

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Real-time Augmented Reality Three-dimensional Guided Robotic Radical Prostatectomy: Preliminary Experience and Evaluation of the Impact on Surgical Planning

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Schiavina, R., Bianchi, L., Lodi, S., Cercenelli, L., Chessa, F., Bortolani, B., et al. (2021). Real-time Augmented Reality Three-dimensional Guided Robotic Radical Prostatectomy: Preliminary Experience and Evaluation of the Impact on Surgical Planning. EUROPEAN UROLOGY FOCUS, 7(6), 1260-1267 [10.1016/j.euf.2020.08.004].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/787247> since: 2021-01-07

*Published:*

DOI: <http://doi.org/10.1016/j.euf.2020.08.004>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

***Real-time Augmented Reality 3D-guided Robotic Radical Prostatectomy: preliminary experience and evaluation of the impact on surgical planning***

***Running title: Augmented reality robotic radical prostatectomy***

*Schiavina R<sup>1-2</sup>, Bianchi L<sup>\*1-2</sup>, Lodi S<sup>3</sup>, Cercenelli L<sup>4-5</sup>, Chessa F<sup>1-2</sup>, Bortolani B<sup>4</sup>, Gaudiano C<sup>6</sup>, Casablanca C<sup>1</sup>, Droghetti M<sup>1</sup>, Porreca A<sup>7</sup>, Romagnoli D<sup>7</sup>, Golfieri R<sup>6</sup>, Giunchi F<sup>8</sup>, Fiorentino M<sup>8</sup>, Marcelli E<sup>3</sup>, Diciotti S<sup>5</sup>, Brunocilla E<sup>1-2</sup>*

<sup>1</sup> Department of Urology, University of Bologna, Bologna, Italy

<sup>2</sup>Department of Experimental, Diagnostic and Specialty Medicine (DIMES), Cardio-Nephro-Thoracic Sciences Doctorate, University of Bologna, Bologna, Italy

<sup>3</sup>Department of Electrical, Electronic and Information Engineering “Guglielmo Marconi”, University of Bologna, Bologna, Italy

<sup>4</sup>Department of Experimental, Diagnostic and Specialty Medicine (DIMES), Laboratory of Bioengineering, University of Bologna, Bologna, Italy

<sup>5</sup>Department of Biomedical and Neuromotor Sciences (DIBINEM), University of Bologna, Bologna, Italy

<sup>6</sup>Radiology Unit, Department of Diagnostic Medicine and Prevention, St. Orsola-Malpighi Hospital, Bologna, Italy

<sup>7</sup>Department of Urology, Abano Terme Hospital, Padua, Italy

<sup>8</sup> Department of Pathology, University of Bologna, Bologna, Italy

1    **\* Corresponding author:**

Lorenzo Bianchi, MD

2    Department of Urology, University of Bologna, Bologna, Italy

Tel. +32 0512142334

Fax. +32 0512142750

E-mail: lorenzo.bianchi3@gmail.com

3

4

**Word count: 2500; Abstract word count: 289; Tables: 3; Figures: 3; Supplementary  
Tables: 3; Supplementary Figures: 3; Supplementary video: 1; References: 30; Pages:  
25**

5

6    **Keywords:** augmented reality; 3D reconstruction; real-time guided surgery; robot assisted  
7    radical prostatectomy; index lesion

8

9

1 **ABSTRACT**

2 **Background:** Augmented Reality (AR) is a novel technology adopted in prostatic surgery.

3 **Objective:** to evaluate the impact on surgical planning of 3D model with AR (AR-3D model)  
4 to guide nerve sparing (NS) during robot-assisted radical prostatectomy (RARP)

5 **Design, Setting, and Participants:** Twenty-six consecutive patients with diagnosis of  
6 prostate cancer (PCa) and multiparametric resonance imaging (mpMRI) available were  
7 scheduled for AR-3D NS RARP.

8 **Intervention:** Segmentation of mpMRI and creation of 3D virtual models were achieved. To  
9 develop the AR guidance, the surgical DaVinci video stream was sent to an AR-dedicated  
10 personal computer and the 3D virtual model was superimposed and manipulated in real-  
11 time on the robotic console.

12 **Outcome Measurements and Statistical Analysis:** The concordance of localization of the  
13 index lesion between the 3D model and the pathologic specimen was evaluated using a  
14 prostate map of 32 specific areas. A preliminary surgical plan to determinate the extent of  
15 NS approach was recorded basing on mpMRI. The final surgical plan was re-assessed  
16 during surgery by implementation of the AR-3D model guidance.

17 **Results and limitations:** PSMs rate were 15.4% in the overall patients' population; 3  
18 patients (11.5%) had PSMs at level of the index lesion. AR-3D technology changed the NS  
19 surgical plan in 38.5% of men on patient-based and in 34.6% of sides on side-based  
20 analysis, resulting an overall appropriateness of 94.4%. The 3D model revealed 70%, 100%  
21 and 92% of sensitivity, specificity and accuracy at 32-areas map analysis.

22 **Conclusion:** AR-3D guided surgery is useful to improve the real-time identification of the  
23 index lesion and allows to change the NS approach in approximately one out three cases,  
24 with overall appropriateness of 94.4%.

- 1 **Patient Summary:** AR-3D guided RARP allows to identify the index PCa during surgery, to
- 2 tailor the surgical dissection to the index lesion and to change the extent of NS dissection.
- 3

## 1 **Introduction**

2 Major technological innovations applied to robotic surgery have set the “precision  
3 medicine” as a novel standard tailored to each patient for the surgical treatment of prostate  
4 cancer (PCa)<sup>1</sup>. The ideal nerve sparing (NS) approach during robotic radical prostatectomy  
5 (RARP) should obtain the optimal compromise between the preservation of neurovascular  
6 bundles (NVBs) and the higher risk of positive surgical margins (PSMs)<sup>2</sup>. As consequence,  
7 different surgical techniques were introduced to reduce PSMs<sup>3</sup> and several methods have  
8 been proposed for “real-time” evaluation of surgical margins<sup>4</sup>.

9 Moreover, multi-parametric Magnetic Resonance Imaging (mpMRI) may modify the  
10 NVB dissection in one out of three individuals<sup>5,6,7</sup> with overall appropriateness of 75%<sup>2</sup>.  
11 Recently, some authors reported that 3D models elaborated from 2D conventional imaging  
12 can be used as additional tools to guide surgery<sup>8,9,10,11,12</sup>. Similarly, Augmented Reality (AR)  
13 technology is increasingly adopted in the surgical field<sup>13</sup> and it has been introduced also in  
14 prostatic surgery<sup>9,14</sup>. Of note, AR technology could increase the understanding of surgical  
15 anatomy and facilitate intraoperative navigation during RARP.

16 Thus, we aimed to evaluate the impact on surgical planning of AR-3D model to guide  
17 surgical dissection during NS RARP for real-time assessment of surgical anatomy and to  
18 test the concordance of the reconstructed anatomical 3D models with the final pathologic  
19 evaluations.

20

21

22

## 1 **Materials and methods**

### 2 *Study design and participants*

3 We prospectively enrolled 26 consecutive patients with diagnosis of PCa basing on positive  
4 MRI-targeted Fusion biopsy at the index lesion<sup>15</sup> with systematic prostate biopsy and  
5 preoperative preserved erectile function (International Index of Erectile Function-5 [IIEF-5]  
6 questionnaire >21<sup>16</sup>) scheduled for RARP at our institution between April 2019 and  
7 September 2019. Participants signed a written informed consent document. The study was  
8 conducted after Institutional Ethics Committee (IRB approval 4325/2017). Exclusion criteria  
9 were contraindications for RARP, undetectable index lesion at preoperative mpMRI or  
10 mpMRI **not** available. Prior to intervention, patients were addressed to undergo 3D virtual  
11 model reconstruction based on preoperative mpMRI images. Finally, the surgeon performed  
12 RARP with the help of the 3D model projected in AR inside the robotic console (AR-3D  
13 guided RARP).

14

### 15 *Preoperative imaging*

16 All the mpMRI examinations were performed with a 1.5-T whole body scanner (Signa HDxt;  
17 GE Healthcare, Milwaukee, WI, USA) and a standard 8-channel pelvic phased-array surface  
18 coil combined with a disposable endorectal coil (MedRad, Indianola, Pa) as previously  
19 described<sup>2</sup>. All lesions were scored using the PI-RADS-v2 score<sup>17</sup>.

20

### 21 *3D Model reconstruction*

22 All 3D virtual model reconstructions based on preoperative mpMRI, were carried out by the  
23 Laboratory of Bioengineering of the University of Bologna (S. Orsola-Malpighi Hospital) and  
24 the development of AR technology was performed by bioengineers of the University of

1 Bologna, Department of Electrical, Electronic and Information Engineering “Guglielmo  
2 Marconi”.

3 Segmentation, i.e. the labelling of each structure of interest in each mpMRI slice was  
4 achieved using D2P™ software (‘DICOM to PRINT’; 3D Systems Inc., Rock Hill, SC; Figure  
5 1). Semi-automatic tools (*multi-slice interpolation* and *threshold segmentation*) of D2P™  
6 software were adopted to segment the healthy prostatic gland, the index lesion, the urinary  
7 sphincter and the NVBs. D2P™ was also used to obtain the 3D virtual models which are  
8 navigable, by converting the segmented structures to 3D triangulated surface mesh file  
9 (STL), using mesh creation methods of D2P™ (*contour* or *gridbase*; Figure 2).

10 To replicate the prostate map used by the pathologist in the whole mount prostate to localize  
11 PCa, the 3D model was stratified in the 3 dimensional space, using Meshmixer software  
12 (Autodesk Inc, San Rafael, CA, US), from base to apex, from right to left and from anterior  
13 to posterior, leading to 32 specific areas including 3-4 macro-sections in which the seminal  
14 vesicles, the base and the apex were analysed separately (Supplementary Figure 1).

15

### 16 *Augmented Reality technology*

17 The augmented reality was delivered through a dedicated hardware and software set-up  
18 (Figure 3). The surgical video stream has been acquired from DaVinci video cart via a frame  
19 grabber (USB3HD, Startech, London, Ontario, Canada) and sent to an AR-dedicated PC  
20 (equipped with an Intel i7 CPU, 8 GB RAM and NVIDIA GeForce 840M video card). A 3D  
21 view of the virtual model obtained using Meshmixer software (Autodesk Inc, San Rafael,  
22 CA, US), was superimposed on the operatory field with vMIX (StudioCoast Pty Ltd, Robina,  
23 Queensland, Australia). During the intervention, the DaVinci TilePro has been activated in  
24 the console and real-time manual alignment has been carried out by a biomedical engineer  
25 using a 6 degrees of freedom (3D) mouse (SpaceMouse, 3D Connexion, Munich, Germany).



1 The resulting augmented video stream was also sent to a second external monitor for quality  
2 control by the surgeon.

3

#### 4 *Surgical technique*

5 All patients underwent RARP using four-arm DaVinci Xi Surgical System (Intuitive Surgical,  
6 Sunnyvale, California, USA), as previous described<sup>18,19,20</sup>. The NS approaches were  
7 classified on patient-based level (considering 26 patients) as bilateral NS, unilateral NS, or  
8 non-NS. Indeed, the extent of neurovascular bundles (NVBs) preservation was recorded on  
9 side-based level (considering 52 sides) as Grade 1, Grade 2, Grade 3-4 according to  
10 incremental NS classification as described by Tewari et al<sup>21</sup>. The main surgeon, when  
11 needed, switched to the TilePro visualization, using the AR technology in which the phantom  
12 of the 3D model was overlaid to the surgical field (Supplementary Video).

13

#### 14 *Surgical planning*

15 The surgical plan to evaluate the extent of NS approach, was planned by surgeons basing  
16 on mpMRI findings together with clinical data. Moreover, the effective intraoperative surgical  
17 plan regarding the level of NVBs preservation<sup>2</sup>, both on patient-based and side-based level  
18 was assessed through a combination of clinical parameters, mpMRI results and the  
19 implemented AR-3D model in the robotic views. At the end of the procedure, the effective  
20 intraoperative surgical approach was compared with the preoperative intended planning in  
21 order to evaluate the potential success of AR-3D guided technology in the management of  
22 surgical dissection.

23

#### 24 *Histopathological examination*

1 Pathologic examinations were performed by a single dedicated uro-pathologist, following a  
2 prostate map in which the gland was divided into 32 specific areas (stratified in the 3-  
3 dimensional space from base to apex, from right to left and from anterior to posterior;  
4 Supplementary Figure 2). The whole mount histological examinations from the RARP  
5 specimens were used as the reference standard, as previously described<sup>22</sup>.

6

### 7 *Statistical analyses*

8 The McNemar-Bowker test was used to compare the surgical plan regarding NS surgery,  
9 before and after its revision following the AR-3D guidance during surgery. The proportion of  
10 surgical plan change was recorded both on patient-based and side-based level. The  
11 appropriateness of surgical plan change was assessed on side-based level and was based  
12 on the presence of ECE or PSMs in the NVBs area at final pathological examination<sup>2</sup>: a less  
13 oncologically radical approach, leading to a grade 1 NS, was considered appropriate in case  
14 of pT2 with negative surgical margins; a less oncologically radical or a more radical  
15 approach, leading to a grade 2 NVBs preservation, was considered appropriate in case of  
16 pT2 or pT3a with negative surgical margins; a more oncologically radical approach leading  
17 to a grade 3-4 NVB preservation was considered appropriate in case of pT3a/pT3b  
18 regardless surgical margins status. Finally, to evaluate the accuracy to localize the index  
19 lesion within the 3D model, the presence of suspected index PCa in each of the 32 prostatic  
20 areas obtained from model stratification was assessed by visual inspection, and it was  
21 compared with the presence of PCa in the corresponding 32 prostatic areas evaluated  
22 separately at pathologic examination, following the prostate map analysis (Supplementary  
23 Figure 3). A p value of <0.05 was considered statistically significant. All statistical tests were  
24 performed using SPSS 22.0.

25

## 1 **Results**

2 Demographic and clinical characteristics are summarized in Table 1. Preoperative mpMRI  
3 reported organ confined index lesion in 23 patients and suspected ECE in 3 individuals;  
4 index lesions were classified as PIRADS 3, 4 and 5 in 9 (34.6%), 9 (34.6%) and 8 (30.8%)  
5 men, respectively. Overall, PSMs rate was 15.4%: 1 (5.3%) PSM in pT2, 2 (33.3%) PSMs  
6 in pT3a and 1 (100%) PSMs in pT3b. Only 3 patients (11.5%) had PSMs at level of the index  
7 lesion detected with 3D-AR model (Table 2). Overall, 2 men experienced postoperative  
8 complication (Clavien<3). The continence recovery rate (0/1 safety PAD) was 57.7%,  
9 73.1%, 88.5% and 92.3% at catheter removal, 1, 3 and 6 months, respectively. The erectile  
10 function recovery rate (IIEF-5>21 with or without Phosphodiesterase 5 inhibitor) was 23.1%  
11 at 1 months, 53.8% at 3 months and 65% at 6 months. No patient had biochemical  
12 recurrence at median follow up of 8 months (Supplementary Table 1). On side-based level,  
13 Grade 1, 2 and 3-4 NS would have been performed in 23 (44.2%), 6 (11.5%) and 23 (44.2%)  
14 sides without 3D-AR model; however, Grade 1, 2 and 3-4 NS was finally performed in 19  
15 (36.5%), 17 (32.7%) and 16 (30.8%) sides with the use of 3D-AR technology, respectively  
16 (p=0.02; Supplementary Table 2). The initial surgical plan of NS techniques was changed  
17 by the use of AR-3D model in 38.5% of men. In 3 (30%) cases, surgery was changed to  
18 more radical approaches, while in 7 (70%) cases the NS approach was attempted (less  
19 radical approach; Table 3a).

20 The use of AR-3D technology induced the surgeon to change the NS surgical plan in 18  
21 (34.6%) sides with overall appropriateness of 94.4% (Table 3b). In 50% of cases surgery  
22 was changed towards a more radical approaches (appropriateness of 77.8%); in the other  
23 half of cases, surgical plan was changed into a less radical approach, (appropriateness of  
24 88.9%). Finally, Supplementary Figure 3 depicts the concordance between the 3D model  
25 and the whole mount specimen according to the 32 prostatic areas to localize the index

1 lesion: the 3D model revealed 70%, 100% and 92% of sensitivity, specificity and accuracy  
2 in the prostate map analysis (Supplementary Table 3).

3

## 1 **Discussion**

2 In the era of precision-medicine, the evolution of real-time imaging-guided technology is an  
3 increasing need to improve the surgical dissection for a tailored surgery of PCa<sup>14</sup>. Of note,  
4 continuous efforts and technique's modifications aiming to reduce PSMs and to preserve  
5 both the periprostatic NVBs<sup>21,23</sup> and the periapical tissue<sup>3</sup> are proposed to obtain better  
6 surgical outcomes<sup>24,25,26</sup>. Despite the use of mpMRI represents nowadays a routine practice  
7 to guide NS surgery<sup>2</sup>, it is not a "real-time" method to evaluate the surgical anatomy during  
8 RARP. Thus, "in vivo" assessment of prostatic gland and periprostatic tissue has gained  
9 special interest in PCa surgery<sup>4</sup>. High fidelity 3D models represent one of the most appealing  
10 method for better understanding of surgical anatomy and guiding surgical planning in  
11 different fields<sup>10,11,14,27</sup>. As consequence, the use of 3D models in urology, showed a  
12 significant impact on surgical planning and decision-making process that may influence  
13 patients' outcomes<sup>28</sup>. Porpiglia and coworkers<sup>14,29</sup> have reported their preliminary  
14 experience with the use of AR technology. Thus, the AR-3D guided surgery is proposed to  
15 facilitate the intraoperative "real-time" navigation and dissection in crucial steps.

16 Several findings are noteworthy in our study. First, the 3D virtual model overlapped to the  
17 DaVinci console through TilePro connection proved to be feasible to define the anatomy of  
18 the prostatic glands and the index lesion, as demonstrated by the good concordance with  
19 the whole-mount pathological analysis. Previously, Porpiglia et al. showed that AR-3D  
20 models were able to identify the index lesion in 100% of patients<sup>14,30</sup>, with a mismatch  
21 between the 3D reconstruction and the scanned prostate recorded from 1 to 5 mm<sup>14</sup>. In our  
22 cohort we used a simple method (a prostate map stratified into 32 specific areas) to validate  
23 the 3D model in terms of index lesion's localization, considering the whole-mount pathologic  
24 specimens as reference: overall, the 3D model revealed 92% of accuracy to localize the  
25 index lesion. The proposed prostate map analysis has two main advantages: firstly, the 32-  
26 areas prostate map on the 3D model is easy to obtain using a free software (Meshmixer);

1 secondly, the proposed prostate map analysis to compare the 32 areas on the 3D model  
2 and the pathologic specimen is able to assess the concordance/non-concordance to localize  
3 the index lesion in the same areas or the presence of PCa in areas nearby the index lesion.  
4 Second, for the first time ever, we evaluated the impact of AR-3D guided surgery to modify  
5 the real-time intraoperative approach to NS both on patient-based and side-based analyses.  
6 The surgical plan was changed through real-time AR-3D visualization, in 38.5% patients and  
7 in 34.6% sides with safe results in terms of PSMs (15.4% overall). Third, the NS plan  
8 changed due to AR-3D was appropriate in the vast majority of cases: overall, in 50% of sides  
9 the AR-3D lead a more radical NS (appropriateness of 77.8%); while in the remaining half  
10 of cases, the AR-3D lead a less radical NS (appropriateness of 88.9%). Fourth, the AR-3D  
11 guided approach to NVBs allows to modulate the NS approach tailored to the index lesion,  
12 by resecting more tissue nearby the lesion and preserving more tissue outside the lesion,  
13 aiming to maximize both oncologic and functional outcomes. Thus, PSMs at level of index  
14 lesion were 11.5% and the erectile function recovery was 65% at 6 months in patients  
15 referred to NS RARP. These findings suggest that the accuracy offered by AR-3D  
16 technology could allow for “real-time” robotic procedure tailored to the patient-specific  
17 anatomy and to the specific cancer location. Future applications of intraoperative frozen  
18 section targeted to the index lesion and guided by AR-3D models could allow further  
19 improvement of surgical outcomes.

20 Despite several strengths, our study is not devoid of limitations. First, in spite of the  
21 prospective nature of the study, the number of patients included is limited. Second, we used  
22 rigid 3D prostate models, that do not represent the tissue deformation during surgical  
23 dissection and the biological realism necessary to create more functional and dynamic  
24 overlapping. Third, the lack of a control group of patients submitted to RARP without the use  
25 of AR-3D model, did not allow to assess the real impact of this technology to modify the  
26 surgical approach. Furthermore, the adoption of different definitions of appropriateness of

1 surgical plan would affect the final results. Moreover, the superimposed screen with the 3D  
2 model can create some problems during the dissection, since the surgeons need to switch  
3 from conventional to the TilePro view that consist in a double and smaller screen. Fourth,  
4 we included also the first patients treated with this novel technology, with possible effect to  
5 prolong the surgical time due to the learning curve process. Indeed, we found that 2 to 5  
6 cases are needed both for the surgeon and the bioengineer to improve the overlap of the  
7 3D model in the surgical field. Finally, the major limitations of AR-assisted surgery consist of  
8 possible registration inaccuracy, translating into a poor navigation precision and the need of  
9 manual external adjustments of the 3D model overlapping on the surgical field<sup>9</sup>. Thus,  
10 automatic registration of the 3D model to the surgical view, based on artificial intelligence,  
11 could be the further implementation to obtain an automated overlapping of the 3D virtual  
12 model inside the DaVinci console liable with the organ movements during surgery.

13

14

1 **Conclusions**

2 The use of AR-3D guided surgery can be a feasible tool to improve the real-time  
3 identification of the index lesion and it could be useful to modulate the NS approach during  
4 RARP. The surgical plan change after the intraoperative adoption of AR-3D guidance  
5 technology was recorded in approximately one out of three cases, with overall  
6 appropriateness of 94.4%.

7



**Acknowledgment:** None

1 **Author Disclosure Statement:** No competing financial interests exist

2

1 **Conflict of interest**

2 The project was supported by a Technology Research Grant by Intuitive Surgical for the  
3 development of augmented reality technology in robotic surgery.

4

1 **Figure legend**

2 **Figure 1.** Example of process to obtain the 3D virtual anatomical model starting from patient  
3 mpMRI (a), by the segmentation (b) of the anatomical regions of interest (prostatic gland,  
4 index lesion, urinary sphincter and neurovascular bundles) to the final 3D model (c).

5 **Figure 2.** Example of the definitive 3D virtual model that can be explored by the surgeon for  
6 surgical planning: the model could be rotated and zoomed and is navigable, i.e. the surgeon  
7 can interact with it changing the transparency of each structure in the 3D rendering, as well  
8 as creating a detailed view of the interaction among the different anatomical structures.

9 **Figure 3.** A schematic diagram of the hardware and software required to implement the  
10 intra-operative use of AR to guide robotic surgery.

11

12

1 **Supplementary Figures**

2 **Supplementary Figure 1.** Three-dimensional stratification of the prostate 3D model (from  
3 base to apex, from right to left and from anterior to posterior) leading to 32 specific areas  
4 including 3-4 macro-sections used to evaluate the concordance of the reconstructed 3D  
5 model with the prostate map stratified in the same 32 areas by pathologist on the whole  
6 mount specimen.

7 **Supplementary Figure 2.** Prostate map used to divide the gland into 32 specific areas  
8 (stratified in the 3-dimensional space from base to apex, from right to left and from anterior  
9 to posterior) on the whole mount specimen.

10 **Supplementary Figure 3.** Prostate map analysis: concordance (green) and discordance  
11 (red) to localize the index lesion between the 3D model and the pathologic whole mount  
12 specimen by visual assessment in each patient (n=26) after stratification of the whole gland  
13 (both on 3D model and on pathologic specimen) according to the 32 prostatic areas.

14

1 **Supplementary Video**

2 Example of AR-3D guided surgical dissection for real-time assessment of the index lesion  
3 during the RARP through the TilePro visualization: the 3D model is overlapped inside the  
4 surgical view during nerve sparing approach and helps the surgeon to modulate the surgical  
5 dissection of the neurovascular bundle.

6

7

8

## 1 References

- 2 1. Checcucci E, Amparore D, Fiori C, et al. 3D imaging applications for robotic urologic  
3 surgery: an ESUT YAUWP review. *World J Urol.* August 2019. doi:10.1007/s00345-  
4 019-02922-4
- 5 2. Schiavina R, Bianchi L, Borghesi M, et al. MRI Displays the Prostatic Cancer  
6 Anatomy and Improves the Bundles Management before Robot-Assisted Radical  
7 Prostatectomy. *J Endourol.* 2018;32(4). doi:10.1089/end.2017.0701
- 8 3. Bianchi L, Turri FM, Larcher A, et al. A Novel Approach for Apical Dissection During  
9 Robot-assisted Radical Prostatectomy: The “Collar” Technique. *Eur Urol Focus.*  
10 2018. doi:10.1016/j.euf.2018.01.004
- 11 4. Eissa A, Zoeir A, Sighinolfi MC, et al. “Real-time” Assessment of Surgical Margins  
12 During Radical Prostatectomy: State-of-the-Art. *Clin Genitourin Cancer.* July 2019.  
13 doi:10.1016/j.clgc.2019.07.012
- 14 5. McClure TD, Margolis DJA, Reiter RE, et al. Use of MR imaging to determine  
15 preservation of the neurovascular bundles at robotic-assisted laparoscopic  
16 prostatectomy. *Radiology.* 2012;262(3):874-883. doi:10.1148/radiol.11103504
- 17 6. Park BH, Jeon HG, Jeong BC, et al. Influence of magnetic resonance imaging in the  
18 decision to preserve or resect neurovascular bundles at robotic assisted  
19 laparoscopic radical prostatectomy. *J Urol.* 2014;192(1):82-88.  
20 doi:10.1016/j.juro.2014.01.005
- 21 7. Kozikowski M, Malewski W, Michalak W, Dobruch J. Clinical utility of MRI in the  
22 decision-making process before radical prostatectomy: Systematic review and meta-  
23 analysis. *PLoS One.* 2019;14(1):e0210194. doi:10.1371/journal.pone.0210194
- 24 8. Porpiglia F, Fiori C, Checcucci E, Amparore D, Bertolo R. Hyperaccuracy Three-

- 1 dimensional Reconstruction Is Able to Maximize the Efficacy of Selective Clamping  
2 During Robot-assisted Partial Nephrectomy for Complex Renal Masses. *Eur Urol.*  
3 2018;74(5):651-660. doi:10.1016/j.eururo.2017.12.027
- 4 9. Bertolo R, Hung A, Porpiglia F, Bove P, Schleicher M, Dasgupta P. Systematic  
5 review of augmented reality in urological interventions: the evidences of an impact  
6 on surgical outcomes are yet to come. *World J Urol.* March 2019.  
7 doi:10.1007/s00345-019-02711-z
- 8 10. Schiavina R, Bianchi L, Borghesi M, et al. Three-dimensional digital reconstruction of  
9 renal model to guide preoperative planning of robot-assisted partial nephrectomy. *Int*  
10 *J Urol.* June 2019. doi:10.1111/iju.14038
- 11 11. Bianchi L, Schiavina R, Barbaresi U, et al. *3D Reconstruction and Physical Renal*  
12 *Model to Improve Percutaneous Puncture during PNL.* Vol 45. Brazil; 2019.  
13 doi:10.1590/S1677-5538.IBJU.2018.0799
- 14 12. Borghesi M, Schiavina R, Gan M, Novara G, Mottrie A, Ficarra V. Expanding  
15 utilization of robotic partial nephrectomy for clinical T1b and complex T1a renal  
16 masses. *World J Urol.* 2013;31(3):499-504. doi:10.1007/s00345-013-1095-2
- 17 13. G. Badiali, L.Cercenelli, S. Battaglia, E. Marcelli, C. Marchetti, V. Ferrari FC. Review  
18 on Augmented Reality in Oral and Cranio-Maxillofacial Surgery: Toward "Surgery-  
19 Specific" Head-Up Displays. *IEEE Access.* 2020;8(February 2020).
- 20 14. Porpiglia F, Fiori C, Checcucci C, Amparore S, Bertolo R, Augmented Reality  
21 Robot-assisted Radical Prostatectomy: Preliminary Experience." *Urology.* 2018  
22 May;115:184. doi: 10.1016/j.urology.2018.01.028.
- 23 15. Ahmed HU. The index lesion and the origin of prostate cancer. *N Engl J Med.*  
24 2009;361(17):1704-1706. doi:10.1056/NEJMcibr0905562

- 1 16. Cappelleri JC, Rosen RC, Smith MD, Mishra A, Osterloh IH. Diagnostic evaluation of  
2 the erectile function domain of the International Index of Erectile Function. *Urology*.  
3 1999;54(2):346-351. doi:10.1016/s0090-4295(99)00099-0
- 4 17. Weinreb JC, Barentsz JO, Choyke PL, et al. PI-RADS Prostate Imaging - Reporting  
5 and Data System: 2015, Version 2. *Eur Urol*. 2016;69(1):16-40.  
6 doi:10.1016/j.eururo.2015.08.052
- 7 18. Mottrie A, De Naeyer G, Schatteman P, Frumenzio E, Rossanese M, Ficarra V.  
8 Robot-assisted radical prostatectomy: tips, tricks and pitfalls. *Minerva Urol Nefrol*.  
9 2012;64(2):89-96.
- 10 19. Porreca A, D'agostino D, Dandrea M, et al. Bidirectional barbed suture for posterior  
11 musculofascial reconstruction and knotless vesicourethral anastomosis during  
12 robot-assisted radical prostatectomy. *Minerva Urol Nefrol*. 2018;70(3):319-325.  
13 doi:10.23736/S0393-2249.18.02969-7
- 14 20. Porreca A, Salvaggio A, Dandrea M, et al. Robotic-Assisted Radical Prostatectomy  
15 with the Use of Barbed Sutures. *Surg Technol Int*. 2017;30:39-43.
- 16 21. Tewari AK, Ali A, Metgud S, et al. Functional outcomes following robotic  
17 prostatectomy using athermal, traction free risk-stratified grades of nerve sparing.  
18 *World J Urol*. 2013;31(3):471-480. doi:10.1007/s00345-012-1018-7
- 19 22. Bianchi L, Schiavina R, Borghesi M, et al. Patterns of positive surgical margins after  
20 open radical prostatectomy and their association with clinical recurrence. *Minerva*  
21 *Urol Nefrol*. May 2019. doi:10.23736/S0393-2249.19.03269-7
- 22 23. Patel VR, Schatloff O, Chauhan S, et al. The role of the prostatic vasculature as a  
23 landmark for nerve sparing during robot-assisted radical prostatectomy. *Eur Urol*.  
24 2012;61(3):571-576. doi:10.1016/j.eururo.2011.12.047



- 1 24. Schiavina R, Borghesi M, Dababneh H, et al. Survival, Continence and Potency  
2 (SCP) recovery after radical retropubic prostatectomy: a long-term combined  
3 evaluation of surgical outcomes. *Eur J Surg Oncol.* 2014;40(12):1716-1723.  
4 doi:10.1016/j.ejso.2014.06.015
- 5 25. Schiavina R, Bertaccini A, Franceschelli A, et al. The impact of the extent of lymph-  
6 node dissection on biochemical relapse after radical prostatectomy in node-negative  
7 patients. *Anticancer Res.* 2010;30(6):2297-2302.
- 8 26. Farolfi A, Ceci F, Castellucci P, et al. (68)Ga-PSMA-11 PET/CT in prostate cancer  
9 patients with biochemical recurrence after radical prostatectomy and PSA <0.5  
10 ng/ml. Efficacy and impact on treatment strategy. *Eur J Nucl Med Mol Imaging.*  
11 2019;46(1):11-19. doi:10.1007/s00259-018-4066-4
- 12 27. Shirk JD, Thiel DD, Wallen EM, et al. Effect of 3-Dimensional Virtual Reality Models  
13 for Surgical Planning of Robotic-Assisted Partial Nephrectomy on Surgical  
14 Outcomes: A Randomized Clinical Trial. *JAMA Netw open.* 2019;2(9):e1911598.  
15 doi:10.1001/jamanetworkopen.2019.11598
- 16 28. Cacciamani GE, Okhunov Z, Meneses AD, et al. Impact of Three-dimensional  
17 Printing in Urology: State of the Art and Future Perspectives. A Systematic Review  
18 by ESUT-YAUWP Group. *Eur Urol.* 2019;76(2):209-221.  
19 doi:10.1016/j.eururo.2019.04.044
- 20 29. Porpiglia F, Bertolo R, Amparore D, et al. Augmented reality during robot-assisted  
21 radical prostatectomy: expert robotic surgeons' on-the-spot insights after live  
22 surgery. *Minerva Urol Nefrol.* 2018;70(2):226-229. doi:10.23736/S0393-  
23 2249.18.03143-0
- 24 30. Porpiglia F, Checcucci E, Amparore D, et al. Three-dimensional Elastic Augmented-

1 reality Robot-assisted Radical Prostatectomy Using Hyperaccuracy Three-  
2 dimensional Reconstruction Technology: A Step Further in the Identification of  
3 Capsular Involvement. *Eur Urol.* April 2019. doi:10.1016/j.eururo.2019.03.037

4

5