

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Distribution, virulence, genotypic characteristics and antibiotic resistance of *Listeria monocytogenes* isolated over one-year monitoring from two pig slaughterhouses and processing plants and their fresh hams

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Rugna, G., Carra, E., Bergamini, F., Franzini, G., Faccini, S., Gattuso, A., et al. (2021). Distribution, virulence, genotypic characteristics and antibiotic resistance of *Listeria monocytogenes* isolated over one-year monitoring from two pig slaughterhouses and processing plants and their fresh hams. INTERNATIONAL JOURNAL OF FOOD MICROBIOLOGY, 336, 1-11 [10.1016/j.ijfoodmicro.2020.108912].

Availability:

This version is available at: <https://hdl.handle.net/11585/789134> since: 2021-01-17

Published:

DOI: <http://doi.org/10.1016/j.ijfoodmicro.2020.108912>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1 **Distribution, virulence, genotypic characteristics and antibiotic resistance of *Listeria monocytogenes* isolated over one-year monitoring from**
2 **two pig slaughterhouses and processing plants and their fresh hams**

3
4 Rugna Gianluca¹, Carra Elena¹, Bergamini Federica¹, Franzini Giuliana¹, Faccini Silvia¹, Gattuso Antonietta², Morganti Marina¹, Baldi Deborah¹,
5 Naldi Simona¹, Serraino Andrea³, Piva Silvia³, Merialdi Giuseppe¹, Giacometti Federica^{3*}

6
7 ¹ Experimental Zooprophyllactic Institute in Lombardy and Emilia Romagna, Brescia, Italy

8 ² Istituto Superiore di Sanità, Department of Food Safety, Nutrition and Veterinary Public Health, Rome, Italy

9 ³ Department of Veterinary Medical Sciences, University of Bologna, Bologna, Italy

10

11 *Correspondence to: Federica Giacometti, Department of Veterinary Medical Sciences, University of Bologna, Bologna, Italy. Email address:

12 federica.giacometti3@unibo.it (F. Giacometti)

13

14

15 **Abstract**

16 *Listeria monocytogenes* contamination in raw pork and ready to eat foods is an important food safety concern, also for the increasing detection of
17 antimicrobial-resistant isolates. Data on *L. monocytogenes* **occurrence**, persistence, distribution and genetic characterization in two different plants,

18 namely in continuum from slaughtered pigs, environment and unfinished products (fresh hams) were observed by one-year monitoring and were
19 integrated with their antimicrobial resistance patterns. A total of 98 samples out of the overall 1,131 (8.7%) were positive for *L. monocytogenes*,
20 respectively 2.6% and 13.2% in plants A and B: only three serotypes were identified, 1/2c (50%), 1/2b (36.7%) and 1/2a (13.27%), and strains were
21 classified in 35 pulsotypes and 16 clusters by PFGE; a unique P-type was highlighted according to the detection of virulence genes. The contamination
22 flow of *L. monocytogenes* has a low occurrence in slaughterhouse (Plant A=1.1%, Plant B: 3.1%; $p>0.05$) and increased throughout the processing
23 chain with trimming area as the most contaminated (Plant A: 25%, Plant B: 57%; $p<0.05$), both in the environment and in unfinished products (80%
24 in hams before trimming in plant B). The dominant role of environmental contamination in post-slaughter processing is confirmed to be a significant
25 cause of meat contamination by *L. monocytogenes*. Very high levels of resistance were observed for clindamycin (57%) and high resistance levels
26 (>20 -50%) to ciprofloxacin, oxacillin, levofloxacin and daptomycin, confirming the *L. monocytogenes* resistance trend to a wide range of antimicrobial
27 agents. A total of 11 *L. monocytogenes* isolates were multidrug resistant and 7 out of them were isolated from slaughtered pigs. An interesting
28 significant ($p<0.05$) statistical correlation has been found between resistance to some antimicrobial agents and lineage/serotypes. Microbiological
29 sampling of food and environments after sanitization are commonly used as verification procedure for the absence of *L. monocytogenes* in food plants
30 and to give assurance of food safety, but strains characterization is necessary for industries to target specific control measures, like the enforcement
31 of the hygiene program and of the control of operator activities, at least for permanent strains. The only presence of *L. monocytogenes* could not be
32 considered as the conclusive assessment of a potential risk for public health, also in terms of emerging and emerged antimicrobial resistances.

33

34 **Keywords**

35 Foodborne pathogen, pig industry, pork product, molecular epidemiology, antimicrobial resistance
36

37

38

39

40

41 **1. Introduction**

42 Pork is the most frequently consumed meat in the European Union (Devine, 2003). In Italy, dry cured hams, in particular those designed by the
43 European Union as PDO (Protected Designation Origin) like Parma and San Daniele, represent a significant source of income for the pork industry,
44 being exported worldwide, even if raw meat and other pork products are produced with carcass parts other than thighs. *Listeria monocytogenes*
45 contamination in raw pork is an important food safety concern (Li et al., 2018), and the management of microbiological hazards transmitted to humans
46 by pork consumption is of major health and economic significance (De Cesare et al., 2018). *L. monocytogenes* is a ubiquitous foodborne pathogen
47 which causes listeriosis mainly among the so called YOPIs (young, old, pregnant, immunocompromised) sensitive groups. Although still relatively
48 rare, in Europe human listeriosis is one of the most serious food-borne diseases under surveillance causing hospitalization, high morbidity and high
49 mortality (EFSA, 2019); treatment with antibiotics is usually needed for the control of the infection caused by this bacterium.

50 *L. monocytogenes* grows and survives in different types of habitats, even in adverse conditions, such as dry environment, high salt concentrations
51 (10%, wt/vol), at refrigeration temperatures and a wide pH range (4.7-9.2). This ability to persist and multiply in the food environment, even by
52 biofilm formation, makes the genus *Listeria* hard to control in food processing plants (Conficoni et al., 2016). In the pig industry, the contamination

53 routes of *L. monocytogenes* are well reported and its incidence in pig carcasses and slaughter plant equipment is usually low (Meloni et al., 2013;
54 Ortiz et al., 2010), whereas a relatively common contamination of raw pork products and ready to eat (RTE) food with *L. monocytogenes* is observed
55 (Camargo et al., 2017; EFSA, 2018). Persistence of *L. monocytogenes* in food processing environments is still considered the major source of RTE
56 food contamination and this persistence appears to be the result both of improper hygiene conditions and of the high adaptive capacity of this bacterium
57 (EFSA, 2018).

58 For these reasons *L. monocytogenes* poses a significant risk to the food industry, particularly producers of RTE foods, and meat products continue to
59 be one of the three most RTE food categories typically associated with human listeriosis (EFSA, 2018). In addition, further concern is due to the
60 increasing detection of antimicrobial-resistant *L. monocytogenes* isolates, mainly for antibiotics commonly used for the treatment of listeriosis, namely
61 a combination of penicillin or ampicillin with aminoglycosides (gentamicin) and trimethoprim-sulfamethoxazole (Sosnowski et al., 2019). Future
62 outbreaks may be more difficult to manage because of the emergence of antimicrobial resistance among *L. monocytogenes* strains isolated from food
63 products (Olaimat et al., 2018).

64 In literature, there are several studies worldwide on *L. monocytogenes* presence, persistence, distribution and genetic characterization in pig
65 slaughterhouses and processing plants, as well as in fresh hams and pork meat products. It is well recognized that, on the one hand, raw pork meat has
66 rarely been implicated in foodborne illness, and perhaps it could be considered a potential source of domestic cross contamination of other food
67 (Thevenot et al., 2006), and that, on the other, the level of *L. monocytogenes* contamination increases along the pork supply chain (López et al., 2008).

68 When contaminated surfaces are involved in food contamination, the risk of the cross-contamination of pork by *L. monocytogenes* could be assessed
69 by predictive models able to predict sporadic event affecting the number of contaminated food samples and the influence of food processing factors

70 and the indirect mechanisms involved in cross-contamination (Jiang et al., 2018; Møller et al., 2016; Possas et al., 2017). Being the role of the after-
71 slaughter phases, such as cooling and/or cutting, crucial in the contamination of fresh hams (Camargo et al., 2017; Larivière-Gauthier et al., 2014;
72 Thevenot et al., 2006), the reduction of *L. monocytogenes* prevalence in the processing plant environment and, as a consequence, the decrease of the
73 initial *L. monocytogenes* load in intermediate fresh hams may lead to a significant reduction in the likelihood of *L. monocytogenes* contamination
74 downstream the meat production chain.

75 Moreover, resistance of *L. monocytogenes* to many antimicrobial agents has emerged and evolved during the past few decades (Olaimat et al., 2018).
76 Resistance is increasingly observed from humans, food, plants and the environment: in humans, *L. monocytogenes* isolates showed resistance for
77 tetracycline and ciprofloxacin; in food, high prevalence of oxacillin and clindamycin resistance was described from meat and fish production chains
78 and significant percentages of resistance against ampicillin, penicillin G, and tetracycline were reported in *L. monocytogenes* strains isolated from
79 meat, fish, and dairy production chains (Caruso et al., 2019). However, among this increasing number of studies on the emergence of antibiotic
80 resistance in *L. monocytogenes* from food products (Camargo et al., 2015; Caruso et al., 2019; Chen et al., 2019; Escolar et al., 2017; Gómez et al.,
81 2014; Li et al., 2016; Moreno et al., 2014; Sala et al., 2016; Sereno et al., 2019; Sosnowski et al., 2019), it is noteworthy that the investigated isolates
82 usually belong to strain collections or to different food sources considered all together, with any correlation to a specific scenario and/or phase within
83 the food chains. No correlation of these antimicrobial resistance levels with information about presence, distribution and genomic characterization is
84 performed along pork chains, let alone food chains in general. Thus, the aim of this study was to integrate these scattered information by one-year of
85 *L. monocytogenes* monitoring in two Italian plants in order to: i) describe the occurrence of *L. monocytogenes* in continuum in two slaughterhouses
86 with annexed cutting and trimming plants, from slaughtered pigs, environment and unfinished products (fresh hams); ii) trace the relevant sources of

87 contamination for fresh hams by PFGE typing method; iii) characterize the *L. monocytogenes* isolates in relation to their serotyping, profiles of
88 virulence genes and antimicrobial resistance patterns.

89

90 **2. Material and methods**

91 **2.1. Sampling in the two plants**

92 Two ham processing facilities, located respectively in the Lombardia and Emilia-Romagna Regions, in Italy, were investigated. Plants A and B
93 comprise two large scale (>400pigs/hour) slaughterhouses authorized to export pork meat products to the Unites States, each annexed to a processing
94 plant, with cutting and trimming areas, that produce fresh hams for the “Parma ham” production and other fresh pork products. Plants A and B were
95 monthly sampled from April 2014 to September 2015 with a sampling plan including food and environment samples collected in continuum at
96 slaughtering, cutting, cooling, trimming areas and storage chill room, respectively. One variable day within the week for each sampling was scheduled.
97 The sampled animals belonged to different batches of about 135 heavy pigs (160-180 kg weight) 9-10 months old. The animals received by the
98 slaughterhouses came for most part (88.9% of pigs and 89.4% of batches) from Emilia–Romagna or from nearby northern Italian regions (Piedmont
99 and Lombardy), within a radius of about 250 km from the slaughterhouses. Travel duration resulted always under 8 hours, and about 60% of batches
100 with less than 90 min. Overall, plants A and B were sampled 10 and 9 times, respectively. At each sampling time point, the following samples were
101 collected from different areas of the processing plant: i) from slaughtering area, cecal content and amygdalae (and ileo-cecal lymph nodes only for
102 plant B) were collected after evisceration from 15 pigs randomly selected; environment and facilities, namely splitting saws (2 swabs in pool) and
103 gloves, bib, and surfaces in contact with the carcasses (4 swabs in pool); ii) from both cutting and trimming areas, hams after cutting (HAC) (4 swabs

104 in pool) and hams before trimming (HBT) (4 swabs in pool) were randomly sampled; environment and facilities, namely meat contact surface swabs
105 (MCS) (4 swabs in pool for hams conveyor, gloves and knives, respectively) and non-meat contact surface swabs (NMCS) (drain); iii) in cooling
106 room, room walls (4 swabs in pool); iv) in storage chill room, ham post-trimming and before curing (HBC) (20 swabs examined in single) were
107 collected from hams belonging to 10 different batches. Plants A and B used different cleaning and disinfection protocols: i) a preliminary cleaning
108 aimed at achieving “visual cleanliness” of the workplace was performed by both plants by washing down with a low-pressure hose, but using cold
109 water in plant A and hot water (between 45°C and 55°C) in plant B; ii) a cleaning stage was performed with alkaline foaming detergent (at a dilution
110 rate of 3% in plant A and 5% in plant B), alternating with acids products at the week-ends only in plant A; in this plant, a specific biofilm-disruptor
111 product was used weekly; iv) intermediate rinsing with hot water (<45°C) at low pressure in plant A and with hot water (between 45°C and 55°C) at
112 high pressure in plant B; v) disinfection performed by rotating products on a daily basis in both plants: chlorine products or equivalents in plant A and
113 citric acid and hydrogen peroxide or quaternary ammonium compound in plant B. In the latter plant, an amphoteric disinfectant based on acetic acid
114 and propanediamine derivatives was used once a fortnight; vi) final rinsing with water at very low pressure in plant A and with cold water at low
115 pressure in plant B.

116 The environmental samples were collected by swabbing 30x30 cm of the surface, during processing. Overall, a total of 1,131 samples were collected,
117 namely 487 in plant A and 644 in plant B. More details are reported in Table 1. All samples were carried under chilled conditions to the laboratory of
118 the Experimental Institute for Zooprophyaxis of Lombardy and Emilia Romagna, and processed within 12 h after collection.

119

120 2.2. Isolation and identification of *Listeria monocytogenes*

121 For the microbiological analysis, the ileo-cecal lymph nodes and amygdalae were separated from the fat and surrounding tissues, respectively, and
122 both were processed using 95% alcohol to flame on specimen surface and then disrupted using a hammer to expose the interior before analysis. Fecal
123 and swab samples were homogenized 1/10 with Fraser broth base (Biolife, Milan, Italy) in a Stomacher Lab Blender 400 (PBI, Italy) for 2 min.
124 The detection of *L. monocytogenes* was performed according to the international standard method ISO 11290-1:1996\Amd 1:2004. Only culture-
125 confirmed samples by biochemical tests (Gram staining, catalase, hemolysis and CAMP tests, carbohydrate utilization test for xylose, rhamnose and
126 mannitol) were deemed positive.
127 One *L. monocytogenes* isolate for each positive sample was considered and preserved in brain heart infusion broth (Biolife) with glycerol (15%,
128 vol/vol) at -80°C. All the considered isolates underwent their characterization by serotyping and by the presence of putative virulence genes. Bacterial
129 DNA extracted from the selected collection of *L. monocytogenes* isolates was subjected to multiplex PCR-based serotyping assay targeting genes
130 *lmo0737*, *lmo1118*, ORF2819, ORF2110, and *prs*, as previously described (Doumith et al., 2004), and confirmed with classical agglutination
131 serotyping according to the method described in the Bacteriological Analytical Manual using commercial anti O and H antisera (Denka Seiken, Tokyo,
132 Japan).

133

134 **2.3. Antibiotics Susceptibility Testing**

135 All the isolates were tested for their antimicrobial resistance using a commercial microbroth dilution method, namely Sensititre™ Gram Positive Plate
136 Format (GPN3F plates, Thermo Scientific, USA) containing 18 antimicrobials (dilution range in mg/l) were used: ampicillin (AMP; 0.12–16),
137 ceftriaxone (AXO; 8–64), ciprofloxacin (CIP; 0.5–2), clindamycin (CLI; 0.12–2), daptomycin (DAP; 1–8), erythromycin (ERY; 0.25–4), gatifloxacin

138 (GAT; 1–8), gentamicin (GEN; 2–16), levofloxacin (LEVO; 0.25–8), linezolid (LZD; 0.5–8), oxacillin (OXA; 0.25–8), penicillin (PEN; 0.06–8),
139 quinupristin/dalfopristin (SYN; 0.12–4), rifampin (RIF; 0.5–4), streptomycin (STR; 1000), tetracycline (TET; 2–16), trimethoprim/sulfamethoxazole
140 (SXT; 0.5/9.5–4/76), vancomycin (VAN; 1–128). The antimicrobials were selected among antimicrobials used for treatment of human *Listeria*
141 infections or for antibiotic therapy that is usually applied against Gram-positive bacteria. The minimal inhibitory concentration (MIC) records were
142 manually read. Antimicrobial resistance of the isolates was determined as first choice according to epidemiological cut-off values for *L.*
143 *monocytogenes* proposed by the European Committee on Antimicrobial Susceptibility Testing (EUCAST), following by EUCAST clinical breakpoints
144 for *L. monocytogenes* and *Staphylococcus* spp., and lastly by Clinical and Laboratory Standards Institute (CLSI) breakpoints for *L. monocytogenes*
145 and *Staphylococcus* spp. (CLSI, 2012; CLSI, 2016). Isolates were considered multidrug resistant (MDR) when showing resistance to three or more
146 antimicrobial classes (Schwarz et al., 2010).

147

148 **2.4. Typing of *Listeria monocytogenes* isolates and detection of virulence genes**

149 Pulsed-field gel electrophoresis (PFGE) was performed according to the Pulse-Net standardized protocol for *L. monocytogenes*
150 (<https://www.cdc.gov/pulsenet/pdf/listeria-pfge-protocol-508c.pdf>) using *AscI* and *ApaI* restriction enzymes. *Salmonella enterica* serovar Braenderup
151 strain H9812 restricted with *XbaI* was used as standard according to the above protocol. The PFGE profiles were analyzed using BioNumerics software
152 (version 6.7; Applied Maths, Sint-Martens-Latem, Belgium); similarity between PFGE profiles was determined using the Dice coefficient with 1%
153 value of tolerance limit and optimization. Pulsotypes were obtained combining *AscI* and *ApaI* profiles. Dendrograms were obtained by cluster analysis
154 of the pulsotypes, set at a 0.90% similarity level, by the Unweighted Pair Group Method Analysis (UPGMA). Isolates showing a PFGE similarity

level $\geq 90\%$ were assigned to the same pulsotype. Isolates showing pulsotypes with a similarity level of $>80\%$ were grouped in the same “PFGE cluster” and were identified by progressive Roman numbers. The definition of the clusters was performed by evaluating the two plants separately. The reliability of cluster analysis was evaluated by calculating the cophenetic correlation coefficient. The discriminatory index (DI) of the PFGE analysis was calculated using the Simpson’s diversity index described by Hunter and Gaston (Hunter and Gaston 1988), considering the combined PFGE *AscI* and *ApaI* profiles, and using the Comparing Partitions Website (<http://www.comparingpartitions.info>). For the identification of virulence genes, DNA was amplified by PCR to detect the *inlA*, *inlB*, *inlC*, and *inlJ* genes (Liu et al., 2007), which code for internalin proteins A, B, C, J, respectively, and of *plcA*, *hlyA*, *actA*, and *iap* genes (Rawool et al., 2007). According to the presence of these genes, different pathotypes (P-types) were determined.

163

164 2.5. Statistical analysis

Descriptive statistics (absolute frequencies and percentages) have been provided regarding the occurrence and characteristics (serotype and P-type) of *L. monocytogenes* isolates. Findings were presented for the two plants by sampling area, sample types and sources. Descriptive statistics (absolute frequencies and percentages) have been also provided for their antimicrobial resistance patterns, namely the number of isolates resistant to, at least, one AA and MDR isolates. Findings were presented by lineage, origin, sources and clusters. Chi-square test or Fisher’s exact test were used to compare nominal variables between different sampling area, origin and sources, as well as serotypes and lineages. Significance was set at a P value of <0.05 . PRISM 5.0 software was used.

171

172 3. Results

173 3.1. Isolation and identification of *Listeria monocytogenes*

174 A total of 98 samples out of the overall 1,131 (8.7%) were positive for *L. monocytogenes* by culture examination, 13 (2.6%) and 85 (13.2%) in plants
175 A and B, respectively. For both plants a low and a relatively low contamination rates were reported in the slaughterhouse area (1.1% and 3.1% for
176 plants A and B, respectively; $p>0.05$), whereas the trimming area resulted the most contaminated (25% and 57% for plants A and B, respectively;
177 $p<0.05$), reaching a maximum level of occurrence of 80% for HBT samples. For the other areas, not negligible differences in *L. monocytogenes*
178 occurrence between the plants were found: in plant A, *L. monocytogenes* was isolated in 2.2 % of samples from cutting area, whereas it was not found
179 in cooling areas and in HBC; on the other hand, in plant B *L. monocytogenes* was isolated in 40%, 10% and 27% of samples collected from cutting
180 area, cooling area and in HBC, respectively. The environment resulted more contaminated than food, with the only exception of the slaughterhouse
181 area of both plants, in which the contamination rate in slaughtered pigs, even if low, resulted higher than in environment. More details in relation to
182 the occurrence, distribution of *L. monocytogenes* in the two plants and their statistical comparisons are reported in Table 1.
183 Only three serotypes were identified in the overall 98 *L. monocytogenes* isolates, namely 1/2c (50%; n=49), 1/2b (36.7%; n=36) and 1/2a (13.27%;
184 n=13). Three lineages were identified in the population of *L. monocytogenes*: the lineage I includes isolates of serotype 4b/4e, 1/2b, 3b, 4b/4e and 3c,
185 the lineage II isolates of serotype 1/2a, 1/2c and 3a, and the lineage III isolates of serotype 4a and 4c. Of the overall 98 strains, 63.2% (62/98) belonged
186 to the lineage II and 36.7% (36/98) to the lineage I. Occurrence, distribution in the two plants and characteristics of the 98 *L. monocytogenes* isolates
187 in relation to lineage distribution, serotyping, and their source are summarized in Table 1.

188

189 3.2. Antibiotics Susceptibility Testing

190 The MIC evaluation of the 98 isolates is reported in Table 2. In total, all the isolates were susceptible to 8 out of 18 tested antimicrobials, namely,
191 quinupristin/dalfopristin, vancomycin, ampicillin, gentamicin, rifampin, penicillin, streptomycin and gatifloxacin. Ceftriaxone was not considered
192 because *L. monocytogenes* is intrinsically resistant to cephalosporins (CRAB, 2020). A total of 93 isolates (94.9%) showed resistance to at least one
193 antimicrobial agent and 11 isolates (11.2%) were MDR, and it is worth noting that 7 (64%) of the overall 11 MDR isolates were from slaughtered
194 pigs. The percentage of MDR isolates was significant higher ($p<0.05$) in slaughtered pigs in comparison to environment and food, and significant
195 differences were ($p<0.05$) in the lineages and serotypes (see Table 3).

196 The most common resistances of *L. monocytogenes* isolates were to clindamycin, with very high level of resistance (n=56/98 isolates; 57.1%), followed
197 by high resistance levels (>20-50%) to ciprofloxacin (n=42/98 isolates; 42.9%), oxacillin (n=35/98 isolates; 35.7%), levofloxacin (n= 34/98 isolates;
198 34.7%) and daptomycin (23/98 isolates; 23.5%). Lastly, low resistance levels (>1-10%) were observed for linezolid (n=4/98 isolates; 4%), tetracycline
199 and trimethoprim/sulfamethoxazole (both n=3/98 isolates; 3.1%) and erythromycin (n=2/98 isolates; 2.04%). Tables 3 and 4 detail antimicrobial
200 susceptibility findings of *L. monocytogenes* isolates.

201 Different resistance levels were observed in relation to the lineage and serotype, and in relation to the origin: i) the percentage of resistant *L.*
202 *monocytogenes* was significantly higher ($p<0.05$) in isolates of lineage II for clindamycin, levofloxacin and ciprofloxacin, whereas for daptomycin
203 and oxacillin was significantly higher ($p<0.05$) in isolates of lineage I, and no differences were found for the other antimicrobial agents (see Table 3);
204 ii) the percentage of resistant *L. monocytogenes* was significantly higher ($p<0.05$) in isolates of serotype 1/2c for clindamycin, levofloxacin and
205 ciprofloxacin, and of serotype 1/2a for daptomycin and oxacillin (see Table 3); iii) although the percentage of MDR isolates was significantly higher

206 ($p<0.05$) in slaughtered pigs, no significant differences were found between the percentage of resistant *L. monocytogenes* isolated from different
207 sources (slaughtered pigs, environment and food) for all the antimicrobial agents, with the exception of clindamycin for which the percentage of
208 resistant *L. monocytogenes* was significantly lower ($p<0.05$) in isolates from food source (see Table 4).

209

210 3.3. Typing of *Listeria monocytogenes* isolates and detection of virulence genes

211 The 98 *L. monocytogenes* isolates characterized using PFGE with *AscI* and *ApaI* were classified in 35 pulsotypes and 16 clusters. The pulsotypes
212 indicated as P16, P01, P04, P13, P15, P26, P03, P12, P18, P08, P14 and P07 were shared between 2 and 29 isolates, whereas the other 23 pulsotypes
213 were identified in single isolates. More details were reported in Figures 1-2. Pulsotypes with >10% of isolates were arbitrarily considered as
214 predominant, namely pulsotypes P16, P01, P04. The most common pulsotype was that indicated as P16, exclusively associated with 29 isolates of
215 serotype 1/2b and with environmental sources, followed by pulsotypes P01, P04, P13 and P15. The only pulsotype identified from all the investigated
216 sources, namely slaughtered pigs, food and environment, was pulsotype P04, even if the unique strain isolated from food shows a different serotype,
217 namely 1/2b, versus serotype 1/2c of all the other 8 isolates of this pulsotype. The resistant isolates were spread in all clusters (except in cluster XIV)
218 even if the *L. monocytogenes* isolates showing resistance to the highest number of antimicrobial agents (>4 antimicrobial agents) were grouped in
219 cluster V, IV and I. For each antimicrobial agent, the percentage of resistant isolates varies between the different clusters, with some clusters gathering
220 the great majority of resistant isolates: in cluster VI were included 32 out of the overall 35 (91%) and 16 of the overall 23 (70%) isolates resistant to
221 oxacillin and daptomycin, respectively. Cluster IV showed 18/34 (53%) and 20/42 (48%) resistant isolates to levofloxacin and ciprofloxacin, as well

222 as 26/56 (46%) of clindamycin resistant isolates were observed in cluster III. MDR isolates belong to 5 clusters (IV, V, VII, IX and XI). See Table 5
223 for more details.

224 The DI of PFGE method ranged between 0.883 (C.I. 95%: 0.843-0.923) for plant A and 0.971 (C.I. 95%: 0.906-1.000) for plant B.

225 All the genes were detected among the investigated 98 isolates *L. monocytogenes* strains, thus a unique P-type was highlighted according to the
226 detection of virulence genes, hindering comparison and evaluation on virulence genotypes, but highlighting their potential pathogenicity.

227

228 4. Discussion

229 The control of *L. monocytogenes* in meat processing facilities continues to be an important challenge consistent with the continuous, even if
230 intermittent, introduction and/or reintroduction of the microorganism in the processing plants, and with the presence of persistent strains in the same
231 environment and the parallel complexity of proper sanitization practices. The first objective of this study was to describe the occurrence and
232 distribution of *L. monocytogenes* in different areas of two different pig industries, from slaughterhouses to fresh hams. Our study confirms the
233 contamination flow of *L. monocytogenes* observed in different pork processing plants with a low occurrence in slaughterhouse that increased
234 throughout the processing chain, both in the environment and in unfinished products (Larivière-Gauthier et al., 2014; Ortiz et al., 2010; Prencipe et
235 al., 2012; Thevenot et al., 2006). Indeed, the low level of contamination at slaughterhouse resulted in line with most of the previous studies reporting
236 an occurrence ranging from 0 to 16.7% in pork carcasses (Larivière-Gauthier et al., 2014; Ortiz et al., 2010; Prencipe et al., 2012; Thevenot et al.,
237 2006) and 0 to 3 % on slaughter equipment (Ortiz et al., 2010) but lower than the study of Meloni and Colleagues (2013) reporting prevalence of 33%
238 in slaughtered pigs. Our findings showed that few batches of pigs entering the slaughterhouse were contaminated with *L. monocytogenes* (from 3 out

239 of 9 samplings for plant A to 5 out of 10 samplings for plant B, with higher frequencies in amygdalae, followed by ileo-cecal lymph nodes and feces)
240 but with a high strain diversity (13 pulsotypes among 17 overall isolates) supporting the observations of Larivière-Gauthier and Colleague (2014) of
241 an infrequent entry of positive batches with a great variety of strains. Furthermore, these strains could be classified as not persistent strains because
242 they were never detected on consecutive visits and they have different pulsotypes, with the exception of pulsotype (P04) observed in the only one
243 splitting saw resulted positive, ileo-cecal lymph nodes, knives and gloves in cutting and trimming area, drains in the trimming area as well as HBC,
244 the latter isolated more than one year later (see P04 in cluster IV of dendrogram, Figure 2). P04 resulted the only pulsotype identified from all the
245 investigated sources: this event demonstrated that these isolates were in-house strains of this plant environment, that a contamination of a plant with
246 *L. monocytogenes* could originate from primary production and that, reciprocally, utensils (splitting saw) could be responsible for cross contamination
247 of the fresh hams, in line with observations of Larivière-Gauthier and Colleagues (2014). Among the debate on the possible origin of contaminations
248 of fresh hams, and as a consequence of the end products, namely if contaminations were incoming from earlier phases of the processing chain or were
249 rather directly originating from the processing environment, our study highlighted the marginality of transfers of *L. monocytogenes* from primary
250 production and, at the same time, the dominant role of environmental contamination. Post-slaughter processing confirmed to be a significant cause of
251 meat contamination, namely in cutting and trimming environment where this contamination is amplified (see Table 1, and cluster III and VI of plant
252 B in Figure 2), even if the initial microbial load as well as the environmental conditions favoring the microbial growth (nutrients from meat and
253 temperature) could have also played an increase in the *L. monocytogenes* level of contamination in both plants (Camargo et al., 2017; Larivière-
254 Gauthier et al., 2014; Thevenot et al., 2006).

255 Only five pulsotypes (P01, P04, P12, P13 and P16) persisted in the plant during the 1-year period of this study but they represented the 60% of the
256 overall *L. monocytogenes* isolates and included all the three predominant pulsotypes (P16, P01, P04). This is a common scenario for food processing
257 plants and a significant risk factor for the application and selection of the appropriate food plant sanitization procedures. The presence of *L.*
258 *monocytogenes* belonging to these predominant pulsotypes in cutting, trimming and storage chill room areas, and their circulation on plant B through
259 the transmission of indistinguishable or closely related pulsotypes between food and the environment (both MCS and NMCS), probably represent the
260 mechanism for *L. monocytogenes* maintenance on the plant. Indeed, differently from slaughter area and cooling room, significant differences ($p<0.05$)
261 in the occurrence between plants A and B were observed in the other investigated areas, in which, even with wide fluctuations, the frequencies of *L.*
262 *monocytogenes* isolation were higher and affected all the samples types and sources. Moreover, the overall occurrence in fresh hams of both plants
263 are in line with literature and the overall occurrence in fresh hams observed in plant B was similar to the worst findings among 13 Italian
264 slaughterhouses (Prencipe et al., 2012) as well as in an Iberian pig slaughterhouse and processing plant underwent to a 3-year surveillance (Ortiz et
265 al., 2010). This difference in contamination between the two plants as well as the varied contamination levels across the processing suggests the
266 existence of limitations to cross-contamination between compartments within the same plant and reflects the operative features of a specific plant,
267 also in terms of sanitization procedures and behavior of workers. Based on the findings observed in cutting and trimming areas in plant B, on one
268 hand, gloves and knives of workers were frequently contaminated and could be considered carriers for HBT and HBC contamination, and on the other,
269 MCS (e.g. ham conveyors) as well as NMCS (e.g. drain) are significant niches indicative of the tendency of *L. monocytogenes* to persist in processing
270 facilities or in general inside the plant. In this context, it is essential the adoption of proper sanitization practices trying to avoid cross-contamination
271 to the products by environmental contamination, a good training of operators on working procedures like movement of materials, and cleaning and

272 disinfection (C&D) procedures. For example, no preventing crossover circuits for equipment and forklift, and the improper use of high-pressure hoses,
273 both observed only in plant B (data not shown), could have respectively represented a possible source of environmental bacterial contamination and
274 provoked the spreading of aerosol particles belonging to drainage water, as observed by Conficoni et al., 2016. These observations could justify the
275 high number of *L. monocytogenes* strains isolated in HBT and HBC in plant B. Finally, besides within-plant strains similarity, pulsotype P01 (namely
276 represented by isolate MN28 in plant A) was found in both the establishments, in line with two studies reporting the presence of indistinguishable
277 strains in different plants, namely for the contamination pattern of *L. monocytogenes* in the environment of Cured Ham food chain (Morganti et al.,
278 2016) and in finished pork-meat products of several unrelated factories (Autio et al., 2002).

279 For serotyping, in line with literature, 1/2a, 1/2b and 1/2c *L. monocytogenes* serotypes were observed in the pig slaughterhouse and processing plants,
280 in slaughtered pigs and from fresh meat (unfinished products). Serotypes 4b, 1/2b, and 1/2a are described as pathogenic and have been shown to be
281 predominant in human listeriosis cases (Orsi et al., 2011): the 4b is responsible for the majority of human listeriosis outbreaks, while sporadic cases
282 of *L. monocytogenes* gastroenteritis are typically caused by consumption of food products contaminated with high bacterial loads of serotype 1/2a and
283 1/2b (Halbedel et al., 2019). Differently from previous studies on pork meat industry showing that serotype 1/2a is the dominant serotype and the
284 most prevalent in food (Li et al., 2016; López et al., 2008; Ortiz et al., 2010; Thevenot et al., 2006), our findings agree with other Italian studies
285 (Meloni et al., 2013; Prencipe et al., 2012) in which serotype 1/2c resulted the most dominant in plants, whereas serotype 1/2a seems to be the
286 prevalent, but not exclusive, in pig animals, in which serotypes 1/2b and 1/2c were also isolated. Although the reasons for dominance of the different
287 serotypes are unclear, strains belonging to serotype 1/2a and 1/2c have demonstrated to be more capable of biofilm formation when compared to
288 lineage I (Borucki et al., 2003; Kalmokoff et al., 2001). In particular, serotype 1/2c adheres significantly more to stainless steel, demonstrating highest

289 degree of adsorption (Ortiz et al., 2010). The *L. monocytogenes* serovars depend upon peptidoglycan-anchored cell wall teichoic acids (WTAs)
290 diversity within the cell wall that defines the O-antigens; WTAs are known to be involved in regulation of cell morphology and division, autolytic
291 activity, ion homeostasis, protection from host defenses and antibiotics, and may mediate host cell invasion and colonization. Recently, the study of
292 Sumrall and Colleagues (2019) demonstrated a switch from serovar 4b to 4d by bacteriophages with a loss of WTA galactosylation, which is involved
293 in cells adhesion; this leads to the loss of biofilm formation capacity of *L. monocytogenes* cells, consequently allowing their easy elimination during
294 sanitization. No hypothesis could be performed in this study for the dominance of serotype 1/2c, but both the mechanisms of serotype selection as a
295 result of sanitization conditions or change in serotype could be taken into account. In our study, all the serotypes are distributed in all sources with
296 some differences: serotype 1/2a, 1/2b and 1/2c isolates were mostly present in slaughtered pigs (61.5%), food (58.3%) and in the environment (59.2%),
297 respectively. In similar studies, serotype 1/2a isolates were present in the environment and equipment, and in different product categories, whereas
298 serotype 1/2b was mostly presented in environmental sites and raw products (Larivière-Gauthier et al., 2014; López et al., 2008; Ortiz et al., 2010;)
299 or even serotype 1/2c in fresh hams (Prencipe et al., 2012).

300 From source attribution studies, it was evidenced that not all *L. monocytogenes* strains are equally capable of causing invasive disease: overall, several
301 studies have shown that *L. monocytogenes* strains belonging to lineage I are on average more virulent and more frequently associated with human
302 clinical cases than lineage II strains (Filipello et al., 2020). Additionally, a recent study showed that a significant proportion of *L. monocytogenes*
303 isolated from food production environments have reduced virulence (Van Stelten et al., 2016) but our findings otherwise showed that all tested *L.*
304 *monocytogenes* strains had all detected virulence genes, and this suggests that *L. monocytogenes* strains isolated from food and/or environment are
305 potentially pathogenic and consequently may play an important role in epidemics, independently from the sources.

306 In relation to data on antimicrobial resistance in *L. monocytogenes*, some studies are available in literature but most of them reported data generically
307 from different food products or meat and human; very few studies investigated the pork chain, and only one study considered the continuum from
308 slaughterhouses, environment and fresh hams in the same plants (Sereno et al., 2019). Furthermore, comparisons are arduous due to differences in
309 antimicrobial agents and breakpoints used. In most studies, the most common resistances were observed for oxacillin, clindamycin, tetracycline,
310 ampicillin and trimethoprim-sulfamethoxazole and MDR isolates varied between zero to 27% (Camargo et al., 2015; Caruso et al., 2019; Chen et al.,
311 2019; Escolar et al., 2017; Gómez et al., 2014; Li et al., 2016; Moreno et al., 2014; Sala et al., 2016; Sereno et al., 2019; Sosnowski et al., 2019). In
312 our study very low resistance or full susceptibility were observed for ampicillin (treatment of choice for listeriosis), tetracycline, and trimethoprim-
313 sulfamethoxazole (used as second choice therapy or as alternative therapy for penicillin-allergic patients). However, the presence of very high and
314 high level of resistances to clindamycin, ciprofloxacin, oxacillin, levofloxacin and daptomycin, is noteworthy and of important concern for public
315 health, since most of these antibiotics are widely used in hospitals to treat Gram-positive infections. Our findings confirm the *L. monocytogenes*
316 increasing trend of resistance to a wide range of antimicrobial agents and the fact that the genus *Listeria* spp. could no longer be reported as susceptible
317 to almost all antimicrobials. In addition, attention should be demanded for the risk of increasing multidrug resistance in *Listeria* and the possibility of
318 its transfer to other bacteria (Moreno et al., 2014).

319 Slaughtered pigs were the only source of strains that resulted resistant to all the 9 antimicrobial agents for which at least a resistance was observed
320 and for which the highest resistance was reported for all these antimicrobials, notwithstanding no significant differences were observed between the
321 different origins (see table 4), whereas resulted the source with significant higher percentage of MDR strains. This finding could be due to the fact
322 that emergence and spread of antimicrobial-resistant *Listeria* spp. has been attributed to the overuse of antibiotics in disease treatments and growth

323 promotion in domestic livestock (Charpentier et al., 1995; Walsh et al., 2001). The most common antibiotic classes worldly used in global pig
324 production are penicillins and tetracyclines. However, the use of Critically Important Antimicrobials for humans is also reported, even at different
325 levels across countries, differently across countries: macrolides were reported at 20% and at 7.4% of total use in France and Austria, fluoroquinolones
326 at 2.4% and 5% and third and fourth generation cephalosporins at 2.2% and 11% of total use in Austria and Belgium (Lekagul et al., 2019). In one
327 study performed in Italy on evaluation of antibiotic usage in swine reproduction farms, penicillins, macrolides, tetracyclines, and polymyxins (colistin)
328 were the most common antimicrobials used; third-generation cephalosporins, penicillins, colistin and fluoroquinolones had the highest prevalence,
329 while sulphonamides and trimethoprim, tetracyclines had the highest intensity of use (DDDs/animals) (Scoppetta et al., 2017).

330 An interesting statistical correlation ($p<0.05$) has been found between resistance and lineage/serotypes. Clindamycin, levofloxacin and ciprofloxacin
331 and, similarly, oxacillin and daptomycin resistances were significantly ($p<0.05$) more frequently identified respectively in lineage II and in particular
332 in serotype 1/2c, and, in lineage I and in particular in serotype 1/2b. This correlation has been previously reported for oxacillin and lineage I, but in
333 particular with serotype 4b/4e, whereas intermediate clindamycin or ciprofloxacin resistances resulted higher in serotype 4b/4e (Caruso et al., 2019)
334 or 1/2a (Kovacevic et al., 2013; Safdar and Armstrong, 2003), but in several other cases no correlations were found (Safdar and Armstrong, 2003).

335 No observations for these correlations have been proposed yet except that serotypes originating from different countries have different antibiotic
336 resistance profiles (Ayaz and Erol, 2010).

337 In conclusion, microbiological sampling of food and environments after sanitization are commonly used as verification procedure for the absence of
338 *L. monocytogenes* in food plants and to give assurance of food safety, but the findings of this study underlined that the presence without recognizing
339 of the real pattern of contamination and the characteristics of the strains could not be considered as the conclusive assessment of a potential risk for

340 public health. Genotyping *L. monocytogenes* strains, mostly in cases in which a previous identification of the pathogen occurred within the plant or
341 in unfinished products, is necessary for industries to target specific control measures, for example the enforcement of the hygiene program and of the
342 control of operator activities, and may help reducing the risk of cross-contamination at the consumer level. *L. monocytogenes* microbiological sampling
343 and its conjunction with a specific evaluation of virulence and antimicrobial resistance of the *L. monocytogenes* strains, at least for permanent strains,
344 are necessary since it should result in a more comprehensive food safety vision and control, also in terms of emerging and emerged antimicrobial
345 resistances.

346

347 **Funding**

348 This work was supported by the Italian Ministry of Health (Project n. IZS LER 09/12 RC).

349

350 **Declaration of competing interest**

351 The authors declare that they have no conflict of interest.

352

353 **References**

354 Devine, R., 200. Meat consumption trends in the world and the European Union. INRA Prod. Anim.16, 325-327.

355 Li, H., Wang, P., Lan, R., Luo, L., Cao, X., Wang, Y., Wang, Y., Li, H., Zhang, L., Ji, S., Ye, C., 2018. Risk factors and level of *Listeria monocytogenes*
356 contamination of raw pork in retail markets in China. Front. Microbiol. 29, 9:1090. <https://doi.org/10.3389/fmicb.2018.01090>

- 357 De Cesare, A., Doménech, E., Comin, D., Meluzzi, A., Manfreda, G., 2018. Impact of cooking procedures and storage practices at home on
358 consumer exposure to *Listeria monocytogenes* and *Salmonella* due to the consumption of pork meat. Risk Anal. 38(4), 638-652.
359 <https://doi.org/10.1111/risa.12882>
- 360 EFSA (European Food Safety Authority), 2019. Scientific report on the European Union One Health 2018 Zoonoses Report. EFSA J. 17(12), 5926.
361 <https://doi.org/10.2903/j.efsa.2019.5926>
- 362 Conficoni, D., Losasso, C., Cortini, E., Di Cesare, A., Cibir, V., Giaccone, V., Corno, G., Ricci, A., 2016. Resistance to biocides in *Listeria*
363 *monocytogenes* collected in meat-processing environments. Front. Microbiol. 19, 7:1627. <https://doi.org/10.3389/fmicb.2016.01627>
- 364 Meloni, D., Piras, F., Mureddu, A., Fois, F., Consolati, S.G., Lamon, S., Mazzette, R., 2013. *Listeria monocytogenes* in five sardinian swine
365 slaughterhouses: prevalence, serotype, and genotype characterization. J. Food Prot. 76(11), 1863-1867. [https://doi.org/10.4315/0362-](https://doi.org/10.4315/0362-028X.JFP-12-505)
366 [028X.JFP-12-505](https://doi.org/10.4315/0362-028X.JFP-12-505)
- 367 Ortiz, S., López, V., Villatoro, D., López, P., Dávila, J.C., Martínez-Suárez, J.N., 2010. A 3-year surveillance of the genetic diversity and
368 persistence of *Listeria monocytogenes* in an Iberian pig slaughterhouse and processing plant. Foodborne Pathog. Dis. 7(10), 1177-1784.
369 <https://doi.org/10.1089/fpd.2010.0535>
- 370 Camargo, A.C., Woodward, J.J., Call, D.R., Nero, L.A., 2017. *Listeria monocytogenes* in food-processing facilities, food contamination, and human
371 listeriosis: the brazilian scenario. Foodborne Pathog Dis. 14(11), 623-636. <https://doi.org/10.1089/fpd.2016.2274>

372 EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), Ricci, A., Allende, A., Bolton, D., Chemaly, M., Davies, R., Fernández Escámez, P.S.,
 373 Girones, R., Herman, L., Koutsoumanis, K., Nørrung, B., Robertson, L., Ru, G., Sanaa, M., Simmons, M., Skandamis, P., Snary, E.,
 374 Speybroeck, N., Ter Kuile, B., Threlfall, J., Wahlström, H., Takkinen, J., Wagner, M., Arcella, D., Da Silva Felicio, M.T., Georgiadis, M.,
 375 Messens, W., Lindqvist, R., 2018. Scientific Opinion on the *Listeria monocytogenes* contamination of ready-to-eat foods and the risk for
 376 human health in the EU. EFSA J. 16(1), 5134. <https://doi.org/10.2903/j.efsa.2018.5134>

377 Sosnowski, M., Lachtara, B., Wieczorek, K., Osek, J., 2019. Antimicrobial resistance and genotypic characteristics of *Listeria monocytogenes*
 378 isolated from food in Poland. Int. J. Food Microbiol. 16(289), 1-6. <https://doi.org/10.1016/j.ijfoodmicro.2018.08.029>

379 Olaimat, A.N., Al-Holy, M.A., Shahbaz, H.M., Al-Nabulsi, A.A., Abu Ghoush, M.H., Osaili, T.M., Ayyash, M.M., Holley, R.A., 2018. Emergence
 380 of antibiotic resistance in *Listeria monocytogenes* isolated from food products: a comprehensive review. Compr. Rev. Food Sci. F. 17, 5.
 381 <https://doi.org/10.1111/1541-4337.12387>

382 Thévenot, D., Dernburg, A., Vernozzy-Rozand, C., 2006. An updated review of *Listeria monocytogenes* in the pork meat industry and its products. J.
 383 Appl. Microbiol. 101, 7-17. <https://doi.org/10.1111/j.1365-2672.2006.02962.x>

384 López, V., Villatoro, D., Ortiz, S., López, P., Navas, J., Dávila, J.C., Martínez-Suárez, J.V. 2008. Molecular tracking of *Listeria monocytogenes* in an
 385 Iberian pig abattoir and processing plant. Meat Sci. 78(1-2), 130-4. <https://doi.org/10.1016/j.meatsci.2007.05.002>

386 Jiang, R.H., Wang, X., Wang, W., Liu, Y.T., Du, J.P., Cui, Y., Zhang, C.Y., Dong, Q.L., 2018. Modelling the cross-contamination of *Listeria*
 387 *monocytogenes* in pork during bowl chopping. Int J Food Sci Tech. 53, 837-846. <https://doi.org/10.1111/ijfs.13660>

388 Møller, C.O., Sant'Ana, A.S., Hansen, S.K., Nauta, M.J., Silva, L.P., Alvarenga, V.O., Maffei, D., Silva, F.F., Lopes, J.T., Franco, B.D., Aabo, S.,
 389 Hansen, T.B., 2016. Evaluation of a cross contamination model describing transfer of *Salmonella* spp. and *Listeria monocytogenes* during
 390 grinding of pork and beef. Int. J. Food Microbiol. 226, 42-52. <https://doi.org/10.1016/j.ijfoodmicro.2016.03.016>
 391 Possas, A., Carrasco, E., Garcia-Gimeno, R.M., Valero, A., 2017. Models of microbial cross-contamination dynamics. Curr. Opin. Food Sci. 14, 43-
 392 49. <http://dx.doi.org/10.1016/j.cofs.2017.01.006>
 393 Larivière-Gauthier, G., Letellier, A., Kérouanton, A., Bekal, S., Quessy, S., Fournaise, S., Fravallo, P., 2014. Analysis of *Listeria monocytogenes* strain
 394 distribution in a pork slaughter and cutting plant in the Province of Quebec. J. Food Prot. 77(12), 2121-8. [https://doi.org/10.4315/0362-](https://doi.org/10.4315/0362-028X.JFP-14-192)
 395 [028X.JFP-14-192](https://doi.org/10.4315/0362-028X.JFP-14-192)
 396 Caruso, M., Fraccalvieri, R., Pasquali, F., Santagada, G., Latorre, L.M., Difato, L.M., Miccolupo, A., Normanno, G., Parisi, A., 2019. Antimicrobial
 397 Susceptibility and Multilocus Sequence Typing of *Listeria monocytogenes* isolated over 11 years from food, humans, and the environment in
 398 Italy. Foodborne Pathog. Dis. 17(4), 284-294. <https://doi.org/10.1089/fpd.2019.2723>
 399 Camargo, A.C., Dias, M.R., Cossi, M.V.C., Lanna, F.G.P.A., Cavicchioli, V.Q., Valim, D.C., Pinto, P.S.A., Hofer, E., Nero, L.A., 2015. Serotypes
 400 and pulsotypes diversity of *Listeria monocytogenes* in a beef-processing environment. Foodb. Pathog. Dis. 12, 323-326.
 401 <http://doi.org/10.1089/fpd.2014.1875>

402 Chen, M., Cheng, J., Zhang, J., Chen, Y., Zeng, H., Xue, L., Lei, T., Pang, R., Wu, S., Wu, H., Zhang, S., Wei, X., Zhang, Y., Ding, Y., Wu, Q., 2019.
 403 Isolation, potential virulence, and population diversity of *Listeria monocytogenes* from meat and meat products in China. Front. Microbiol.
 404 10, 946. <https://www.frontiersin.org/article/10.3389/fmicb.2019.00946>

405 Escolar, C., Gómez, D., Del Carmen Rota García, M., Conchello, P., Herrera, A., 2017. Antimicrobial resistance profiles of *Listeria monocytogenes*
 406 and *Listeria innocua* isolated from ready-to-eat products of animal origin in Spain. Foodborne Pathog. Dis. 14(6), 357-363.
 407 <https://doi.org/10.1089/fpd.2016.2248>

408 Gómez, D., Azón, E., Marco, N., Carramiñana, J.J., Rota, C., Ariño, A., Yangüela, J., 2014. Antimicrobial resistance of *Listeria monocytogenes* and
 409 *Listeria innocua* from meat products and meat-processing environment. Food Microbiol. 42, 61-65. <https://doi.org/10.1016/j.fm.2014.02.017>

410 Li, L., Olsen, R.H., Ye, L., Wang, W., Shi, L., Yan, H., Meng, H., 2016. Characterization of antimicrobial resistance of *Listeria monocytogenes* strains
 411 isolated from a pork processing plant and its respective meat markets in southern China. Foodborne Pathog. Dis. 13(5), 262-268.
 412 <https://doi.org/10.1089/fpd.2015.2087>

413 Moreno, L.Z., Paixão, R., Gobbi, D.D.S., Raimundo, D.C., Ferreira, T.P., Moreno, A.M., Hofer, E., Reis, C.M.F., Matté, G.R., Matté, M.H., 2014.
 414 Characterization of antibiotic resistance in *Listeria* spp. isolated from slaughterhouse environments, pork and human infections. J. Infect. Dev.
 415 Ctries. 8(4), 416-23. <https://jids.org/index.php/journal/article/view/24727506>

416 Sala, C., Morar, A., Tîrziu, E., Nichita, I., Imre, M., Imre, K., 2016. Environmental Occurrence and Antibiotic susceptibility profile of *Listeria*
 417 *monocytogenes* at a slaughterhouse raw processing plant in Romania. J. Food Protect. 79(10), 1794-1797. [https://doi.org/10.4315/0362-](https://doi.org/10.4315/0362-028X.JFP-16-052)
 418 [028X.JFP-16-052](https://doi.org/10.4315/0362-028X.JFP-16-052)

419 Sereno, M.J., Viana, C., Pegoraro, K., da Silva, D.A.L., Yamatogi, R.S., Nero, L.A., Bersot, L.D.S., 2019. Distribution, adhesion, virulence and
 420 antibiotic resistance of persistent *Listeria monocytogenes* in a pig slaughterhouse in Brazil. Food Microbiol. 84, 103234.
 421 <https://doi.org/10.1016/j.fm.2019.05.018>

422 ISO, 2004. Microbiology of food and animal feeding stuffs — Horizontal method for the detection and enumeration of *Listeria monocytogenes* —
 423 Part 1: Detection method (ISO 11290-1:1996/AMD 1:2004). International Organization for Standardization, Geneva, Switzerland.

424 Doumith, M., Buchrieser, C., Glaser, P., Jacquet, C., Martin, P., 2004. Differentiation of the major *Listeria monocytogenes* serovars by multiplex
 425 PCR. J. Clin. Microbiol. 42(8), 3819-22. <https://dx.doi.org/10.1128%2FJCM.42.8.3819-3822.2004>

426 CLSI (Clinical and Laboratory Standards Institute), 2012. Performance Standards for Antimicrobial Susceptibility Testing Twenty Second
 427 Informational Supplement. vol. 33, no. 3 Clinical and Laboratory Standards Institute, Wayne, PA, USA (Document M100-S30-20).

428 CLSI (Clinical and Laboratory Standards Institute), 2016. Methods for Antimicrobial Dilution and Disk Susceptibility Testing of Infrequently
 429 Isolated or Fastidious Bacteria. Clinical and Laboratory Standards Institute, Wayne, PA, USA (Document M45).

430 Schwarz, S., Silley, P., Simjee, S., Woodford, N., van Duijkeren, E., Johnson, A. P., Gastra, W., 2010. Editorial: assessing the antimicrobial
 431 susceptibility of bacteria obtained from animals. J. Antimicrob. Chemother. 65(4):601-604. <https://doi.org/10.1093/jac/dkq037>

- 432 Hunter, P.R. and Gaston, M.A., 1988. Numerical index of the discriminatory ability of typing systems: an application of Simpson's index of
433 diversity. J. Clin. Microbiol. 26(11), 2465-6.
- 434 Liu, D., Lawrence, M.L., Austin, F.W., Ainsworth, A.J., 2007. A multiplex PCR for species- and virulence-specific determination of *Listeria*
435 *monocytogenes*. J. Microbiol. Methods 71, 133-140. <https://doi.org/10.1016/j.mimet.2007.08.007>
- 436 Rawool, D.B., Malik, S.V.S., Barbuddhe, S.B., Shakuntala, I., Aurora, R.A., 2007. A multiplex PCR for detection of virulence associated genes in
437 *Listeria monocytogenes*. Int. J. Food Safety 9, 56–62.
- 438 CRAB, 2020. Centro di Referenza Nazionale per l'Antibioticoresistenza. Table 1. URL: [http://www.izslt.it/crab/wp-](http://www.izslt.it/crab/wp-content/uploads/sites/8/2018/08/Tabelle-Resistenze-intrinseche-in-batteri-di-interesse-veterinario.pdf)
439 [content/uploads/sites/8/2018/08/Tabelle-Resistenze-intrinseche-in-batteri-di-interesse-veterinario.pdf](http://www.izslt.it/crab/wp-content/uploads/sites/8/2018/08/Tabelle-Resistenze-intrinseche-in-batteri-di-interesse-veterinario.pdf) (Accessed 08.01. 2020).
- 440 Prencipe, V.A., Rizzi, V., Acciari, V., Iannetti, L., Giovannini, A., Serraino, A., Calderone, D., Rossi, A., Morelli, D., Marino, L., Migliorati, G.,
441 Caporale, V., 2012. *Listeria monocytogenes* prevalence, contamination levels and strains characterization throughout the Parma ham
442 processing chain. Food Control, 25(1), 150-158. <https://doi.org/10.1016/j.foodcont.2011.10.018>
- 443 Morganti, M., Scaltriti, E., Cozzolino, P., Bolzoni, L., Casadei, G., Pierantoni, M., Foni, E., Pongolini, S., 2016. Processing-Dependent and Clonal
444 Contamination Patterns of *Listeria monocytogenes* in the Cured Ham Food Chain Revealed by Genetic Analysis. Appl. Environ. Microbiol.
445 82(3), 822-831. <https://doi.org/10.1128/AEM.03103-15>.
- 446 Autio, T., Lundén, J., Fredriksson-Ahomaa, M., Björkroth, J., Sjöberg, A.M., Korkeala, H., 2002. Similar *Listeria monocytogenes* pulsotypes detected
447 in several foods originating from different sources. Int. J. Food Microbiol. 77(1-2), 83-90. [https://doi.org/10.1016/S0168-1605\(02\)00055-7](https://doi.org/10.1016/S0168-1605(02)00055-7)

448 Orsi, R.H., den Bakker, H.C., Wiedmann, M., 2011. *Listeria monocytogenes* lineages: Genomics, evolution, ecology, and phenotypic characteristics.
 449 Int. J. Medical Microbiol. 301(2), 79-96. <https://doi.org/10.1016/j.ijmm.2010.05.002>

450 Halbedel, S., Prager, R., Banerji, S., Kleta, S., Trost, E., Nishanth, G., Alles, G., Hölzel, C., Schlesiger, F., Pietzka, A., Schlüter, D., Flieger, A., 2019.
 451 A *Listeria monocytogenes* ST2 clone lacking chitinase ChiB from an outbreak of non-invasive gastroenteritis. Emerg. Microbes Infect. 8(1),
 452 17-28. <https://doi.org/10.1080/22221751.2018.1558960>

453 Borucki, M.K., Peppin, J.D., White, D., Loge, F., Call, D.R., 2003. Variation in biofilm formation among strains of *Listeria monocytogenes*. Appl.
 454 Environ. Microbiol. 69, 7336-7342. <https://doi.org/10.1128/AEM.69.12.7336-7342.2003>

455 Kalmokoff, M.L., Austin, J.W., Wan, X.D., Sanders, G., Banerjee, S., Farber, J.M., 2001. Adsorption, attachment and biofilm formation among
 456 isolates of *Listeria monocytogenes* using model conditions. J. Appl. Microbiol. 91, 725-734. <https://doi.org/10.1046/j.1365-2672.2001.01419.x>

457 Sumrall, E.T., Shen, Y., Keller, A.P., Rismondo, J., Pavlou, M., Eugster, M.R., Boulos, S., Disson, O., Thouvenot, P., Kilcher, S., Wollscheid, B.,
 458 Cabanes, D., Lecuit, M., Gründling, A., Loessner, M.J., (2019). Phage resistance at the cost of virulence: *Listeria monocytogenes* serovar 4b
 459 requires galactosylated teichoic acids for InlB-mediated invasion. PLoS Pathog. 15(10), e1008032.
 460 <https://doi.org/10.1371/journal.ppat.1008032>

461 Filipello, V., Mughini-Gras, L., Gallina, S., Vitale, N., Mannelli, A., Pontello, M., Decastelli, L., Allard, M.W., Brown, E.W., Lomonaco, S., 2020.
 462 Attribution of *Listeria monocytogenes* human infections to food and animal sources in Northern Italy. Food Microbiol. 89, 103433.
 463 <https://doi.org/10.1016/j.fm.2020.103433>

464 Van Stelten, A., Roberts, A.R., Manuel, C.S., Nightingale, K.K., 2016. *Listeria monocytogenes* isolates carrying virulence-attenuating mutations in
 465 internalin A are commonly isolated from ready-to-eat food processing plant and retail environments. J. Food Prot. 79 (10), 1733-1740.
 466 <https://doi.org/10.4315/0362-028X.JFP-16-145>

467 Charpentier, E., Gerbaud, G., Jacquet, C., Rocourt, J., Courvalin, P., 1995. Incidence of antibiotic resistance in *Listeria* species. J. Infect. Dis. 172,
 468 277-281. <https://doi.org/10.1093/infdis/172.1.277>

469 Walsh, D., Duffy, G., Sheridan, J.J., Blair, I.S., McDowell, D.A., 2001. Antibiotic resistance among *Listeria*, including *Listeria monocytogenes*, in
 470 retail foods. J. Appl. Microbiol. 90, 517-522. <https://doi.org/10.1046/j.1365-2672.2001.01273.x>

471 Lekagul, A., Tangcharoensathien, V., Yeung, S., 2019. Patterns of antibiotic use in global pig production: A systematic review. Vet. An. Science 7,
 472 100058. <https://doi.org/10.1016/j.vas.2019.100058>

473 Scoppetta, F., Sensi, M., Franciosini, M.P., Capuccella, M., 2017. Evaluation of antibiotic usage in swine reproduction farms in Umbria region based
 474 on the quantitative analysis of antimicrobial consumption. Ital J. Food Saf. 6:6886. <https://doi.org/10.4081/ijfs.2017.6886>

475 Kovacevic, J., Sagert, J., Wozniak, A., Gilmour, M.W., Allen, K.J., 2013. Antimicrobial resistance and co-selection phenomenon in *Listeria* spp.
 476 recovered from food and food production environments. Food Microbiol. 34(2), 319-327. <https://doi.org/10.1016/j.fm.2013.01.002>

477 Safdar, A., Armstrong, D., 2003. Listeriosis in patients at a comprehensive cancer center, 1955-1997. Clin. Infect. Dis. 37, 359-364.
 478 <https://doi.org/10.1086/376631>

479 Ayaz, N.D., Erol, I., 2010. Relation between serotype distribution and antibiotic resistance profiles of *Listeria monocytogenes* isolated from ground
480 turkey. J.Food Protect. 73(5), 967-972. <https://doi.org/10.4315/0362-028X-73.5.967>

481 **Table 1. Occurrence of *Listeria monocytogenes* in the different areas, typology and sources investigated in two plants considered**
482 **in this study and serotype and pulsed-field gel electrophoresis results of the 98 isolates included in this study.**
483

Sampling area	Sample types	Sources	Plant A			Plant B		
			No. pos/tot samples(%)	Serotype(n)	P-type	No. pos/tot samples(%)	Serotype(n)	P-type
Slaughterhouse	Slaughtered pigs	faeces	0/135(0) ^a	1/2a(1);1/2b(2)	P27;P28;P29	2/150(1.3) ^a	1/2a(1);1/2c(1)	P03;P05
		amygdales	3/135(2.2) ^a			7/150(4.6) ^a	1/2a(5);1/2c(2)	P02;P03;P06;P07;P08;
		ileo-cecal lymphonodes	n.p.			4/114(3.5)	1/2a(1);1/2c(3)	P04;P09;P20
		total	3/270(1.1) ^a			13/414(3.1) ^a		
	environment	MCS	0/9(0) ^a			0/10(0) ^a		
		gloves, knives, bib, splitting saw	0/18(0) ^a			1/20(5) ^a	1/2c(1)	P04
		total	0/27(0) ^a			1/30(3.3) ^a		
Cutting	total		3/297(1) ^a			14/444(3.1) ^a		
	food	HAC	0/9(0) ^a			4/10(40) ^a	1/2b(3);1/2c(1)	P15;P16
	environment	MCS	1/18(5.5) ^a	1/2a(1)	P35	6/20(30) ^a	1/2b(3);1/2c(3)	P01;P15;P16
		gloves, knives	0/9(0) ^a			7/10(70) ^b	1/2b(3);1/2c(4)	P01;P04;P12;P16
		NMCS	0/9(0) ^a			3/10(30) ^a	1/2c(3)	P01;P23
		total	1/36(2.7) ^a			16/40(40) ^b		
	total		1/45(2.2) ^a			20/50(40) ^b		
Cooling Room	environment	NMCS	0/9(0) ^a			1/10(10) ^a	1/2c(1)	P13
	food	HBT	0/9(0) ^a			8/10(80) ^b	1/2b(4);1/2c(4)	P01;P13;P16
Trimming	environment	MCS	3/9(33.3) ^a	1/2c(3)	P26;P31;P33	6/10(60) ^a	1/2b(3);1/2c(3)	P01;P13;P16;P19
		gloves and knives	3/9(33.3) ^a	1/2c(3)	P26;P32;P34	5/10(50) ^a	1/2b(3);1/2c(3)	P04;P16
		NMCS	3/9(33.3) ^a	1/2c(3)	P01;P26;P30	4/10(40) ^a	1/2b(1);1/2c(3)	P01;P04;P13;P16
		total	9/27(33.3) ^a			15/30(50) ^a		
	total		9/36(25) ^a			23/40(57.5) ^b		
Storage Chill Room	food	HBC	0/100 ^a			27/100(27) ^b	1/2a(4);	P01;P04;P10;P11;P12;P14;

		1/2b(14);1/2c(9 P16;P17;P18;P21;P22;P24;P2 5
Total	13/487(2.6) ^a	85/644(13.2) b

484

485

486

487

488

489

490

491

492

493

n.p.: not performed; MCS: meat contact surface swabs, namely ham conveyor for cutting and trimming areas, and hams rack for cooling room; NMCS: non-meat contact surface swabs, namely drain; HAC: hams after cutting; HBT: hams before trimming; HBC: ham post-trimming and before curing. Number of isolates in rows bearing different letters are significantly different (p< 0.05) between plants A and B.

Table 2. Minimum Inhibitory Concentration distribution of the 98 *Listeria monocytogenes* strains isolated in the present study.

Antimicrobials	MIC												
	0,06	0,12	0,25	0,5	1	2	4	8	16	32	64	128	1000
Erythromycin		94*		2				2*					
Clindamycin	1*		10	31	36	18	2*						
Quinupristin/dalfopristin			1	44	50	3							
Daptomycin					1	17	57	23					
Vancomycin				98*									
Tetracycline					95*					3*			
Ampicillin	56*		42										
Gentamicin					98*								
Levofloxacin				6	58	33			1*				
Linezolid					21	73	4						
Ceftriaxone							9*		32	35	13	9*	
Streptomycin												98*	
Penicillin	5	18	70	5									
Rifampin			98*										
Gatifloxacin				98*									
Ciprofloxacin			26*		30	41	1*						
Trimethoprim/sulfamethoxazole			95*		3								
Oxacillin				1	3	59	35						

494 Black vertical lines indicate breakpoints for resistance. Gray shading indicates that the isolates were not tested for susceptibility to those concentrations of a given
495 antimicrobial agent. Asterisked number indicates the number of isolates exhibiting MIC values equal to or higher or lower than concentration of the test range. No
496 usable breakpoint was for ceftriaxone.

497
498
499
500
501
502
503
504
505
506
507
508
509

Table 3. Antimicrobial resistance, lineage, serotypes and origin of the 98 *Listeria monocytogenes* strains isolated in the present study.

Lineage	Serotype	Origin	N. of isolate s	No. of resistant isolates (%)									R at least one AA	MDR
				Macrolides	Lincosamides	Lipopeptides	Fluoroquinolones	Tetracycline	Folate Pathway Inhibitors	Oxazolidinone s	Penicillins			
				ERY	CLI	DAP	LEVO	CIP	TET	SXT	LZD	OXA		
I	1/2b	Slaughtered pigs	2	-	1(50)	-	1(50)	1(50)	-	-	-	-	2(100)	-
		Environmen t	13	-	-	5(38.5)	-	-	-	-	-	13(100)	13(100)	-
		Food	21	-	4(19)	11(52.4)	1(4.8)	3(14.3)	-	-	-	15(71.4)	20(95)	1(4.8)
	Total	36	-	5(13.8) ^{a,1}	16(44.4) ^{a,1}	2(5.5) ^{a,1}	4(11) ^{a,1}	-	-	-	28(77.7) ^{a,1}	35(97.2) ^{a,1}	1(2.7) ^{a,1}	
II	1/2a	Slaughtered pigs	8	2(25)	6(75)	3(37.5)	4(50)	3(37.5)	2(25)	2(25)	1(12.5)	1(12.5)	8(100)	3(37.5)
		Environmen t	1	-	-	-	-	-	-	-	-	-	-	-
		Food	4	-	-	1(25)	-	-	-	-	-	4(100)	4(100)	-
		Total	13	2(15.3)	6(46.1) ^b	4(30.7) ^a	4(30.7) ^a	3(23) ^b	2(15.3) ^a	2(15.3) ^a	1(7.6) ^a	5(38.4) ^a	12(92.3) ^a	3(23) ^b
	1/2c	Slaughtered pigs	6	-	6(100)	2(33.3)	3(50)	5(83.3)	-	-	2(33.3)	2(33.3)	6(100)	4(66.7)
		Environmen t	29	-	25(86)	1(3.4)	17(58.6)	20(68.9)	1(3.4)	-	1(3.4)	-	27(90)	2(6.7)
		Food	14	-	14(100)	-	8(57.1)	10(71.4)	-	1(7.1)	-	-	14(100)	1(7.1)

	Total	49	-	45(91.8) ^c	3(6.1) ^b	28(57.1) _b	35(71.4) _c	1(2) ^a	1(2) ^a	3(6.1) ^a	2(4) ^b	47(95.9) ^a	7(14.2) ^b
	Total	62	2(3.2)	51(82.2) ²	7(11.2) ²	32(51.6) ₂	38(61.2) ₂	3(4.8)	3(4.8)	4(6.4)	7(11.2) ¹	59(95.1) ¹	10(16.1) ₂
	Total	98	2(2.04)	56(57.1)	23(23.5)	34(34.7)	42(42.9)	3(3.1)	3(3.1)	4(4)	35(35.7)	93(94.9)	11(11.2)

510 ERY: erythromycin; DAP: daptomycin; CLI: clindamycin; LEVO: levofloxacin; CIP: ciprofloxacin; TET: tetracycline; SXT: trimethoprim/sulfamethoxazole;
511 LZD: linezolid; OXA: oxacillin; AA: antimicrobial agent; MDR: multidrug resistant strains; - : not detected; number of isolates in columns bearing different
512 numbers and letters are significantly different (p< 0.05) in relationship to respectively lineage and serotypes
513
514
515
516
517
518
519
520
521
522
523

524 **Table 4. Antimicrobial resistance and origin of the 98 *Listeria monocytogenes* strains isolated in the present study.**
525

AA	Slaughtered pigs	Environment				Food
		MCS	Utensil	NMCS	Total	
Erythromycin	2(12.59)	-	-	-	-	-
Clindamycin	13(81.3)	10(58.8)	8(50)	7(70)	25(58.1)	18(46.2)*
Daptomycin	5(31.3)	1(5.9)	3(18.8)	2(20)	6(14)	12(30.8)
Levofloxacin	8(50)	7(41.2)	7(43.8)	3(30)	17(39.5)	9(23.1)
Ciprofloxacin	9(56.3)	7(41.2)	9(56.3)	4(40)	20(46.5)	13(33.3)
Tetracycline	2(12.5)	-	1(6.3)	-	1(2.3)	-
Trimethoprim/sulfamethoxazole	2(12.5)	-	-	-	-	1(7.1)
Linezolid	3(18.8)	-	1(6.3)	-	1(2.3)	-
Oxacillin	3(18.8)	6(35.3)	6(37.5)	1(10)	13(30.2)	19(48.7)
R at least one AA	16(100)	16 (94)	16(100)	8(80)	40(93)	38(97.4)
MDR	7(63)*	-	1(9)	1(9)	2(18)	2(18)

526 MCS: meat contact surface swabs, namely ham conveyor for cutting and trimming areas, and hams rack for chilling room; NMCS: non-meat contact
527 surface swabs, namely drain; AA: antimicrobial agent; - : not detected; number of isolates bearing * are significantly different (p< 0.05) in relationship to
528 the source

529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551

Table 5. Antimicrobial resistance and pulsed-field gel electrophoresis results of the 98 *Listeria monocytogenes* strains isolated in the present study.

Cluster	Isolates	No. of resistant isolates (%)									R at least one AA	MDR
		ERY	CLI	DAP	LEVO	CIP	TET	SXT	LZD	OXA		
1	2	-	2(3.6)	1(4.3)	2(5.8)	1(2.3)	-	-	1(25)	-	2	1
2	2	-	1(1.7)	-	-	-	-	-	-	-	1	-
3	27	-	26(46.4)	-	18(52.9)	20(47.6)	-	-	-	-	26	-
4	13	-	12(21.4)	-	10(29.4)	13(30)	-	1(33.3)	2(50)	-	13	3
5	2	2(100)	2(3.6)	1(4.3)	2(5.8)	2(4.7)	2(66.7)	2(66.7)	-	-	2	2
6	34	-	-	16(69.6)	-	-	-	-	-	32(91.4)	33	-
7	3	-	3(5.4)	2(8.6)	-	2(4.7)	-	-	-	-	3	2
8	1	-	-	1(4.3)	-	-	-	-	-	1(2.8)	1	-
9	2	-	2(3.6)	2(8.6)	-	1(2.3)	-	-	-	2(5.7)	2	2
10	2	-	2(3.6)	-	-	-	-	-	-	-	2	-
11	1	-	-	-	1(2.9)	1(2.3)	1(33.3)	-	1(25)	-	1	1

12	3	-	2(3.6)	-	-	1(2.3)	-	-	-	-	2	-
13	2	-	2(3.6)	-	-	-	-	-	-	-	2	-
14	1	-	-	-	-	-	-	-	-	-	-	-
15	2	-	2(3.6)	-	-	-	-	-	-	-	2	-
16	1	-	-	-	1(2.9)	1(2.3)	-	-	-	-	1	-
98		2	56	23	34	42	3	3	4	34	93	11

552 ERY: erythromycin; DAP: daptomycin; CLI: clindamycin; LEVO: levofloxacin; CIP: ciprofloxacin; TET: tetracycline; SXT: trimethoprim/sulfamethoxazole;
553 LZD: linezolid; OXA: oxacillin
554

555

556

557

558