

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Modeling the behavior of *Listeria innocua* in Italian salami during the production and high-pressure validation of processes for exportation to the U.S

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Bonilauri, P., Meriardi, G., Ramini, M., Bardasi, L., Taddei, R., Grisenti, M.S., et al. (2021). Modeling the behavior of *Listeria innocua* in Italian salami during the production and high-pressure validation of processes for exportation to the U.S. *MEAT SCIENCE*, 172, 1-9 [10.1016/j.meatsci.2020.108315].

Availability:

This version is available at: <https://hdl.handle.net/11585/789136> since: 2021-01-17

Published:

DOI: <http://doi.org/10.1016/j.meatsci.2020.108315>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1 **Modeling the behavior of *Listeria innocua* in Italian salami during the production and High-**
2 **Pressure validation of processes for exportation to the U.S.**

3

4 Paolo Bonilauri¹, Giuseppe Merialdi², Mattia Ramini², Lia Bardasi², Roberta Taddei², Maria Silvia
5 Grisenti³, Paolo Daminelli⁴, Elena Cosciani-Cunico⁴, Elena Dalzini⁵, Maria Angela Frustoli³,
6 Federica Giacometti^{6*}, Federico Tomasello⁶, Silvia Piva⁶, Andrea Serraino⁶.

7 1 Institute for Zooprophyllaxis in Lombardy and Emilia Romagna- Via Pitagora 2 - 42100 Reggio
8 Emilia – Italy

9 2 Experimental Institute for Zooprophyllaxis in Lombardy and Emilia Romagna, Via Fiorini 5 40127
10 Bologna – Italy

11 3 SSICA Stazione Sperimentale per l’Industria delle Conserve Alimentari V.le Tanara 31/a 43121
12 Parma.

13 4 Experimental Institute for Zooprophyllaxis in Lombardy and Emilia-Romagna - Via Bianchi 7/9 -
14 25124 Brescia – Italy

15 5 National Reference Centre for Emerging Risks in Food Safety, Experimental Institute for
16 Zooprophyllaxis in Lombardy and Emilia-Romagna - Via Celoria 12, 20133, Milan – Italy

17 6 Department of Veterinary Medical Sciences - University of Bologna - Via Tolara di Sopra 50 -
18 40064 Ozzano Emilia (BO) – Italy. Tel. +39 51 2097320; fax +39 51 2097346;

19

20 * Corresponding author: federica.giacometti3@unibo.it (F. Giacometti).

21

22 **Declaration of interest**

23 None

24 **Funding**

25 This research did not receive any grant from funding agencies in the public, commercial, or not-for-
26 profit sectors.

27

28 **Abstract**

29 The study evaluated 51 Italian commercial salami processes in association with High-Pressure
30 Processing (HPP) to assess *Listeria innocua* lethality. We used data to model the decrease of *Listeria*
31 *monocytogenes* according to process parameters. At the end of the process, the total *L. innocua*
32 reduction always resulted in > 1 Log₁₀ CFU/g (1.04-5.68). In the univariate analysis, we observed a
33 significant association between the decrease of *L. innocua* count, a_w at the end of the
34 acidification/drying process, a_w decrease during seasoning, duration of the seasoning, trimming and
35 caliber. HPP further reduced the *L. innocua* count by 0.48-3.47 Log₁₀ CFU/g. The model represents
36 a useful tool for enterprises and Authorities to evaluate the efficacy of processes to reduce *L.*
37 *monocytogenes*, predicting its load at the end of the process and the need for a process modification
38 or for the addition of a final lethal process. The model of HPP treatment predicts treatment efficacy
39 based on pH and a_w of the product.

40

41 **Keywords:** fermented sausages, Italian salami, *Listeria innocua*, *Listeria monocytogenes*, challenge
42 test, linear regression model.

43

44 **1. Introduction**

45 *Listeria monocytogenes* is the causative agent of foodborne listeriosis; in healthy individuals,
46 infection is usually mild, but it is possible to observe severe consequences in fetuses, infants,
47 immunocompromised patients, elderly, and pregnant women with a high mortality rate (Tolvanen et
48 al., 2008).

49 *L. monocytogenes* frequently contaminates meat and meat products, including fermented sausages.
50 Such contamination occurs via raw materials, ingredients, and processing equipment. Furthermore, it
51 can also be related to the post-processing phase (Hwang et al., 2009; Lin et al., 2006; Prencipe et al.,
52 2012). *L. monocytogenes* is often isolated from fermented meat products due to its capability to
53 survive their adverse conditions (Degenhardt & Sant'Anna, 2007), and the presence of *L.*
54 *monocytogenes* in ready-to-eat (RTE) products, like salami, poses a risk to the consumer. Several
55 studies have reported the presence of *L. monocytogenes* in fermented sausages worldwide, with
56 prevalence reaching up to 60% (Bohaychuk et al., 2006; De Cesare, Mioni, & Manfreda, 2007;
57 Doménech, Jimenez -Belenguer, Amoros, Ferrus, & Escriche, 2015; Ferreira et al., 2007; Glass &
58 Doyle, 1989; Gounadaki, Skandamis, Drosinos, & Nychas, 2008; Martin, Garriga, & Aymerich,
59 2011; Meloni, 2015; Meloni et al., 2012, 2014).

60 The European Union food safety Regulations admit the presence of *L. monocytogenes* in RTE
61 products that do not sustain its growth, to a concentration not exceeding 100 CFU/g throughout the
62 defined shelf life. Regardless of whether or not they support the growth of the pathogen, the U.S.
63 legislation applies a zero-tolerance approach for RTE products. According to the guidance document
64 of the U.S. Department of Agriculture Food Safety and Inspection Service (FSIS & USDA, 2017),
65 establishments producing dry-cured meat products should implement a process addressing lethality
66 of *L. monocytogenes*, to ensure the safety of products for consumption. In this context, lethality is
67 defined as the process or combination of processes, ensuring a specific and significant reduction in
68 the number of pathogens in products.

69 Although literature frequently reported the presence of *L. monocytogenes* in dry and semi-dry
70 fermented sausages, these products are considered at low risk for foodborne listeriosis due to their
71 intrinsic properties (Barmpalia-Davis, Geornaras, Kendall, & Sofos, 2008; Simpson et al., 2008). On
72 the other hand, fermented sausages were traced as a possible source of a listeriosis outbreak that
73 involved 36 individuals in Philadelphia with four cases of death (Schwartz et al., 1989), while in
74 2010, two cases were linked to salami consumption in Ontario
75 ([http://outbreakdatabase.com/details/siena-foods-salame-](http://outbreakdatabase.com/details/siena-foods-salame-2010/?organism=Listeria+monocytogenes&vehicle=sausage)
76 [2010/?organism=Listeria+monocytogenes&vehicle=sausage](http://outbreakdatabase.com/details/siena-foods-salame-2010/?organism=Listeria+monocytogenes&vehicle=sausage)). Besides, the economic impact due to
77 the recall of meat products contaminated with *L. monocytogenes*, despite their ability to cause
78 illnesses, should be considered.

79 Several studies evaluated the fate of *L. monocytogenes* in dry sausage manufacturing processes, but
80 their results vary considerably. Although most studies observed a reduction of *L. monocytogenes*
81 contamination, others demonstrated the ability of *L. monocytogenes* to survive when the initial load
82 of *L. monocytogenes* in raw materials was high (Nightingale, Thippareddi, Phebus, Marsden, &
83 Nutsch, 2006); moreover, in some cases, growth was shown during processing (Campanini,
84 Pedrazzoni, Barbuti, & Baldini, 1993; Nissen & Holck, 1998). Literature indicates that *L.*
85 *monocytogenes* can contaminate fermented sausages and may not be completely eliminated during
86 processing. Consequently, to comply with the zero-tolerance policy requested by the U.S. Federal
87 Authorities, additional treatments must be necessary, including the implementation of post-
88 processing operations (Hereu, Bover-Cid, Garriga, & Aymerich, 2012). High-Pressure Processing
89 (HPP) can be used on several products to improve their microbiological characteristics without a
90 significant modification of their organoleptic features (Hayman, Baxter, O’Riordan, & Stewart,
91 2004).

92 The variability of process procedures and parameters in salami production highlights the need for
93 process validation research to reduce the prevalence of *L. monocytogenes* contamination and the risk

94 for consumers (Nightingale, Thippareddi, Phebus, Marsden, & Nutsch, 2006). The use of predictive
95 models explaining the effect of a wide range of food variables can help producers to plan their
96 formulae or process to achieve the goal of controlling *L. monocytogenes* contamination (Novelli et
97 al., 2017). The objective of this study was, through the evaluation of 51 Italian commercial salami
98 production processes, also in association with HPP, to assess the reduction of *Listeria innocua* during
99 manufacturing; we used processing data, and *L. innocua* count to develop a model that describes the
100 evolution of *Listeria* population according to process parameters.

101

102 **2. Materials and methods**

103 **2.1 Salami production and inoculation**

104 Data on ingredients and processing parameters of 51 different Italian salami were collected from
105 thirteen manufacturers as previously described by Bonilauri et al. (2019); briefly, thirteen enterprises
106 producing fermented sausages for the European market and exportation to the U.S. were involved in
107 the study, and they were requested to supply the laboratory with the ingredient mix (sausage mix) to
108 be inoculated and stuffed. We performed a total of 51 different challenge tests, and table 1 shows the
109 details of ingredients and process parameters. The sausage mix was stored, refrigerated, and
110 transported to the laboratory within 24 hours after production.

111 The sausage mix was inoculated with a blend of five strains of *Listeria innocua* isolates used as the
112 surrogate of *L. monocytogenes* (Hu & Gurtler, 2017; Lebow et al., 2017; Merialdi, Ramini, Ravanetti,
113 Gherri, & Bonilauri, 2015). More specifically, the laboratory used the following isolates: IZSLER
114 111373/1 and IZSLER 111373/2 isolated from environmental swabs collected in a pork meat
115 transformation plant, IZSLER 257529/1 isolated from fresh pork sausages, IZSLER 257529/2
116 isolated from fresh pork meat and a collection strain ATCC 33090. The bacterial cultures were
117 prepared following what indicated by Bonilauri et al. (2019). The enumeration of *L. innocua* count

118 was performed by serial decimal dilution and inoculation on Agar *Listeria* Ottaviani & Agosti
119 (Biolife, Milan, Italy) plates, according to 11290-2 (ISO, 2017a). Following the study of Bonilauri
120 and Colleagues (2019), we prepared a multi-strain cocktail; then, we stuffed the mix, and sausages
121 were acidified/dried and seasoned according to the instructions of each enterprise. At the end of
122 seasoning, 5 inoculated sausages were vacuum packed and used for HPP at 600 MPa for 300'' (Bover-
123 Cid, Belletti, Aymerich, & Garriga, 2015; Meriardi, Ramini, Ravanetti, Gherri & Bonilauri, 2015;
124 Rubio, Possas, Rincón, García-Gímeno & Martínez, 2018) by commercial apparatus. At the moment
125 of HPP, the temperature of fermented sausages and treatment water was 4 and 14°C, respectively.

126 **2.2 Sampling, physicochemical and microbiological examination**

127 Five samples (of 25 g each) for each sausage mix were tested for the presence/absence of *Listeria*
128 spp. according to according to ISO 11290-1 (ISO, 2017b) to evaluate the natural contamination of
129 meat.

130 To determine pH and a_w values and to evaluate *L. innocua* count, for each challenge test, we collected
131 and analyzed after the inoculation with *L. innocua* strains a total of 3 samples per sampling time
132 namely; before stuffing (1), at the end of the acidification step (2) and at the end of seasoning (3),
133 and, after HPP treatment, a total of 5 samples. Physicochemical analyses (pH and a_w) were performed
134 as stated by Bonilauri and Colleagues (2019). *L. innocua* count was performed by serial dilution and
135 direct surface plating onto Agar *Listeria* Ottaviani & Agosti (Biolife, Milan, Italy) plates according
136 to ISO 11290-2 (ISO, 2017a). Whenever the count showed a result equal to < 10 CFU/g of *L. innocua*,
137 we analyzed samples to detect *L. innocua* (presence/absence) 25g of each. *L. innocua* counts were
138 expressed in CFU/g and converted into Log₁₀ CFU/g. For the statistical elaboration of negative
139 *Listeria* counts, a presence/absence test was performed and the result was reported respectively as 5
140 CFU/g (Log₁₀ = 0.70) and 0.04 CFU/g (Log₁₀ = -1.40) in case of presence or absence of the
141 pathogen.

142

143 **2.3 Data collection and the linear regression model**

144 Data (mean of the 3 or 5 samples) on *L. innocua* count, pH, and a_w values were collected and used
145 for the linear regression model as detailed in the study of Bonilauri and Colleagues (2019) and as
146 outlined in Figure 1. Briefly, standard deviation (SD) of the *L. innocua* count obtained from the 3 or
147 5 samples of each challenge test was calculated; we considered the SD acceptable when $<0.5 \text{ Log}_{10}$
148 CFU/g. We calculated the Δ_a difference as well as Δ_s difference between *L. innocua* concentrations
149 between the different process steps as extensively described in the study of Bonilauri and
150 Colleagues (2019) and as summarized in Figure 1.

151 We used a linear regression model to describe the *L. innocua* count decrease observed during the
152 production process. In such a model, we reported the dependent and the independent variables in
153 Figure 1. Then, we used the variables that in the univariate model had a p-value <0.10 in a
154 multivariate stepwise linear regression with backward elimination. We considered second- and
155 third-order models (polynomial regression), and the polynomial regression model was kept only in
156 case of a significant increase in the coefficient of determination (R²). We tested a pair of
157 independent variables to detect co-linearity, while interactions were tested in those cases where two
158 or more factors were significant. Intercooled STATA 7.0 (Statacorp) was used to calculate the
159 coefficient of determination (R²) and significance of the model (F-statistic) ($p < 0.05$).

160

161 **3. Results**

162 **3.1 Salami productions experiment**

163 We observed no association between the decrease of *L. innocua* count after the acidification/drying
164 phase and other independent variables; t_1 and t_2 , maximum, minimum, and mean temperature during
165 acidification/drying and seasoning phase, caliber, and trimming. The concentration of NaCl, nitrate,
166 and nitrite used by the different enterprises was very similar, and thus, it was not possible to show an
167 association with *L. innocua* count.

168 Table 2 reports the results of the *L. innocua* count decrease after acidification/drying, seasoning, and
169 HPP and the associated pH and a_w measurements (before stuffing, after acidification/drying, and after
170 seasoning). Each type of salami has very different process parameters, and also, within the same type
171 of salami, a high variability of process parameters between the different manufacturers has to be
172 mentioned.

173 Before stuffing, a_w and pH values ranged respectively from 5.57 to 6.11 and from 0.963 to 0.979 (see
174 Table 2). The acidification/drying phase had a variable length between 5 and 17 days and led the pH
175 and a_w values, respectively, to 4.78 and 5.41, and 0.936 and 0.971 (see Table 2).

176 After seasoning, it varied from a minimum of 20 to a maximum of 90 days, and pH values showed a
177 general increase that reached values ranging from 4.99 to 6.53, whereas a_w values were reduced by
178 0.880 to 0.936 (see Table 2).

179 The observed *L. innocua* counts always resulted in an SD <0.5 Log₁₀ CFU/g, and therefore, in the
180 text and tables, data are reported exclusively as mean. A decrease of *L. innocua* count could be
181 observed during acidification/drying by 0.02-2.20 Log₁₀ CFU/g (median 0.63), and, this decrease
182 resulted statistically significant (p<0.05) in the univariate analysis reporting an association between
183 the decrease of *L. innocua* count and the pH value after acidification/drying (pH₂), ΔpH₁-pH₂, water
184 activity at the end of the acidification/drying process (a_w 2), and trimming (mm). Even pH₂ and Δ
185 pH₁-pH₂ results were correlated, and thus, in the multivariate analysis, it was used Δ pH₁-pH₂
186 exclusively.

187 During acidification/drying phase, the observed *L. innocua* decrease could be outlined by a highly
188 significant equation with about 50% of the variability that could be explicated by the model (R² =
189 0.48; p<0.00001):

190
$$\Delta_a = 10.107 + 1.564 \Delta \text{pH1-pH2} - 10.607 a_w 2 - 0.052 \text{ trimming (mm)}$$

191 The White general test (White, 1980) is used for heteroscedastic errors in the regression by the model
192 (White general test statistic: 11.10227 Chi-sq(9) P -value = 0.27), and in the independence test,
193 residuals resulted uncorrelated with all the previous inputs.

194 In Figure 2, the *L. innocua* decay (Δa) observed at the end of the acidification/drying process was
195 reported by a graph of residuals against data fitted into the multivariate model.

196 Considering polynomial regression or the interaction between variables, no significant increase of the
197 model was observed.

198 During seasoning, it was observed a *L. innocua* reduction, varying from 0.02 to 3.4 Log₁₀ CFU/g
199 (median 0.53); only in three cases (see tests 6, 23, 43 in Table 2) the *L. innocua* count slightly
200 increased during seasoning, but this apparent growth did not reach 0.5 Log₁₀ CFU/g and, therefore,
201 these three cases were not included in the regression analysis (EURL Lm, 2014).

202 In the univariate analysis, the decrease of *L. innocua* count (Δ_s) resulted significantly ($p < 0.05$)
203 associated with water activity at the end of the acidification/drying process ($a_w 2$), $\Delta a_w 2 - a_w 3$,
204 duration of the seasoning in days (t_s), trimming (mm), and caliber (mm). $a_w 2$ and $\Delta a_w 2 - a_w 3$,
205 seasoning in days (t_s) and caliber (mm) results autocorrelated, so only the $\Delta a_w 2 - a_w 3$ and caliber
206 (mm) were used in the multivariate analysis.

207 This highly significant equation here reported describes the reduction of *L. innocua* during the
208 seasoning phase even if it only explains about 30% of the variability ($R^2 = 0.32$; $p < 0.0007$):

$$209 \Delta_s = 22.87 \Delta a_w 2 - a_w 3 + 0.008 \text{ caliber (mm)} - 10.607 a_w 2 - 0.078 \text{ trimming(mm)} - 0.092.$$

210 Again, the White general test (White, 1980) is used for heteroscedastic errors in the regression by the
211 model (White general test statistic: 15.89082 Chi-sq(9) P -value = .07), and in the independence test,
212 residuals resulted not correlated with all the past inputs. In Figure 3, the graph of residuals against
213 fitted values of the multivariate model reports the *L. innocua* decay (Δ_s) at the end of the seasoning

214 process. However, when polynomial regression or interaction between variables was considered, the
215 model did not show a significant increase.

216

217 **3.2 HPP of Salami at the end of the seasoning phase**

218 We observed a further decrease of *L. innocua* count after HPP (0.48-3.47 Log₁₀ CFU/g). The efficacy
219 of HPP resulted associated with pH₂, pH₃, a_w 2, a_w 3, caliber (mm), and duration of
220 acidification/drying phase (days) in the univariate analysis, while only a_w 2, a_w 3 and pH₂ remained
221 significant in multivariate analysis. In this way, the reduction of *L. innocua* after HPP process could
222 be described by the model with almost 36% of the variability resulted by the following highly
223 significant equation (R² = 0.357; p=0.0001):

$$224 \Delta h = 2.069 \text{ pH}_2 + 27.188 a_w 2 + 24.431 a_w 3 - 57.246$$

225 In the model, the test for heteroscedasticity of the regression (White, 1980) (White general test
226 statistic: 9.416029 Chi-sq(9) *P*-value = .40) was passed; as before, residuals resulted not correlated
227 with all the past inputs in the independence test.

228 The graph of residuals against fitted values of the multivariate model, which describes the *L. innocua*
229 decay (Δ s) observed at the end of the s acidification/drying process, is reported in Figure 4. No
230 significant increase of the model was observed by polynomial regression or interaction between
231 variables. This relation means that the higher are pH and a_w after the acidification/drying and
232 seasoning phases, respectively, the higher will be the reduction of *L. innocua* after HPP.

233

234 **3.4 Global model**

235 The reduction of *L. innocua* globally observed during the validation experiment (acidification/drying,
236 seasoning and HPP treatment) could be described by the combination of the three models:

237 $Dt = D_a + D_s + D_h = 2.069 \text{ pH}_2 + 1.564 \Delta \text{ pH}_1\text{-pH}_2 + 16.5808 a_w^2 + 24.431 a_w^3 +$
238 $22.8704 \Delta a_w^2 - a_w^3 - 0.1292 \text{ trimming(mm)} + 0.00793 \text{ caliber (mm)} - 47.2304$, in Figure 5, it is
239 possible to observe the graph of residuals against fitted values of the global model that describes
240 approximately 65% of the total variability ($R^2=0.648$). Finally, *L. innocua* count reduction always
241 resulted > 1 Log₁₀ CFU/g, varying from 1.04 to of 5.68.

242

243 4. Discussion

244 Data on the reduction of *L. innocua* during the processing of fifty-one challenge studies and after
245 HPP were collected: we used the overall data to define a model that could be utilized and could be
246 useful to predict the fate of *L. monocytogenes* in other types of salami or other enterprises.

247 In our study, we noted an overall reduction of *L. innocua* count during processing
248 (acidification/drying plus seasoning) in all the challenge tests performed, ranging from 0.34-4.32
249 Log₁₀ CFU/g. These results are roughly in agreement with previously reported studies on fermented
250 sausages for example: in Soudjouk style fermented sausage a 0-1.86 Log CFU/g reduction of *L.*
251 *monocytogenes* during production was reported (Hwang et al., 2009); Porto-Fett et al., 2010 reported
252 a 1.1 to 2.2 reduction during fermentation and drying of Genoa salami; and Johnson et al. (Johnson,
253 Doyle, Cassens, & Schoeni, 1988) reported a 1.2-1.8 reduction in salami; in Chouriço de Vinho
254 Garcia Diez & Patarata (2013) it was observed a reduction of about 2 Log CFU/g, higher when
255 *Lactobacillus sakei* was present as a starter culture. In Italian salami (Cacciatore and Felino), a < 1 -
256 log reduction was observed during the production process (Mataragas et al., 2015). Some authors
257 (Degenhardt & Sant'Anna, 2007) observed a slightly higher reduction of *L. monocytogenes* in Italian
258 sausages (2.57-3.81 log at final pH 5.10-5.16 and a_w at 0.883-0.897) and, in some cases, a growth
259 during processing was shown (Campanini, Pedrazzoni, Barbuti, & Baldini, 1993; Nissen & Holck,
260 1998).

261 In traditional dry-cured fermented sausages, salt concentration, pH, and a_w are the main parameters
262 controlling pathogenic microorganisms (Messier, Smith, & Tittiger, 1989) and also *L. monocytogenes*
263 (Lindqvist & Lindblad, 2009). Additional hurdles that should be considered are nitrite, and another
264 preservative addition, the temperature of fermentation, length of the fermentation and drying phases,
265 smoking, starter addition; also an influence of polyphosphate addition was reported (Gonzales-Barron
266 et al., 2015); many of these last variables are more or less related to the pH and a_w reduction during
267 processing.

268 In our study, we did not show any association between the concentration of NaCl, nitrate, and nitrite
269 in the sausage mix, and the decrease of *L. innocua* count to be quite similar in the production processes
270 of different types of salami. All the investigated enterprises used starter cultures. The duration and
271 temperature of the acidification/drying process, as well as the temperature of the seasoning process,
272 resulted not to be related to the reduction of *L. innocua* in the multivariate analysis. The reduction of
273 *L. innocua* during processing turned out to be related to the pH value at the end of acidification/drying
274 (pH₂ or Δ pH₁-pH₂), to water activity at the end of the seasoning process (a_{w3} or $\Delta a_{w2} - a_{w3}$), to the
275 caliber of salami, and the duration of the seasoning in days (ts) which autocorrelated. In none of the
276 challenge tests performed during acidification/drying phase, pH and a_w reached the no-growth limits
277 indicated by EU Regulation 2073/2005 (Regulation (EC) 2073/2005/EC): pH \leq 4.4 or $a_w \leq$ 0.92 or pH
278 \leq 5.0 and \leq 0.94; despite this, in all the tests carried out, a reduction of *L. innocua* count was observed
279 in the acidification/drying phase (0.02-2.20 Log₁₀ CFU/g median 0.63). A further reduction was
280 observed during seasoning (0.02-3.4 Log₁₀ CFU/g median 0.53).

281 These results are generally in agreement with the results reported in the literature. In Linguíçola, a
282 Portuguese traditional dry fermented sausage, at least three hurdles determined the evolution of *L.*
283 *monocytogenes* population (low a_w , low pH and nitrite at an input level of about 150 ppm), and other
284 factors contributed to its control like a more prolonged ripening and maceration period, a_w at the end
285 of smoking (Gonzales-Barron et al., 2015). Hwang et al. (2009) reported that the decrease of *L.*

286 *monocytogenes* was significantly related to pH and a_w during the production of Soudjouk style
287 fermented sausage and that *L. monocytogenes* decrease occurred after pH was lowered to pH 5.1; a
288 significant lethality was observed at $a_w < 0.92$. Garriga et al. (2005) highlighted the importance of the
289 decrease of pH during the first 7 days of ripening for the prevention of *L. monocytogenes*
290 multiplication, suggesting the use of starter cultures in Mediterranean fermented sausages; also
291 Chikthimmah, Guyer, & Knabel (2001) reported that fermentation alone at pH 4.7 allowed a 2.4-log
292 reduction of *L. monocytogenes* load.

293 Based on an analysis of studies in literature carried out by Mataragas et al. (2015) relating the
294 inactivation rate of *L. monocytogenes* in fermented sausages, to the temperature of fermentation, pH
295 decrease in the fermentation phase and a_w decrease; temperatures above 20°C, especially in the first
296 48 hours of fermentation, turned out to be necessary for rapid inactivation of *L. monocytogenes*. The
297 same authors reported that the relation of pH and a_w with *L. monocytogenes* decrease appears to be
298 temperature-related and that temperature is related to *L. monocytogenes* reduction when pH and a_w
299 values are in the range that prevents *L. monocytogenes* growth. In our study the decrease of *L. innocua*
300 count resulted not related to the temperature of fermentation: we should consider that pH values at
301 the end of fermentation never resulted lower than the no-growth limit for *L. monocytogenes* (4.4)
302 resulting in 12 tests < 5.0 (4.78 – 4.98) and in 39 tests > 5.00 . A fundamental aspect that should be
303 mentioned is the relation of higher trimming and higher *L. innocua* load reduction; it is known that
304 physical status can influence bacterial behavior in food: in our study, we observed better acidification
305 in Milano type salami with their typical thin grain.

306 Italian salami has a higher pH (Garriga et al., 2005) in comparison to northern European sausages,
307 which are characterized by a sharper and faster pH reduction (Holck et al., 2011). In Mediterranean
308 sausages, the a_w reduction and the increase in aging can be more critical than in reducing *L.*
309 *monocytogenes* contamination (Mataragas et al., 2015; Nightingale, Thippareddi, Phebus, Marsden,
310 & Nutsch, 2006). In our study, the a_w reached values between 0.88-0.952 in different types of salami

311 at the end of seasoning with $a_w > 0.92$ in only three challenge tests (median of 51 challenge tests 0.92).
312 The importance of a_w in controlling *L. monocytogenes* was reported in almost all studies on fermented
313 sausages (Degenhardt & Sant'Anna, 2007; Gonzales-Barron et al., 2015; Mataragas et al., 2015;
314 Nightingale, Thippareddi, Phebus, Marsden, & Nutsch, 2006; Novelli et al., 2017; Porto-Fett et al.,
315 2010; Roccatto et al., 2017).

316 HPP was able to reduce further the *L. innocua* count by 0.48-3.47 Log₁₀ CFU/g in all the challenge
317 tests with efficacy that resulted directly associated with a_w 3 (a_w at the end of seasoning) and pH2 (pH
318 at the end of acidification/drying). The few references available on meat products, and dried
319 fermented sausages showed similar results: in cooked ham Jofré et al. (2008) showed a 3.5 Log CFU/g
320 reduction through treatment at 600 MPa for 5 minutes and a 3.4 log (400 MPa for 10 minutes)
321 reduction was observed by Marcos et al. (2008). Bover-Cid et al. (2015) showed a reduction of *L.*
322 *monocytogenes* in dry-cured ham at 600MPa for 5 minutes ranging from 2.24 log to 6.82 Log CFU/g
323 depending on the a_w and fat content of the ham.

324 Regarding fermented, dried products, the following studies reported a 1.6 – 5.0 Log CFU/g reduction
325 at 600MPa for 5 min in Genoa salami (Porto-Fett et al., 2010), a 1.79-3.15 Log CFU/g reduction in
326 Spanish chorizo at 600 MPa for 5-10 minutes (Rubio, Possas, Rincón, García-Gímeno, & Martínez,
327 2018), and a 0.9 Log CFU/g reduction in slightly fermented sausages at 400MPa for 10 minutes
328 (Garriga et al., 2005).

329 Several studies have shown that *L. monocytogenes* baroresistance increases when meat products
330 present a low a_w . Findings demonstrated that water activity could influence the efficacy of HPP
331 treatment and that low water activity protects microorganisms from environmental stresses. Bover-
332 Cid et al. (2015) showed that a_w affected the efficacy of HPP in *L. monocytogenes* inactivation in dry-
333 cured ham at different pressures; for example, a difference of more than 4 Log CFU/g of inactivation
334 was registered with an a_w of 0.960 vs an a_w of 0.860 at 600 MPa. An increase of a_w from 0.79 to 0.92
335 raised the reduction of *L. monocytogenes* from 0.07 to 2.17 Log in Spanish Chorizo sausage at 475

336 MPa (Rubio, Possas, Rincón, García-Gímeno, & Martínez, 2018); similarly, an increase of a_w from
337 0.88 to about 0.92 decreased *L. monocytogenes* significantly in Genoa salami (Porto-Fett et al., 2010).

338 To our knowledge, no data are available on the influence of pH on the baroresistance of *L.*
339 *monocytogenes* in fermented meat products, but in other food low pH resulted in a noticeable
340 synergistic effect with pressure on the inactivation of *L. monocytogenes* (Dogan & Erkmen, 2004;
341 Gao, Ju, & Wu-Ding, 2007; Xu, Hyeon-Yong, & Ahn, 2009). The combination of low pH values with
342 HHP processing resulted in a higher efficacy of this technology for reducing *L. monocytogenes* levels
343 (Possas, Pérez-Rodríguez, Valero, & García-Gimeno, 2017); this relation should be further
344 investigated in subsequent studies to improve the accuracy models on the survival of *L.*
345 *monocytogenes* in fermented meat products undergoing HPP.

346 Usually, *L. monocytogenes* contamination in raw meat is approximately < 100 CFU (Farber &
347 Peterkin, 1991). Whereas some processes are adequate to ensure the absence of *L. monocytogenes*
348 others may permit the survival of *L. monocytogenes* in the final product, notably when raw materials
349 are highly contaminated (Glass & Doyle, 1989; Johnson, Doyle, Cassens, & Schoeni, 1988) and
350 additional treatment could be necessary. In this work, based on 51 challenge tests, we demonstrated
351 the ability of Italian type salami production process with a final HPP treatment aimed at addressing
352 lethality of *L. monocytogenes*, in order to validate products for export to the U.S.

353 Control of *L. monocytogenes* during the Italian Salami production process was attributed mainly to
354 the decrease of pH during the fermentation and low a_w at the end of seasoning. Differences in
355 fermentation and drying parameters in Italian style salami studied by different authors, together with
356 strain resistance variation, may explain the different results reported in the literature (Nightingale,
357 Thippareddi, Phebus, Marsden, & Nutsch, 2006). In this study, we performed 51 challenge tests by
358 using the same 5 *L. innocua* strains, avoiding different strains behavior.

359 The highly significant equations of the model allow us to predict the Log CFU/g reduction of *L.*
360 *monocytogenes* during the processing of Italian Salami based on the process parameters; the model

361 enables us to assess the further reduction of a commercial HPP process depending on the intrinsic
362 characteristics of the product. By evaluating raw materials to determine the initial level of
363 contamination, processors and Authorities may assess the ability of the process to control incoming
364 pathogens, to predict the *L. monocytogenes* load at the end of the process, and to evaluate the need
365 for a process modification or for the addition of a final lethal process.

366

367 5. References

- 368 Barmpalia-Davis, I. M., Geornaras, I., Kendall, P. A., & Sofos, J. N. (2008). Survival of *Listeria*
369 *monocytogenes* in a simulated dynamic gastrointestinal model during storage of inoculated
370 bologna and salami slices in vacuum packages. *Journal of Food Protection*, *71*(10), 2014–
371 2023. <https://doi.org/10.4315/0362-028X-71.10.2014>
- 372 Bohaychuk, V. M., Gensler, G. E., King, R. K., Manninen, K. I., Sorensen, O., Wu, J. T., Stiles, M.
373 E., & McMullen, L. M. (2006). Occurrence of pathogens in raw and ready-to-eat meat and
374 poultry products collected from the retail marketplace in Edmonton, Alberta, Canada. *Journal*
375 *of Food Protection*, *69*(9), 2176–2182. <https://doi.org/10.4315/0362-028X-69.9.2176>
- 376 Bonilauri, P., Grisenti, M.S., Daminelli, P., Merialdi, G., Ramini, M., Bardasi, L., Taddei, R.,
377 Cosciani-Cunico, E., Dalzini, E., Frustoli, M.A., Giacometti, F., Piva, S., & Serraino, A.
378 (2019). Reduction of *Salmonella* spp. populations in Italian salami during production process
379 and high pressure processing treatment: Validation of processes to export to the U.S. *Meat*
380 *Science*, *157*:107869. <https://doi.org/10.1016/j.meatsci.2019.06.005>
- 381 Bover-Cid, S., Belletti, N., Aymerich, T., & Garriga, M. (2015). Modeling the protective effect of a
382 w and fat content on the high pressure resistance of *Listeria monocytogenes* in dry-cured ham.
383 *Food Research International*, *75*, 194–199. <https://doi.org/10.1016/j.foodres.2015.05.052>
- 384 Campanini, M., Pedrazzoni, I., Barbuti, S., & Baldini, P. (1993). Behaviour of *Listeria*

385 monocytogenes during the maturation of naturally and artificially contaminated salami: effect
386 of lactic-acid bacteria starter cultures. *International Journal of Food Microbiology*, 20(3),
387 169–175. [https://doi.org/10.1016/0168-1605\(93\)90109-T](https://doi.org/10.1016/0168-1605(93)90109-T)

388 Chikthimmah, N., Guyer, R. B., & Knabel, S. J. (2001). Validation of a 5-log₁₀ reduction of
389 *Listeria monocytogenes* following simulated commercial processing of Lebanon bologna in a
390 model system. *Journal of Food Protection*, 64(6), 873–876. [https://doi.org/10.4315/0362-](https://doi.org/10.4315/0362-028X-64.6.873)
391 028X-64.6.873

392 De Cesare, A., Mioni, R., & Manfreda, G. (2007). Prevalence of *Listeria monocytogenes* in fresh
393 and fermented Italian sausages and ribotyping of contaminating strains. *International Journal*
394 *of Food Microbiology*, 120(1–2), 124–130. <https://doi.org/10.1016/j.ijfoodmicro.2007.06.009>

395 Degenhardt, R., & Sant’Anna, E. S. (2007). Survival of *Listeria monocytogenes* in low acid Italian
396 sausage produced under Brazilian conditions. *Brazilian Journal of Microbiology*, 38(2), 309–
397 314. <https://doi.org/10.1590/S1517-83822007000200024>

398 Dogan, C., & Erkmen, O. (2004). High pressure inactivation kinetics of *Listeria monocytogenes*
399 inactivation in broth, milk, and peach and orange juices. *Journal of Food Engineering*, 62(1),
400 47–52. [https://doi.org/10.1016/S0260-8774\(03\)00170-5](https://doi.org/10.1016/S0260-8774(03)00170-5)

401 Doménech, E., Jimenez -Belenguer, A., Amoros, J. A., Ferrus, M. A., & Escriche, I. (2015).
402 Prevalence and antimicrobial resistance of *Listeria monocytogenes* and *Salmonella* strains
403 isolated in ready-to-eat foods in Eastern Spain. *Food Control*, 47, 120–125.
404 <https://doi.org/10.1016/j.foodcont.2014.06.043>

405 European Commission. (2005). Commission Regulation (EC) No 2073/2005 of 15 November 2005
406 on microbiological criteria for foodstuffs. *Off J Eur Union*, L338, 1–26.

407 European Union Reference Laboratory for *Listeria monocytogenes*. (2014). Technical guidance
408 document for conducting shelf-life studies on *Listeria monocytogenes* in ready-to-eat foods.

409 *EURL Lm-ANSES*, 1–47.

410 Farber, J. M., & Peterkin, P. I. (1991). *Listeria monocytogenes*, a food-borne pathogen. In
411 *Microbiological Reviews* (Vol. 55, Issue 3, pp. 476–511).
412 <https://doi.org/10.1128/membr.55.3.476-511.1991>

413 Ferreira, V., Barbosa, J., Silva, J., Felício, M. T., Mena, C., Hogg, T., Gibbs, P., & Teixeira, P.
414 (2007). Characterisation of alheiras, traditional sausages produced in the North of Portugal,
415 with respect to their microbiological safety. *Food Control*, 18(5), 436–440.
416 <https://doi.org/10.1016/j.foodcont.2005.11.011>

417 Food Safety and Inspection Service/ United States Department of Agricultural (USDA/FSIS).
418 (2017). *FSIS Salmonella Compliance Guidelines for Small and Very Small Meat and Poultry*
419 *Establishments that Produce Ready-to-Eat (RTE) Products and Revised Appendix A*.

420 Gao, Y. L., Ju, X. R., & Wu-Ding. (2007). A predictive model for the influence of food components
421 on survival of *Listeria monocytogenes* LM 54004 under high hydrostatic pressure and mild
422 heat conditions. *International Journal of Food Microbiology*, 117(3), 287–294.
423 <https://doi.org/10.1016/j.ijfoodmicro.2007.04.008>

424 Garriga, M., Marcos, B., Martín, B., Veciana-Nogués, M. T., Bover-Cid, S., Hugas, M., &
425 Aymerich, T. (2005). Starter cultures and high-pressure processing to improve the hygiene and
426 safety of slightly fermented sausages. *Journal of Food Protection*, 68(11), 2341–2348.
427 <https://doi.org/10.4315/0362-028X-68.11.2341>

428 Glass, K. A., & Doyle, M. P. (1989). Fate of *Listeria monocytogenes* in processed meat products
429 during refrigerated storage. *Applied and Environmental Microbiology*, 55(6), 1565–1569.
430 <https://doi.org/10.1128/aem.55.6.1565-1569.1989>

431 Gonzales-Barron, U., Cadavez, V., Pereira, A. P., Gomes, A., Araújo, J. P., Saavedra, M. J.,
432 Estevinho, L., Butler, F., Pires, P., & Dias, T. (2015). Relating physicochemical and

433 microbiological safety indicators during processing of linguiça , a Portuguese traditional dry-
434 fermented sausage. *Food Research International*, 78, 50–61.
435 <https://doi.org/10.1016/j.foodres.2015.11.007>

436 Gounadaki, A. S., Skandamis, P. N., Drosinos, E. H., & Nychas, G. J. E. (2008). Microbial ecology
437 of food contact surfaces and products of small-scale facilities producing traditional sausages.
438 *Food Microbiology*, 25(2), 313–323. <https://doi.org/10.1016/j.fm.2007.10.001>

439 Hayman, M. M., Baxter, I., O’Riordan, P. J., & Stewart, C. M. (2004). Effects of high-pressure
440 processing on the safety, quality, and shelf life of ready-to-eat meats. *Journal of Food*
441 *Protection*, 67(8), 1709–1718. <https://doi.org/10.4315/0362-028X-67.8.1709>

442 Hereu, A., Bover-Cid, S., Garriga, M., & Aymerich, T. (2012). High hydrostatic pressure and
443 biopreservation of dry-cured ham to meet the Food Safety Objectives for *Listeria*
444 *monocytogenes*. *International Journal of Food Microbiology*, 154(3), 107–112.
445 <https://doi.org/10.1016/j.ijfoodmicro.2011.02.027>

446 Holck, A. L., Axelsson, L., Rode, T. M., Høy, M., Måge, I., Alvseike, O., L’Abée-Lund, T. M.,
447 Omer, M. K., Granum, P. E., & Heir, E. (2011). Reduction of verotoxigenic *Escherichia coli* in
448 production of fermented sausages. In *Meat Science* (Vol. 89, Issue 3, pp. 286–295). Elsevier.
449 <https://doi.org/10.1016/j.meatsci.2011.04.031>

450 Hu, M., & Gurtler, J. B. (2017). Selection of surrogate bacteria for use in food safety challenge
451 studies: A review. In *Journal of Food Protection* (Vol. 80, Issue 9, pp. 1506–1536).
452 International Association for Food Protection. <https://doi.org/10.4315/0362-028X.JFP-16-536>

453 Hwang, C. A., Porto-Fett, A. C. S., Juneja, V. K., Ingham, S. C., Ingham, B. H., & Luchansky, J. B.
454 (2009). Modeling the survival of *Escherichia coli* O157:H7, *Listeria monocytogenes*, and
455 *Salmonella Typhimurium* during fermentation, drying, and storage of soudjouk-style
456 fermented sausage. *International Journal of Food Microbiology*, 129(3), 244–252.

457 <https://doi.org/10.1016/j.ijfoodmicro.2008.12.003>

458 International Organization for Standardization (ISO). (2017a). *Microbiology of the food chain.*
459 *Horizontal method for the detection and enumeration of Listeria monocytogenes and other*
460 *Listeria spp.-Part 2: enumeration method.* (ISO standard 11290-2:2017)

461 International Organization for Standardization (ISO). (2017b). *Microbiology of the food chain.*
462 *Horizontal method for the detection and enumeration of Listeria monocytogenes and other*
463 *Listeria spp.-Part 1: detection method.* (ISO standard 11290-1:2017)

464 Jofré, A., Garriga, M., & Aymerich, T. (2008). Inhibition of Salmonella sp. Listeria monocytogenes
465 and Staphylococcus aureus in cooked ham by combining antimicrobials, high hydrostatic
466 pressure and refrigeration. *Meat Science*, 78(1–2), 53–59.
467 <https://doi.org/10.1016/j.meatsci.2007.06.015>

468 Johnson, J. L., Doyle, M. P., Cassens, R. G., & Schoeni, J. L. (1988). Fate of Listeria
469 monocytogenes in tissues of experimentally infected cattle and in hard salami. *Applied and*
470 *Environmental Microbiology*, 54(2), 497–501. <http://www.ncbi.nlm.nih.gov/pubmed/3128165>

471 Lebow, N. K., DesRocher, L. D., Younce, F. L., Zhu, M.-J., Ross, C. F., & Smith, D. M. (2017).
472 Influence of High-Pressure Processing at Low Temperature and Nisin on *Listeria innocua*
473 Survival and Sensory Preference of Dry-Cured Cold-Smoked Salmon. *Journal of Food*
474 *Science*, 82(12), 2977–2986. <https://doi.org/10.1111/1750-3841.13957>

475 Lin, C. M., Takeuchi, K., Zhang, L., Dohm, C. B., Meyer, J. D., Hall, P. A., & Doyle, M. P. (2006).
476 Cross-contamination between processing equipment and deli meats by Listeria
477 monocytogenes. *Journal of Food Protection*, 69(1), 71–79. [https://doi.org/10.4315/0362-](https://doi.org/10.4315/0362-028X-69.1.71)
478 [028X-69.1.71](https://doi.org/10.4315/0362-028X-69.1.71)

479 Lindqvist, R., & Lindblad, M. (2009). Inactivation of Escherichia coli, Listeria monocytogenes and
480 Yersinia enterocolitica in fermented sausages during maturation/storage. *International Journal*

481 *of Food Microbiology*, 129(1), 59–67. <https://doi.org/10.1016/j.ijfoodmicro.2008.11.011>

482 Marcos, B., Aymerich, T., Monfort, J. M., & Garriga, M. (2008). High-pressure processing and
483 antimicrobial biodegradable packaging to control *Listeria monocytogenes* during storage of
484 cooked ham. *Food Microbiology*, 25(1), 177–182. <https://doi.org/10.1016/j.fm.2007.05.002>

485 Martin, B., Garriga, M., & Aymerich, T. (2011). Prevalence of salmonella spp. and listeria
486 monocytogenes at small-scale spanish factories producing traditional fermented sausages.
487 *Journal of Food Protection*, 74(5), 812–815. <https://doi.org/10.4315/0362-028X.JFP-10-437>

488 Mataragas, M., Bellio, A., Rovetto, F., Astegiano, S., Decastelli, L., & Cocolin, L. (2015). Risk-
489 based control of food-borne pathogens *Listeria monocytogenes* and *Salmonella enterica* in the
490 Italian fermented sausages Cacciatore and Felino. *Meat Science*, 103, 39–45.
491 <https://doi.org/10.1016/j.meatsci.2015.01.002>

492 Meloni, D. (2015). Presence of *Listeria monocytogenes* in Mediterranean-Style Dry Fermented
493 Sausages. *Foods*, 4(4), 34–50. <https://doi.org/10.3390/foods4010034>

494 Meloni, D., Consolati, S. G., Mazza, R., Mureddu, A., Fois, F., Piras, F., & Mazzette, R. (2014).
495 Presence and molecular characterization of the major serovars of *Listeria monocytogenes* in
496 ten Sardinian fermented sausage processing plants. *Meat Science*, 97(4), 443–450.
497 <https://doi.org/10.1016/j.meatsci.2014.02.012>

498 Meloni, D., Piras, F., Mureddu, A., Mazza, R., Nucera, D., & Mazzette, R. (2012). Sources of
499 *Listeria monocytogenes* contamination in traditional fermented sausage processing plants in
500 Italy. *Italian Journal of Food Science*, 24(3), 214–222.

501 Merialdi, G., Ramini, M., Ravanetti, E., Gherri, G., & Bonilauri, P. (2015). Reduction of *Listeria*
502 *innocua* contamination in vacuum-pack-aged dry-cured italian pork products after high
503 hydrostatic pressure treatment. *Italian Journal of Food Safety*, 4(2), 101–103.
504 <https://doi.org/10.4081/ijfs.2015.4515>

505 Messier, S., Smith, H. J., & Tittiger, F. (1989). Survival of *Salmonella typhimurium* and
506 *Staphylococcus aureus* in Genoa salami of varying salt concentration. *Canadian Journal of*
507 *Veterinary Research*, 53(1), 84.

508 Nightingale, K. K., Thippareddi, H., Phebus, R. K., Marsden, J. L., & Nutsch, A. L. (2006).
509 Validation of a Traditional Italian-Style Salami Manufacturing Process for Control of
510 *Salmonella* and *Listeria monocytogenes* †. In *Journal of Food Protection* (Vol. 69, Issue 4).
511 http://meridian.allenpress.com/jfp/article-pdf/69/4/794/1679233/0362-028x-69_4_794.pdf

512 Nissen, H., & Holck, A. (1998). Survival of *Escherichia coli* O157:H7, *Listeria monocytogenes* and
513 *Salmonella kentucky* in Norwegian fermented, dry sausage. *Food Microbiology*, 15(3), 273–
514 279. <https://doi.org/10.1006/fmic.1997.0165>

515 Novelli, E., Santo, L. D., Balzan, S., Cardazzo, B., Spolaor, D., Lombardi, A., Carraro, L., &
516 Fasolato, L. (2017). Analysis of process factors of dry fermented salami to control *Listeria*
517 *monocytogenes*. *Italian Journal of Food Safety*, 6(1), 28–32.
518 <https://doi.org/10.4081/ijfs.2017.6184>

519 Porto-Fett, A. C. S., Call, J. E., Shoyer, B. E., Hill, D. E., Pshebniski, C., Cocoma, G. J., &
520 Luchansky, J. B. (2010). Evaluation of fermentation, drying, and/or high pressure processing
521 on viability of *Listeria monocytogenes*, *Escherichia coli* O157:H7, *Salmonella* spp., and
522 *Trichinella spiralis* in raw pork and Genoa salami. *International Journal of Food*
523 *Microbiology*, 140(1), 61–75. <https://doi.org/10.1016/j.ijfoodmicro.2010.02.008>

524 Possas, A., Pérez-Rodríguez, F., Valero, A., & García-Gimeno, R. M. (2017). Modelling the
525 inactivation of *Listeria monocytogenes* by high hydrostatic pressure processing in foods: A
526 review. In *Trends in Food Science and Technology* (Vol. 70, pp. 45–55). Elsevier Ltd.
527 <https://doi.org/10.1016/j.tifs.2017.10.006>

528 Prencipe, V. A., Rizzi, V., Acciari, V., Iannetti, L., Giovannini, A., Serraino, A., Calderone, D.,

529 Rossi, A., Morelli, D., Marino, L., Migliorati, G., & Caporale, V. (2012). *Listeria*
530 *monocytogenes* prevalence, contamination levels and strains characterization throughout the
531 Parma ham processing chain. *Food Control*, *25*(1), 150–158.
532 <https://doi.org/10.1016/j.foodcont.2011.10.018>

533 Roccato, A., Uyttendaele, M., Barrucci, F., Cibin, V., Favretti, M., Cereser, A., Cin, M. D.,
534 Pezzuto, A., Piovesana, A., Longo, A., Ramon, E., De Rui, S., & Ricci, A. (2017). Artisanal
535 Italian salami and sopresse: Identification of control strategies to manage microbiological
536 hazards. *Food Microbiology*, *61*, 5–13. <https://doi.org/10.1016/j.fm.2016.07.010>

537 Rubio, B., Possas, A., Rincón, F., García-Gímeno, R. M., & Martínez, B. (2018). Model for *Listeria*
538 *monocytogenes* inactivation by high hydrostatic pressure processing in Spanish chorizo
539 sausage. *Food Microbiology*, *69*, 18–24. <https://doi.org/10.1016/j.fm.2017.07.012>

540 Schwartz, B., Hexter, D., Broome, C. V., Hightower, A. W., Hirschhorn, R. B., Porter, J. D., Hayes,
541 P. S., Bibb, W. F., Lorber, B., & Faris, D. G. (1989). Investigation of an Outbreak of
542 Listeriosis: New Hypotheses for the Etiology of Epidemic *Listeria monocytogenes* Infections.
543 *Journal of Infectious Diseases*, *159*(4), 680–685. <https://doi.org/10.1093/infdis/159.4.680>

544 Simpson, C. A., Geornaras, I., Yoon, Y., Scanga, J. A., Kendall, P. A., & Sofos, J. N. (2008). Effect
545 of inoculum preparation procedure and storage time and temperature on the fate of *Listeria*
546 *monocytogenes* on inoculated salami. *Journal of Food Protection*, *71*(3), 494–501.
547 <https://doi.org/10.4315/0362-028X-71.3.494>

548 Tolvanen, R., Hellström, S., Elsser, D., Morgenstern, H., Björkroth, J., & Korkeala, H. (2008).
549 Survival of *Listeria monocytogenes* strains in a dry sausage model. *Journal of Food*
550 *Protection*, *71*(8), 1550–1555. <https://doi.org/10.4315/0362-028X-71.8.1550>

551 White, H. (1980). A Heteroskedasticity-Consistent Covariance Matrix Estimator and a Direct Test
552 for Heteroskedasticity. *Econometrica*, *48*(4), 817. <https://doi.org/10.2307/1912934>

553 Xu, H., Hyeon-Yong, L., & Ahn, J. (2009). High pressure inactivation kinetics of Salmonella
554 enterica and Listeria monocytogenes in milk, orange juice, and tomato juice. *Food Science and*
555 *Biotechnology*, 18(4), 861–866.

556

Table 1. Process parameters and ingredients of the different types of Italian salami used for challenge tests.

Test	Salami type	Enterprise	Acidification/drying		Seasoning		Characteristics of different salami					
			Time (days)	Temperature °C ¹	Time (days)	Temperature °C ¹	NaCl %	Nitrate	Nitrite	Caliber (mm)	Trimming (mm)	Type of envelope
1	Low fat	A	7	18-22	31	12-15	2.3	150 ppm	50 ppm	80	5.0	synthetic
2	Milano	B	11	7-19	65	7	2.7	125 ppm	67 ppm	80	3.0	synthetic
3	Finocchiona	B	10	15-20	68	10	2.7	125 ppm	67 ppm	120	6.0	synthetic
4	Napoli	B	7	15-20	28	10	3.0	125 ppm	67 ppm	90	12.0	synthetic
5	Felino	C	5	14-22	40	13-14	2.5	70 ppm	120 ppm	60	5.0	synthetic
6	Felino	C	7	14-22	55	12-13	2.5	70 ppm	70 ppm	80	7.0	synthetic
7	Pepperoni	C	5	14-22	30	13-14	2.5	70 ppm	120 ppm	60	8.0	synthetic
8	Strolghino	C	5	14-22	21	13-14	2.5	70 ppm	120 ppm	38	8.0	synthetic
9	Felino large caliber	D	5	15-25	69	10-15	2.3	150ppm	75ppm	80	6.0	synthetic
10	Genoa	E	8	18-27	41	16-18	2.3	50 ppm	150 ppm	60	6.0	synthetic
11	Milano	F	5	16-24	70	9-11	2.5	150 ppm	0 ppm	110	3.5	synthetic
12	Milano	F	5	16-24	90	9-11	2.5	150 ppm	0 ppm	110	3.5	synthetic
13	Cacciatore	G	6	18-26	21	18-19	2.5	150 ppm	0 ppm	38	5.0	synthetic
14	Cacciatore	G	6	18-26	28	18-19	2.6	150 ppm	0 ppm	38	5.0	synthetic
15	Napoli	H	3	19-24	35	14-16	2.6	150 ppm	0 ppm	65	6.5	synthetic
16	Napoli	H	3	19-24	31	14-16	2.3	150 ppm	0 ppm	65	6.5	synthetic
17	Napoli	H	3	19-24	27	14-16	2.3	150 ppm	0 ppm	65	6.5	synthetic
19	Varzi	I	6	15-25	43	12-19	2.3	150 ppm	0 ppm	65	13.0	natural
18	Varzi	I	6	15-25	43	12-19	2.5	150 ppm	0 ppm	65	13.0	natural
20	Milano	J	7	18-22	63	10-14	2.5	150 ppm	0 ppm	100	3.5	synthetic
21	Cacciatore	B	6	12.5-21	28	10-16	2,4	150 ppm	70 ppm	45	8.0	synthetic
22	Pepperoni small caliber	B	6	12.5-21	28	10-16	2.4	150 ppm	70 ppm	45	8.0	synthetic
23	Flattened pepperoni	B	17	9-21	43	8-12.5	2.7	150 ppm	70 ppm	60	13.0	synthetic
24	Flattened pepperoni	B	17	9-21	43	13	2.7	150 ppm	70 ppm	60	13.0	synthetic
25	Flattened pepperoni	B	17	9-21	72	8-12.5	2.7	150 ppm	70 ppm	60	13.0	synthetic
26	Cacciatore	B	8	12.5-21	31	10-16	2.4	150 ppm	70 ppm	45	8.0	synthetic
27	Milano small caliber	B	8	12.5-21	35	10-16	2.4	150 ppm	70 ppm	45	4.5	synthetic
28	Hungarian small caliber	B	6	12.5-21	25	10-16	2.4	150 ppm	70 ppm	45	3.0	synthetic
29	Pepperoni small caliber	B	6	12.5-21	24	10-16	2.4	150 ppm	70 ppm	45	8.0	synthetic
30	Pepperoni small caliber	B	8	12.5-21	29	10-16	2.4	150 ppm	70 ppm	45	8.0	synthetic
31	Felino	G	8	14-24	57	14-15	2.5	0 ppm	0 ppm	44	7.0	synthetic
32	Felino	L	4	15-24	33	10.5-17	2.3	150 ppm	50 ppm	65	7.0	natural
33	Garlic	M	4	16-23	37	13-15	2.4	120 ppm	30 ppm	65	8.0	synthetic
34	Genoa	D	8	14-24	57	14-15	2.5	150 ppm	0 ppm	44	7.0	synthetic

35	Genoa	I	6	17-26	47	17	2.3	150 ppm	50 ppm	95	6.0	synthetic
36	Felino	I	9	6-22	38	12-13	2.4	0 ppm	0 ppm	65	8.0	synthetic
37	Felino	I	5	19-24	20	19	2.5	150 ppm	50 ppm	45	3.5	synthetic
38	Felino	N	5	26-18	42	14-16	2.4	150 ppm	0 ppm	75	7.0	synthetic
39	Felino	O	6	16-23	37	13-15	2.4	150 ppm	30 ppm	43	8.0	synthetic
40	Milano	O	6	8-19	20	12-13	2.6	150 ppm	30 ppm	43	3.0	synthetic
41	Pepperoni	O	6	8-19	26	12-13	2.6	150 ppm	30 ppm	43	5.0	synthetic
42	Pepperoni	O	6	8-19	31	12-13	2.6	150 ppm	30 ppm	50	5.0	synthetic
43	Pepperoni	O	6	8-19	29	12-13	2.6	150 ppm	30 ppm	50	3.0	synthetic
44	Milano	O	7	8-16	42	14-16	2.6	150 ppm	30 ppm	90	6.0	synthetic
45	Milano	O	5	16-24	69	12-14	2.5	150 ppm	30 ppm	100	3.0	synthetic
46	Pepperoni	O	7	8-19	26	12-13	2.6	150 ppm	30 ppm	43	5.0	synthetic
47	Pepperoni	O	5	8-19	24	12-13	2.6	150 ppm	30 ppm	43	5.0	synthetic
48	Pepperoni	O	5	8-19	34	12-13	2.6	150 ppm	30 ppm	50	5.0	synthetic
49	Pepperoni	O	5	8-19	37	12-13	2.6	150 ppm	30 ppm	50	5.0	synthetic
50	Felino	O	5	8-19	24	12-13	2.6	150 ppm	30 ppm	40	5.0	synthetic
51	Felino	O	5	8-19	31	12-13	2.6	150 ppm	30 ppm	50	5.0	synthetic

¹= range of temperature: a variable temperature can be applied during the phase.

Table 2. *Listeria innocua* Log CFU reduction and pH, a_w values during manufacturing and after HPP treatment of Italian salami.

Test	Salami type	Salami before stuffing		Salami after acidification/drying			Salami after seasoning			Process + HPP		
		pH1	a_w 1	pH2	a_w 2	$-\Delta_a$	pH3	a_w 3	Δ_s	Δp	Δ HPP treatment	Total reduction
1	Low fat	6.00	0.969	5.12	0.971	0.25	5.11	0.951	0.31	0.56	3.01	3.57
2	Milano	5.75	0.970	5.24	0.966	0.49	5.47	0.930	0.47	0.96	3.00	3.96
3	Finocchiona	5.91	0.974	5.41	0.960	0.61	5.20	0.940	1.32	1.93	1.79	3.72
4	Napoli	5.79	0.968	5.25	0.957	0.78	5.38	0.933	0.43	1.21	2.69	3.90
5	Felino	5.57	0.970	5.27	0.962	0.30	5.74	0.920	0.80	1.10	3.46	4.56
6	Felino	6.11	0.965	5.29	0.963	0.84	5.42	0.916	-0.02	0.82	2.50	3.32
7	Pepperoni	5.65	0.966	5.28	0.971	0.37	5.56	0.880	1.54	1.91	1.43	3.34
8	Strolghino	5.87	0.970	5.10	0.963	0.77	5.83	0.922	0.10	0.87	2.55	3.42
9	Felino large caliber	5.99	0.973	5.06	0.970	1.51	5.62	0.910	1.53	3.04	1.17	4.21
10	Genoa	5.89	0.973	4.98	0.967	1.65	6.11	0.917	1.70	3.35	1.31	4.66
11	Milano	6.02	0.966	5.34	0.953	0.41	5.45	0.922	0.51	0.92	3.03	3.95
12	Milano	6.02	0.966	5.34	0.953	0.41	5.47	0.898	1.09	1.50	2.29	3.79
13	Cacciatore	5.73	0.971	5.38	0.948	0.64	5.40	0.910	0.22	0.86	2.65	3.51
14	Cacciatore	5.73	0.971	5.38	0.948	0.64	5.52	0.896	0.36	1.00	1.60	2.60
15	Napoli	6.10	0.974	5.11	0.947	1.33	5.58	0.898	0.73	2.06	1.81	3.87
16	Napoli	6.10	0.974	5.11	0.947	1.33	5.63	0.923	0.27	1.60	1.57	3.17
17	Napoli	6.10	0.974	5.11	0.947	1.33	5.44	0.932	0.09	1.42	3.47	4.89
18	Varzi	5.80	0.972	5.20	0.967	0.06	5.96	0.913	0.53	0.59	1.78	2.37
19	Varzi	5.80	0.972	5.20	0.967	0.06	5.96	0.913	0.53	0.59	2.20	2.79
20	Milano	5.65	0.967	4.78	0.953	1.32	5.16	0.905	0.97	2.29	1.03	3.32
21	Cacciatore	5.93	0.970	5.22	0.950	0.60	5.79	0.908	0.02	0.62	0.49	1.11
22	Pepperoni small caliber	5.89	0.970	4.86	0.950	0.32	5.33	0.910	0.13	0.45	0.59	1.04
23	Flattened pepperoni	5.74	0.967	4.97	0.956	0.58	5.46	0.921	-0.24	0.34	1.03	1.37
24	Flattened pepperoni	5.74	0.967	5.05	0.956	0.58	5.12	0.915	0.23	0.81	0.48	1.29
25	Flattened pepperoni	5.76	0.976	4.87	0.957	0.22	4.99	0.908	0.26	0.48	0.69	1.17
26	Cacciatore	5.69	0.970	5.21	0.960	0.40	5.40	0.917	1.26	1.66	1.13	2.79
27	Milano small caliber	5.71	0.970	5.15	0.960	0.17	4.99	0.917	2.14	2.31	1.35	3.66
28	Hungarian small caliber	5.95	0.970	5.31	0.960	0.51	5.26	0.911	2.22	2.73	0.73	3.46
29	Pepperoni small caliber	5.71	0.970	5.16	0.960	0.11	5.22	0.913	1.25	1.36	1.00	2.36
30	Pepperoni small caliber	5.66	0.977	5.18	0.957	0.02	5.52	0.920	0.71	0.73	1.46	2.19
31	Felino	5.92	0.975	5.09	0.953	1.08	6.17	0.921	1.20	2.28	1.73	4.01
32	Felino	5.76	0.975	5.28	0.969	0.04	5.34	0.920	0.58	0.62	3.20	3.82
33	Garlic	5.91	0.969	5.16	0.950	0.59	5.38	0.919	1.51	2.10	1.11	3.21
34	Genoa	5.90	0.979	5.26	0.959	0.53	6.33	0.920	0.57	1.10	2.52	3.62
35	Genoa	5.77	0.965	4.88	0.960	1.36	5.28	0.912	1.20	2.56	2.54	5.10

36	Felino	5.88	0.963	5.32	0.956	0.05	5.64	0.915	0.48	0.53	2.43	2.96
37	Felino	5.75	0.971	5.10	0.960	1.16	6.53	0.921	0.63	1.79	2.93	4.72
38	Felino	5.89	0.971	5.05	0.968	0.6	6.16	0.921	1.64	2.24	1.96	4.20
39	Felino	5.76	0.971	5.14	0.952	0.92	5.89	0.919	0.65	1.57	2.36	3.93
40	Milano	5.75	0.969	5.06	0.954	0.81	5.85	0.921	0.47	1.28	1.86	3.14
41	Pepperoni	5.82	0.970	4.91	0.944	1.85	5.76	0.918	0.72	2.57	1.06	3.63
42	Pepperoni	5.82	0.970	4.91	0.949	2.2	6.10	0.923	0.21	2.41	1.37	3.78
43	Pepperoni	5.75	0.969	4.97	0.955	1.43	5.75	0.921	-0.42	1.01	2.59	3.60
44	Milano	5.68	0.971	4.98	0.960	0.75	5.12	0.918	0.95	1.70	2.30	4.00
45	Milano	5.88	0.970	5.04	0.966	0.92	6.08	0.921	3.40	4.32	1.36	5.68
46	Pepperoni	5.72	0.971	4.97	0.952	0.68	5.37	0.920	0.49	1.17	0.99	2.16
47	Pepperoni	5.80	0.972	5.14	0.936	1.03	5.21	0.921	0.26	1.29	0.85	2.14
48	Pepperoni	5.80	0.972	5.10	0.941	1.040	5.12	0.906	0.07	1.11	0.62	1.73
49	Pepperoni	5.72	0.971	4.97	0.956	0.580	5.29	0.920	0.84	1.42	0.55	1.97
50	Felino	5.77	0.973	5.13	0.938	0.680	5.16	0.925	0.22	0.90	0.75	1.65
51	Felino	5.77	0.973	4.95	0.945	0.460	5.06	0.903	0.17	0.63	0.81	1.44

Data are expressed as mean of three or five (only for HPP) samples; SD was calculated and considered acceptable when $<0.5 \text{ Log}_{10} \text{ CFU/g}$.

$-\Delta_a$: reduction of *Listeria innocua* after acidification/drying; Δ_s : reduction of *Listeria innocua* at the end of seasoning; Δ_s : reduction of *Listeria innocua* as sum of Δ_a and Δ_s ; Δ HPP: reduction of *Listeria innocua* after HPP; Total Log reduction: reduction of *Listeria innocua* as sum of Δ_s and Δ HPP