



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

ON FINITE-BY-NILPOTENT GROUPS

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Detomi E., Donadze G., Morigi M., Shumyatsky P. (2021). ON FINITE-BY-NILPOTENT GROUPS. GLASGOW MATHEMATICAL JOURNAL, 63(1), 54-58 [10.1017/S0017089519000508].

Availability:

This version is available at: <https://hdl.handle.net/11585/789414> since: 2023-07-11

Published:

DOI: <http://doi.org/10.1017/S0017089519000508>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

DETOMI, E., DONADZE, G., MORIGI, M., & SHUMYATSKY, P. (2021). ON FINITE-BY-NILPOTENT GROUPS. *Glasgow Mathematical Journal*, 63(1), 54-58

The final published version is available online at
<https://dx.doi.org/10.1017/S0017089519000508>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

On finite-by-nilpotent groups

Eloisa Detomi, Guram Donadze, Marta Morigi, and Pavel Shumyatsky

ABSTRACT. Let $\gamma_n = [x_1, \dots, x_n]$ be the n th lower central word. Denote by X_n the set of γ_n -values in a group G and suppose that there is a number m such that $|g^{X_n}| \leq m$ for each $g \in G$. We prove that $\gamma_{n+1}(G)$ has finite (m, n) -bounded order. This generalizes the much celebrated theorem of B. H. Neumann that says that the commutator subgroup of a BFC-group is finite.

1. Introduction

Given a group G and an element $x \in G$, we write x^G for the conjugacy class containing x . Of course, if the number of elements in x^G is finite, we have $|x^G| = [G : C_G(x)]$. A group is said to be a BFC-group if its conjugacy classes are finite and of bounded size. One of the most famous of B. H. Neumann's theorems says that in a BFC-group the commutator subgroup G' is finite [6]. It follows that if $|x^G| \leq m$ for each $x \in G$, then the order of G' is bounded by a number depending only on m . A first explicit bound for the order of G' was found by J. Wiegold [10], and the best known was obtained in [5] (see also [7] and [9]).

The recent articles [3] and [2] deal with groups G in which conjugacy classes containing commutators are bounded. Recall that multilinear commutator words are words which are obtained by nesting commutators, but using always different variables. More formally, the group-word $w(x) = x$ in one variable is a multilinear commutator; if u and v are multilinear commutators involving different variables then the word $w = [u, v]$ is a multilinear commutator, and all multilinear commutators are obtained in this way. Examples of multilinear commutators

2010 *Mathematics Subject Classification.* 20E45; 20F12; 20F24.

Key words and phrases. Conjugacy classes, commutators.

The first and third authors are members of INDAM. The fourth author was supported by CNPq-Brazil.

include the familiar lower central words $\gamma_n(x_1, \dots, x_n) = [x_1, \dots, x_n]$ and derived words δ_n , on 2^n variables, defined recursively by

$$\delta_0 = x_1, \quad \delta_n = [\delta_{n-1}(x_1, \dots, x_{2^{n-1}}), \delta_{n-1}(x_{2^{n-1}+1}, \dots, x_{2^n})].$$

We let $w(G)$ denote the verbal subgroup of G generated by all w -values. Of course, $\gamma_n(G)$ is the n th term of the lower central series of G while $\delta_n(G) = G^{(n)}$ is the n th term of the derived series.

The following theorem was established in [2].

THEOREM 1.1. *Let m be a positive integer and w a multilinear commutator word. Suppose that G is a group in which $|x^G| \leq m$ for any w -value x . Then the order of the commutator subgroup of $w(G)$ is finite and m -bounded.*

Throughout the article we use the expression “ (a, b, \dots) -bounded” to mean that a quantity is finite and bounded by a certain number depending only on the parameters a, b, \dots .

The present article grew out of the observation that a modification of the techniques developed in [3] and [2] can be used to deduce that if $|x^{G'}| \leq m$ for each $x \in G$, then $\gamma_3(G)$ has finite m -bounded order. Naturally, one expects that a similar phenomenon holds for other terms of the lower central series of G . This is indeed the case.

THEOREM 1.2. *Let m, n be positive integers and G a group. If $|x^{\gamma_n(G)}| \leq m$ for any $x \in G$, then $\gamma_{n+1}(G)$ has finite (m, n) -bounded order.*

Using the concept of verbal conjugacy classes, introduced in [4], one can obtain a generalization of Theorem 1.2. Let $X_n = X_n(G)$ denote the set of γ_n -values in a group G . It was shown in [1] that if $|x^{X_n}| \leq m$ for each $x \in G$, then $|x^{\gamma_n(G)}|$ is (m, n) -bounded. Hence, we have

COROLLARY 1.3. *Let m, n be positive integers and G a group. If $|x^{X_n(G)}| \leq m$ for any $x \in G$, then $\gamma_{n+1}(G)$ has finite (m, n) -bounded order.*

Observe that Neumann’s theorem can be obtained from Corollary 1.3 by specializing $n = 1$. Another result which is straightforward from Corollary 1.3 is the following characterization of finite-by-nilpotent groups.

THEOREM 1.4. *A group G is finite-by-nilpotent if and only if there are positive integers m, n such that $|x^{X_n}| \leq m$ for any $x \in G$.*

2. Preliminary results

Recall that in any group G the following “standard commutator identities” hold, when $x, y, z \in G$.

- (1) $[xy, z] = [x, z]^y [y, z]$
- (2) $[x, yz] = [x, z][x, y]^z$
- (3) $[x, y^{-1}, z]^y [y, z^{-1}, x]^z [z, x^{-1}, y]^x = 1$ (Hall-Witt identity);
- (4) $[x, y, z^x][z, x, y^z][y, z, x^y] = 1$.

Note that the fourth identity follows from the third one. Indeed, we have

$$[x^y, y^{-1}, z^y][y^z, z^{-1}, x^z][z^x, x^{-1}, y^x] = 1.$$

Since $[x^y, y^{-1}] = [y, x]$, it follows that

$$[y, x, z^y][z, y, x^z][x, z, y^x] = 1.$$

Recall that X_i denote the set of γ_i -values in a group G .

LEMMA 2.1. *Let k, n be integers with $2 \leq k \leq n$ be integers and let G be a group such that $[\gamma_k(G), \gamma_n(G)]$ is finite and $|x^{\gamma_n(G)}| \leq m$ for any $x \in G$. Then for every $g \in X_n$ we have*

$$|g^{\gamma_{k-1}(G)}| \leq m^{n-k+2} |[\gamma_k(G), \gamma_n(G)]|.$$

PROOF. Let $N = [\gamma_k(G), \gamma_n(G)]$. It is sufficient to prove that in the quotient group G/N , for every integer d with $k-1 \leq d \leq n$

$$|(gN)^{\gamma_d(G/N)}| \leq m^{n-d+1} \quad \text{for every } \gamma_{n-d+1}\text{-value } gN \in G/N,$$

since this implies that $g^{\gamma_d(G)}$ is contained at most m^{n-d+1} cosets of N , whenever $g \in X_{n-d+1}$.

So in what follows we assume that $N = 1$. The proof is by induction on $n-d$. The case $d = n$ is immediate from the hypotheses.

Let $c = n-d+1$. Choose $g \in X_c$ and write $g = [x, y]$ with $x \in X_{c-1}$ and $y \in G$. Let $z \in \gamma_d(G)$. We have

$$[x, y, z^x][z, x, y^z][y, z, x^y] = 1.$$

Note that

$$[z, x] \in [\gamma_d(G), \gamma_{c-1}(G)] \leq \gamma_{d-1+c}(G) = \gamma_n(G)$$

and

$$[y, z] \in \gamma_{d+1}(G) \leq \gamma_k(G),$$

whence $[z, x, y^z] = [z, x, y[y, z]] = [z, x, y]$. Thus,

$$\begin{aligned} 1 = [x, y, z^x][z, x, y^z][y, z, x^y] &= [x, y]^{-1} [x, y]^{z^x} [z, x, y][y, z, x^y] \\ &= [x, y]^{-1} [x, y]^{z^x} (y^{-1})^{[z, x]} y ((x^y)^{-1})^{[y, z]} x^y. \end{aligned}$$

It follows that

$$[x, y]^{z^x} = [x, y](x^{-1})^y(x^y)^{[y, z]}y^{-1}y^{[z, x]}.$$

Since $x^y \in X_{c-1}$ and $[y, z] \in \gamma_{d+1}(G)$, by induction

$$|\{(x^y)^{[y, z]} \mid z \in \gamma_d(G)\}| \leq m^{n-d-1+1}.$$

Moreover, $[z, x] \in \gamma_n(G)$ and so $|\{y^{[z, x]} \mid z \in \gamma_d(G)\}| \leq m$. Thus,

$$|\{[x, y]^{z^x} \mid z \in \gamma_d(G)\}| = |\{[x, y]^z \mid z \in \gamma_d(G)\}| \leq mm^{n-d} = m^{n-d+1}$$

as claimed. \square

Let H be a group generated by a set X such that $X = X^{-1}$. Given an element $g \in H$, we write $l_X(g)$ for the minimal number l with the property that g can be written as a product of l elements of X . Clearly, $l_X(g) = 0$ if and only if $g = 1$. We call $l_X(g)$ the length of g with respect to X . The following result is Lemma 2.1 in [3].

LEMMA 2.2. *Let H be a group generated by a set $X = X^{-1}$ and let K be a subgroup of finite index m in H . Then each coset Kb contains an element g such that $l_X(g) \leq m - 1$.*

In the sequel the above lemma will be used in the situation where $H = \gamma_n(G)$ and $X = X_n$ is the set of γ_n -values in G . Therefore we will write $l(g)$ to denote the smallest number such that the element $g \in \gamma_n(G)$ can be written as a product of as many γ_n -values.

Recall that if G is a group, $a \in G$ and H is a subgroup of G , then $[H, a]$ denotes the subgroup of G generated by all commutators of the form $[h, a]$, where $h \in H$. It is well-known that $[H, a]$ is normalized by a and H .

LEMMA 2.3. *Let $k, m, n \geq 2$ and let G be a group in which $|x^{\gamma_n(G)}| \leq m$ for any $x \in G$. Suppose that $[\gamma_k(G), \gamma_n(G)]$ is finite. Then for every $x \in \gamma_{k-1}(G)$ the order of $[\gamma_n(G), x]$ is bounded in terms of m, n and $[[\gamma_k(G), \gamma_n(G)]]$ only.*

PROOF. By Neumann's theorem $\gamma_n(G)'$ has m -bounded order, so the statement is true for $k \geq n + 1$. Therefore we deal with the case $k \leq n$. Without loss of generality we can assume that $[\gamma_k(G), \gamma_n(G)] = 1$.

Let $x \in \gamma_{k-1}(G)$. Since $|x^{\gamma_n(G)}| \leq m$, the index of $C_{\gamma_n(G)}(x)$ in $\gamma_n(G)$ is at most m and by Lemma 2.2 we can choose elements $y_1, \dots, y_m \in X_n$ such that $l(y_i) \leq m - 1$ and $[\gamma_n(G), x]$ is generated by the commutators $[y_i, x]$. For each $i = 1, \dots, m$ write $y_i = y_{i1} \cdots y_{im-1}$, where $y_{ij} \in X_n$. The standard commutator identities show that $[y_i, x]$ can be written

as a product of conjugates in $\gamma_n(G)$ of the commutators $[y_{ij}, x]$. Since $[y_{ij}, x] \in \gamma_k(G)$, for any $z \in \gamma_n(G)$ we have that

$$[[y_{ij}, x], z] \in [\gamma_k(G), \gamma_n(G)] = 1.$$

Therefore $[y_i, x]$ can be written as a product of the commutators $[y_{ij}, x]$.

Let $T = \langle x, y_{ij} \mid 1 \leq i, j \leq m \rangle$. It is clear that $[\gamma_n(G), x] \leq T'$ and so it is sufficient to show that T' has finite (m, n) -bounded order. Observe that $T \leq \gamma_{k-1}(G)$. By Lemma 2.1, $C_{\gamma_{k-1}(G)}(y_{ij})$ has (m, n) -bounded index in $\gamma_{k-1}(G)$. It follows that $C_T(\{y_{ij} \mid 1 \leq i, j \leq m\})$ has (m, n) -bounded index in T . Moreover, $T \leq \langle x \rangle \gamma_n(G)$ and $|x^{\gamma_n(G)}| \leq m$, whence $|T : C_T(x)| \leq m$. Therefore the centre of T has (m, n) -bounded index in T . Thus, Schur's theorem [8, 10.1.4] tells us that T' has finite (m, n) -bounded order, as required. \square

The next lemma can be seen as a development related to Lemma 2.4 in [3] and Lemma 4.5 in [10]. It plays a central role in our arguments.

LEMMA 2.4. *Let $k, n \geq 2$. Assume that $|x^{\gamma_n(G)}| \leq m$ for any $x \in G$. Suppose that $[\gamma_k(G), \gamma_n(G)]$ is finite. Then the order of $[\gamma_{k-1}(G), \gamma_n(G)]$ is bounded in terms of m, n and $|[\gamma_k(G), \gamma_n(G)]|$ only.*

PROOF. Without loss of generality we can assume that $[\gamma_k(G), \gamma_n(G)] = 1$. Let $W = \gamma_n(G)$. Choose an element $a \in X_{k-1}$ such that the number of conjugates of a in W is maximal possible, that is, $r = |a^W| \geq |g^W|$ for all $g \in X_{k-1}$.

By Lemma 2.2 we can choose $b_1, \dots, b_r \in W$ such that $l(b_i) \leq m - 1$ and $a^W = \{a^{b_i} \mid i = 1, \dots, r\}$. Let $K = \gamma_{k-1}(G)$. Set $M = (C_K(\langle b_1, \dots, b_r \rangle))_K$ (i.e. M is the intersection of all K -conjugates of $C_K(\langle b_1, \dots, b_r \rangle)$). Since $l(b_i) \leq m - 1$ and, by Lemma 2.1, $C_K(x)$ has (m, n) -bounded index in K for each $x \in X_n$, the subgroup $C_K(\langle b_1, \dots, b_r \rangle)$ has (m, n) -bounded index in K , so also M has (m, n) -bounded index in K .

Let $v \in M$. Note that $(va)^{b_i} = va^{b_i}$ for each $i = 1, \dots, r$. Therefore the elements va^{b_i} form the conjugacy class $(va)^W$ because they are all different and their number is the allowed maximum. So, for an arbitrary element $h \in W$ there exists $b \in \{b_1, \dots, b_r\}$ such that $(va)^h = va^b$ and hence $v^h a^h = va^b$. Therefore $[h, v] = v^{-h} v = a^h a^{-b}$ and so $[h, v]^a = a^{-1} a^h a^{-b} a = [a, h][b, a] \in [W, a]$. Thus $[W, v]^a \leq [W, a]$ and so $[W, M] \leq [W, a]$.

Let x_1, \dots, x_s be a set of coset representatives of M in K . As $[W, x_i]$ is normalized by W for each i , it follows that

$$[W, K] \leq [W, x_1] \cdots [W, x_s][W, M] \leq [W, x_1] \cdots [W, x_s][W, a].$$

Since s is (m, n) -bounded and by Lemma 2.3 the orders of all subgroups $[W, x_i]$ and $[W, a]$ are bounded in terms of m and n only, the result follows. \square

PROOF OF THEOREM 1.2. Let G be a group in which $|x^{\gamma_n(G)}| \leq m$ for any $x \in G$. We need to show that $\gamma_{n+1}(G)$ has finite (m, n) -bounded order. We will show that the order of $[\gamma_k(G), \gamma_n(G)]$ is finite and (m, n) -bounded for $k = n, n-1, \dots, 1$. This is sufficient for our purposes since $[\gamma_1(G), \gamma_n(G)] = \gamma_{n+1}(G)$. We argue by backward induction on k . The case $k = n$ is immediate from Neumann's theorem so we assume that $k \leq n-1$ and the order of $[\gamma_{k+1}(G), \gamma_n(G)]$ is finite and (m, n) -bounded. Lemma 2.4 now shows that also the order of $[\gamma_k(G), \gamma_n(G)]$ is finite and (m, n) -bounded, as required. \square

PROOF OF COROLLARY 1.3. Let G be a group in which $|x^{X_n(G)}| \leq m$ for any $x \in G$. We wish to show that $\gamma_{n+1}(G)$ has finite (m, n) -bounded order. Theorem 1.2 of [1] tells us that $|x^{\gamma_n(G)}|$ is (m, n) -bounded. The result is now immediate from Theorem 1.2. \square

PROOF OF THEOREM 1.4. In view of Corollary 1.3 the theorem is self-evident since a group G is finite-by-nilpotent if and only if some term of the lower central series of G is finite. \square

References

- [1] S. Brazil, A. Krasilnikov and P. Shumyatsky, *Groups with bounded verbal conjugacy classes*, J. Group Theory **9** (2006), 127–137.
- [2] E. Detomi, M. Morigi, P. Shumyatsky, *BFC-theorems for higher commutator subgroups*, Q. J. Math., doi: 10.1093/qmath/hay068.
- [3] G. Dierings, P. Shumyatsky, *Groups with Boundedly Finite Conjugacy Classes of Commutators*, Q. J. Math., **69** (2018), 1047–1051, doi: 10.1093/qmath/hay014.
- [4] S. Franciosi, F. de Giovanni and P. Shumyatsky, *On groups with finite verbal conjugacy classes*, Houston J. Math. **28** (2002), 683–689.
- [5] R. M. Guralnick, A. Maroti, *Average dimension of fixed point spaces with applications*, J. Algebra, **226** (2011), 298–308.
- [6] B. H. Neumann, *Groups covered by permutable subsets*. J. London Math. Soc. **29**, (1954). 236–248.
- [7] P. M. Neumann, M.R. Vaughan-Lee, *An essay on BFC groups*, Proc. Lond. Math. Soc. **35** (1977), 213–237.
- [8] D. J. S. Robinson, *A course in the theory of groups*, Second edition. Graduate Texts in Mathematics, **80**. Springer-Verlag, New York, 1996.
- [9] D. Segal, A. Shalev, *On groups with bounded conjugacy classes*, Quart. J. Math. Oxford **50** (1999), 505–516.
- [10] J. Wiegold, *Groups with boundedly finite classes of conjugate elements*, Proc. Roy. Soc. London Ser. A **238** (1957), 389–401.

DIPARTIMENTO DI INGEGNERIA DELL'INFORMAZIONE - DEI, UNIVERSITÀ DI
PADOVA, VIA G. GRADENIGO 6/B, 35121 PADOVA, ITALY

E-mail address: `eloisa.detomi@unipd.it`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRASILIA, BRASILIA-DF,
70910-900 BRAZIL AND INSTITUTE OF CYBERNETICS OF THE GEORGIAN TECH-
NICAL UNIVERSITY, SANDRO EULI STR. 5, 0186, TBILISI, GEORGIA

E-mail address: `gdonad@gmail.com`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, PIAZZA DI PORTA
SAN DONATO 5, 40126 BOLOGNA, ITALY

E-mail address: `marta.morigi@unibo.it`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRASILIA, BRASILIA-DF,
70910-900 BRAZIL

E-mail address: `pavel2040@gmail.com`