



ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

The CUORE Detector and Results

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

The CUORE Detector and Results / Nutini I.; Adams D.Q.; Alduino C.; Alfonso K.; Avignone F.T.; Azzolini O.; Bari G.; Bellini F.; Benato G.; Biassoni M.; Branca A.; Brofferio C.; Bucci C.; Caminata A.; Campani A.; Canonica L.; Cao X.G.; Capelli S.; Cappelli L.; Cardani L.; Carniti P.; Casali N.; Chiesa D.; Chott N.; Clemenza M.; Copello S.; Cosmelli C.; Cremonesi O.; Creswick R.J.; D'Addabbo A.; D'Aguzzo D.; Dafinei I.; Davis C.J.; Dell'Oro S.; Domizio S.D.; Dompe V.; Fang D.Q.; Fantini G.; Faverzani M.; Ferri E.; Ferroni F.; Fiorini E.; Franceschi M.A.; Freedman S.J.; Fujikawa B.K.; Giachero A.; Gironi L.; Giuliani A.; Gorla P.; Gotti C.; Gutierrez T.D.; Han K.; Heeger K.M.; Huang R.G.; Huang H.Z.; Johnston J.; Keppel G.; Kolomensky Y.G.; Ligi C.; Ma Y.G.; Ma L.; Marini L.; Maruyama R.H.; Mei Y.; Moggi N.; Morganti S.; Napolitano T.; Nastasi M.; Nikkel J.; Nones C.; Norman E.B.; Novati V.; Nucciotti A.; O'Donnell T.; Ouellet J.L.; Pagliarone C.E.; Pagnanini L.; Pallavicini M.; Pattavina L.; Pavan M.; Pessina G.; Pettinacci V.; Pira C.; Pirro S.; Pozzi S.; Previtali E.; Puiu A.; Rosenfeld C.; Rusconi C.; Sakai M.; Sangiorgio S.; Schmidt B.; Scielzo N.D.; Singh V.; Sisti M.; Speller D.; Taffarello L.; Terranova F.; Tomei C.; Vignati M.; Wagaarachchi S.L.; Wang B.S.; Welliver B.; Wilson J.; Wilson K.; Winslow L.A.; Zanotti L.; Zimmermann S.; Zucchelli S. - In: JOURNAL OF LOW TEMPERATURE PHYSICS. - ISSN 0022-2291. - STAMPA. - 199:1-2(2020), pp. 519-528. [10.1007/s10909-020-02402-9]

This version is available at: <https://hdl.handle.net/11585/790208> since: 2021-01-21

Published:

DOI: <http://doi.org/10.1007/s10909-020-02402-9>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Nutini, I., Adams, D.Q., Alduino, C. et al. *The CUORE Detector and Results*. J Low Temp Phys 199, 519–528 (2020).

The final published version is available online at: <https://doi.org/10.1007/s10909-020-02402-9>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

The CUORE detector and results

I. Nutini^{1,2}, D. Q. Adams³, C. Alduino³,
 K. Alfonso⁴, F. T. Avignone III³,
 O. Azzolini⁵, G. Bari⁶, F. Bellini^{7,8},
 G. Benato⁹, M. Biassoni², A. Branca^{1,2},
 C. Brofferio^{1,2}, C. Bucci¹⁰, A. Caminata¹¹,
 A. Campani^{12,11}, L. Canonica^{13,10},
 X. G. Cao¹⁴, S. Capelli^{1,2}, L. Cappelli^{10,9,15},
 L. Cardani⁸, P. Carniti^{1,2}, N. Casali⁸,
 D. Chiesa^{1,2}, N. Chott³, M. Clemenza^{1,2},
 S. Copello^{16,10}, C. Cosmelli^{7,8},
 O. Cremonesi², R. J. Creswick³,
 A. D'Addabbo¹⁰, D. D'Aguzzo^{10,17},
 I. Dafinei⁸, C. J. Davis¹⁸, S. Dell'Oro¹⁹,
 S. Di Domizio^{12,11}, V. Dompè^{10,16},
 D. Q. Fang¹⁴, G. Fantini^{10,16},
 M. Faverzani^{1,2}, E. Ferri^{1,2}, F. Ferroni^{16,8},
 E. Fiorini^{2,1}, M. A. Franceschi²⁰,
 S. J. Freedman^{15,9,fn1}, B. K. Fujikawa¹⁵,
 A. Giachero^{1,2}, L. Gironi^{1,2}, A. Giuliani²¹,
 P. Gorla¹⁰, C. Gotti^{1,2}, T. D. Gutierrez²²,
 K. Han²³, K. M. Heeger¹⁸, R. G. Huang⁹,
 H. Z. Huang⁴, J. Johnston¹³, G. Keppel⁵,
 Yu. G. Kolomensky^{9,15}, C. Ligi²⁰,
 Y. G. Ma¹⁴, L. Ma⁴, L. Marini^{9,15},
 R. H. Maruyama¹⁸, Y. Mei¹⁵, N. Moggi^{24,6},
 S. Morganti⁸, T. Napolitano²⁰,
 M. Nastasi^{1,2}, J. Nikkel¹⁸, C. Nones²⁵,
 E. B. Norman^{26,27}, V. Novati²¹,
 A. Nucciotti^{1,2}, T. O'Donnell¹⁹,
 J. L. Ouellet¹³, C. E. Pagliarone^{10,17},
 L. Pagnanini^{1,2}, M. Pallavicini^{12,11},
 L. Pattavina¹⁰, M. Pavan^{1,2}, G. Pessina²,
 V. Pettinacci⁸, C. Pira⁵, S. Pirro¹⁰,
 S. Pozzi^{1,2}, E. Previtali², A. Puiu^{1,2},
 C. Rosenfeld³, C. Rusconi^{3,10},
 M. Sakai⁹, S. Sangiorgio²⁶, B. Schmidt¹⁵,
 N. D. Scielzo²⁶, V. Singh⁹, M. Sisti^{1,2},
 D. Speller¹⁸, L. Taffarello²⁸,
 F. Terranova^{1,2}, C. Tomei⁸, M. Vignati⁸,
 S. L. Wagaarachchi^{9,15}, B. S. Wang^{26,27},
 B. Welliver¹⁵, J. Wilson³, K. Wilson³,
 L. A. Winslow¹³, L. Zanotti^{1,2},
 S. Zimmermann²⁹, S. Zucchelli^{24,6}

the date of receipt and acceptance should be inserted later

Abstract The Cryogenic Underground Observatory for Rare Events (CUORE) is a cryogenic experiment searching for neutrino-less double beta decay ($0\nu\beta\beta$) of ^{130}Te . The detector consists of an array of 988 TeO_2 crystals arranged in a compact cylindrical structure of 19 towers. We report the CUORE initial operations and optimization campaigns. We then present the CUORE results on $0\nu\beta\beta$ and $2\nu\beta\beta$ decay of ^{130}Te obtained from the analysis of the physics data acquired in 2017.

Keywords Neutrinos, $0\nu\beta\beta$, Macro-calorimeters, Cryogenics, CUORE, ...

1 Double beta decay

Double beta decay is a second-order weak decay where a nucleus (A, Z) undergoes two beta decays to its isobar ($A, Z+2$) in a single step, emitting two electrons in the process. According to the Standard Model (SM), the process observes lepton number conservation and two electron anti-neutrinos are simultaneously emitted ($2\nu\beta\beta$). The SM extensions that accommodate for a Majorana neutrino nature predict a lepton-number violating neutrino-less double beta decay channel in which only the two electrons are emitted ($0\nu\beta\beta$). The experimental signature of $0\nu\beta\beta$ decay is a peak in the summed energy spectrum of the final state electrons at the Q-value of the decay ($Q_{\beta\beta}$).

For the $2\nu\beta\beta$ decay, several measurements of the half-lives have been made for few isotopes, giving values of the order of $T_{2\nu\beta\beta}^{1/2} \sim 10^{18}\text{-}10^{24} \text{ yr}^1$. At the same time,

E-mail: irene.nutini@mib.infn.it

¹Dipartimento di Fisica, Università di Milano-Bicocca, Milano I-20126, Italy · ²INFN – Sezione di Milano Bicocca, Milano I-20126, Italy · ³Department of Physics and Astronomy, University of South Carolina, Columbia, SC 29208, USA · ⁴Department of Physics and Astronomy, University of California, Los Angeles, CA 90095, USA · ⁵INFN – Laboratori Nazionali di Legnaro, Legnaro (Padova) I-35020, Italy · ⁶INFN – Sezione di Bologna, Bologna I-40127, Italy · ⁷Dipartimento di Fisica, Sapienza Università di Roma, Roma I-00185, Italy · ⁸INFN – Sezione di Roma, Roma I-00185, Italy · ⁹Department of Physics, University of California, Berkeley, CA 94720, USA · ¹⁰INFN – Laboratori Nazionali del Gran Sasso, Assergi (L'Aquila) I-67100, Italy · ¹¹INFN – Sezione di Genova, Genova I-16146, Italy · ¹²Dipartimento di Fisica, Università di Genova, Genova I-16146, Italy · ¹³Massachusetts Institute of Technology, Cambridge, MA 02139, USA · ¹⁴Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800, China · ¹⁵Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA · ¹⁶INFN – Gran Sasso Science Institute, L'Aquila I-67100, Italy · ¹⁷Dipartimento di Ingegneria Civile e Meccanica, Università degli Studi di Cassino e del Lazio Meridionale, Cassino I-03043, Italy · ¹⁸Wright Laboratory, Department of Physics, Yale University, New Haven, CT 06520, USA · ¹⁹Center for Neutrino Physics, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, USA · ²⁰INFN – Laboratori Nazionali di Frascati, Frascati (Roma) I-00044, Italy · ²¹CSNSM, Univ. Paris-Sud, CNRS/IN2P3, Universitris-Saclay, 91405 Orsay, France · ²²Physics Department, California Polytechnic State University, San Luis Obispo, CA 93407, USA · ²³INPAC and School of Physics and Astronomy, Shanghai Jiao Tong University; Shanghai Laboratory for Particle Physics and Cosmology, Shanghai 200240, China · ²⁴Dipartimento di Fisica e Astronomia, Alma Mater Studiorum – Università di Bologna, Bologna I-40127, Italy · ²⁵Service de Physique des Particules, CEA / Saclay, 91191 Gif-sur-Yvette, France · ²⁶Lawrence Livermore National Laboratory, Livermore, CA 94550, USA · ²⁷Department of Nuclear Engineering, University of California, Berkeley, CA 94720, USA · ²⁸INFN – Sezione di Padova, Padova I-35131, Italy · ²⁹Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

there are several experiments hunting for $0\nu\beta\beta$ process, but no evidence for it has been uncovered yet; thus only lower limits on half-life of $0\nu\beta\beta$ are provided, $T_{0\nu\beta\beta}^{1/2} > 10^{24}\text{-}10^{26}$ yr². In case of a light Majorana neutrino exchange mechanism for the $\beta\beta$ decay, the half-life of the $0\nu\beta\beta$ process can be expressed as a function of the neutrino mass,

$$[T_{0\nu\beta\beta}^{1/2}]^{-1} \propto G_{0\nu} |M_{0\nu\beta\beta}|^2 |m_{\beta\beta}|^2$$

where $m_{\beta\beta}$ is usually referred to as the "effective Majorana mass", $G_{0\nu}$ is the phase-space factor and $|M_{0\nu\beta\beta}|^2$ is the nuclear matrix element³.

2 The CUORE experiment

CUORE (Cryogenic Underground Observatory for Rare Events) is a ton-scale cryogenic experiment. The main goal of the experiment is the investigation of the $0\nu\beta\beta$ decay of ¹³⁰Te^{4,5}. The CUORE detector, shown in Fig.1[left], is composed by 988 TeO₂ crystals arranged in a structure of 19 towers. The total mass of TeO₂ is 742 kg, corresponding to 206 kg of ¹³⁰Te.

2.1 TeO₂ macro-calorimeters

The CUORE choice of using the ¹³⁰Te isotope to search for the $0\nu\beta\beta$ process has been driven by several factors. This isotope has large transition energy, $Q_{\beta\beta} = 2527.518$ keV, and this helps in order to have a reduced background from natural radioactivity in the region of interest (ROI). It has also the highest natural isotopic abundance, $\eta = 34.167\%$, among the $\beta\beta$ emitters. This avoids the need for a costly and time-consuming isotopic enrichment.

The CUORE experiment employs a calorimetric phonon energy detection method, and operates at temperatures of the order of ~ 10 mK. This method is characterized by an extremely good energy resolution, which allows to consider a narrower ROI, and fewer background events will contend with the $0\nu\beta\beta$ signal. However it has to be taken into account the technical difficulty of cooling a large mass detector to such a low temperature.

Cryogenic calorimeters made of materials containing $\beta\beta$ -emitters, were developed by the group of E.Fiorini⁶, that mainly focused on the search for the $0\nu\beta\beta$ of ¹³⁰Te using tellurium dioxide (TeO₂) crystals, which allows to have the ¹³⁰Te source within the detector absorber itself.

The CUORE TeO₂ crystals⁷ are $5 \times 5 \times 5$ cm³ each and weigh an average of 750 g (see Fig.1). They are made with ^{nat}Te; each crystal contains 208 g of ¹³⁰Te. Each TeO₂ crystal is instrumented with a Neutron Transmutation Doped germanium (NTD-Ge) thermistor, which acts as a voltage transducer; the thermal phonons, produced by the energy release in the TeO₂ crystals, are converted into a voltage pulse at the thermistor ends. Each crystal absorber is also instrumented with a silicon Joule heater, which is designed to periodically provide a fixed amount of energy in the detector for gain stabilization purposes. Copper strips connected to the NTD and heaters by gold wires are utilized for the signal readout. A dedicated low-noise circuit with two large load resistors maintained at room temperature provides the bias voltage to each NTD. High precision calibration pulses are sent periodically to the heaters from pulser boards to study possible thermal instabilities of the system⁸.




Fig. 1 [Left] The complete CUORE detector: the 19 towers hosting the 988 TeO₂ crystals. [Right] CUORE instrumented detectors: TeO₂ crystals held in the copper structure, Ge-NTD and Si-heater chips are glued on the crystals.

2.2 Experimental setup

The CUORE experiment is hosted at the Laboratori Nazionali del Gran Sasso of the Istituto Nazionale di Fisica Nucleare (INFN) in central Italy. The Gran Sasso mountain provides the detector with an overburden of 1400 m of rock. The cosmic muon rate is reduced by 6 orders of magnitude relative to the surface; this creates appropriate conditions for rare event physics experiments, such as the search for $0\nu\beta\beta$.

The CUORE TeO₂ crystals have to be operated at a temperature of ~ 10 mK in order to behave as efficient calorimeters. A custom cryostat has been designed to house the ton-scale CUORE detector^{9,10}. Moreover, to avoid radioactive background, only a few construction materials were acceptable and lead shields were integrated in the structure. The cryogenic system implemented in CUORE has to cool down the TeO₂ detectors, the cryostat components and the Pb shields and to maintain the detector volume stably at a temperature around 10 mK for at least 5 years. A cryogen free refrigerator system is utilized for the cooling of the detector; it consists of three subsystem: the Fast Cooling System (FCS), five Pulse Tube cryocoolers (PTs), and a powerful custom ³He/⁴He dilution refrigerator (DU).

3 CUORE initial operations and optimization

The CUORE experiment started taking data in 2017. Detector optimization campaigns were performed in order to characterize and improve the detectors and overall system performance. This required a lot of effort since it was the first time that such a large number of macro-calorimeters were simultaneously operated in a completely new and unique cryogenic system.

3.1 Noise reduction

The spectral shape of the noise in the CUORE detectors is the sum of several contributions, mainly vibration transmission from the cooling system: Pulse Tubes induced vibrations (peaks at 1.4 Hz and its harmonics), residual mechanical vibrations and oscillations related to the detector suspension and support structure.^{11,12}

A system to control the relative phases of the pressure oscillations of the PT cryocoolers was developed. Indeed pressure oscillations induce mechanical vibrations which are transmitted to the cryostat and eventually to the detectors. By driving the relative phases of the PTs it is possible to find the configuration that maximizes the noise cancellation profiting of the interference between the PTs.

3.2 Setting the detector operating conditions

The choice of the correct bias voltage is crucial for the operation of the CUORE NTD sensors. Therefore, special runs were acquired, in which the variation of the NTD resistance, signal amplitude and baseline noise were analyzed as a function of the applied voltage. The characteristic I-V curve (called "load curve") of each detector was measured and the best bias voltage, the so called "working point" (WP), was then identified along the load curve. The optimal WP is usually a compromise between a large signal amplitude and the lowest noise condition, while still ensuring a linear and uniform behavior of the detector for small temperature variations. The load curve measurements and the identification of the optimal working points are, in general, procedures that can be performed manually in cryogenic experiments with a limited number of channels, like it was for CUORE-0. In CUORE, the number of channels to be optimized is almost one thousand, moreover the NTD thermistors have resistances values higher and with a wider spread than small-size experiments. Therefore it was necessary to develop new procedures and algorithms to automatize the load curve measurement and the working point identification for each operating temperature¹³.

The system temperature has to be carefully regulated in order to successfully operate the CUORE detector array. Indeed it is crucial to identify the best operating temperature in order to achieve the highest quality readout with a good signal-to-noise ratio. Moreover it is necessary to maintain this temperature stable over the course of a prolonged run. Temperature scans were performed in 2017 to select the best operational temperature that, together with the noise reduction, contributes to reach the CUORE goal of 5 keV nominal energy resolution.

The working temperature was initially set at 15 mK for the CUORE initial data-taking in 2017. Based on the results of the temperature scans, it appeared that lower temperatures lead to a better energy resolution in the detector response to pulser events and particle signals. For this reason for the data taking from 2018, it was decided to set an operational temperature of 11 mK. Since the NTD working resistances at 11 mK were higher than the nominal values proposed for CUORE, special care was taken in setting NTD operating conditions which could ensure a stable and uniform response in a wide energy range.

4 CUORE physics data taking

The CUORE first science runs were carried out by mid-2017 at 15 mK temperature. After the initial CUORE physics data taking, dedicated runs devoted to further characterization and optimization of the detector were performed in late 2017. In early 2018, the system underwent a period of maintenance, during which the gate valves for the internal detector calibration system were replaced and an external system for calibration was installed. The physics data taking re-started in spring 2018 at 11 mK operating temperature. The data-taking proceeded up to late 2018 when the experiment experienced another warm-up at 100 K, followed by the investigation of the leak in the cryostat causing the issue. A dedicated maintenance of the cryogenic system was performed in early 2019, in order to ensure a stable and long-term operation of the system. The data-taking restarted in March 2019 and it is ongoing. CUORE acquired a total raw exposure (before analysis cuts) of more than 400 kg(TeO₂)·yr up to July 2019.

The CUORE experiment released physics results come from the unblinded data from 2017. The first performed analysis consisted in the search of $0\nu\beta\beta$ decay of TeO₂ and afterwards the analysis was focused on the evaluation of the $2\nu\beta\beta$ half-life and background model reconstruction^{14,18}. The total collected exposure (after the analysis selections) was 86.3 kg(TeO₂)·yr for the 2017 data.

4.1 Search for $0\nu\beta\beta$ of ¹³⁰Te

The CUORE physics energy spectra, in which to look for a possible $0\nu\beta\beta$ signal of ¹³⁰Te, were obtained after applying basic quality cuts, rejection of spurious signals by pulse shape analysis and anti-coincidence selections to the raw data. Science data were blinded until all the selections and fit procedures had been fixed.

The average energy resolution at $Q_{\beta\beta}$ in the science runs was (7.7 ± 0.5) keV FWHM, almost reaching the 5 keV CUORE goal.

The background index in the ROI was obtained via fitting the region with no $0\nu\beta\beta$ component in the model: $b = (0.014 \pm 0.002)$ counts/(keV·kg·yr), which was consistent with the expectations given the CUORE background budget.

The complete model for the ROI including of the $0\nu\beta\beta$ component was utilized for performing a fit in the region around $Q_{\beta\beta}$: there was no evidence for $0\nu\beta\beta$ decay. Including the systematic uncertainties, the 90% lower limit on the half-life was extracted: $T_{1/2}^{0\nu}(^{130}\text{Te}) > 1.3 \times 10^{25}$ yr (90% C.L.).

The CUORE result was combined with the CUORE-0^{15,16} and Cuoricino¹⁷ results: $T_{1/2}^{0\nu}(^{130}\text{Te}) > 1.5 \times 10^{25}$ yr (at 90% C.L.)¹⁴. A limit on the effective Majorana neutrino mass ($m_{\beta\beta}$) in $0\nu\beta\beta$ decay models mediated by a light Majorana neutrino exchange, can be extracted from the half-life limits: $m_{\beta\beta} < 110 - 520$ meV (at 90% C.L.). In Fig.2, the current experimental limits on $m_{\beta\beta}$ are shown. The regions of $m_{\beta\beta}$ allowed by oscillations are shown both for inverted and normal hierarchies of neutrino mass. The horizontal bands with arrows indicate the most stringent upper limits on $m_{\beta\beta}$ coming from the experimental searches of $0\nu\beta\beta$ with several isotopes, including the results for ¹³⁰Te from CUORE combined with CUORE-0 and Cuoricino; moreover the CUORE sensitivity on $m_{\beta\beta}$ for 5 years of data taking is shown.




Fig. 2 Experimental limits on $m_{\beta\beta}$ and current CUORE result and sensitivity¹⁴.

4.2 Measurement of $2\nu\beta\beta$ half-life of ^{130}Te

Given the physics data acquired in 2017, the analysis to reconstruct the observed CUORE background and extract the ^{130}Te $2\nu\beta\beta$ half-life was also performed¹⁸.

The physics data were split into four types of spectra: by-multiplicity and by-layer. Single-detector events fill the multiplicity 1 (M1) spectrum, while double-hit events are assigned with multiplicity 2 (M2). The M1 spectrum is also split in two geometrical layers: the 12 outer towers are more sensitive to contaminations on the shields (outer layer, L1), the 9 inner towers to contaminations on the crystals (inner layer, L0).

A set of 60 sources, considering different isotopes and locations, was sufficient to reconstruct the CUORE observed physics data, via a Markov-Chain Monte-Carlo simultaneous fit across the four energy spectra¹⁹. Many contaminations were identified and located, others were still utilized by the fit but only a limit on their activity was obtained. The model was able to reproduce nearly all the major features of the observed spectra. The $2\nu\beta\beta$ decay is the dominant component of the observed spectrum for single-site events between 1 and 2 MeV (see Fig.3). The $2\nu\beta\beta$ half-life of ^{130}Te was obtained: $T_{1/2}^{2\nu}(^{130}\text{Te}) = [7.9 \pm 0.1 \text{ (stat)} \pm 0.2 \text{ (syst)}] \times 10^{20} \text{ yr}$. The result is consistent with previous measurements of the same half-life^{20,21}. It represents the most precise measurement of ^{130}Te $2\nu\beta\beta$ half-life.

5 Conclusions

CUORE is the first tonne-scale operating cryogenic calorimetric $0\nu\beta\beta$ detector. The first CUORE physics results of $0\nu\beta\beta$ and $2\nu\beta\beta$ in ^{130}Te were obtained from the analysis of physics data collected in 2017 for 86.3 kg(TeO_2)·yr exposure. A total raw exposure of more than 400 kg·yr was already available in July 2019; updated results for 372.5 kg(TeO_2)·yr exposure (after the analysis selections) were released in December 2019²².

The CUORE data taking is currently underway to collect 5 years of run time. While proceeding with the physics data-taking, CUORE keeps studying and testing new




Fig. 3 The CUORE observed spectrum (black) of multiplicity 1 events in the inner layer (M1L0), with a reconstruction of the $2\nu\beta\beta$ component (cyan) of the background as well as the ^{40}K background (yellow).¹⁸

strategies to improve the detector energy resolution and to better understand the sources of background and their effect of the measured energy spectra.

Acknowledgements The author thanks the CUORE collaboration, the directors and staff of the Laboratori Nazionali del Gran Sasso and the technical staff of the laboratories. This work was supported by the Istituto Nazionale di Fisica Nucleare (INFN); the European Research Council; the National Science Foundation (NSF); the US Department of Energy (DOE) Office of Science and by the DOE Office of Nuclear Physics.

References

1. Tanabashi M. et al. (Particle Data Group), *Phys. Rev. D*, **98**, 030001, (2018)
2. Dolinski M. J. et al., *Annu. Rev. Nucl. Part. Sci.*, **69**, 219, (2019)
3. Dell’Oro S. et al., *Adv. High Energy Phys.*, **2016**, 2162659, (2016)
4. Artusa D.R. et al. (CUORE collaboration), *Adv. High Energy Phys.*, **2015**, 879871, (2015)
5. Alduino C. et al. (CUORE collaboration), *Eur. Phys. J. C* **77**, 532, (2017)
6. Fiorini E. and Niinikoski T., *Nucl. Instrum. Meth.* **224**, 83, (1984)
7. Alduino C. et al. (CUORE collaboration), *J. Inst.* **11(07)**, P07009, (2016)
8. Carniti P. et al., *J. Instrum.* **13**, P02029, (2018)
9. D’Addabbo et al., *J. Low Temp. Phys* **193**, 867-875, (2018)
10. Alduino C. et al., *Cryogenics* **102**, 9, (2019)
11. D’Addabbo A. et al., *Cryogenics* **93**, 55-56, (2018)
12. Bersani A. et al., "The detector suspension system of the CUORE experiment", In preparation.
13. Nutini I. et al. "An highly automated system to set working points for large bolometric arrays", In preparation
14. Alduino C. et al. (CUORE collaboration), *Phys. Rev. Lett.* **120**, 132501, (2018)
15. Alfonso K. et al. (CUORE collaboration), *Phys. Rev. Lett.* **115**, 102502, (2015)
16. Alduino C. et al. (CUORE collaboration), *Phys. Rev. C* **93**, 045503, (2016)
17. Andreotti E. et al., *Astropart. Phys.* **34**, 822, (2011)

-
18. Adams D. Q. et al. (CUORE collaboration), "Update on the recent progress of the CUORE experiment", arXiv:1808.10342, (2018)
 19. Alduino C. et al.(CUORE collaboration), *Eur. Phys. J.* **C77**, 8, p. 543, (2017)
 20. Alduino C. et al.(CUORE collaboration), *Eur. Phys. J.* **C77**, 1, p.13, (2017).
 21. Arnold R. et al. (NEMO-3 collaboration), *Phys. Rev. Lett.* **107**, 062504, (2011)
 22. Adams D. Q. et al. (CUORE collaboration), "Improved Limit on Neutrinoless Double-Beta Decay in ^{130}Te with CUORE", arXiv:1912.10966, (2019)

Author accepted manuscript