

### Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

In vivo femorotibial kinematics of medial-stabilized total knee arthroplasty correlates to post-operative clinical outcomes

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Alesi D., Marcheggiani Muccioli G.M., Roberti di Sarsina T., Bontempi M., Pizza N., Zinno R., et al. (2021). In vivo femorotibial kinematics of medial-stabilized total knee arthroplasty correlates to postoperative clinical outcomes. KNEE SURGERY, SPORTS TRAUMATOLOGY, ARTHROSCOPY, 29(2), 491-497 [10.1007/s00167-020-05975-w].

Availability:

This version is available at: https://hdl.handle.net/11585/795865 since: 2024-08-02

Published:

DOI: http://doi.org/10.1007/s00167-020-05975-w

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

## In-vivo femorotibial kinematics of medial-stabilized total knee arthroplasty correlates to postoperative clinical outcomes

Domenico Alesi, M.D.*	domenicoalesi@ymail.com
Giulio Maria Marcheggiani Muccioli, M.D., Ph.D. *#	marcheggianimuccioli@me.com
Marco Bontempi, Ph.D. *	m.bontempi@biomec.it
Nicola Pizza *	nickpiz90@gmail.com
Raffaele Zinno MSc *	raffaele.zinno.ufficio@gmail.com
Stefano Di Paolo Eng *	stefano.dipaolo@ior.it
Stefano Zaffagnini, M.D., Full Professor *#	stefano.zaffagnini@unibo.it
Laura Bragonzoni Ph.D., Senior Assistant Professor #	laura.bragonzoni4@unibo.it

\* 2nd Orthopaedic and Traumatologic Clinic – IRCCS – Istituto Ortopedico Rizzoli – Via G.B. Pupilli 1, 40136,

Bologna, Italy

# University of Bologna - Italy

Corresponding Author: Giulio Maria Marcheggiani Muccioli, M.D., Ph.D.

Orthopaedic Surgeon - IRCCS - Istituto Ortopedico Rizzoli

tel: +39 0516366509

mail: marcheggianimuccioli@me.com

**Funding Statement** 

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

#### **Conflict of interest**

Authors declare that they have no conflict of interest.

#### **Blinded Manuscript**

# In-vivo femorotibial kinematics of medial-stabilized total knee arthroplasty correlates to postoperative clinical outcomes

3

#### 4 ABSTRACT

**Purpose:** To evaluate if there was a correlation between in vivo kinematics of a medial-stabilized (MS) total knee
arthroplasty (TKA) and post-operative clinical scores. We hypothesized that (1) a MS-TKA would produce a medial pivot
movement and that (2) this specific pattern would be correlated with higher clinical scores.

8 Methods: 18 patients were evaluated trough clinical and functional scores evaluation (Knee Society Score clinical and

9 functional, Womac, Oxford), and kinematically through dynamic radiostereometric analysis (RSA) at 9 months after MS-

10 TKA, during the execution of a sit-to-stand and a lunge motor task. The anteroposterior (AP) Low Point translation of

12 scores and kinematics was performed through the Pearson's correlation coefficient r.

Results: A significantly greater (p<0.0001) anterior translation of the lateral compartment with respect to the medial one</li>
was found in both sit-to-stand (medial 2.9 mm ± 0.7 mm, lateral 7.1 mm ± 0.6 mm) and lunge (medial 5.3 mm ± 0.9 mm,
lateral 10.9 mm ± 0.7 mm) motor tasks, thus resulting in a medial pivot pattern in about 70% of patients. Significant
positive correlation in sit-to-stand was found between the peak of AP translation in the lateral compartment and clinical
scores (r=0.59 for Knee Society Score clinical and r=0.61 for Oxford). Moreover, we found that the higher peak of AP

18 translation of the medial compartment correlated with lower clinical scores (r=-0.55 for Knee Society Score clinical, r=-

- 19 0.61 for Womac and r=-0.53 for Oxford) in the lunge. A negative correlation was found between Knee Society Score
- 20 clinical and VV laxity during sit-to-stand (r=-0.56) and peak of external rotation in the lunge motor task (r=-0.66).

21 Conclusions: The MS-TKA investigated produced in-vivo a medial pivot movement in about 70% of patients in both

22 examined motor tasks. There was a correlation between the presence of medial pivot and higher post-operative scores.

- 23 Level of evidence: IV
- 24 Keywords: medial-stabilized, TKA, dynamic RSA, medial pivot
- 25
- 26

#### 27 Introduction

28

29 Nowadays, about 20% of patients who underwent total knee arthroplasty (TKA) is not fully satisfied in terms of pain

- 30 relief and functional outcomes [11]. Despite good coronal and sagittal alignment, the restoration of normal knee
- 31 kinematics, which is supposed to be a crucial factor, is not always reached [2, 9]. Many studies focusing on the motion of
- 32 the native knee joint revealed that the lateral compartment performs a wider anterior-posterior (AP) translation during
- 33 flexion-extension compared to the medial one. This type of motion is known as "medial pivoting" movement of the knee
- **34** [9, 13].
- 35 Several TKA designs have been developed with the aim of reproducing this pattern as much as possible. A new concept
- of TKA, the medial-stabilized (MS) knee prosthesis, has been developed trying to mimic distal femur and tibial plateau
   geometry, thus reproducing the natural knee motion. The MS-TKA guarantees anteroposterior (AP) stability of the medial
- 38 compartment thanks to the "ball-in-socket" geometry, coupling with a greater AP translation of the external compartment
- during knee flexion allowed by the flat lateral surface of the insert [18].
- 40 Multiple studies reported favourable outcomes for the MS-TKA, with satisfactory results at medium to long-term follow-
- 41 up [5, 8, 12, 14]. Nevertheless, to authors' knowledge, there is lack of evidence of better clinical outcomes in presence of
- 42 specific kinematic patterns. In particular, just few investigations have been focused on the correlation between in-vivo
- 43 kinematics of MS prosthesis and post-operative clinical outcomes [19].
- 44 Hence, the aim of the present study was to evaluate if the in vivo kinematics of a MS total knee arthroplasty, designed to
- 45 produce medial pivot pattern, was somehow correlated with post-operative clinical scores. The hypotheses were that (1)
- the MS-TKA would be able to produce medial pivot movement during the examined motor tasks, that (2) this specific
- 47 pattern would be correlated with higher clinical scores.
- 48

#### 49 Materials and Methods

- 50 A cohort of 18 randomly selected not consecutive patients in waiting list for TKA at our institution was enrolled in this 51 study after providing an informed consent. The randomization was carried out using a random number generator, available 52 on the web site of National Health System, which generated 18 casual numbers within the interval (1, 45). These numbers 53 were used to select the patients from the waiting list for TKA. The inclusion criteria were: (1) age (50-85 years old); (2) 54 severe radiographic primary osteoarthritis (Kellgren-Lawrence grade 3 and grade 4); (3) patients scheduled for a primary 55 TKA. The exclusion criteria were: (1) previous corrective osteotomy on the affected lower limb; (2) post-traumatic 56 arthritis; (3) severe preoperative varus-valgus deformity (Hip Knee Ankle angle >  $10^{\circ}$ ); Body Mass Index >  $40 \text{ kg/m}^2$ ; 57 (4) rheumatoid arthritis; (5) chronic inflammatory joint diseases; (6) patients with a pre-pathological abnormal gait 58 (amputated, neuromuscular disorders, poliomyelitis, developmental dysplasia of the hip); (7) severe ankle osteoarthritis 59 (Kellgren-Lawrence grade 3 and grade 4); (8) severe hip osteoarthritis (Kellgren-Lawrence grade 3 and grade 4); (9) 60 previous total hip or ankle replacement; (10) unwillingness to take part in this study and providing Health Insurance
- 61 Portability and Accountability Act (HIPAA) authorisation.
- The mean age of the patients was 72.1 ± 7.4 years old (95% CI). The cohort included 7 right legs and 11 left legs, with 9
  females and 9 males. All patients were operated with the standard technique (medial parapatellar approach, adjusted
  mechanical alignment)[21] receiving a cemented MS-TKA (GMK Sphere, Medacta International, Switzerland) with
- 65 patellar resurfacing. Patients were evaluated after a minimum 9-month follow-up by Model-based dynamic

- 66 radiostereometric analysis (RSA). The RSA set-up was analogous to the ones already published in previous articles from
- 67 the same study group [3, 4, 6, 15].
- 68 All patients performed two motor tasks that represent common daily activities, a sit-to-stand and a lunge movement:
- 69 Sit-to-stand: from the sitting position, the patient had to stand up, without support for the upper limbs. The chair
  70 was made of a radiolucent material and was 40 cm high. The chair rising movement was not standardised to best
  71 simulate the natural movement that the patient would have done in daily life.
- Lunge: from the standing position, the patient made a bending with the operated limb, reaching their maximum
   flexion according to their possibilities, then returned in the standing position. Patients were not allowed to use
   supports for the upper limbs. Also, this motor task was not strictly standardised.
- 75

The global accuracy of dynamic RSA in model positioning and orientation, evaluated in terms of "trueness  $\pm$  precision", resulted to be sub-millimetric, respectively 0.2 mm  $\pm$  0.5 mm and 0.3°  $\pm$  0.2°. Data are presented as mean  $\pm$  standard error.

Moreover, pre-operative and post-operative visit at 9 months, patients were clinically evaluated by one single expert
orthopaedic surgeon which calculated the clinical and functional Knee Society Score (KSSC and KSSF), Womac and
Oxford scores. These scores are commonly used to assess patients' clinical status in terms of pain and functionality. For
the KSSC, KSSF and Womac, the scores range from 0 to 100, and is considered good over 69. The Oxford score ranges

- 83 from 0 to 48 and is considered good over 39.
- This study obtained the approval from Institutional Review Board (IRB) of the IRCCS Rizzoli Orthopaedic Institute (ID:
   003603 February 16<sup>th</sup>, 2016).

#### 86 Statistical Analysis

87 Statistical analysis was performed with MATLAB®. For each of the two movements, the AP translation of medial and

- 88 lateral compartments was compared in order to assess the presence of medial pivot pattern: a "pivoting ratio" was
  89 assessed within (-1, 1) for each task. In all tasks with ratio between 0 and 1, prostheses were considered to perform
- 90 consistent medial pivot.
- 91 The matched pairs Student's t-test was used to assess the statistical differences, with p < 0.05. Furthermore, a ratio
- 92 between medial and lateral AP translation was calculated to evaluate how many knees showed the medial pivot pattern.

A correlation analysis between post-operative clinical scores and kinematic parameters was performed for each of the
 two tasks through the Pearson's correlation coefficient r (p<0.05). A correlation coefficient r<0.4 was considered poor, r</li>
 between 0.4 and 0.75 was considered fair, r >0.75 was considered good.

- 96 In order to calculate the sample size, an a priori power analysis was performed. The two sample T-test was used with p <97 0.05. Based on previously published studies with the same technology, the true difference of means between medial and
- 57 0.05. Dased on previously published studies with the same technology, the true difference of means between mediat and
- 98 lateral compartment was set to 3 mm for both motor tasks [6, 15], while the standard deviation was considered 3 mm.
- **99** The power value was 0.95. Based on this analysis, the number of patients resulted to be 16.
- 100

#### 101 Results

102 All the patients were able to perform the sit-to-stand motor task, while all the patients except one were able to perform 103 the lunge motor task. The average time of follow-up evaluation was  $10.0 \pm 1.8$  months. The sex distribution was not found 104 to statistically influence the kinematic and outcome analyses (p<0.05).

- 105 Regarding the sit-to-stand, the kinematic analysis showed the presence of medial pivot, with a significantly greater
- 106 (p<0.0001) anterior translation of the lateral condyle (7.1 mm  $\pm$  0.6 mm) compared to the medial one (2.9 mm  $\pm$  0.7 mm)
- on average (Fig.1 and Tab.1). A consistent medial pivot pattern was found in 12 patients out of 18 (67%). Thus, the
   kinematic analysis was repeated in the sub-group of patients performing medial pivot and in the sub-group of patients not
- 109 performing it. A significantly greater anterior translation of lateral condyle (p<0.0001) was found in the sub-group of
- 110 patients performing medial pivot, while no statistical difference was found in the sub-group of patients not performing
- medial pivot. Also, regarding the lunge motor task, the lateral condyle showed a significantly larger (p<0.0001) anterior translation (10.9 mm  $\pm$  0.7 mm) with respect to the medial one (5.3 mm  $\pm$  0.9 mm) (Fig.2 and Tab.1) on average. A
- 113 consistent medial pivot pattern was found in 11 patients out of 17 (65%). As for the sit-to-stand, the initial cohort was
- split in two sub-groups, and the same trend was found (Tab.1). The Freeman graphics regarding the kinematical results
- split in two groups of patients, performing a consistent medial pivot or not, are provided in supplementary material A.
- 116 Regarding the post-operative clinical outcomes, significant correlations (p < 0.05) were found between clinical outcomes 117 and knee kinematics (Tab.2). In particular, a positive correlation was found between the peak of AP translation in the 118 lateral compartment and both the KSSC (r = 0.59) and the Oxford (r = 0.61) scores in the sit-to-stand motor task. In 119 addition, we found that the higher the peak of AP translation in the medial compartment the lowest the KSSC score (r = 120 -0.55 lunge), the WOMAC (r = -0.60 lunge), and the Oxford (r = -0.53 lunge). We found also a negative correlation 121 between KSSC score and VV laxity regarding the sit-to-stand (r = -0.56) motor task and between the peak of external 122 rotation and the KSSC (r = -0.66) regarding the lunge motor task. A specific post-hoc power analysis for the clinical 123 outcomes was performed. Using a sample size of 18 patients and the correlation coefficient with the lowest absolute 124 magnitude, i.e. r=0.53, the power resulted 0.81. The complete list of patients' outcomes in relation to the medial pivot 125 pattern is provided in the supplementary material B.
- 126

#### 127 Discussion

The most important findings of the present study were that (1) the MS TKA produced medial pivot movement during both sit-to-stand and lunge motor tasks in the 70% of patients; that (2) higher values of AP translation of the lateral compartment correlated with higher clinical scores, while the higher values of AP translation of the medial compartment correlated with lower clinical scores. Moreover, higher values of VV laxity during sit-to-stand and peak of external rotation during the lunge motor task were correlated with lower clinical scores.

- 133 The medial pivot motion of the healthy knee joint has been widely described in literature [7, 10, 13, 16]. The restoration
- 134 of normal knee biomechanics is a crucial point of TKA surgery, but it can be highly related to implant design and patients
- activity [1]. The MS prosthesis evaluated in this study (GMK Sphere, Medacta International, Switzerland) demonstrated
- a consistent medial pivot movement both in the sit-to-stand and in the lunge motor tasks, two common activities of daily
- 137 life with different levels of engagement. This prosthetic design is intended to reproduce posterior femoral rollback and

- medial pivot motion, avoiding paradoxical anterior sliding or rolling forward of the femoral component with respect tothe tibia during flexion-extension.
- 140 The medial pivot pattern was found in about 70% of patients for both motor tasks, thus confirming the average trend. Our 141 kinematic results are in line with the ones of different authors, reproduced in either an in-vivo or in-vitro setting. 142 Steinbrück A. et al. [18] tested the GMK Sphere MS TKA in an in-vitro setting during loaded squat from 20° to 120° of 143 flexion. They found a combination of anterior translation of the lateral compartment (14.7 mm ± 6 mm) with a lower
- 144 anterior translation on the medial one (3.5 mm  $\pm$  4.8 mm), concluding that the specific TKA was able to mimic the
- physiological knee kinematics. Also Scott G. et al.[17] reported a greater AP translation on the lateral side (8 mm  $\pm$  4
- 146 mm) with respect to the medial one  $(2 \text{ mm} \pm 3 \text{ mm})$  in 16 knees implanted with the same prosthesis during in-vivo lunge
- 147 movement. The range of AP translation of the present study resulted lower than the one presented by Steinbrück A. et al.
- and similar to the one of Scott G. et al. This difference could be explained by the higher range of knee flexion (20-120°)
- reached in-vitro and in the absence of patients' active muscle control [18].
- 150 On average, the cohort of patients reported good clinical and functional outcomes. Regarding the correlation between the 151 clinical outcomes and AP Low Point translations of medial and lateral compartments, it seems that the medial pivot 152 movement has a positive effect on the clinical and functional outcomes of the patients. Analogue results were found in a recent paper by Van Onsem et al. The authors reported that patients with poor post-operative patient-reported outcomes 153 154 (PROMs), after a Posterior-Stabilized (PS) TKA implantation experienced more anterior translation on the medial side 155 and less posterior translation on the lateral side during deep flexion activities. Meanwhile the patients who reported good 156 clinical outcomes showed a greater AP stability on the medial compartment with respect to the lateral one [21]. It must 157 be outlined that the TKA model examined by Van Onsem is substantially different from the one of the present study, 158 since AP stability of PS-TKA relies more on the post-cam mechanism. However, the concept of higher congruency of the 159 medial compartment with respect to the lateral one seems to be equally important to ensure AP stability of the implant.
- 160 Furthermore, the negative correlation between VV laxity and peak of external rotation with clinical outcomes suggests 161 that excessive coronal and rotational laxity could play a role in knee function after TKA. To our knowledge, no article 162 has been published about correlation between dynamic rotational or coronal laxity with clinical outcomes. In addition, 163 this correlation seems to highlight the importance of soft tissue balancing during surgery. Medial compartment stability 164 and lateral compartment wider AP translation allow proper implant function. On the medial side, the stability is provided 165 by the high congruency of the inlay with the medial condyle and by the integrity of medial soft tissues (superficial medial 166 collateral ligament). On the lateral side, the flat surface of the inlay facilitates the rollback, thus, lateral soft tissue release 167 is more likely to be done. However, to avoid excessive lateral compartment laxity, the anterior fibers of the ileo-tibial 168 band should remain intact, as suggested also by Van Overschelde et al. [20]. However, the ranges of VV laxity found for 169 the patients of our study were always below  $5^{\circ}$ , thus suggesting that intraoperative stability was achieved properly. As 170 recently confirmed by Watanabe et al., valgus laxity averaging 3.6° correlates with good PROMs after PS TKA, while 171 increasing joint laxity reversely affected some of the clinical outcomes [22].
- 172 The present study has some limitations. First of all, the RSA evaluation was performed only after surgery, as specifically 173 required by our IRB in order to reduce x-rays exposure. Thus, it was not possible to compare joint kinematics before and 174 after implantation. Anyway, joint kinematics of an arthritic knee cannot be assumed as the one of the native joint. Second, 175 a strict standardization of patients' movement in terms of movement magnitude and velocity was not performed. Patients 176 were required to perform the motor tasks in the most natural way as possible, in order to mimic daily life conditions.

- 177 However, to generate comparable data, all the movements were normalized in the post processing (see Methods for
- details). Furthermore, the correlation analysis between the clinical outcomes and the kinematical data did not allow us to
- define a cut-off for laxity, over which the clinical outcomes becomes poor. This aspect is more likely to be investigated
- 180 in a wider cohort of patients. Finally, all the pre-operative and post-operative clinical evaluations was carried out by one
- 181 single expert surgeon (who did not participate to surgeries nor to dynamic RSA analysis) to avoid inter-rater variability.
- 182 The results of this study have an important clinical relevance. In fact, the knowledge that a medial pivot kinematics after
- 183 the implantation of a MS-TKA leads to better clinical results will push research in obtaining it in a higher ratio of patients.
- 184

#### 185 Conclusions

- 186 The MS-TKA investigated produced in-vivo a medial pivot movement in about 70% of patients in both examined motor
- tasks. There was a correlation between the presence of medial pivot and higher post-operative scores.

188

189

#### 190 References

Banks SA, Hodge WA (2004) 2003 Hap Paul Award Paper of the International Society for Technology in
 Arthroplasty. Design and activity dependence of kinematics in fixed and mobile-bearing knee arthroplasties. J
 Arthroplasty 19:809–816

Behrend H, Giesinger K, Giesinger JM, Kuster MS (2012) The "forgotten joint" as the ultimate goal in joint
 arthroplasty: validation of a new patient-reported outcome measure. J Arthroplasty 27:430-436.e1

Bontempi M, Roberti di Sarsina T, Marcheggiani Muccioli GM, Pizza N, Cardinale U, Bragonzoni L, Zaffagnini
 S (2019) J-curve design total knee arthroplasty: the posterior stabilized shows wider medial pivot compared to the cruciate
 retaining during chair raising. Knee Surg Sports Traumatol Arthrosc doi: 10.1007/s00167-019-05645-6.

Bragonzoni L, Marcheggiani Muccioli GM, Bontempi M, Roberti di Sarsina T, Cardinale U, Alesi D, Iacono F,
 Neri MP, Zaffagnini S (2018) New design total knee arthroplasty shows medial pivoting movement under weight-bearing
 conditions. Knee Surg Sports Traumatol Arthrosc 27(4):1049-1056

Brinkman J-M, Bubra PS, Walker P, Walsh WR, Bruce WJM (2014) Midterm results using a medial pivot total
 knee replacement compared with the Australian National Joint Replacement Registry data. ANZ J Surg 84:172–176

Cardinale U, Bragonzoni L, Bontempi M, Alesi D, Roberti di Sarsina T, Lo Presti M, Zaffagnini S, Marcheggiani
 Muccioli GM, Iacono F (2019) Knee kinematics after cruciate retaining highly congruent mobile bearing total knee
 arthroplasty: An in vivo dynamic RSA study. Knee pii: S0968-0160(19)30258-3. doi: 10.1016/j.knee.2019.11.003

207 7. Dennis DA, Mahfouz MR, Komistek RD, Hoff W (2005) In vivo determination of normal and anterior cruciate
208 ligament-deficient knee kinematics. J Biomech 38:241–253

8. Fan C-Y, Hsieh JT-S, Hsieh M-S, Shih Y-C, Lee C-H (2010) Primitive results after medial-pivot knee
arthroplasties: a minimum 5-year follow-up study. J Arthroplasty 25:492–496

9. Freeman M a. R, Pinskerova V (2005) The movement of the normal tibio-femoral joint. J Biomech 38:197–208

Freeman M a. R, Pinskerova V (2003) The movement of the knee studied by magnetic resonance imaging. Clin
 Orthop Relat Res (410):35–43

Gunaratne R, Pratt DN, Banda J, Fick DP, Khan RJK, Robertson BW (2017) Patient Dissatisfaction Following
 Total Knee Arthroplasty: A Systematic Review of the Literature. J Arthroplasty 32:3854–3860

Hossain F, Patel S, Rhee S-J, Haddad FS (2011) Knee arthroplasty with a medially conforming ball-and-socket
 tibiofemoral articulation provides better function. Clin Orthop Relat Res 469:55–63

218 13. Johal P, Williams A, Wragg P, Hunt D, Gedroyc W (2005) Tibio-femoral movement in the living knee. A study
219 of weight bearing and non-weight bearing knee kinematics using 'interventional' MRI. Journal of Biomechanics 38:269–
220 276

14. Karachalios T, Roidis N, Giotikas D, Bargiotas K, Varitimidis S, Malizos KN (2009) A mid-term clinical
 outcome study of the Advance Medial Pivot knee arthroplasty. Knee 16:484–488

Marcheggiani Muccioli GM, Pizza N, Di Paolo S, Zinno R, Alesi D, Roberti Di Sarsina T, Bontempi M,
 Zaffagnini S, Bragonzoni L (2020) Multi-radius posterior-stabilized mobile-bearing total knee arthroplasty partially
 produces in-vivo medial pivot during activity of daily living and high demanding motor task. Knee Surg Sports Traumatol
 Arthrosc doi: 10.1007/s00167-020-05846-4

16. Nakagawa S, Kadoya Y, Todo S, Kobayashi A, Sakamoto H, Freeman MA, Yamano Y (2000) Tibiofemoral
 movement 3: full flexion in the living knee studied by MRI. J Bone Joint Surg Br 82:1199–1200

17. Scott G, Imam MA, Eifert A, Freeman M a. R, Pinskerova V, Field RE, Skinner J, Banks SA (2016) Can a total
knee arthroplasty be both rotationally unconstrained and anteroposteriorly stabilised? A pulsed fluoroscopic investigation.
Bone Joint Res 5:80–86

18. Steinbrück A, Schröder C, Woiczinski M, Fottner A, Pinskerova V, Müller PE, Jansson V (2016) Femorotibial
kinematics and load patterns after total knee arthroplasty: An in vitro comparison of posterior-stabilized versus medialstabilized design. Clin Biomech (Bristol, Avon) 33:42–48

Van Onsem S, Verstraete M, Van Eenoo W, Van Der Straeten C, Victor J (2019) Are TKA Kinematics During
 Closed Kinetic Chain Exercises Associated with Patient-reported Outcomes? A Preliminary Analysis. Clin Orthop Relat
 Res 478(2):255-263

238 20. Van Overschelde P, Pinskerova V, Koch PP, Fornasieri C, Fucentese S (2017) Redefining Knee Balance in a
239 Medially Stabilized Prosthesis: An In-Vitro Study. Open Orthop J 11:1165–1172

240 21. Vanlommel L, Vanlommel J, Claes S, Bellemans J (2013) Slight undercorrection following total knee
241 arthroplasty results in superior clinical outcomes in varus knees. Knee Surg Sports Traumatol Arthrosc 21:2325–2330

242 22. Watanabe T, Koga H, Katagiri H, Otabe K, Nakagawa Y, Muneta T, Sekiya I, Jinno T (2019) Coronal and
243 sagittal laxity affects clinical outcomes in posterior-stabilized total knee arthroplasty: assessment of well-functioning
244 knees. Knee Surg Sports Traumatol Arthrosc doi: 10.1007/s00167-019-05500-8

## Figure captions246

Fig.1 Average of the AP translation of the Low Point of medial and lateral compartment during sit-to-stand over the entire
 cohort of patients, represented as Freeman graphic. On the right are represented the knee flexion angles.

249 Fig.2 Average of the AP translation of the Low Point of medial and lateral compartment during lunge over the entire

250 cohort of patients, represented as Freeman graphic. On the right are represented the knee flexion angles.





LOW POINT	LOW POINT KINEMATICS		Lateral Condyle AP translation (mm)	p-value	
	Entire cohort (n=18)	$2.9\pm0.7$	$7.1 \pm 0.6$	< 0.0001 *	
Sit to stand	Medial pivot pt (n=12)	$1.2\pm0.8$	8.2 ± 1.1	< 0.0001 *	
	No medial pivot pt (n=6)	$5.1\pm1.4$	4.7 ± 1.7	n.s	
	Entire cohort (n=17)	$5.3 \pm 0.9$	$10.9\pm0.7$	< 0.0001 *	
Lunge	Medial pivot pt (n=12)	$5.7\pm0.8$	$11.9\pm0.8$	< 0.0001 *	
	No medial pivot pt (n=5)	$6.3 \pm 2.4$	$6.6\pm1.9$	n.s	

Table 1 – Anteroposterior (AP) Low Point translation of medial and lateral condyles during sit to stand (top) and lunge (bottom) evaluated through dynamic RSA. Data are reported as entire cohort, patients performing consistent medial pivot and patients not performing consistent medial pivot. \* represent statistically significant differences (p<0.05).

	Sit to Stand			Lunge				
CORRELATION COEFFICIENT [r]	KSSC	KSSF	WOMAC	OXFORD	KSSC	KSSF	WOMAC	OXFORD
Peak of AP translation lateral compartment	0.59 *	n.s.	n.s.	0.61 *	n.s.	n.s.	n.s.	n.s.
Peak of AP translation medial compartment	n.s.	n.s.	n.s.	n.s.	-0.55 *	n.s.	-0.60 *	-0.53 *
VV laxity	-0.56 *	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
Peak Ext-rotation	n.s.	n.s.	n.s.	n.s.	-0.66 *	n.s.	n.s.	n.s.

Table 2 – Pearson's correlation coefficients (r) between kinematical parameters and patient-reported outcomes. \* represent statistically significant differences (p<0.05).