



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Ultrasonographic measurement of the adrenal gland in neonatal foals: reliability of the technique and assessment of variation in healthy foals during the first five days of life

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Lauteri, E., Mariella, J., Beccati, F., De Graaf-Roelfsema, E., Castagnetti, C., Pepe, M., et al. (2020). Ultrasonographic measurement of the adrenal gland in neonatal foals: reliability of the technique and assessment of variation in healthy foals during the first five days of life. *THE VETERINARY RECORD*, 187(12), E117-E23 [10.1136/vr.106027].

Availability:

This version is available at: <https://hdl.handle.net/11585/797929> since: 2021-03-09

Published:

DOI: <http://doi.org/10.1136/vr.106027>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Adrenal glands ultrasonographic measurements in neonatal foals: reliability of the technique and assessment of variation in healthy foals during the first 5 days of life.

The final published version is available online at: <https://doi.org/10.1136/vr.106027>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Adrenal glands ultrasonographic measurements in neonatal foals: reliability of the technique and assessment of variation in healthy foals during the first 5 days of life.

E. Lauteri^{a1*}, J. Mariella^b, F. Beccati^a, E. Roelfsema^c, C. Castagnetti^{b,e}, M. Pepe^a, T. Peric^d, O. Barbato^a, M. Montillo^d, S. Rouge^a, F. Freccero^b

^a Department of Veterinary Medicine, University of Perugia, Perugia, Italy, Via San Costanzo 4, 06126, Perugia, Italy

^b Department of Veterinary Medical Sciences, University of Bologna, via Tolara di Sopra 50, 40064 Ozzano dell'Emilia, Bologna, Italy

^d Department of Food Sciences, University of Udine, Piazzale Osoppo 1, 33100 Udine, Italy

^c Department of Equine Sciences, Faculty of Veterinary Medicine, Utrecht University, Yalelaan 1, 3584 CL Utrecht, The Netherlands

^e Health Science and Technologies Interdepartmental Center for Industrial Research (HST-ICIR), University of Bologna, Bologna, Italy

¹ Present address: Clinèquine, Vetagro-sup, University of Lyon, 1 Avenue Bourgelat, Marcy-l'Etoile, Lyon, France

*Corresponding author. Tel: +39 3409249803

E-mail address: leonoralauteri@gmail.com (E. Lauteri)

¹ Actual address: Equine Clinic, University of Lyon, VetAgroSup, 1 Avenue Bourgelat, Marcy L'Etoile, France

Keywords: Adrenal glands, equine neonates, ultrasound, size, inter-observer

Word count: 2716

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Abstract

Background: Adrenal gland ultrasonographic measurements are useful in clinical evaluation of patients with adrenal dysfunction in several species. In human healthy neonates, ultrasonographic size of the adrenal glands decreases during the first days of life. Ultrasonography of adrenal glands was demonstrated to be feasible in neonatal foals.

The aims of this study were to describe a technique for ultrasonographic measurement of adrenal gland size, to test its reliability in neonatal foals, to assess any variation of ultrasonographic measurements during the first 5 days of life in healthy foals.

Methods: First, measurements of the adrenal glands were retrospectively obtained by three observers in 26 adrenal gland images of 13 healthy and sick neonatal foals. The inter- and intra-observer agreement were tested. Later, adrenal gland ultrasonographic images and measurements were acquired by one operator in 11 healthy neonatal foals at 1, 3 and 5 days of life and differences among the measurements obtained at the different time points were assessed.

Results: Inter-observer agreement ranged from fair to excellent (0.48–0.92), except for cortex width (<0.4); intra-observer agreement ranged from good to excellent (0.52–0.98). No significant differences were found among the measurements obtained at 1, 3 and 5 days of life.

Conclusion: Adrenal glands ultrasonographic measurements can be obtained consistently in equine neonates and, in contrast to humans, they do not vary during the first 5 days of life in healthy foals.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Introduction

Adrenal glands and hypothalamic-pituitary (HP) axis are involved in many vital processes in healthy neonate and adult horses. Hypothalamic-pituitary-adrenal (HPA) axis is involved in the homeostasis of metabolism, cardiovascular function, growth and immune response and it plays a central role in the organism response to stress and illness¹⁻³. Furthermore, emerging evidence in equine medicine suggests that adrenocortical dysfunction, resulting in cortisol insufficiency, frequently develops during critical illness in neonatal foals contributing substantially to morbidity and mortality associated with the primary disease. These findings strongly encourage the evaluation of the adrenal gland function in neonatal equine patients⁴.

Adrenal glands had been considered not detectable by ultrasound examination in neonatal foals for a long time^{5,6}. Hoffman et al obtained and described adrenal gland ultrasonographic images in neonatal foals⁷ for the first time. Recently, Beccati et al, described a simple and non-invasive ultrasonographic technique to image both adrenal glands in a transverse-oblique plan⁸. The structures and the shape of the adrenal glands were assessed and described, although only subjectively⁸. As far as we know, no data are reported on the objective or quantitative evaluation of the ultrasonographic size of the adrenal glands in neonatal foals.

Ultrasonographic examination has become the method of choice for evaluation of the appearance and size of the adrenal glands in men, dogs and cats; it is considered pivotal in clinical practice in the diagnosis of hyperadrenocorticism in dogs and in the diagnosis and treatment of congenital adrenal disorders in humans⁹⁻¹³. In human neonates, the normal variation in ultrasonographic measurements of the adrenal glands has been assessed, demonstrating a physiological decrease in ultrasonographic adrenal gland size in neonates throughout the first days of life¹⁴.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

The aims of this cohort study were: i) to describe an ultrasonographic measurement technique for adrenal glands in neonatal foals; ii) to test the reliability of the technique; iii) to assess any variation of the ultrasonographic size of the adrenal glands in the first 5 days of life in healthy neonatal foals.

Materials and Methods

Measurement technique and assessment of measurement reliability

A total of 26 two-dimensional (2D) ultrasonographic images of the left and right adrenal glands of 13 neonatal foals (5 healthy, 8 sick; 8 colts and 5 fillies, aging from 12 hours to 7 days) of various breeds were chosen from a pool of ultrasonographic images previously obtained during the foaling seasons 2014-2016 using the technique described by Beccati et al⁸.

All the images had been obtained by two different operators (F.B; E.L) with two different ultrasonographic machines (Logic C5, GE Healthcare, Milan, Italy; M-Turbo, FUJIFILM Sonosite, Bothell, USA), using a convex electronic transducer (4–5.5 MHz). The less expert operator (E.L) had obtained the images after a training period of 5 adrenal glands ultrasonographic examination under the guidance of the more expert one (F.B.). The depth setting had been fixed at 11 cm and maintained for all the foals; other parameters such as focus position and gains were adjusted individually for image optimization. The ultrasound examinations had been performed with the foals in the standing or sternal position and the haircoat over the region of the kidney was not clipped. Aorta and the left kidney and caudal vena cava and the right kidney had been used as landmarks for the correct identification of the left and right adrenal gland, respectively. Both images and videos had been obtained and stored in DICOM format. All the acquired images and frames from the video were reviewed by a single observer who selected one image for each adrenal gland of each foal. The

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

selection was based on the ability of the image or frame to allow the entire visualisation of the short and long axis of the adrenal gland

To assess the inter-observer agreement, the ultrasonographic images were then reviewed off-line, retrospectively, in a blinded fashion and independently by three different observers. To assess the intra-observer agreement, one observer repeated all the measurements twice at least one week apart.

Each observer measured (in cm) the following dimensions using the electronic calliper of a DICOM viewer:

- height and width of the left (Figure.1A) and the right (Figure.1B) whole adrenal gland;
- width of the cortex and medulla of the left (Figure. 2A) and the right (Figure. 2B) adrenal gland;
- height of the cortex and medulla of the left (Figure.3A) and the right (Figure. 3B) adrenal gland.

The height was defined as the maximum dorsoventral diameter and the measurement was made along the long axis of the gland; the width was defined as the maximum lateromedial diameter and the measurement was made perpendicular to the long axis of the gland. The width of the right adrenal gland was obtained in two sites, dorsal and ventral, because of its “peanut” shape (Figure 1B; Figure 2B; Figure 3B).

Each observer was allowed to record additional free comments to the images.

Assessment of variation in adrenal gland ultrasonographic size in the first 5 days of life

Definition criteria for foals

Eleven healthy neonatal foals (2 Warmbloods and 9 Standardbreds; 2 colts and 9 fillies) were included. The foals were privately owned, and were born from mares referred for attended delivery to the Equine Perinatology Unit of the University of Bologna, during the 2017–2018 foaling seasons.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

All procedures on the animals were carried out with the approval of the Ethical Committee of the University of Bologna. The owners gave oral informed consent for their foals' inclusion in the study. The foals were included in the study when all the following criteria were met: at term birth (>320 days) and eutocic delivery, APGAR score ≥ 9 at birth¹⁵, normal clinical examination, haematology (white blood cell count and leukogram, red blood cell and platelet counts, packed cell volume, haemoglobin concentration) and biochemistry (total proteins, albumin, total and direct bilirubin, glucose, urea, creatinine, alkaline phosphatase, lactate dehydrogenase, creatine kinase, gamma glutamyl transferase, aspartate aminotransferase, electrolyte concentrations and fibrinogen) within the normal range, and serum immunoglobulin G (IgG) concentrations >800 mg/dl at 18–24 hours of life.

Ultrasonographic protocol

Ultrasonographic examinations were performed after informed consent from horse owners. Two-D ultrasonographic images of the right and left adrenal glands were obtained at three different time-points: 1 (T1), 3 (T3) and 5 (T5) days of life. A single operator (F.F.) acquired all the images and videos using a convex transducer (1-8 MHz), (MyLabAlpha, ESAOTE, Genova, Italy). The technique described by Beccati et al⁸ was employed after a training period of five adrenal gland ultrasound examinations under the guidance of an expert operator. The depth setting had been fixed at 11 cm and maintained for all the foals; other parameters such as focus position and gains were adjusted individually for image optimization.

The ultrasound examinations were performed with the foals in the standing or sternal position and the haircoat over the region of the kidney was not clipped. Aorta and the left kidney and caudal vena cava and the right kidney were used as landmarks for the correct identification of the left and right adrenal gland, respectively. All the images and frames of the videos were stored in DICOM format.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

From the acquired images only the image allowing the best visualisation of the adrenal gland was chosen the left and right adrenal gland of each foal at each time point.

Adrenal gland measurements were obtained off-line, retrospectively, in a blinded fashion by a single observer (E.L.) that, analysed the images in a random order at the end of the season.

The following measurements were obtained as previously described: height of the left and the right whole adrenal gland, width of the left and the right whole adrenal gland, and height and width of the left and the right adrenal medulla.

Statistical analysis

All measurements were tabulated in a Microsoft Excel worksheet and transferred to R Studio (version 3.4.3) and/or JASP (version 0.8.6.0) statistical software for analyses.

Inter- and intra-observer agreement for each measurement were assessed by calculating the intraclass correlation coefficient (ICC); a value of ICC less than 0.40 was considered poor, 0.40 to 0.59 fair, 0.60 to 0.74 good and 0.75 to 1.00 excellent¹⁶.

The homoscedasticity of the variables was tested using the Shapiro–Wilk test (normal distribution) and Levene’s test (homogeneity of the variance) and subsequent tests chosen as appropriate. Differences in ultrasonographic measurements of the adrenal glands at the different time points (T1, T3, T5) were tested using a one-way ANOVA for repeated measurements.

The mean value and the standard deviation were calculated for the measurements of the adrenal glands obtained at 1, 3 and 5 days of life.

Significance was set at $p < 0.05$.

Results

Measurement technique and assessment of measurement reliability

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

All the observers judged the dorsoventral diameter of the adrenal cortex not easy to assess, especially in the left adrenal gland, because of a blurred demarcation between the cortex and the medulla. For this reason, the inter- and intra-observer reliability for the height of the cortex and medulla, were not calculated.

The overall inter-observer ICC ranged from poor (10%) to excellent (85%) depending on the measurements evaluated; specifically, the inter-observer ICC was poor for the left cortex width, right dorsal cortex width and right ventral cortex width, fair for the whole right adrenal gland height, good for the whole left adrenal gland height and left medulla width, excellent for all the other measurements considered. The overall intra-observer ICC varied from fair to excellent depending on the measurements evaluated; specifically, the ICC was fair for the right dorsal cortex width, good for the whole right ventral lobe and excellent for all the other measurements. Details of the ICC values are presented in Table 1.

Assessment of variation in adrenal gland ultrasonographic size in the first 5 days of life

The measurements of the adrenal glands in healthy foals did not change in the first 5 days of life ($p > 0.05$). Measurements obtained at each time-point are presented in Table 2.

Discussion

The results of this study demonstrate that objective ultrasonographic measurements of the adrenal glands in neonatal foals can be consistently obtained as previously described in dogs, cats and men^{10,14,17}. However, in contrast to dogs, cats and men, where measurements are usually obtained in both the transverse and longitudinal plane, in this study the adrenal glands were visualized and measured using an ultrasonographic transverse-oblique plane that has been recently suggested as the

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

easiest approach for adrenal gland visualisation in foals since it offers clearer landmarks for gland identification and it avoids the acoustic shadowing due to the ribs^{8,10,14,17}. In the transverse-oblique plane, it is not possible to assess the thickness and the length of the glands. For this reason, the measurement of the long axis corresponded to the height of the gland, and the measurement of the short axis had been defined as the width of the gland, in line with previous reports for kidney measurements in the transverse-oblique plane in the horse⁷.

For both adrenal glands, the width of the whole gland, medulla and cortex, and the height of the whole gland were successfully obtained. The height of the cortex and the medulla was not assessed because of the absence of a clear-cut corticomedullary demarcation in the long axis of the left gland due to a blurred image appearance. Image blurring is often unavoidable in ultrasound images and, as reported in previous studies, it could be a consequence of the poor lateral resolution secondary to an excessive divergence of the ultrasonographic beam. In this study the image blurring was noticed especially in the left adrenal gland, possibly because of the greater distance between the left adrenal gland and the transducer in comparison to the right gland¹⁸⁻²⁰.

The values of the ICC demonstrate that the technique can be considered reliable for all measurements considered in the study except for the cortex measurements, and that the technique should be able to be easily applied by other clinicians with a short training. The poor agreement obtained for the cortex is likely to be due to its small size, which increases the error in positioning the electronic calliper used for the measurements. With this consideration, we excluded the cortex from subsequent measurements. However, in the authors' opinion, the lack of cortex measurement does not affect the utility of the assessment of the adrenal gland size since the whole adrenal gland measurements have been demonstrated to be useful markers of the gland and cortex function in other species^{9,17}. The fair agreement obtained for the height of the whole right adrenal gland, in contrast with the good to

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

excellent agreement obtained for the other measurements, can be explained by the fact that movements of the lung could make it difficult to obtain high-quality images of the right gland⁸.

In contrast to reports in newborn children, in which a rapid decrease in the size of the adrenal glands was demonstrated between day 1 and day 5 of life, in this study no changes in adrenal size were found in neonatal foals¹⁴. These differences can reflect the different histological development of the adrenal glands in horses and men. In fact, the human fetal adrenal gland is characterized by the presence of a fetal zone, which is responsible for DHEA synthesis and androgen production in general, throughout the gestation period^{21,22}. The fetal zone occupies about 80% of the gland in newborn children and it is demonstrated to involute rapidly in the first few weeks of life¹⁴. The fetal zone characterizes the fetal adrenal glands of men and other primates, it has been described in sheep and cats, but is not present in the horse. In horses, steroidogenic enzymes are all expressed in the cortical cells and their expression increases significantly just before or after birth, corresponding to the increase in plasma cortisol that occurs very close to birth^{22,23}.

In this study, the cross-section of the aorta and the caudal vena cava represented two important markers in the identification of the left and right adrenal gland, respectively. Accordingly with the previous study by Beccati et al⁸, left adrenal gland was always identified ventrally to the aorta and medially to kidney, while the right adrenal gland was identified located between the right kidney and the caudal vena cava, confirming that the aorta and caudal vena cava represent two important ultrasonographic markers in the identification of the left and right adrenal gland, respectively, in the transversal-oblique plane.

Two main limitations are present: first of all, the repeatability of the image acquisition has not been tested; the operators, the variability intrinsic to image collection and the employ of two different ultrasonographic machines could have potentially affected the results. However, the operators involved in the study underwent the same training and, furthermore, videos, in addition to images,

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

were obtained; videos were subsequently reviewed by one single observer with the purpose of selecting images allowing the entire visualisation of the short and long axis of the gland. In light of this, in authors' opinion, the variability linked to images collection and its effect on final results are limited. The second limitation is represented by the fact that no comparison was made between ultrasonographic size of the adrenal glands and morphometric measurements on gross specimen. The technique and the peculiar anatomical localisation of the adrenal glands in the body could affect the correspondence between ultrasonographic and gross anatomical size of the adrenal glands. Small changes in dimensions could have been missed and, although not detected by the ultrasonographic examination, changes in gross dimensions of the adrenal glands in the first 5 days of life in healthy foals cannot be excluded. The assessment of variation in gross anatomical size of the adrenal glands, was, however, beyond the purpose of this study.

Conclusions

This study demonstrates that 2D ultrasonographic measurements of the adrenal glands in neonatal foals can be reliably obtained and that the ultrasonographic size of both the adrenal glands does not vary significantly during the first 5 days of life in healthy neonatal foals. This finding suggests that the same measurement ranges could be used as reference for healthy neonatal foals indistinctly from birth to the 5th day of life. In line to what previously reported in other species, further studies are recommended to investigate the correlation between measurements and sex, age and weight of the foals and to further investigate the applicability of adrenal glands ultrasonographic measurements in the clinical evaluation of foals.

Conflict of Interest Statement

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

None of the authors has any financial or personal relationships that could inappropriately influence or bias the content of the paper.

Acknowledgements

We would like to thank all the students and the staff of the Equine Teaching Hospital of the University of Perugia and of the Equine Clinical Service of University of Bologna for the help with procedures and data collection.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflicts of interest statement:

No competing interests have been declared.

References

1. Dembek KA, Onasch K, Hurcombe SDA et al. Renin-Angiotensin-aldosterone system and hypothalamic-pituitary-adrenal axis in hospitalized newborn foals. *J Vet Intern Med* 2013;27, 331-338.
2. Hurcombe SDA. Hypothalamic-pituitary gland axis function and dysfunction in horses. *Vet Clin N Am-Equine* 2011;27, 1-17.
3. Toribio RE. Endocrine dysregulation in critically ill foals and horses. *Vet Clin N Am-Equine* 2011;27, 35-47

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

4. Hart KA, Barton MH. Adrenocortical insufficiency in horses and foals. *Vet Clin N Am-Equine* 2011;27, 19-34.
5. Bain TF. Ultrasound of the thorax and abdomen in the foal. Proceeding of the 58th American Association of Equine Practitioners Conference; December 1-5, 2012; Anaheim, California, USA.
6. Rantanen NW. Disease of the abdomen. *Vet Clin N Am-Equine* 1986;2, 67-88.
7. Hoffmann KL, Wood AKW, McCarthy PH. Ultrasonography of the equine neonatal kidney. *Equine Vet J* 2000;32, 109-113.
8. Beccati F, Lauteri E, Cercone M et al. Ultrasonographic technique and appearance of adrenal gland in neonatal foals: a pilot study. *J Equine Vet Sci* 2018;61, 13-17.
9. Sandhya Bhavani M, Thirunavukkarasu PS, Kovitha S et al. Ultrasonography of adrenal glands in normal and hyperadrenocorticoid dogs. *Int J Curr Res* 2015;7, 16119-16122.
10. Cartee RE, Finn Bodner ST, Gray BW. Ultrasound examination of the feline adrenal gland. *J Diag Med Son* 1993;9, 327-330.
11. Grooters AM, Biller DS, Theisen SK et al. Ultrasonographic characteristics of the adrenal glands in dogs with pituitary-dependent hyperadrenocorticism: comparison with normal dogs. *J Vet Intern Med* 1996;10, 110-115.
12. Karaguzel G, Eyuboglu I, Ozdem S et al. Reference intervals for serum 17 α -hydroxyprogesterone and ultrasonographic adrenal gland sizes in healthy newborns. *J Matern-Fetal Neo M* 2019.
13. Wenger M, Mueller C, Kook PH et al. Ultrasonographic evaluation of adrenal glands in dogs with primary hypoadrenocorticism or mimicking diseases. *Vet Rec* 2010;167, 207-210.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

14. Scott EM, Thomas A, McGarrigle HHG et al. Serial Adrenal Ultrasonography in Normal Neonates. *J Med Ultrasound* 1990;9, 279-283.
15. Vaala WE, House JK, Madigan JE. Initial management and physical examination of the neonate. In: Large Animal Internal Medicine, Smith B.P. (Ed). Mosby: St. Louis, 2002:277-293.
16. Rutherford L, Beever L, Bruce et al. Assessment of computed tomography derived cricoid cartilage and tracheal dimensions to evaluate degree of cricoid narrowing in brachycephalic dogs. *Vet Radiol Ultrasound* 2017;59(6), 634-646.
17. Douglass JP, Clifford RB, James S. Ultrasonographic adrenal gland measurements in dogs without evidence of adrenal disease. *Vet Radiol Ultrasound* 1997;38, 124-130.
18. Alexander NG, Justiaan Swanevelder MB. Resolution in ultrasound imaging. *Anaesth Crit Care Pa* 2011;11, 186-192.
19. Goldstein A. Errors in ultrasound digital image distance measurements. *Ultrasound Med Biol* 2000;26(7):1125-1132.
20. Soulshy SN, Holland M, Hudson JA et al. Ultrasonographic evaluation of adrenal gland size compared to body weight in normal dogs. *Vet Radiol and Ultrasound* 2015;56, 317-326.
21. Mesiano S, Coulter CL, Jaffe RB. Localization of cytochrome P450 cholesterol side-chain cleavage, cytochrome P450 17 alpha-hydroxylase/17, 20-lyase, and 3 beta-hydroxysteroid dehydrogenase isomerases steroidogenic enzymes in human and rhesus monkey fetal adrenal glands: reappraisal of functional zonation. *J Clin Endocrinol Metab* 1993;77, 1184-1189.
22. Mesiano S, Jaffe RB. Developmental and functional biology of the primate fetal adrenal cortex. *Endocr Rev* 1997;18, 378-403.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

23. Han X, Fowden AL, Silver M et al. 1995. Immunohistochemical localisation of steroidogenic enzymes and phenylethanolamine-N-methyl-transferase (PNMT) in the adrenal gland of the fetal and newborn foal. *Equine Vet J* 1995;27, 140-146.

Tables

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Table 1. Intra- and inter-observer reliability for ultrasonographic measurements obtained, retrospectively, in 13 healthy and sick neonatal foals.

Measurement	Intraobserver ICC (95% CIs)	Interobserver ICC (95% CIs)
Whole left adrenal gland height	0.91 (0.72–0.97)	0.60 (0.27–0.84)
Whole left adrenal gland width	0.91 (0.72–0.97)	0.81 (0.59–0.93)
Left cortex width	0.89 (0.66–0.97)	0.27 (-0.067–0.65)
Left medulla width	0.98 (0.93–0.99)	0.63 (0.31–0.86)
Whole right adrenal gland height	0.93 (0.79–0.98)	0.48 (0.14–0.77)
Whole right dorsal lobe width	0.97 (0.90–0.99)	0.76 (0.52–0.91)
Right dorsal cortex width	0.52 (-0.02–0.82)	0.17 (-0.13–0.56)
Right dorsal medulla width	0.86 (0.60–0.96)	0.82 (0.62–0.93)
Whole right ventral lobe width	0.73 (0.32–0.91)	0.85 (0.68–0.94)
Right ventral cortex width	0.86 (0.60–0.96)	0.22 (-0.09–0.60)
Right ventral medulla width	0.84 (0.56–0.95)	0.84 (0.65–0.94)

ICC, intraclass correlation coefficient; CI, confidence intervals.

Table 2. Serial ultrasound measurements obtained in healthy neonatal foals (n = 11) at T1, T3 and T5. Data are presented as mean value and standard deviation.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Measurement (cm)	T1 (mean ± SD)	T3 (mean ± SD)	T5 (mean ± SD)	P-value
Whole left adrenal gland height	2.5 ± 0.48	2.6 ± 0.58	2.86 ± 0.52	0.2
Whole left adrenal gland width	0.69 ± 0.21	0.63 ± 0.25	0.73 ± 0.19	0.2
Left medulla width	0.52 ± 0.17	0.40 ± 0.18	0.49 ± 0.13	0.1
Whole right adrenal gland height	2.78 ± 0.35	3.08 ± 0.4	3.28 ± 0.58	0.6
Whole right dorsal lobe width	0.77 ± 0.26	0.75 ± 0.09	0.74 ± 0.15	0.2
Right dorsal medulla width	0.55 ± 0.19	0.53 ± 0.15	0.49 ± 0.13	0.1
Whole right ventral lobe width	0.79 ± 0.24	0.79 ± 0.14	0.83 ± 0.23	0.3
Right ventral medulla width	0.67 ± 0.26	0.54 ± 0.14	0.56 ± 0.21	0.1
* significative difference in row. SD = standard deviation				

Figure legends

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Figure 1. Ultrasound image obtained in a transverse-oblique plane in a standing neonatal foal. D = dorsal; M = medial. The figure illustrates the measurement of the width (white solid line) and the height (white dashed line) of the left (Figure 1A) and the right (Figure 1B) whole adrenal gland.

Figure 2. Ultrasound image of the adrenal gland obtained in a transverse-oblique plane in a standing neonatal foal. D = dorsal; M = medial. The figure illustrates the measurement of the width of the cortex (white asterisks) and the medulla (white solid line) of the left (Figure 2A) and the right (Figure 2B) adrenal gland.

Figure 3. Ultrasound image obtained in a transverse-oblique plane in a standing neonatal foal. D = dorsal; M = medial. The figure illustrates the measurement of the height of the cortex (white asterisks) and the medulla (white dashed line) of the left (Figure 3A) and the right (Figure 3B) adrenal gland.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.