

# ARCHIVIO ISTITUZIONALE DELLA RICERCA

## Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Optical spectroscopic observations of gamma-ray blazar candidates. X. Results from the 2018–2019 SOAR and OAN-SPM observations of blazar candidates of uncertain type

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

### Published Version:

Optical spectroscopic observations of gamma-ray blazar candidates. X. Results from the 2018–2019 SOAR and OAN-SPM observations of blazar candidates of uncertain type / de Menezes R.; Amaya-Almazan R. A.; Marchesini E. J.; Pena-Herazo H. A.; Massaro F.; Chavushyan V.; Paggi A.; Landoni M.; Masetti N.; Ricci F.; D'Abrusco R.; La Franca F.; Smith H. A.; Milisavljevic D.; Tosti G.; Jimenez-Bailon E.; Cheung C. C.. - In: ASTROPHYSICS AND SPACE SCIENCE. - ISSN 0004-640X. - STAMPA. - 365:1(2020), pp. 12.1-12.23. [10.1007/s10509-020-3727-5]

This version is available at: https://hdl.handle.net/11585/803898 since: 2021-02-22

Published:

DOI: http://doi.org/10.1007/s10509-020-3727-5

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

(Article begins on next page)

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version. This is the final peer-reviewed accepted manuscript of:

de Menezes, R., Amaya-Almazán, R.A., Marchesini, E.J. et al. Optical spectroscopic observations of gamma-ray blazar candidates. X. Results from the 2018–2019 SOAR and OAN-SPM observations of blazar candidates of uncertain type. Astrophys Space Sci 365, 12 (2020).

The final published version is available at: <u>http://dx.doi.org/10.1007/s10509-020-</u> 3727-5

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<u>https://cris.unibo.it/</u>)

When citing, please refer to the published version.

### Optical spectroscopic observations of gamma-ray blazar candidates. X. Results from the 2018–2019 SOAR and OAN-SPM observations of blazar candidates of uncertain type

R. de Menezes <sup>1,2</sup>, R. A. Amaya-Almazán<sup>3</sup>, E. J. Marchesini<sup>2,4,7,8,9</sup>, H. A. Peña-Herazo<sup>2,3,4,5</sup>, F. Massaro<sup>2,4,5,6</sup>, V. Chavushyan<sup>3</sup>, A. Paggi<sup>4,5</sup>, M. Landoni<sup>10</sup>, N. Masetti<sup>9,11</sup>, F. Ricci<sup>12</sup>, R. D'Abrusco<sup>13</sup>, F. La Franca<sup>15</sup>, Howard A. Smith<sup>13</sup>, D. Milisavljevic<sup>16</sup>, G. Tosti<sup>17</sup>, E. Jiménez-Bailón<sup>18</sup>, C.C. Cheung<sup>14</sup>

R. de Menezes, R. A. Amaya-Almazán, E. J. Marchesini, H. A. Peña-Herazo, F. Massaro, V. Chavushyan, A. Paggi, M. Landoni, N. Masetti, F. Ricci, R. D'Abrusco, F. La Franca, Howard A. Smith, D. Milisavljevic, G. Tosti, E. Jiménez-Bailón, C.C. Cheung

R. de Menezes

raniere.m.menezes@gmail.com

<sup>1</sup>Universidade de São Paulo, Departamento de Astronomia, São Paulo, SP 05508-090, Brazil

<sup>2</sup>Dipartimento di Fisica, Università degli Studi di Torino, via Pietro Giuria 1, I-10125 Torino, Italy

<sup>3</sup>Instituto Nacional de Astrofísica, Óptica y Electrónica, Apartado Postal 51-216, 72000 Puebla, México

<sup>4</sup>Istituto Nazionale di Fisica Nucleare, Sezione di Torino, I-10125 Torino, Italy

<sup>5</sup>INAF-Osservatorio Astrofisico di Torino, via Osservatorio 20, 10025 Pino Torinese, Italy.

<sup>6</sup>Consorzio Interuniversitario per la Fisica Spaziale (CIFS), via Pietro Giuria 1, I-10125, Torino, Italy.

<sup>7</sup>Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata, La Plata, Argentina.

<sup>8</sup>Instituto de Astrofísica de La Plata, CONICET-UNLP, CCT La Plata, La Plata, Argentina.

<sup>9</sup>INAF – Osservatorio di Astrofisica e Scienza dello Spazio, via Gobetti 93/3, I-40129, Bologna, Italy

<sup>10</sup>INAF-Osservatorio Astronomico di Brera, Via Emilio Bianchi 46, I-23807 Merate, Italy

<sup>11</sup>Departamento de Ciencias Físicas, Universidad Andrés Bello, Fernández Concha 700, Las Condes, Santiago, Chile

<sup>12</sup>Instituto de Astrofísica and Centro de Astroingeniería, Facultad de Física, Pontificia Universidad Catolica de Chile, Casilla 306, Santiago 22, Chile

<sup>13</sup>Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA

 $^{14}\mbox{Naval}$  Research Laboratory, Space Science Division, Code 7650, Washington, DC 20375, USA

<sup>15</sup>Dipartimento di Matematica e Fisica, Università degli Studi Roma Tre, Via della Vasca Navale 84, I-00146, Roma, Italy

<sup>16</sup>Department of Physics and Astronomy, Purdue University, 525 Northwestern Avenue, West Lafayette, IN 47907, USA

<sup>17</sup>Dipartimento di Fisica, Università degli Studi di Perugia, 06123 Perugia, Italy

<sup>18</sup>Instituto de Astronomía, Universidad Nacional Autónoma de México, Apdo. Postal 877, Ensenada, 22800 Baja California, México

#### Received: 01 Nov 2019 / Accepted: 09 Jan 2020

#### Abstract

The fourth Fermi Large Area Telescope Source Catalog (4FGL) lists over 5000  $\gamma$ -ray sources with statistical significance above  $4\sigma$ . About 23% of the sources listed in this catalog are unidentified/unassociated  $\gamma$ -ray sources while  $\sim 26\%$  of the sources are classified as blazar candidates of uncertain type (BCUs), lacking optical spectroscopic information. To probe the blazar nature of candidate counterparts of UGSs and BCUs, we started our optical spectroscopic follow up campaign in 2012, which up to date account for more than 350 observed sources. In this paper, the tenth of our campaign, we report on the spectroscopic observations of 37 sources, mostly BCUs, whose observations were carried out predominantly at the Observatorio Astronómico Nacional San Pedro Mártir and the Southern Astrophysical Research Observatory between August 2018 and September 2019. We confirm the BL Lac nature of 27 sources and the flat spectrum radio quasar nature of three sources. The remaining ones are classified as six BL Lacs galaxydominated and one normal galaxy. We were also able to measure the redshifts for 20 sources, including 10 BL Lacs. As in previous analyses, the largest fraction of BCUs revealed to be BL Lac objects.

**Keywords** galaxies: active - galaxies: BL Lacertae objects - quasars: general - surveys - radiation mechanisms: nonthermal

#### 1 Introduction

Blazars are one of the most peculiar and among the rarest class (Abdo et al. 2010) of active galactic nuclei (AGNs) whose emission, over the whole electromagnetic spectrum, is mainly arising from relativistic particles accelerated in a jet closely aligned to the line of sight (Blandford & Rees 1978;

Urry & Padovani 1995). Together with radio galaxies they constitute the main class of AGNs emitting at MeV-to-TeV energies, with a few exceptions due to a small subset of nearby galaxies (Abdollahi et al. 2019).

There are two main sub-classes of blazars, distinguished on the basis of their optical spectra: BL Lac objects and flat spectrum radio quasars. The former subclass presents weak emission/absorption lines in their optical spectra with equivalent width (EW) smaller than 5 Å (Stickel et al. 1991) or even completely featureless spectra, while sources belonging to the latter show quasar-like optical spectra with broad emission lines.

In the last decade we initiated and carried out an optical spectroscopic campaign to unveil the nature of potential counterparts of unidentified/unassociated  $\gamma$ -ray sources (UGSs, D'Abrusco et al. 2013; Massaro et al. 2013a; Massaro discovered with the *Fermi* Large Area Telescope (LAT) and to confirm the classification of blazar candidates of uncertain type (BCUs; Ackermann et al. 2015a).

Given the continuously increasing number of discovered UGSs by *Fermi*-LAT, our observations are mainly focused on searching blazar-like sources within the positional uncertainty of the UGSs (Acero et al. 2013; Massaro et al. 2015a; Paggi et al. 2014) and/or eventually BCUs (Massaro et al. 2017) Álvarez-Crespo et al. 2016). Given the large positional uncertainty of  $\gamma$ -ray sources, of the order of ~4 arcmin in the third and fourth releases of the *Fermi*-LAT point source catalogs (3FGL and 4FGL, Acero et al. 2015; Abdollahi et al. 201 potential counterparts, targets of our spectroscopic observations, were selected on the basis of several criteria based, for example, on their mid-infrared colors (MIR; Massaro et al. 2011; D'Abrusco et al. 2012), presence of X-ray emission (Paggi et al. 2013; Takeuchi et al. 2013) and/or presence of radio sources (Schinzel et al. 2015).

Here, in the tenth paper of a series devoted to our optical spectroscopic campaign, we focused on the observations of a selected sample of BCUs, that being already associated with *Fermi*-LAT sources, needed an optical spectroscopic classification (Ackermann et al. 2015a) to confirm their nature. In the present analysis, for the blazar classification, we adopted the nomenclature of the Roma-BZCAT (Massaro et al. 2015b) distinguishing between BL Lac objects (i.e., BZBs), flat spectrum radio quasars (labelled as BZQs), as well as blazars of galaxy type (BZGs) whereas the spectrum is dominated by the emission from its host galaxy but presents a non-thermal blue continuum likely originated in a relativistic jet (see Section 3).

The main goal of our optical spectroscopic campaign is to "hunt" blazars among UGSs and BCUs, since they constitute the largest known population of  $\gamma$ -ray emitters (Abdollahi et al. 2019). If we confirm a blazar lies within the positional uncertainty region of a UGS, it bolsters its likelihood of being the counterpart of the UGS, due to the scarcity of such sources in the sky. It is thus a valuable information, that has been so far used in every subsequent release of the *Fermi*-LAT catalogs (Massaro et al. 2015a). Even though there has been continued efforts to develop new and alternative methods to increase the association probability (e.g., D'Abrusco et al. 2019), still optical spectroscopy remains an efficient method that provides a precise classification of the potential counterpart of a  $\gamma$ -ray source.

The state of the art of our campaign can be briefly summarized as follows. We collected spectra for 337 sources to date and classified them as 255 BZBs, 39 BZQs and 20 BZGs, while additional 23 sources have been classified as normal quasars, due to the lack of multifrequency observations, particularly in the radio band, that could confirm the presence of a jet. In particular, 167 sources out of 337 etver 201 BGUs for which the largest fraction (i.e., 114 out of 167,  $\sim$ 70%) were identified as BZBs, confirming that they are the most elusive subclass of blazars. During our campaign we extensively and continuously searched in the optical databases to exclude targets for which an optical spectrum was already available (see e.g., Massaro et al. 2014; Massaro et al. 2016b) and we also compared our results with those present in the literature, due to other spectro-2bs opic campaigns (Landoni et al. 2018; Paiano et al. 2019), to maintain an updated database. The largest fraction of the sources classified to date come from the archival search in Sloan Digital Sky Survey (SDSS) data, which counts for a total of 127 spectra; and then more than 100 observations were also carried out in the southern hemisphere with the Southern Astrophysical Research Telescope (SOAR) telescope. All remaining ones were performed with other 2 m- to 6 m-class telescopes as the Kitt Peak National Observatory (KPNO), the William Herschel Telescope (WHT), Telescopio Nazionale Galileo (TNG), Nordic Optical Telescope (NOT), Observatorio Astrofísico Guillermo Haro (OAGH), Observatorio Astronómico Nacional San Pedro Mártir (OAN-SPM) and Multiple Mirror Telescope (MMT) to name a few.

Our spectroscopic identifications were used by other groups to (i) build the luminosity function of BL Lacs (Ajello et al. 2014); (ii) select potential targets for the Cherenkov Telescope Array (CTA; Massaro et al. 2013b; Arsioli et al. 2015); (iii) obtain stringent limits on the dark matter annihilation in sub-halos (e.g., Zechlin & Horns 2012; Berlin & Hooper 2014); (iv) search for counterparts of new flaring  $\gamma$ -ray sources (Bernieri et al. 2013); (v) test new  $\gamma$ ray detection algorithms (Campana et al. 2015; Campana et al. 2016); (vi) perform population studies on the UGSs (e.g., Acero et al. 2013) and (vii) discover the new subclass of radio weak BL Lacs (e.g., Massaro et al. 2017). All the above studies confirm the legacy value of our results thus motivating the prosecution of our BCU follow up campaign.

The present paper is organized as follows. In § 2 we present the selection criteria for the the BCU sample anal-

ysed here, while in § 3 the data reduction procedures for SOAR and OAN-SPM observations are described. In § 4 we discuss results obtained, and § 5 is devoted to our summary and conclusions.

#### 2 Sample selection

Our sample consists of 37 sources predominantly listed as counterparts of  $\gamma$ -ray sources in the 4FGL catalog (35 sources), as well as in the Fermi-LAT 8-year Point Source List<sup>1</sup> (FL8Y; 2 sources), a preliminary release of 4FGL based on the same 8-years of data. We highlight that the two sources selected from FL8Y do not appear in 4FGL. The 37 targets are listed as 33 BCUs and 4 BZBs in these  $\gamma$ -ray catalogs.

Sources classified as BCUs have unknown blazar nature, as all BCUs i) lack an optical spectra sensitive enough to classify them as BZBs or BZQs, or ii) are classified as blazars of uncertain or transitional types in Roma-BZCAT. However, all BCUs have multiwavelength data indicating a typical blazar-like behaviour (see Ackermann et al. 2015a; Abdollahi et al. 2019) and thus, as previously stated, our spectroscopic analysis has been carried out to confirm their real nature.

In 4FGL, there are 1312 associated sources classified as BCUs (i.e., 26% of the entire catalog), a fraction significantly higher than on its predecessor 3FGL, for which the BCUs accounted for 19% of the sample (Acero et al. 2015). All targets selected in our sample have optical magnitude in the r band brighter than 18.5, as listed in the USNO-B catalog (Monet et al. 2003). During our observing nights we were able to observe 33 BCUs and 4 sources already classified as BZBs for which we searched for a spectroscopic confirmation and potentially a redshift estimate.

We also verified which of our targets are listed in the latest releases of the Wide-field Infrared Survey Explorer (WISE) Blazar-Like Radio-Loud Sources catalog (WIBRaLS; D'Abrusco et al. 2019) and the Kernel Density Estimation selected candidate BL Lacs catalog (KDE-BLLACS; D'Abrusco et al. 2019), both known to present a

We found a total of 19 matches with WIBRaLS (14 classified here as BZBs, 2 BZGs and 3 BZQs; see § 4) and 3 more matches with KDEBLLACS (all classified here as BZBs, see § 4). Additionally, two of our targets, 4FGL J1208.4+6121 and 4FGL J1239.4+0728, are listed in Roma-BZCAT and classified there as blazars of uncertain type.

Although we selected the BCUs mainly from 4FGL, in our previous papers we tend to select sources based on MIR 6

1.6

2.5

W3-W4 (mag)

3.0

3.5

4.0

2.0

1.0

1.5

<sup>∿</sup> ⊼∄ BCUs 1.4 BZCAT BZB BZCAT BZQ 1.2 1.0 W1-W2 (mag) 0.8 0.6 0.4 0.2 0.0 -0.2 L 2 3 4 5 1 W2-W3 (mag) ★ BCUs BZCAT BZB 4.0 **BZCAT BZQ** 3.5 (mage) 3.0 (mag) 2.5 (mage) 2.0 ( 3.0 1.5 1.0 0.5

high number of spectroscopically confirmed blazars (de Menezes et al. 2019) Fig. 1 WISE color-color diagrams. Top: distribution of Roma-BZCAT  $\gamma$ -ray BZBs (blue circles), BZQs (red circles) and our targets (stars) in the W1-W2 × W2-W3 diagram. Here, black stars represent the targets detected in WISE W4 band, while the white stars lack such a detection. Bottom: same as upper panel but for  $W2-W3 \times W3-W4.$ 

methods, as the *Fermi*-LAT  $\gamma$ -ray blazars are known to occupy a distinct area in the WISE MIR color-color-color diagram (D'Abrusco et al. 2012; D'Abrusco et al. 2019). Here we compare the distribution of our targets in the WISE MIR color-color diagrams W1-W2 vs. W2-W3 and W2-W3 vs. W3-W4, using the magnitudes measured at the four WISE bands W1[3.4  $\mu$ m], W2[4.6  $\mu$ m], W3[12  $\mu$ m] and W4[22  $\mu$ m], with the distribution of MIR colors for  $\gamma$ -ray blazars listed in Roma-BZCAT, finding that all of them have MIR colors consistent with  $\gamma$ -ray blazars (Figure 1), except perhaps for 4FGL J1612.2+2828/WISE J161217.62+282546.3, the black star in the bottom of both panels, classified as the only normal galaxy in our sample. Among our sources, 27 are detected in WISE W4 band and plotted as black stars in Figure 1. The remaining targets are plotted as white stars. The distribution of our targets in these diagrams suggest that their infrared emission is dominated by non-thermal radiation, as expected for  $\gamma$ -ray blazars.

*WISE* magnitudes used here are in the Vega system and are not corrected for the Galactic extinction. As shown in our previous analyses (D'Abrusco et al. 2013; D'Abrusco et al. 2014), such correction affect only the magnitude measures at  $3.4\mu$ m for sources lying at low Galactic latitudes, and it ranges between 2% and 5%, thus being almost negligible.

#### **3** Observations and data reduction

We observed a total of 25 targets with the 2.1 m telescope at OAN-SPM, in Baja California, México, during several nights in 2018–2019 (see Table 1). The data were collected with the Boller & Chivens spectrograph, with a slit width of 2.5", grating of 300 l/mm, wavelength range from 3800 Å up to 7800 Å, and resolution of ~ 14 Å.

A single source, 4FGL J0434.7+0922/TXS 0431+092, was observed with the 2.1 m telescope at OAGH in Cananea, México, using the Boller & Chivens spectrograph. The collected spectrum has a wavelength range from  $\sim 3600$  to 7300 Å and was acquired with a slit width of 2.5" and grating of 150 l/mm, having a final resolution of  $\sim 14$  Å.

The data for the remaining 11 targets were collected with the 4.1 m SOAR telescope at Cerro Pachón, Chile, in remote observing mode on May 22 and 23 of 2019. We used the single, long slit mode of the Goodman High Throughput Spectrograph (Clemens et al. 2004) with slit width of 1" and a grating of 400 l/mm, giving a dispersion of ~ 3Å per pixel in a spectral range from ~ 4100 Å up to 7900 Å, and resolution of ~ 6Å.

In Table 1 we report the log of all observations, giving the name of the  $\gamma$ -ray source and its associated counterpart in the *Fermi*-LAT catalogs, the associated counterpart in WISE, the telescope used, the average signal to noise ratio (SNR), the exposure time and the date of observation.

The data were reduced using standard procedures, with the images trimmed, bias subtracted, corrected for flat field and stacked using standard IRAF packages (Tody 1986). The cosmic ray removal was performed with Python packages ccdproc (Craig et al. 2015) and astropy (Astropy Collaboration et al. Astropy Collaboration et al. 2018), in the case of SOAR data, and using the L.A. Cosmic IRAF algorithm (van Dokkum 2001) in the case of OAN-SPM data. All spectra were then flux and wavelength calibrated using, respectively, a standard spectrophotometric star and a Hg-Ar (SOAR) or CuNeHeAr (OAN-SPM) lamp. The corrections for Galactic extinction were made using the reddening law of Cardelli et al. (1989) and values of visual extinction  $A_{\nu}$ computed based on Schlafly & Finkbeiner (2011) assuming a visual extinction to reddening ratio  $A_v/E(B-V) = 3.1$ . Finally, we performed a box smoothing for visual representation, normalized the spectra using a 5th degree polynomial fitted to the continuum to highlight possible spectroscopic features, and visually identified the emission/absorption lines (Figure 2).

#### 4 Analysis and results

Thanks to the new optical spectra collected, we are able to confirm the blazar nature for 36 targets in our sample, consisting of 27 BZBs, 6 BZGs, and 3 BZQs. Additionally, we identified a normal galaxy listed in 4FGL as the counterpart of 4FGL J1612.2+2828.

We obtained the redshifts (z) for 20 of the targets, including 10 BZBs, all with relatively low z distributed in the range 0.12–0.28, as expected for BZBs. Particularly we successfully measured redshifts for 3 of the 4 BZBs listed in the 4FGL (see Section 2). They are 4FGL J0420.2+4012/WISE J042013.43+401121.2 with z = 0.132, 4FGL J2220.5+2813/WISE J222028.72+281355.8 with z = 0.148 and 4FGL J2358.3+3830/WISE J235825.19+382856.5 with z = 0.200. For the last source, we observed emission lines with EW > 5 Å, and thus classified it as a BZQ.

All results achieved for the entire sample are summarized in Table 2, where we show the name of the  $\gamma$ -ray sources as listed in the *Fermi*-LAT catalogs, the name of the WISE counterparts, the redshifts when available, the spectral classes and information about the observed absorption/emission lines.

The optical spectra and finding charts for the analysed sources are available in Appendix A, ordered by right ascension. Finding charts were obtained from ESO Online Digitized Sky Survey<sup>2</sup> and have a size of  $5' \times 5'$ . The name of each emission/absorption line is also indicated in the figure. Telluric absorption lines are labelled with  $\bigoplus$ .

<sup>&</sup>lt;sup>2</sup>http://archive.eso.org/dss/dss

Criteria adopted to distinguishing between BZBs, BZGs and normal galaxies are based on the observed Ca II H&K break contrast. We computed the ratio  $C = (F_+ - F_-)/F_+$ where  $F_+$  and  $F_-$  are the mean flux densities measured in small ranges of 200 Å just above (4050-4250 Å) and below (3750-3950 Å) the Ca II H&K break, respectively. A value of C < 0.4 means that the source has a blue continuum emission dominated by non-thermal radiation likely originated in a relativistic jet (Marcha et al.1996; Landt et al. 2002; Massaro et al. 2012a). Then sources with C > 0.4 are classified as normal galaxies, sources with  $0.25 < C \le 0.4$  are classified as BZG, while values of  $C \le 0.25$  lead towards a BZB or BZO classification.

To distinguish between BZBs and BZQs we computed the rest frame equivalent width (EW) of their emission lines (when available) and classified them as BZBs if all lines had  $EW \le 5$ Å. In particular, we found only 3 BCUs having emission lines with EW > 5Å, leading to a BZQ classification.

For those sources for which a redshift estimate has been provided, the redshift uncertainties are of the order of  $\Delta z = 0.001$  for the data collected with SOAR, while those pointed with OAGH and OAN-SPM have  $\Delta z \sim 0.004$ .

#### 5 Summary and Conclusions

We presented the tenth paper of our optical spectroscopic campaign aiming at the identification of BCUs listed in *Fermi*-LAT catalogs. We collected and analysed data for a total of 37 sources, confirming the blazar nature for 36 of them. Our results are summarized as follows:

- Among the 33 sources classified as BCUs in the *Fermi*-LAT catalogs, 24 of them (73%) are found to be BZBs, 6 are BZGs (18%), 2 are BZQs (6%) and 1 is a normal galaxy (3%). We also measured the red-shifts for 17 of them.
- We observed 4 sources classified as BZBs in 4FGL and were able to measure the redshifts for 3 of them: 4FGL J0420.2+4012, 4FGL J2220.5+2813 and 4FGL J2358.3+3830, with z = 0.132, z = 0.148 and z =0.200 respectively. In particular, 4FGL J2358.3+3830 presents emission lines with EW > 5Å, thus being classified as BZQ.
- All sources classified as blazars (36 out of 37) have MIR colors consistent with what is expected for γ-ray blazars (Figure 1), except perhaps for the counterpart of 4FGL J1608.0-2038.

The results here, as in the previous papers of our campaign (see e.g., Marchesini et al. 2019; Peña-Herazo et al. 201 confirm that the largest fraction of BCUs tend to be BZBs. These objects, although very common in the  $\gamma$ -ray sky, continue to be one of the most elusive classes of  $\gamma$ -ray sources.

Our optical spectroscopic campaign is still ongoing with observing nights scheduled for the current and next semesters. New results will be made available as soon as they are collected and analysed. The next paper of the campaign will focus mainly on UGSs (Peña-Herazo et al. 2019) observed with the 4 m Blanco telescope at Cerro Tololo, Chile. We observed a total of 374 targets to date, mostly BCUs and UGSs, but including a small fraction (i.e., of the order of a few percent) of known blazars, especially BL Lac objects, trying to catch them in a low optical state and obtain a *z* estimate. Particularly, out of 200 sources previously classified as BCUs, we found that 141 of them are BZBs.

This work was supported by FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo) under grants 2016/25484-9 and 2018/24801-6 (R.M.). R.A.A-A. and H.P. acknowledge support from the CONACyT program for Ph.D. studies. The work of F.M. and A.P. is partially supported by the "Departments of Excellence 2018-2022" Grant awarded by the Italian Ministry of Education, University and Research (MIUR) (L. 232/2016) and made use of resources provided by the Compagnia di San Paolo for the grant awarded on the BLENV project (S1618\_L1\_MASF\_01) and by the Ministry of Education, Universities and Research for the grant MASF\_FFABR\_17\_01 F.M. also acknowledges financial contribution from the agreement ASI-INAF n.2017-14-H.0 while A.P. the financial support from the Consorzio Interuniversitario per la Fisica Spaziale (CIFS) under the agreement related to the grant MASF\_CONTR\_FIN\_18\_02. F.R. acknowledges support from FONDECYT Postdoctorado 3180506 and CONI-CYT project Basal AFB-170002. C.C.C. at NRL was supported by the Chief of Naval Research.

This research made use of Astropy,<sup>3</sup> a communitydeveloped core Python package for Astronomy (Astropy Collaboration et al. 2 Astropy Collaboration et al. 2018), ccdproc, an Astropy package for image reduction (Craig et al. 2015), and TOP-CAT<sup>4</sup> (Taylor 2005) for the preparation and manipulation of the tabular data.

This project makes use of data products from the Southern Astrophysical Research (SOAR) telescope, which is a joint project of the Ministério da Ciência, Tecnologia, Inovações e Comunicações (MCTIC) da República Federativa do Brasil, the U.S. National Optical Astronomy Observatory (NOAO), the University of North Carolina at Chapel Hill (UNC), and Michigan State University (MSU).

This project makes use of spectroscopic observations acquired at the 2.1 m telescope of the Observatorio Astronómico Nacional San Pedro Mártir (OAN-SPM), Baja California, México. We also thank the staff at the Observatorio Astrofísico Guillermo Haro (OAGH) for all their help during the observation runs.

<sup>&</sup>lt;sup>3</sup>http://www.astropy.org

<sup>&</sup>lt;sup>4</sup>http://www.star.bris.ac.uk/~mbt/topcat/

 Table 1
 Observation log. Col. (1) The name reported in the *Fermi*-LAT catalogs (i.e., FL8Y and 4FGL); Col. (2) the name of the associated counterpart listed in the *Fermi*-LAT catalogs; Col. (3) the WISE name of the associated counterpart; Col. (4) and Col. (5) The coordinates (Equinox J2000) of the pointed source; Col. (6) the telescope used to carry out the observation; Col. (7) The average signal to noise ratio; Col. (8) the exposure time and Col. (9) the observation date. The source tagged with "\*" is found only in the AllWISE Multiepoch Photometry Table.

4FGL J2358.3+3830	4FGL J2236.6+3706	4FGL J2220.5+2813	FL8Y J2201.6+2953	4FGL J1858.7+5708	4FGL J1838.0-5959	4FGL J1831.9+3820	4FGL J1830.0-5225	4FGL J1808.2+3500	4FGL J1741.2+5739	4FGL J1707.5+1649	4FGL J1612.2+2828	4FGL J1608.0-2038	4FGL J1606.9+5919	4FGL J1550.8-0822	4FGL J1546.1-1003	4FGL J1440.0-2343	4FGL J1400.2-4010	4FGL J1351.4-1529	4FGL J1310.6+2449	4FGL J1239.4+0728	4FGL J1213.8-4345	4FGL J1208.4+6121	4FGL J1156.6-2248	4FGL J1042.1-4128	4FGL J1028.3+3108	4FGL J0928.7-3529	4FGL J0842.7+6656	4FGL J0602.7-0007	FL8Y J0516.6+2742	4FGL J0442.7+6142	4FGL J0434.7+0922	4FGL J0420.2+4012	4FGL J0325.3+3332	4FGL J0204.3+2417	4FGL J0119.6+4158	4FGL J0015.9+2440		Fermi-LAT name
B3 2355+382	NVSS J223626+370713	RX J2220.4+2814	2WHSP J220123.8+294934	87GB 185759.9+570427	SUMSS J183806-600033	1RXS J183202.2+382132	SUMSS J183004-522618	MG2 J180813+3501	NVSS J174111+573812	MG1 J170732+1649	TXS 1610+285	NVSS J160756-203942	1RXS J160709.7+592115	NVSS J155053-082247	PMN J1546-1003	PMN J1439-2341	2MASS J14002208-4008235	2MASX J13511746-1530155	MG2 J131037+2447	PKS 1236+077	PMN J1213-4343	RGB J1208+613	NVSS J115633-225004	1RXS J104204.1-412936	TXS 1025+313	NVSS J092849-352947	TXS 0838+671	PMN J0602-0004	VCS J0516+2743	GB6 J0442+6140	TXS 0431+092	1RXS J042008.3+401138	2MASX J03251760+3332435	B2 0201+24	2MASX J01200274+4200139	GB6 J0016+2440		Association
J235825.19+382856.5	J223626.34+370713.5	J222028.72+281355.8	J220123.82+294934.5	J185853.50+570809.5	J183806.74-600032.1	J183200.98+382137.0	J183004.32-522618.8	J180811.52+350118.7	J174111.70+573812.4	J170731.55+164844.6	J161217.62+282546.3	J160756.90-203943.5	J160708.14+592129.5	J155053.26-082246.7	J154611.48-100326.1	J143959.46-234141.0	J140022.08-400823.6	J135117.45-153016.0	J131038.49+244822.7	J123924.58+073017.2	J121350.38-434324.6	J120837.13+612106.4	J115633.22-225004.2	J104203.00-412929.9	J102817.61+310734.4	J092849.82-352948.8	J084243.19+665729.4	J060242.88-000426.7	J051640.48+274310.2	J044240.66+614039.4*	J043440.98+092348.6	J042013.43+401121.2	J032517.58+333243.7	J020421.54+241750.7	J012002.76+420013.8	J001603.62+244014.7		WISE name
23:58:25	22:36:26	22:20:29	22:01:24	18:58:54	18:38:07	18:32:01	18:30:04	18:08:12	17:41:12	17:07:32	16:12:18	16:07:57	16:07:08	15:50:53	15:46:11	14:39:59	14:00:22	13:51:17	13:10:38	12:39:25	12:13:50	12:08:37	11:56:33	10:42:03	10:28:18	09:28:50	08:42:43	06:02:43	05:16:40	04:42:41	04:34:41	04:20:13	03:25:18	02:04:22	01:20:03	00:16:04	(hh:mm:ss)	R.A. (J2000)
+38:28:57	+37:07:14	+28:13:56	+29:49:35	+57:08:10	-60:00:32	+38:21:37	-52:26:19	+35:01:19	+57:38:12	+16:48:45	+28:25:46	-20:39:44	+59:21:30	-08:22:47	-10:03:26	-23:41:41	-40:08:24	-15:30:16	+24:48:23	+07:30:17	-43:43:25	+61:21:06	-22:50:04	-41:29:30	+31:07:34	-35:29:49	+66:57:29	-00:04:27	+27:43:10	+61:40:39	+09:23:49	+40:11:21	+33:32:44	+24:17:51	+42:00:14	+24:40:15	(dd:mm:ss)	Dec. (J2000)
OAN	OAN	OAN	OAN	OAN	SOAR	OAN	SOAR	OAN	OAN	OAN	OAN	SOAR	OAN	OAN	SOAR	SOAR	SOAR	SOAR	OAN	OAN	SOAR	OAN	SOAR	SOAR	OAN	SOAR	OAN	OAN	OAN	OAN	OAGH	OAN	OAN	OAN	OAN	OAN		Telescope
7	6	11	10	7	19	8	13	14	6	8	11	18	10	8	21	19	12	19	7	17	16	9	20	20	6	16	10	9	8	S	S	10	9	8	=	15		SNR
3600	1800	1800	1800	3600	2 x 1200	3600	2 x 1600	1200	1800	1800	1200	2 x 1700	1800	1800	2 x 1200	2 x 1500	2 x 1800	2 x 1300	1800	1800	2 x 1500	1800	2 x 1300	2 x 1500	3600	2 x 1200	1800	3600	3600	1800	1800	1800	1200	1800	1800	1800	(s)	Exposure
04/08/2018	18/10/2018	03/08/2018	05/08/2018	08/08/2018	23/05/2019	07/08/2018	23/05/2019	08/04/2019	04/07/2019	07/08/2018	06/06/2019	22/05/2019	05/06/2019	05/08/2018	23/05/2019	22/05/2019	22/05/2019	23/05/2019	08/04/2019	08/04/2019	23/05/2019	07/04/2019	23/05/2019	23/05/2019	06/04/2019	22/05/2019	07/04/2019	17/10/2018	18/10/2018	17/10/2018	08/11/2018	19/10/2018	17/10/2018	18/10/2018	17/10/2018	17/10/2018	(dd/mm/yyyy)	Obs. date

Fermi-LAT name	WISE counterpart	Class	С	z	Line	EW (Å)	$\lambda_{obs}$ (Å)	Туре
4FGL J0015.9+2440	J001603.62+244014.7	BZB						
4FGL J0119.6+4158	J012002.76+420013.8	BZG	0.30	0.109	[O II]	5	4132	Е
					Ca II H&K	10; 9	4362; 4399	Α
					G band	5	4773	Α
					Mg I	5	5737	Α
					Na I	5	6532	А
4FGL J0204.3+2417	J020421.54+241750.7	BZB	0.21	0.210	Ca II H&K	7; 5	4760; 4803	А
					G band	3	5510	Α
					Mg I	5	6267	А
					Na I	4	7131	Α
4FGL J0325.3+3332	J032517.58+333243.7	BZG	0.36	0.128	Ca II H&K	12; 9	4440; 4477	А
					G band	6	4855	А
					Mg I	6	5838	А
4FGL J0420.2+4012	J042013.43+401121.2	BZB	0.01	0.132	G band	4	4874	A
					Mg I	6	5860	А
					Na I	4	6676	А
4FGL J0434.7+0922	J043440.98+092348.6	BZB						
4FGL J0442.7+6142	J044240.66+614039.4	BZB						
FL8Y J0516.6+2742	J051640.48+274310.2	BZG	0.34	0.060	[O II]	5	3952	E
					Ca II H&K	8;6	4172; 4206	А
					G band	3	4566	А
					Mg I	5	5492	А
					Na I	4	6251	А
					$H_{\alpha}$	5	6962	E
					[N II]	7	6984	E
					[S II]	2	7125	Е
					[S II]	2	7141	E
4FGL J0602.7-0007	J060242.88-000426.7	BZG	0.29	0.118	Ca II H&K	5	4447	A
					G band	6	4810	A
					Mg I	6	5788	А
4FGL J0842.7+6656	J084243.19+665729.4	BZB	0.24	0.121	Ca II H&K	8; 8	4412; 4450	A
					G band	3	4829	A
					Mg I	4	5803	A
					Na I	3	6609	A
4FGL J0928.7-3529	J092849.82-352948.8	BZB						
4FGL J1028.3+3108	J102817.61+310734.4	BZB						
4FGL J1042.1-4128	J104203.00-412929.9	BZB						
4FGL J1156.6-2248	J115633.22-225004.2	BZB		0.860?	Mg II?	2;2	5200; 5214	A
4FGL J1208.4+6121	J120837.13+612106.4	BZB	0.13	0.275	Ca II H&K	3; 3	5018; 5062	A
					G band	4	5486	A
4501 11010 0 4045	1101050 00 404004 (	DZD			Mg I	4	0004	А
4FGL J1213.8-4345	J121350.38-434324.6	BZB						
4FGL J1239.4+0728	J123924.58+073017.2	BZB	0.00	0.226		0.10	4925, 4966	
4FGL J1310.0+2449	J151058.49+244822.7	BZG	0.29	0.220	C hor -	9; 10 5	4823; 4800	A
					G band	3	5209	A
					MgI	4	0342	А

**Table 2** Summary of our spectroscopic identifications. Col. (1) the name reported in the *Fermi*-LAT catalogs (i.e., FL8Y and 4FGL); Col. (2) the WISE name of the associated counterpart; Col. (3) the source class (i.e., BZB for BL Lac objects, BZQ for Blazars of quasar type, BZG for blazars of galaxy type); Col. (4) the measured Ca II H&K contrast; Col. (5) measured redshift; Col. (6) Emission or Absorption lines identified; Col. (7) Measured equivalent width (EW); Col. (8) Observed wavelength of each emission/absorption line identified; Col. (9) type of line: absorption (A) or emission (E).

Fermi-LAT name	WISE counterpart	Class	C	z	Line	EW (Å)	$\lambda_{obs}$ (Å)	Туре
4FGL J1351.4-1529	J135117.45-153016.0	BZB	0.01	0.285	Ca II H&K	1; 2	5061; 5101	A
					G band	2	5530	A
					Mg I	1	6656	A
4FGL J1400.2-4010	J140022.08-400823.6	BZB	0.24	0.203	Ca II H&K	7	4775	A
					G band	4	5177	A
					MσI	5	6226	A
					NaI	3	7090	A
4FGL 11440 0-2343	114395946-2341410	BZB			i u i	5	1090	
4FGL 11546 1-1003	I154611 48-100326 1	BZB						
4FGL 11550 8-0822	1155053 26-082246 7	BZB						
4FGL 11606 9±5919	$1160708 14 \pm 592129 5$	BZB	0.18	0.132	10 11	4	4217	F
41 OL J1000.7 ( 571 )	J100700.14+372127.5	DLD	0.10	0.152		т 1·1	4453-4401	
					G band	т, т 3	4872	
					Mal	2	5855	
					No I	5	5655	
AECL 11609 0 2029	1160756 00 202042 5	D7D			INAI	2	0007	A
4FGL J1008.0-2038	J100/30.90-203943.3	BZB	0.47	0.052	C. HILLEV	10.0	4140. 4102	
4FGL J1612.2+2828	J101217.02+282540.5	galaxy	0.47	0.053	Call H&K	10; 9	4148; 4183	A
					G band	7	4537	A
					MgI	5	5454	A
					Nal	4	6211	A
					$H_{\alpha}$	2	6915	E
						2	6938	E
		-			[S II]?	9	7081	E
4FGL J1707.5+1649	J170731.55+164844.6	BZQ	0.10	0.291	[O II]	7	4817	E
					Ca II H&K	1	5133	A
					G band	1	5558	A
					[O III]	3; 5	6409; 6470	E
4FGL J1741.2+5739	J174111.70+573812.4	BZB						
4FGL J1808.2+3500	J180811.52+350118.7	BZB		0.269?	[O II]?	5	4721	E
4FGL J1830.0-5225	J183004.32-522618.8	BZB						
4FGL J1831.9+3820	J183200.98+382137.0	BZG	0.30	0.216	Ca II H&K	8;7	4785; 4826	A
					G band	4	5234	A
					Mg I	2	6304	A
					Na I	4	7168	A
4FGL J1838.0-5959	J183806.74-600032.1	BZB						
4FGL J1858.7+5708	J185853.50+570809.5	BZQ	-0.02	0.076	[O II]	4	4017	E
					$H_{\beta}$	2	5236	E
					[O III]	4;5	5346; 5393	E
					$H_{\alpha}$	12	7067	E
FL8Y J2201.6+2953	J220123.82+294934.5	BZB	0.22	0.149	Ca II H&K	7;4	4521; 4560	A
					G band	3	4945	A
					Mg I	3	5951	A
4FGL J2220.5+2813	J222028.72+281355.8	BZB	0.24	0.148	Ca II H&K	8; 5	4519; 4558	A
					G band	4	4944	A
					Mg I	4	5945	A
					NaI	4	6768	A
4FGL J2236.6+3706	J223626.34+370713.5	BZB	0.15	0.235	Ca II H&K	6:5	4861: 4899	A
		_			G band	3	5318	A
					MgI	4	6393	A
4FGL 12358 3+3830	J235825 19+382856 5	BZO	0.00	0.200		9	4474	E
	12002010100200000			0.200	H <sub>e</sub>	1	5837	Ē
						4:10	5953: 6011	Ē
					[Fe VIII?	4	6874	A
		l			LIC, III.			_ <b>^ *</b>

**Table 3**Continued from Table 2.

9

This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration.

#### References

- Abdo, A. A., Ackermann, M., Agudo, I. et al. 2010, ApJ, 716, 30
- Abdollahi, S., Acero, F., Ackermann, M. et al. 2019, arXiv e-prints, arXiv:1902.10045v4
- Acero, F., Donato, D., Ojha, R. et al. 2013, ApJ, 779, 133
- Acero, F., Ackermann, M., Ajello, M. et al. 2015, ApJS, 218, 23
- Ackermann, M., Ajello, M., Atwood, W. B. et al. 2015, ApJ, 810, 14
- Ajello, M., Romani, R. W., Gasparrini, D., et al. 2014, ApJ, 780, 73
- Álvarez-Crespo, N., Massaro, F., D'Abrusco, R. et al. 2016, Ap&SS 361, 316
- Arsioli, B. et al. 2015, A&A, 579, A34
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, A&A, 558, A33
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, AJ, 156, 123
- Berlin, A., & Hooper, D. 2014, Phys. Rev. D, 89, 016014
- Bernieri, E. et al. 2013 A&A, 551, L5
- Blandford, R.D. & Rees, M.J. 1978, In: Proc. Pittsburgh Conference on BL Lac Objects, p. 328
- Campana, R. et al. 2015, Ap&SS, 360, 65
- Campana, R., Massaro, E., Bernieri, E. 2016, Ap&SS 361, 183
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, ApJ, 345, 245
- Clemens, J. C., Crain, J. A., & Anderson, R. 2004, Proc. SPIE, 5492, 331
- Craig, M. W., Crawford, S. M., Deil, C., et al. 2015, ccdproc: CCD data reduction software, ascl:1510.007
- D'Abrusco, R., Massaro, F., Ajello, M. et al. 2012, ApJ, 748, 68
- D'Abrusco, R., Massaro, F., Paggi, A. et al. 2013, ApJS, 206, 12
- D'Abrusco, R., Massaro, F., Paggi, A. et al. 2014, ApJS, 215, 14
- D'Abrusco, R., Álvarez Crespo, N., Massaro, F., et al. 2019, ApJS, 242, 4
- de Menezes, R., Peña-Herazo, H. A., Marchesini, E. J., et al. 2019, A&A, 630, A55
- Landoni, M., Paiano, S., Falomo, R. et al. 2018, ApJ, 861, 130
- Landt, H., Padovani, P., & Giommi, P. 2002, MNRAS, 336, 945
- Marcha, M. J. M., Browne, I. W. A., Impey, C. D., et al. 1996, MNRAS, 281, 425
- Marchesini, E. J., Peña-Herazo, H. A., Álvarez Crespo, N. et al. 2019, Ap&SS, 364, 5
- Massaro, F., D'Abrusco, R., Ajello, M. et al. 2011, ApJ, 740, L48
- Massaro, E., Nesci, R. & Piranomonte S. 2012a, MNRAS, 422, 2322
- Massaro, F., D'Abrusco, R., Tosti, G. et al. 2012b, ApJ, 750, 138
- Massaro, F., D'Abrusco, R., Paggi, A. et al. 2013a, ApJS, 206, 13
- Massaro, F., D'Abrusco, R., Paggi, A. et al. 2013b, ApJS, 209, 10
- Massaro, F., Masetti, N., D'Abrusco, R. et al. 2014, AJ, 148, 66
- Massaro, F., D'Abrusco, R., Landoni, M. et al. 2015a, ApJS, 217, 2
- Massaro, E., Maselli, A., Leto, C. et al. 2015b, Ap&SS, 357, 75
- Massaro, F., Thompson, D. J., Ferrara, E. C. 2016a, A&AR, 24, 2 Massaro, F., Álvarez Crespo, N., D'Abrusco, R. et al. 2016b,
- Ap&SS 361, 337
- Massaro, F., Marchesini, E. J., D'Abrusco, R., et al. 2017, ApJ, 834, 113
- Monet, D. G., Levine, S. E., Canzian, B., et al. 2003, AJ, 125, 984
- Paggi, A., Massaro, F., D'Abrusco, R. et al. 2013, ApJS, 209, 9
- Paggi, A., Milisavljevic, D., Masetti, N. et al. 2014, AJ, 147, 112
- Paiano, S., Falomo, R., Treves, A., Franceschini, A., & Scarpa, R. 2019, ApJ, 871, 162

- Peña-Herazo, H. A., Massaro, F., Chavushyan, V., et al. 2019, Ap&SS, 364, 85
- Schinzel, F. K., Petrov, L., Taylor, G. B. 2015, ApJS, 217, 4
- Schlafly, E. F., & Finkbeiner, D. P. 2011, ApJ, 737, 103
- Stickel, M., Padovani, P., Urry, C. M. et al. 1991, ApJ, 374, 431
- Takeuchi, Y., Kataoka, J., Maeda, K. et al. 2013, ApJS, 208, 25
- Taylor, M. B. 2005, ASP Conf. Ser., 347, 29
- Tody, D. 1986, SPIE, 627, 733
- Urry, C. M., & Padovani, P. 1995, PASP, 107, 803
- van Dokkum, P. G. 2001, PASP, 113, 1420V
- Zechlin, H. & Horns, D. 2012, JCAP, 11, 050.

This manuscript was prepared with the AAS LATEX macros v5.2.

Here we present all the spectra collected together with their respective finding charts extracted from ESO Online Digitized Sky Survey (Figure 2).



Fig. 2 Left: flux calibrated and normalized optical spectra. Right: finding charts. The name of the  $\gamma$ -ray sources and their respective counterparts listed in the *Fermi*-LAT catalogs are indicated, together with the spectral class and average SNR. Redshifts and emission/absorption lines are shown whenever possible.



Fig. 3 Continued from Figure 2.



Fig. 4 Continued from Figure 2.



Fig. 5 Continued from Figure 2.



Fig. 6 Continued from Figure 2.



Fig. 7 Continued from Figure 2.



Fig. 8 Continued from Figure 2.



Fig. 9 Continued from Figure 2.



Fig. 10 Continued from Figure 2.



Fig. 11 Continued from Figure 2.



Fig. 12 Continued from Figure 2.



Fig. 13 Continued from Figure 2.



Fig. 14 Continued from Figure 2.