

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Corrigendum to "Orlik-Solomon-type presentations for the cohomology algebra of toric arrangements"

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Callegaro F., D'Adderio M., Delucchi E., Migliorini L., Pagaria R. (2021). Corrigendum to "Orlik-Solomon-type presentations for the cohomology algebra of toric arrangements". TRANSACTIONS OF THE AMERICAN MATHEMATICAL SOCIETY, 374(5), 3779-3781 [10.1090/tran/8262].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/832437> since: 2021-09-16

*Published:*

DOI: <http://doi.org/10.1090/tran/8262>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

**Callegaro, F., D'Adderio, M., Delucchi, E., Migliorini, L., & Pagaria, R. (2021). Corrigendum to “Orlik-Solomon-type presentations for the cohomology algebra of toric arrangements” (trans. amer. math. soc. (2020) 373:3 (1909-1940) DOI: 10.1090/tran/7952). Transactions of the American Mathematical Society, 374(5), 3779-3781**

The final published version is available online at <https://dx.doi.org/10.1090/tran/8262>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

ERRATUM TO “ORLIK-SOLOMON-TYPE PRESENTATIONS  
 FOR THE COHOMOLOGY ALGEBRA OF TORIC  
 ARRANGEMENTS”

FILIPPO CALLEGARO, MICHELE D’ADDERIO, EMANUELE DELUCCHI,  
 LUCA MIGLIORINI, AND ROBERTO PAGARIA

ABSTRACT. In this short note we correct the statement of the main result of [1]. That paper presented the rational cohomology ring of a toric arrangement by generators and relations. One of the series of relations given in [1] is indexed over the set circuits in the arrangement’s arithmetic matroid. That series of relations should however be indexed over all sets  $X$  with  $|X| = \text{rk}(X) + 1$ . Below we give the complete and correct presentation of the rational cohomology ring.

We state the correct version of [1, Theorem 6.13]:

**Theorem 1.** *Let  $\mathcal{A}$  be an essential toric arrangement. The rational cohomology algebra of the complement  $H^*(M(\mathcal{A}), \mathbb{Q})$  is isomorphic to the algebra  $\mathcal{E}$  with*

- *Set of generators  $e_{W,A;B}$ , where  $W$  ranges over all layers of  $\mathcal{A}$ ,  $A$  is a set generating  $W$  and  $B$  is disjoint from  $A$  and such that  $A \sqcup B$  is an independent set; the degree of the generator  $e_{W,A;B}$  is  $|A \sqcup B|$ .*
- *The following types of relations:*
  - *For any two generators  $e_{W,A;B}$ ,  $e_{W',A';B'}$ ,*

$$e_{W,A;B}e_{W',A';B'} = 0$$

*if  $A \sqcup B \sqcup A' \sqcup B'$  is a dependent set, and otherwise*

$$(1) \quad e_{W,A;B}e_{W',A';B'} = (-1)^{\ell(A \cup B, A' \cup B')} \sum_{L \in \pi_0(W \cap W')} e_{L, A \cup A'; B \cup B'}.$$

- *For every linear dependency  $\sum_{i \in E} n_i \chi_i = 0$  with  $n_i \in \mathbb{Z}$ , a relation*

$$(2) \quad \sum_{i \in E} n_i e_{T, \emptyset; \{i\}} = 0.$$

- *For every subset  $X \subseteq E$  where  $\text{rk}(X) = |X| - 1$  write  $X = C \sqcup F$  with  $C$  the unique circuit in  $X$ . Consider the associated (unique) linear dependency  $\sum_{i \in C} n_i \chi_i = 0$  with  $n_i \in \mathbb{Z}$ , and for every connected*

---

2010 *Mathematics Subject Classification.* Primary 14N20, 52C35, 55R80.

The first and fourth authors were supported by PRIN 2015 “Moduli spaces and Lie theory” 2015ZWST2C - PE1.

The third author was supported by the Swiss National Science Foundation professorship grant PP00P2\_150552/1.

component  $L$  of  $\bigcap_{i \in X} H_i$  a relation

$$(3) \quad \sum_{j \in C} \sum_{\substack{A, B \subset X \\ A \supseteq F \\ X = A \sqcup B \sqcup \{j\} \\ |B| \text{ even} \\ W \supseteq L}} (-1)^{|A \leq j|} c_B \frac{m(A)}{m(A \cup B)} e_{W, A; B} = 0$$

where, for all  $i \in C$ ,  $c_i := \text{sgn } n_i$ ,  $c_B = \prod_{i \in B} c_i$ .

The only difference between Theorem 1 and [1, Theorem 6.13] consists in eq. (3) that hold not only for every circuit  $C$  but also for all  $X$  with  $|X| = \text{rk}(X) + 1$ . This difference is important as shown in the following example.

**Example 2.** Consider the central toric arrangement in  $T = (\mathbb{C}^*)^3$  given by the four hypertori  $H_1 = \{x = 1\}$ ,  $H_2 = \{y = 1\}$ ,  $H_3 = \{xy = 1\}$ , and  $H_4 = \{xy^{-1}z^3 = 1\}$ . The zero-dimensional layers are the points  $p = (1, 1, 1)$ ,  $q = (1, 1, \zeta_3)$ , and  $r = (1, 1, \zeta_3^2)$ . Let  $W$  be the layer  $H_1 \cap H_2 \cap H_3$ . The relations given by (3) are the following:

$$\begin{aligned} \bar{\omega}_{W, \{1,2\}} - \bar{\omega}_{W, \{1,3\}} + \bar{\omega}_{W, \{2,3\}} + \psi_1 \psi_2 - \psi_1 \psi_3 - \psi_2 \psi_3 &= 0 \\ \bar{\omega}_{s, \{1,2,4\}} - \bar{\omega}_{s, \{1,3,4\}} + \bar{\omega}_{s, \{2,3,4\}} + \frac{1}{3} \psi_1 \psi_2 \bar{\omega}_4 - \frac{1}{3} \psi_1 \psi_3 \bar{\omega}_4 - \frac{1}{3} \psi_2 \psi_3 \bar{\omega}_4 &= 0 \end{aligned}$$

for all  $s = p, q, r$ . Notice that the relation in degree two does not imply the relations in degree three.

The proof of Theorem 1 is the same as in [1] with the following corrections (see also the preprint arXiv:1806.02195v3).

Let  $X \subseteq E$  with  $|X| = \text{rk}(X) + 1$ , then  $X$  can be written uniquely as  $C \sqcup F$  where  $C$  is a circuit and  $F = X \setminus C$ . Theorem 6.12 of [1] holds in a wider generality: the set  $C$  does not need to be a circuit but can be any  $X \subseteq E$  with  $|X| = \text{rk}(X) + 1$ . It can be proven by choosing a suitable separating cover of  $X$ : for  $i \in C$  define  $a_i = m(X) \prod_{j \in C \setminus \{i\}} m(C \setminus \{j\})$  and  $a_i = m(X)$  for  $i \in F$ . Let  $\Lambda(X)$  be the lattice generated by  $\frac{X_i}{a_i}$  for all  $i \in X$ , it defines a covering  $\pi_U: U \rightarrow T$  of tori (cf. [1, Definition 6.6]).

Lemmas 6.3, 6.4, 6.5, 6.7, 6.8, and 6.10 of [1] should be corrected by changing  $C$  with  $X$  and with minor changes in their proofs. The proof of [1, Theorem 6.12] needs an extra step: let  $L$  be a connected component of  $\bigcap_{i \in X} H_i$  and  $p$  a point in  $L$ , we use the relation (12) of [1] in the torus  $U$  to obtain:

$$\sum_{j \in C} \sum_{\substack{A, B \subset C \\ C = A \sqcup B \sqcup \{j\} \\ |B| \text{ even}}} (-1)^{|A \leq j|} \bar{\eta}_{A, B}^U(q) c_B = 0.$$

for all  $q \in \pi_U^{-1}(p)$ . We multiply this equation by  $\bar{\eta}_{F, \emptyset}^U(q)$ , and using the equality  $\bar{\eta}_{A, B}^U(q) \bar{\eta}_{F, \emptyset}^U(q) = (-1)^{|F \leq j| + \ell(C, F)} \bar{\eta}_{A \sqcup F, B}^U(q)$  we obtain:

$$\sum_{j \in C} \sum_{\substack{A, B \subset X \\ A \supseteq F \\ X = A \sqcup B \sqcup \{j\} \\ |B| \text{ even}}} (-1)^{|A \leq j|} \bar{\eta}_{A, B}^U(q) c_B = 0.$$

This corrects the proof of [1, Theorem 6.12].

In the proof of Theorem 6.13 it was claimed that the old relations allow one to write each  $e_{W,A;B}$  in term of generators with  $A$  a no-broken-circuit set. In the following we prove the claim by using the new relations.

Indeed if  $A$  contains a broken-circuit  $A_1 \subseteq A$ , i.e.  $A_1 = C \setminus \min(C)$  for a circuit  $C$ , consider the relation (3) for  $X = C \cup A$ : it expresses the element  $e_{W,A;B}$  as a linear combination of some  $e_{W',A';B'}$  with  $|A'| < |A|$  or with  $|A'| = |A|$  and  $A'$  lexicographically smaller than  $A$ . We have inductively proved the claim.

## REFERENCES

- [1] Filippo Callegaro, Michele D’Adderio, Emanuele Delucchi, Luca Migliorini, and Pagaria Roberto, *Orlik-Solomon type presentations for the cohomology algebra of toric arrangements*, Trans. Amer. Math. Soc. **373** (2019), no. 3, 1909–1940.

FILIPPO CALLEGARO

UNIVERSITÀ DI PISA, DIPARTIMENTO DI MATEMATICA, LARGO BRUNO PONTECORVO 5, 56127 PISA, ITALY

*Email address:* `callegaro@dm.unipi.it`

MICHELE D’ADDERIO

UNIVERSITÉ LIBRE DE BRUXELLES (ULB), DÉPARTEMENT DE MATHÉMATIQUE, BOULEVARD DU TRIOMPHE, B-1050 BRUXELLES, BELGIUM

*Email address:* `mdadderi@ulb.ac.be`

EMANUELE DELUCCHI

UNIVERSITÉ DE FRIBOURG, DÉPARTEMENT DE MATHÉMATIQUES, CHEMIN DU MUSÉE 23, CH-1700 FRIBOURG, SWITZERLAND

*Email address:* `emanuele.delucchi@unifr.ch`

LUCA MIGLIORINI

UNIVERSITÀ DI BOLOGNA, DIPARTIMENTO DI MATEMATICA, PIAZZA DI PORTA SAN DONATO 5, 40126 BOLOGNA, ITALY

*Email address:* `luca.migliorini@unibo.it`

ROBERTO PAGARIA

UNIVERSITÀ DI BOLOGNA, DIPARTIMENTO DI MATEMATICA, PIAZZA DI PORTA SAN DONATO 5, 40126 BOLOGNA, ITALY

*Email address:* `roberto.pagaria@unibo.it`