



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

## ARCHIVIO ISTITUZIONALE DELLA RICERCA

### Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Nectar chemistry is not only a plant's affair: floral visitors affect nectar sugar and amino acid composition

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Bogo G., Fisogni A., Rabassa-Juventeny J., Bortolotti L., Nepi M., Guarnieri M., et al. (2021). Nectar chemistry is not only a plant's affair: floral visitors affect nectar sugar and amino acid composition. *OIKOS*, 130(7), 1180-1192 [10.1111/oik.08176].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/837802> since: 2021-11-09

*Published:*

DOI: <http://doi.org/10.1111/oik.08176>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

0 **Nectar chemistry is not only a plant's affair: floral visitors affect** 61  
**nectar sugar and amino acid composition**

5 **Gherardo Bogo, Alessandro Fisogni, Joan Rabassa-Juventeny, Laura Bortolotti, Massimo Nepi,** 65  
**Massimo Guarnieri, Lucia Conte and Marta Galloni**

10  
15 *G. Bogo* (<https://orcid.org/0000-0001-7415-2224>) ([gberardo1985@hotmail.com](mailto:gberardo1985@hotmail.com)), *Grupo de Ecología de la Polinización, INIBIOMA (Univ. Nacional del Comahue-CONICET), San Carlos de Bariloche, Argentina.* – *A. Fisogni* (<https://orcid.org/0000-0001-6179-2767>), *Dept of Evolution, Ecology and* 75  
*Organismal Biology, Univ. of California, Riverside, Riverside, CA, USA.* – *AF, L. Conte* (<https://orcid.org/0000-0003-0513-0616>) and *M. Galloni* (<https://orcid.org/0000-0001-5304-7820>), *Dept of Biological, Geological and Environmental Sciences, Univ. of Bologna, Bologna, Italy.* – *J. Rabassa-Juventeny* (<https://orcid.org/0000-0002-5554-2304>), *Faculty of Biology, Univ. of Barcelona, Barcelona, Spain.* – *L. Bortolotti* (<https://orcid.org/0000-0001-7372-1579>), *CREA Research Centre for Agriculture and Environment, Bologna, Italy.* – *M. Nepi* (<https://orcid.org/0000-0002-6765-897X>) and *M. Guarnieri,* 80  
*Dept of Life Sciences, Univ. of Siena, Siena, Italy.*

25 Floral nectar is the primary reward directly consumed by floral visitors and its chemical 85  
composition affects their behaviour and fidelity. In turn, floral visitors are expected to  
alter floral nectar composition directly or indirectly through the introduction of external  
contaminants, such as pollen grains and microorganisms. To understand the effect  
of exogenous factors on nectar chemistry, we investigated the modifications of nectar  
30 sugar and amino acid composition in relation to floral visitors in the perennial herb 90  
*Gentiana lutea*. We quantified nectar contamination by floral visitors through visitor  
exclusion experiments in the field, estimating pollen and yeast concentration in visited  
flowers and in uncontaminated nectar at two different flower stages (flower bud and  
35 older open flowers). We then assessed changes in the composition and concentration 95  
of sugars and both protein and non-protein amino acid in the nectar. We found clear  
differences in the chemical composition of flower nectar in relation to flower stage  
and floral visitation. Nectar sugar and amino acid concentrations were significantly  
40 higher in older flowers than in flower buds. In addition, nectar sugar concentration 100  
was significantly higher in older flowers where visitors were excluded than in visited  
flowers. On the contrary, amino acid concentration was comparable between visited  
and non-visited older flowers. We also found that pollen contamination enriched the  
45 amino acid profile of the host nectar both quantitatively and qualitatively. Finally, 105  
nectar exposed to floral visitors showed a higher number of yeast cells than unvisited  
flowers although too low to cause the observed changes in the glucose:fructose ratio,  
suggesting the presence of fructose-related bacteria. Our results show that the chemical  
50 composition of floral nectar is altered by several endogenous and exogenous factors 110  
in a complex process of ecological relations, and suggest that the modulation of floral  
visitation through nectar is not solely related to the plant itself.

55 **Keywords:** generalist pollination, *Gentiana lutea*, nectar contamination, nectar 115  
microorganisms, nectar yeast, plant–pollinator interactions

60 121

## 0 Introduction

5 Floral nectar is a complex aqueous solution secreted by plants whose chemical composition is dominated by three simple sugars: the disaccharide sucrose and its monosaccharide constituents, fructose and glucose. These three sugars are derived either from sucrose translocated into phloem sap and/or from the hydrolysis of starch accumulated in the developing nectaries (Pacini and Nepi 2007, Heil 2011). The maintenance of the sugar ratio can be regulated by the enzyme invertase, which hydrolyzes sucrose into glucose and fructose before, during and after nectar secretion (Nicolson and Thornburg 2007). In addition to sugars, other substances are present in smaller quantities, such as proteins, protein and non-protein amino acids, lipids, phenols, alcohols, alkaloids and antioxidants (Nicolson and Thornburg 2007, Nepi 2014).

10 The chemistry of floral nectar can be shaped by ecological as well as by phylogenetic constraints (Nepi et al. 2010). Among the first, interaction with specific guilds of pollinators may drive selection towards convergent nectar chemistry in unrelated taxa (Fenster et al. 2004, Pozo et al. 2015). On the other hand, phylogenetic conservatism may result in similar nectar chemistry in related taxa regardless of their pollinators (Nicolson and Thornburg 2007, Nepi et al. 2010). Nectar components have two main functions: to mediate plant interactions with pollinators (attraction) and to protect nectar (protection) from other floral visitors, such as nectar robbers, and from nectar-dwelling microorganisms (Adler 2000, Heil 2011, Chalcoff et al. 2017). Several studies have shown that differences in nectar composition were related to different groups of floral visitors. For example, sucrose-rich nectars are usually preferred by hummingbirds, Megachiroptera, Lepidoptera and long-tongued bees, while hexose-rich nectars are usually preferred by passerine birds, Microchiroptera, short-tongued bees and flies (Baker and Baker 1982, 1983, 1990, Kress 1985, Baker et al. 1998). Moreover, butterflies and flies usually show a preference for nectars with high amino acid concentration (Potter and Bertin 1988, Alm et al. 1990, Erhardt and Rusterholz 1998), honey bees prefer sugar solutions enriched with proline and phenylalanine (Inouye and Waller 1984, Alm et al. 1990, Bertazzini et al. 2010), and bumble bees prefer solutions enriched with  $\beta$ -alanine (Bogo et al. 2019). Recent studies focusing on nectar secondary compounds and their effects on pollinators (Nepi 2014, Stevenson et al. 2017, Mustard 2020) challenge the classical view reported above in which just the alimentary importance of nectar is highlighted. According to the 'manipulation' hypothesis nectar is considered a pollinator manipulator rather than simply an attractant or reward for pollinators (Pyke 2016, Nepi et al. 2018).

15 Quantity and quality of nectar can vary considerably in relation to several abiotic and biotic factors (Baude et al. 2011). Changes can be induced by water availability, ambient humidity and temperature, soil-related factors, light availability and CO<sub>2</sub> concentration (Chalcoff et al. 2017, Parachnowitsch et al. 2019). At the same time, nectar composition can be affected by flower phenology, interactions

61 with herbivores, nectar robbers or floral visitors, and by the presence of bacteria and yeasts (Aizen and Raffaele 1996, Lasso and Naranjo 2003, Vannette and Fukami 2018, Parachnowitsch et al. 2019) resulting in complex determination of nectar chemical profile. 65

The concentration of nectar solutes can also change in relation to flower age, independently of external factors (Pacini and Nepi 2007). Once the flowers open, floral visitors can affect nectar composition directly and indirectly in several ways. Visitors can alter nectar sugar composition by feeding on it, often stimulating further secretion. On the other hand, failure to visit can also affect nectar composition, often causing the reabsorption of unconsumed sugars (Pacini and Nicolson 2007, Nepi and Stpiczyńska 2008). Floral visitors can also modify nectar chemical composition by adding amino acids present in their saliva or by introducing conspecific or heterospecific pollen into the nectar (Willmer 1980, Gottsberger et al. 1990). Moreover, floral visitors can transfer yeasts directly between flowers or indirectly by transporting yeast-contaminated pollen grains (Herrera et al. 2009, Mittelbach et al. 2015, Pozo et al. 2015). In turn, nectar yeasts can cause important variation in the composition and concentration of nectar sugars (Canto and Herrera 2012, Chappell and Fukami 2018), and can decrease the concentration of certain amino acids (Vannette and Fukami 2018) or increase fatty acid derivatives (Yang et al. 2019). 70 75 80 85


In this article we aim to better understand how flower nectar composition is altered by floral visitors and endogenous factors. Specifically, we performed floral visitor exclusion experiments in the field to compare the composition of virgin (i.e. uncontaminated) nectar with that of flower nectar in which visits were allowed, to evaluate the effects of exogenous factors (i.e. insect floral visits and contamination by pollen grains and yeasts) on sugar and amino acid composition. In addition, we analysed the composition of nectar collected from flower buds to evaluate intrinsic changes related to flower age. In this way we can obtain comprehensive information on the alteration of nectar chemistry occurring at different floral stages and after floral visitation. We expected to find differences in sugar and amino acid concentration and in amino acid spectrum richness related to flower age, regardless of exogenous factors. Moreover, based on the potential direct and indirect effects of floral visitors, we also expected an increase in the presence of pollen grains and yeasts in the nectar, which in turn would modify nectar chemistry (Fig. 1). 90 95 100 105

## 110 Material and methods

### 115 Model species and study site

*Gentiana lutea* subsp. *symphyandra* is a long-lived scapose hemicryptophyte. It presents an unbranched stout stem (rarely two), growing up to 2 m tall (Tutin et al. 1972). Flowering occurs between June and July. Flowering stems carry up to ten pseudo-whorls containing about 20 pedicellate flowers each. On average, flower lifespan lasts about three 121

0  
5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60



61  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
121

Figure 1. Concept map of the endogenous and exogenous factors that can directly or indirectly influence each other and flower nectar composition.




Figure 2. Stages of flower development in *Gentiana lutea* subsp. *symphyandra* (a–e) and study site (f). (a) Floral bud, (b) stage I: perianth open, anthers closed and stigma unreceptive, (c) stage II: perianth open, one to four dehiscent anthers, unreceptive stigma undivided or hardly bilamellate (here a small bee of the family Halictidae is collecting pollen from the anthers), (d) stage III: perianth open, anthers completely dehiscent and stigma bilamellate, (e) stage IV: perianth withered.

121



0 days and four stages of flower development can be recognized  
 after floral buds (Fig. 2a) have opened (Rossi 2012): 1) peri-  
 5 anth open, anthers closed and stigma unreceptive (Fig. 2b),  
 2) perianth open, one to four dehiscent anthers, unreceptive  
 stigma undivided or hardly bilamellate (Fig. 2c), 3) perianth  
 10 open, anthers completely dehisced and stigma bilamellate  
 (Fig. 2d) and 4) perianth withered (Fig. 2e). Flowers of *G.*  
*lutea* subsp. *symphyandra* are visited by several insect species  
 belonging to the orders Hymenoptera, Diptera, Coleoptera  
 and Lepidoptera, of which wild bees are among the most  
 15 efficient pollinators (Rossi et al. 2014). Although *G. lutea* is  
 a self-compatible species, partial flower dichogamy reduces  
 within-flower selfing while pollinator-mediated geitonogamy  
 is more likely to occur (Rossi et al. 2014). Seeds developed  
 from self-pollination have lower viability and germination  
 than cross-pollinated seeds (Rossi et al. 2016).

Samplings and field observations were performed in a  
 population situated on the northeast face of Mount Grande  
 (Vidiciatico, Bologna – Italy), within the Habitats Directive  
 (92/43/EEC) Site of Community Importance and Special  
 Protection Area IT4050002 ‘Corno alle Scale’ (44°8’57”N,  
 10°52’10”E, 1380–1460 m a.s.l.). This isolated population  
 (Fig. 2f) occurs in a meadow surrounded by a beech forest  
 and is probably preserved by the steepness of the slope and  
 by the rocky nature of the substrate (Rossi 2012, Rossi et al.  
 2014). In the studied population plants bloom synchronously  
 with a peak occurring in mid-July, and flowers present  
 copious amounts of nectar (up to 10 µl, Bogo et al unpubl.).  
 30 Bumblebees are the most abundant flower visitors (about  
 50% of the total visitors), and present the highest visitation  
 frequency, fidelity and pollinator performance (Rossi et al.  
 2014).

### 35 Test design

We collected nectar samples from four different flower groups  
 (hereafter: flower treatments): 1) flower buds, to investigate  
 the composition of freshly secreted unvisited pure nectar  
 (young virgin nectar, YVN); 2) flowers with open perianth  
 40 emasculated and bagged from the bud stage, to investigate the  
 composition of unvisited nectar a few days old not contami-  
 nated by own or external pollen (old virgin nectar, OVN); 3)  
 45 non-emasculated flowers with open perianth bagged from the  
 bud stage, to investigate the composition of unvisited nectar  
 contaminated by pollen from the same flower (self-contami-  
 nated nectar, SCN); 4) non-emasculated flowers with open

perianth, bagged only three to four hours before sampling  
 to allow floral visits but to avoid nectar depletion (naturally  
 contaminated nectar, NCN) (Table 1). Flowers allocated to  
 treatments OVN, SCN and NCN were bagged individually  
 65 with nonwoven fabric (30 g m<sup>-2</sup>, white, permeable to air,  
 water and light) to exclude floral visitors. We also sampled  
 nectar from flower buds to test for the presence of invertase  
 activity (INV). Finally, we collected pollen from dehiscent  
 anthers to investigate the effect of pollen contamination on  
 70 the nectar amino acid profile under controlled conditions.  
 When possible, we applied all treatments to different flow-  
 ers on the same plant. If the four flower developmental  
 stages required for treatments YVN, OVN, SCN and NCN  
 were not present together, we sampled flowers from differ-  
 75 ent plants (Table 1). The number of treatments allocated per  
 plant ranged between three and six.

### 80 Nectar sampling

Nectar was collected from a total of nine plants and 90 flow-  
 ers of *G. lutea* (Table 1) throughout the study population  
 during two non-consecutive days (July 14 and 17) in 2015.  
 We sampled nectar using Drummond Microcaps of different  
 85 volumes, depending on its amount in the flower (3, 5 and  
 10 µl). For each sample, we measured the nectar level in the  
 microcap using a Vernier calliper and then calculated its exact  
 volume as a proportion of the total microcap volume. Nectar  
 samples were then transferred into separate eppendorf tubes  
 90 filled with 100 µl ethanol. All samples were transported to  
 the laboratory in thermal insulated ice containers on the same  
 day of field sampling, and stored at -20°C until analyses.

### 95 Nectar chemical analyses

Just prior to analyses nectar samples were thawed and air-  
 dried in a Speedvac centrifuge to eliminate the ethanol and  
 diluted 1:50 with distilled water. We analysed nectar sugar  
 composition in all samples from the four flower treatments  
 100 (n=78 flowers belonging to 4–6 different plants per treat-  
 ment, Table 1) by isocratic HPLC, using a Waters Sugar-Pak  
 I ion-exchange column (6.5 × 300 mm) maintained at 90°C  
 and a Waters 2410 refractive index detector. Water (MilliQ,  
 105 pH 7) was used as mobile phase at a flow rate of 0.6 ml min<sup>-1</sup>.  
 The sample and standard solutions containing glucose, fruc-  
 tose and sucrose (20 µl) were injected. The concentration of  
 each single sugar was calculated by comparing the area under

50 Table 1. Summary of the experimental treatments and sample size (n) used in this study.

Treatment	Abbr.	Flower stage	Bagged	Emasculated	n plants	n flowers
Young virgin nectar	YVN	bud	No	No	6	19
Old virgin nectar	OVN	III–IV	Yes	Yes	4	19
Self-contaminated nectar	SCN	III–IV	Yes	No	5	20
Naturally contaminated nectar	NCN	III–IV	No	Yes	6	20
Invertase activity	INV	bud	No	No	7	12
Pollen suspension	–	II–III	No	No	3	5

Abbr.: treatment abbreviation. Flower stage II: perianth open, one to four dehiscent anthers, unreceptive stigma undivided or hardly bilamel-  
 late. Flower stage III: perianth open, anthers completely dehisced and stigma bilamellate, Flower stage IV: perianth withered.

0 the chromatogram peaks with standards using the software  
Clarity (DataApex).

We analysed amino acid composition in a subset of eight  
randomly chosen samples per flower treatment ( $n=32$  flow-  
ers belonging to 3–5 plants per treatment) by gradient HPLC,  
5 using an ion exchange Novapak C18 ( $15 \times 4.6$  mm) cartridge  
with guard column maintained at  $37^\circ\text{C}$  and a Waters 470  
scanning fluorescence detector (excitation at 295 nm, detec-  
tion at 350 nm). A solvent composed of TEA-phosphate buf-  
fer (pH 5.0) mixed with a 6:4 acetonitrile–water solution was  
10 used as mobile phase at a flow rate of  $1.0 \text{ ml min}^{-1}$ . According  
to AccQtag protocol, the selected volume ( $20 \mu\text{l}$ ) of each  
reconstituted sample was amino acid derivatized (Cohen and  
Michaud 1993) with AQC fluorescent reagent and 0.02 M  
borate buffer (pH 8.6). In addition to all the protein amino  
15 acids (PAA), standards of the non-protein amino acids  
(NPAA)  $\beta$ -alanine, citrulline, l-homoserine,  $\alpha$ -aminobutyric  
acid (AABA),  $\gamma$ -aminobutyric acid (GABA), hydroxyproline,  
ornithine and taurine were also used (Nocentini et al. 2012).  
Tryptophan was not detectable with this procedure. The  
20 concentration of each individual amino acid was calculated  
by comparing the area under the chromatogram peaks with  
standards using the software Clarity (DataApex).

### 25 Pollen and yeast detection

Each of the 78 nectar samples was topped up to  $200 \mu\text{l}$  with  
ethanol. We then determined pollen and yeast content by  
30 counting the number of pollen grains and yeast cells present  
in aliquots of  $10 \mu\text{l}$  in ethanol. These aliquots were placed on  
a microscope slide with a glass cover slip and observed under  
an optical microscope for counting the number of pollen  
grains and yeast cells present on the slide. We calculated the  
35 total number of pollen grains and yeast cells in each sample  
using the expression  $N=(X \times 20)/V$  where  $X$  is the aliquot  
count and  $V$  is the nectar sample volume without ethanol.  
Moreover, we discriminated between pollen of *G. lutea* and  
40 pollen of other species based on pollen grain morphology.

### Invertase test

Since the floral nectar of *G. lutea* is almost devoid of sucrose,  
45 we checked for the presence of invertase activity in the secre-  
tion. For this purpose we collected  $100 \mu\text{l}$  of nectar from 12  
flower buds (Table 1, INV), with a micropipette K7501 2–20  
 $\mu\text{l}$ . The nectar sample was diluted 1:20 with a  $4.6 \text{ mg ml}^{-1}$   
sucrose solution. The sugar profile at time zero was deter-  
50 mined soon after the dilution. Subsequently, we incubated  
the solution at  $30^\circ\text{C}$  and repeated the sugar determination  
after 1, 2, 3, 4 and 24 h to quantify the potential hydrolysis of  
sucrose into glucose and fructose. The determination of sugars  
was carried out according to the procedure described above.

### 55 Pollen suspension test

We suspended 1 mg of *G. lutea* pollen collected from five  
60 flowers on three plants (Table 1) in  $100 \text{ mg}$  of a solution  
mimicking the average sugar composition of the young virgin

nectar (glucose  $27.65 \text{ mg ml}^{-1}$ , fructose  $26.59 \text{ mg ml}^{-1}$ ,  
61 Fig. 4a–b). We sampled  $10 \mu\text{l}$  of the suspension after a few  
minutes and after 1, 4 and 24 h from the pollen suspension,  
and measured the amino acid content by gradient HPLC  
65 (same methodology explained above) to evaluate if the amino  
acid profile had changed.

### Data analysis

70 We performed a principal component analysis (PCA) on the  
subset of eight samples for each of the four flower treatments  
to explore similarities in nectar composition, considering  
sugar and amino acid concentration and concentrations of  
pollen grains and yeast cells (component variables are listed in  
75 Supporting information), and a linear discriminant analysis  
(LDA) with jackknife leave-one-out cross validation to gener-  
ate posterior probability scores for each treatment based on  
principal components. Data were scaled and centred around  
the mean, and analyses were performed using the functions  
80 ‘dudi.pca’ and ‘lda’ in the R-packages *ade4* and *MASS*, respec-  
tively (Venables and Ripley 2002, Dray and Dufour 2007).

We estimated the effect of flower treatment on sugar and  
amino acid concentration by fitting linear mixed-effects mod-  
85 els (LMMs) using the function ‘lmer’ in the R-package *lme4*  
(Bates et al. 2015). For sugars, we fitted four separate mod-  
els using the log-transformed glucose concentration, fructose  
concentration, total sugar concentration and glucose:fructose  
ratio as response variable, respectively. For amino acids, we  
90 fitted four separate models using the log-transformed PAA  
concentration, NPAA concentration, total amino acid con-  
centration and PAA:NPAA ratio as response variable, respec-  
tively. In all models we included the four flower treatments  
(YVN, OVN, SCN, NCN) as explanatory variables, and whorl  
95 identity on the plant stem nested within plant identity  
as random effect to account for intra- and inter-plant vari-  
ability. We then performed pairwise comparisons between  
treatments by estimating the marginal means (EMMs) with  
100 Tukey-adjusted p-values using the function ‘emmeans’ in the  
R-package *emmeans* (Lenth et al. 2020).

We analysed sucrose, glucose and fructose concentrations  
among the different time intervals in the test ‘invertase’ using  
105 a Pearson’s chi-squared test.

We measured the diversity of amino acids based on their  
concentration using Hill numbers of order 1, corresponding  
to the exponential of Shannon entropy (Jost 2006). We esti-  
110 mated the effect of flower treatment on amino acid diversity  
by fitting linear models (LMs) using Hill numbers as response  
variable and the four flower treatments (YVN, OVN, SCN,  
NCN) as explanatory variables. Flower treatments were then  
compared between them using EMMs with Tukey-adjusted  
115 p-values.

We estimated the effect of flower treatment on pollen and  
yeast concentration by fitting two separate LMMs includ-  
ing  $\log(x+1)$ -transformed pollen and yeast concentrations  
as response variables, respectively, and the four flower treat-  
120 ments (YVN, OVN, SCN, NCN) as explanatory variables.  
Whorl identity nested within plant identity was included

as random effect to account for inter-flower and inter-individual variability. Flower treatments were then compared by estimating EMMs with Tukey-adjusted p-values. Figures were drawn using the R-package *ggplot2* (Wickham 2016), and all analyses were performed in R ver. 4.0.2 (<www.r-project.org>).

## Results

### Nectar analyses

The PCA showed a clear separation of the flower treatments with regard to nectar composition, with the first two components explaining 73.0% of the variance (Fig. 3). The first component was positively correlated with the protein amino acid and total amino acid concentrations (PC1 loadings=0.44 for both). The second component was positively correlated with the concentration of yeast cells and with the glucose:fructose ratio (PC2 loadings=0.45 and 0.58, respectively; Supporting information). The LDA showed a total correct assignment by cross-validation of 93.75% of samples (Supporting information). The correct assignment was 100% in treatments YVN, OVN and SCN, and 75% in treatment NCN.

### Sugar composition

We excluded sucrose from the analysis of nectar sugar composition because its concentration was always lower than 1.5% of the total sugar concentration. We found significant effects of flower treatment on glucose ( $F_{3,74}=376.97$ ,  $p < 2.2e^{-16}$ ), fructose ( $F_{3,74}=461.38$ ,  $p < 2.2e^{-16}$ ) and total sugar ( $F_{3,74}=422.05$ ,  $p < 2.2e^{-16}$ ) concentration in nectar. Sugar concentrations always showed the same pattern,




Figure 3. Principal components analysis (PCA) of the nectar composition in eight flowers for each of the four flower treatments, based on sugar and amino acid concentration and on concentrations of pollen grains and yeast cells. Yellow dots: young virgin nectar (YVN); blue dots: old virgin nectar (OVN); green dots: self-contaminated nectar (SCN); purple dots: naturally contaminated nectar (NCN).

with the lowest value found for YVN, followed by NCN, OVN and the highest value for SCN (Fig. 4a–c, Supporting information).

Glucose:fructose ratio was significantly higher in the NCN than in the other treatments ( $p < 0.0001$  all tests), while no other differences were observed (Fig. 4d, Supporting information).

### Amino acids composition

We found significant effects of flower treatment on protein ( $F_{3,28}=17.55$ ,  $p < 7.1 \times 10^{-06}$ ), non-protein ( $F_{3,28}=28.32$ ,  $p < 2.9 \times 10^{-07}$ ) and total amino acid ( $F_{3,28}=33.51$ ,  $p < 2.1 \times 10^{-09}$ ) concentration in nectar. YVN always had the lowest concentration (Fig. 5a–c, Supporting information). Both PAA and total amino acid concentrations were similar among the OVN, SCN and NCN treatments (Fig. 5a–c), while NPAA concentration did not differ significantly between the two contaminated treatments with the highest value observed in the OVN treatment (marginally significant difference between OVN and SCN,  $p=0.09$ ; Fig. 5b, Supporting information).

The PAA:NPAA ratio was significantly different among treatments ( $F_{3,28}=6.64$ ,  $p=0.0026$ ). The YVN and OVN treatments had similar low ratios, while SCN and NCN had significantly higher ratios than OVN and marginally higher ratios than YVN ( $p=0.06$  and  $p=0.08$ , respectively; Fig. 5d, Supporting information).

The diversity of amino acids significantly differed among treatments ( $F_{3,28}=10.97$ ,  $p=6.152 \times 10^{-05}$ ). Both uncontaminated nectars (YVN=2.38 ± 0.29, OVN=1.80 ± 0.09) had a significantly lower diversity ( $p \leq 0.01$ , Supporting information) than contaminated nectars (SCN=4.50 ± 0.60, NCN=4.70 ± 0.66).  $\beta$ -Alanine was the most concentrated amino acid in both the uncontaminated (YVN and OVN) and in the NCN treatments, while it was the second most concentrated single amino acid in the SCN treatment (Supporting information, Fig. 6). Proline was the most and the second most concentrated amino acid in the SCN and NCN treatments, respectively. We discarded the arginine/threonine peak in the SCN treatment, since it was not possible to separate the two amino acids in the chromatograms (Supporting information, Fig. 6).

### Pollen and yeast detection

The concentration of pollen grains in the nectar samples significantly differed among flower treatments ( $F_{3,74}=26.55$ ,  $p=6.713 \times 10^{-09}$ ), and was significantly higher in the SCN and NCN than in the YVN and OVN treatments ( $p < 0.0001$  in all significant pairwise comparisons; Fig. 7a, Supporting information).

In a few samples of the uncontaminated nectars we found traces of pollen grains that did not belong to *G. lutea*, likely due to contamination during field manipulations. In the SCN treatment 78.5% of the pollen grains detected in nectar samples belonged to *G. lutea* (self-pollen), while in the




Figure 4. Glucose (a), fructose (b) and total sugar (c) concentration and glucose:fructose ratio (d) detected in the 78 nectar samples of the four flower treatments. Different letters indicate significant differences. YVN: young virgin nectar; OVN: old virgin nectar; SCN: self-contaminated nectar; NCN: naturally contaminated nectar. Note the different scales on the y-axes.

NCN treatment only 31.5% of pollen grains originated from *Gentiana lutea* (self- and conspecific cross-pollen). The concentration of yeast cells in the nectar samples significantly differed among treatments ( $F_{3,74} = 33.80$ ,  $p = 3.322 \times 10^{-13}$ ) and was the highest in the NCN treatment, followed by the SCN, OVN and YVN treatments ( $p < 0.01$  in all pairwise comparisons except OVN–YVN; Fig. 7b, Supporting information).

#### Invertase test

We did not find significant differences among flower treatments and time intervals in the sugar composition ( $\chi^2_{10} = 0.093$ ;  $p = 1$ ). The constant sugar composition through time highlights the absence of the enzyme invertase in the nectar of *G. lutea*, as sucrose was not hydrolysed into glucose and fructose (Supporting information).

#### Pollen suspension test

The total protein and non-protein amino acid concentrations increased considerably (more than tenfold and more than thirtyfold for PAAs and NPAAAs, respectively) after one hour from the addition of pollen grains of *G. lutea* to the sugar

solution mimicking the YVN composition. Total amino acid concentration continued to increase after four hours and after 24 h from the pollen suspension, reaching more than 40 times the initial amino acid concentration. The PAA:NPAA ratio was very high at all intervals after the pollen suspension, and increased through time (Fig. 8).

After one hour from pollen suspension, proline was the most concentrated amino acid (3243.6 nmol/100 mg), representing more than 25% of the total amino acid content. After four hours the concentration of proline decreased considerably (382.9 nmol/100 mg), while the overall amino acid spectrum showed minor variations. After 24 h only a few peaks were detectable, and cysteine represented more than 70% of the total amino acid content (Supporting information).

#### Discussion

We found clear differences in the chemical composition of flower nectar of *Gentiana lutea* subsp. *symphyandra* in relation to flower stage and access by floral visitors. Nectar composition was relatively constant in uncontaminated nectar, especially in floral buds, while it became more variable when nectar was exposed to the external environment and to






Figure 5. Concentration of (a) protein amino acids (PAA), (b) non-protein amino acids (NPAA), (c) total amino acids and (d) protein:non-protein amino acid ratio in the 32 nectar samples from the four flower treatments. Different letters indicate significant differences. YVN: young virgin nectar; OVN: old virgin nectar; SCN: self-contaminated nectar; NCN: naturally contaminated nectar. Note the different scales on the y-axes.

contamination by pollen grains and yeasts. Nectar sugar and amino acid concentration significantly increased with flower age and as a consequence of contamination by pollen grains. Contamination of nectar by pollen, either autonomously from flowers of the same plant or transported by floral visitors, enhanced nectar amino acid diversity and quantity. Nectar exposed to floral visitors showed a higher number of yeast cells than that of unvisited flowers, although yeasts likely did not significantly alter nectar composition because of their low overall density. These modifications are in turn expected to interact with floral visitors to shape plant-insect interactions.

### Nectar sugar composition

The sugar profile of *G. lutea* is dominated by glucose and fructose. The absence of sucrose was not due to the activity of an extracellular invertase but rather to the presence of cell wall-bound invertase such as the one that is required for nectar production in *Arabidopsis* (Ruhlmann et al. 2010) or to other cytoplasmic invertases. Our results showed that the concentrations of glucose, fructose and total sugars followed the same pattern and were significantly different among all

treatments. The freshly secreted and uncontaminated nectar presented the lowest sugar concentration, which markedly increased in older flowers. Nectar sugar concentration is often negatively correlated with flower age (Pacini and Nepi 2007), although opposite patterns can be observed (Petanidou et al. 1996, Nicolson and Nepi 2005, Pacini and Nepi 2007). Our results indicated that nectar sugar concentration in *G. lutea* significantly increases with flower age, regardless of external factors. In addition, although pollen only contains low amounts of sugars that could be released into nectar (Nakamura and Suzuki 1981, Pacini 1996, Pacini and Nepi 2007), the higher concentration of sugars found in nectar contaminated by self-pollen (SCN) than in old virgin nectar (OVN) suggests a significant contribution of sugars released by pollen.

The low sugar concentration in nectar of unbagged flowers (NCN) could be related to sugar consumption by yeasts whose presence was highest in this type of nectar. Lowering the sugar concentration is one of the most common impacts of yeasts on nectar quality (Pozo et al. 2015). However, the yeast cell concentration was very low, ranging between 0 and 481 yeast cells  $\mu\text{l}^{-1}$ , and only small changes in nectar chemistry have been detected when yeast concentration was lower




Figure 6. Mean concentration ( $\pm$  SE) of protein (PAA) and non-protein (NPAA) amino acids detected in the 32 nectar samples of the four nectar treatments. YVN: young virgin nectar; OVN: old virgin nectar; SCN: self-contaminated nectar; NCN: naturally contaminated nectar. \*: concentration of  $\beta$ -alanine (BALA) in the OVN treatment was  $4553 \pm 480.9$  nmol ml<sup>-1</sup>. ARG/THR corresponds to the sum of the two amino acid concentrations, because it was not possible to separate them in the chromatograms.

than  $10^4$  cells  $\mu\text{l}^{-1}$  (Hausmann et al. 2017). The low overall concentration of yeast cells can be related to the hexose-dominant nectar of *G. lutea*, as yeasts are more commonly found in sucrose-dominant nectars (Herrera et al. 2008, Pozo et al. 2015). More likely, the low sugar concentration observed can be related to consumption by floral visitors (Pacini and Nicolson 2007), as nectar removal can decrease further sugar secretion by the plant (Galletto and Bernardello 1992, Rivera et al. 1996, Pacini and Nepi 2007).

The glucose:fructose ratio was close to one in all treatments except in the unbagged flowers. The lower concentration of fructose and the subsequent higher glucose:fructose ratio in visited flowers can be explained by the presence of microorganisms. Although yeasts can significantly alter the proportion of sugars in nectar (Canto and Herrera 2012, Pozo et al. 2015, Borghi and Fernie 2017, Schaeffer et al. 2017), they do so by hydrolyzing sucrose into glucose and fructose (Herrera et al. 2008, De Vega and Herrera 2013, Pozo et al. 2015). Because sucrose is very scarce in nectar of *G. lutea*, yeasts are likely not the main cause of the G:F ratio increase observed in visited flowers (NCN treatment). We therefore hypothesise that the low fructose concentration found in visited flowers (NCN treatment) was caused

by some fructose-related bacteria, another abundant group of nectar-dwelling microorganisms (Vannette et al. 2013, Pozo et al. 2015).

Yeasts are frequently transported by floral visitors during their foraging bouts, either directly via their proboscis or indirectly through the transportation and involuntary release of pollen grains into nectar (Brysch-Herzberg 2004, Canto et al. 2008, Borghi and Fernie 2017, Hausmann et al. 2017). The lower concentration of conspecific pollen recorded in flowers open to visitors, compared to flowers where visitors were excluded, suggests that under natural conditions most of the pollen produced by *G. lutea* is removed by floral visitors, and that only a fraction contaminates its own nectar (< 31.5%). Although there was no difference in the total pollen content between self- (SCN) and naturally contaminated (NCN) nectar, results clearly indicate that most of the pollen found in the SCN treatment originated from *G. lutea* (> 75%), while almost 70% of the pollen in the NCN treatment belonged to different plant species. This suggests that the higher concentration of yeast cells found in the naturally contaminated nectar was mainly driven by floral visitors, either directly through contact with their proboscis or indirectly through contamination by foreign pollen.




Figure 7. Concentration of (a) pollen grains and (b) yeast cells found in the 72 nectar samples of the four flower treatments. Different letters indicate significant differences. YVN: young virgin nectar; OVN: old virgin nectar; SCN: self-contaminated nectar; NCN: naturally contaminated nectar. Note the different scales on the y-axes.

### Nectar amino acid composition

As for sugars, the freshly secreted and uncontaminated nectar had the lowest total amino acid concentration, which significantly increased in older flowers. This difference was mainly driven by a marked increase in non-protein amino acids. Nectar contaminated by self-pollen and by pollen from floral visitors had relatively high and comparable concentrations of amino acids. Moreover, the diversity of amino acids was significantly higher in the contaminated (SCN, NCN) than in the uncontaminated nectar (YVN, OVN). Our results clearly demonstrate a weak effect of yeast contamination and a strong effect of pollen contamination on amino acid diversity and abundance of floral nectar of *G. lutea*, because the total concentration and the profile of amino acids was similar in the SCN (self-pollen contaminated) and NCN (contaminated by pollen and yeasts) treatments. It has also been demonstrated that pollen contamination may increase yeast growth and play an important role in the population dynamics of nectar-dwelling yeasts (Poza and Jacquemyn 2019).




Figure 8. Concentration of total (yellow squares), protein (blue triangles) and non-protein (purple dots) amino acids (AA) at different time intervals after the suspension of pollen grains of *Gentiana lutea* in a solution mimicking nectar sugar composition of young virgin (i.e. uncontaminated) nectar (YVN). t0: after a few minutes, t + 1: after one hour, t + 4: after four hours and t + 24: after 24 h from pollen suspension.

$\beta$ -alanine was the most abundant non-protein amino acid in all nectar treatments, and one of the most abundant amino acids overall. Since  $\beta$ -alanine was detected at high concentrations in both uncontaminated nectar treatments, we assume that it is produced by the plant.  $\beta$ -alanine is the most common non-protein amino acid in nectar, along with GABA (Nepi 2014), and has important roles in the functioning and regulation of plant physiology and metabolism. It mainly functions as a precursor of Coenzyme A and protects plants from several external stresses (Parthasarathy et al. 2019). Recently, it has been suggested that  $\beta$ -alanine may influence plant-pollinator relationships by affecting insects' nervous systems and stimulating the flight muscle function (Nepi 2014, Felicioli et al. 2018, Bogó et al. 2019). The high concentration of  $\beta$ -alanine produced by the nectar of *G. lutea* could therefore play an important role in the attraction of floral visitors.

The protein:non-protein amino acid ratio was higher in nectars contaminated by self-pollen and by pollen from floral visitors than in uncontaminated nectars. The significant amount of pollen grains observed in contaminated nectars is the most likely source of this increase. Although we only performed a single pollen supplementation test, these preliminary results clearly showed an increase of the total amino acids and in particular of protein amino acids. This was supported by the highest total and protein amino acid concentrations found in the self-contaminated nectar treatment (SCN), while the old uncontaminated nectar (OVN) presented a very low total amino acid concentration and protein amino acids were almost absent. The effect of pollen contamination on nectar amino acid concentration and composition needs further systematic examination, since few direct tests have been performed, and these have yielded contrasting results (Gottsberger et al. 1990, Nicolson 2007).

0 Both contaminated nectars presented a high concentra-  
tion of proline, which was absent in uncontaminated nectars.  
Free proline accumulates in extremely high concentrations  
in the pollen of several species (Pálfi et al. 1987, Chiang and  
5 Dandekar 1995), reaching up to 70% of free amino acids  
(Hong-qi et al. 1982), and is usually the primary amino acid  
released in nectar-like solutions (Erhardt and Baker 1990,  
Gottsberger et al. 1990, O'Brien et al. 2003). It is therefore  
likely that the increase of proline content resulted from the  
10 large amount of pollen grains observed in contaminated nec-  
tars, originating either from within-plant pollen transfer or  
from pollen transported by floral visitors.

Nectar contamination can affect the activity of floral visi-  
tors. High proline concentration can increase nectar attractive-  
15 ness towards floral visitors by stimulating insects' salt cells and  
consequently enhancing the intensity of feeding behaviour  
(Hansen et al. 1998, Wacht et al. 2000). Moreover, proline  
is degraded during the initial stages or the lift phase of insect  
flight (Micheu et al. 2000), acting as an efficient short-term  
20 fuel – more efficient than sugar – that results in quick bursts  
of energy production (Carter et al. 2006, Teulier et al. 2016).

In a previous study, Rossi et al. (2014) found similar con-  
centrations of proline in different populations and subspe-  
cies of *G. lutea*, and hypothesised an endogenous production  
of proline rather than a contamination by exogenous pollen.  
25 However, our results indicate that pollen is the main source  
of proline in flower nectar of *G. lutea* subsp. *symphyandra*.  
Since all subspecies of *G. lutea* have generalist pollination  
and show similar pollinator communities (Rossi et al. 2014),  
30 we can expect that the similar patterns observed among nec-  
tars of different subspecies are driven by the same mechanism of  
contamination by floral visitors.

35

## Conclusions

Nectar is the main interface between the plant and its floral  
visitors, which use it as the main source of energy. Our results  
40 show that the floral nectar implicated in this process is not the  
pure uncontaminated nectar secreted by the plant, but rather  
a complex intertwining of numerous endogenous and exog-  
enous factors, partially mediated by visitors themselves. We  
45 found that nectar sugar and amino acid composition depends  
on both flower development and on nectar contamination by  
pollen and microorganisms. While the main impact of micro-  
organism presence is the shift of glucose:fructose ratio, pollen  
contamination mainly increases the amino acid concentra-  
50 tion and diversity. By transferring pollen grains and microor-  
ganisms to flowers while foraging, floral visitors appear to be  
the main indirect cause of major modifications of the nectar  
chemical composition. In turn, all these modifications are  
55 likely to modify the attractiveness of flowers towards floral  
visitors, in a complex process of ecological relationships.

60 *Acknowledgements* – We thank 'Ente di gestione per i Parchi e la  
Biodiversità – Emilia Orientale' for permitting field investigations,

Mariachiara Zanichelli and Natasha de Manincor for help with field  
61 work and Marcelo Aizen for feedback during the writing of this  
manuscript. Joan Rabassa-Juvanteny was supported by an Erasmus+  
traineeship grant by the European Commission.  
65 *Conflict of interest* – The authors declare that they have no conflict  
of interest.

## Author contributions

**Gherardo Bogo** and **Alessandro Fisogni** contributed equally  
70 to this publication. **Gherardo Bogo**: Formal analysis (equal);  
Visualization (equal); Writing – original draft (equal);  
Writing – review and editing (equal). **Alessandro Fisogni**:  
Formal analysis (equal); Investigation (equal); Visualization  
75 (equal); Writing – original draft (equal); Writing – review  
and editing (equal). **Joan Rabassa-Juvanteny**: Investigation  
(equal); Writing – review and editing (equal). **Laura  
Bortolotti**: Investigation (equal); Writing – review and  
80 editing (equal). **Massimo Nepi**: Conceptualization (equal);  
Methodology (equal); Writing – review and editing (equal).  
**Massimo Guarnieri**: Investigation (equal); Writing – review  
and editing (equal). **Lucia Conte**: Project administra-  
tion (equal); Writing – review and editing (equal). **Marta  
85 Galloni**: Conceptualization (equal); Investigation (equal);  
Methodology (equal); Project administration (equal);  
Supervision (equal); Writing – review and editing (equal).

## Data availability statement

Data are available from the Zenodo Digital Repository:  
95 <<http://doi.org/10.5281/zenodo.4265415>>.

## References

- Adler, L. S. 2000. The ecological significance of toxic nectar. –  
100 *Oikos* 91: 409–420.
- Aizen, M. A. and Raffaele, E. 1996. Nectar production and pollina-  
tion in *Alstroemeria aurea*: responses to level and pattern of  
105 flowering shoot defoliation. – *Oikos* 76: 312.
- Alm, J. et al. 1990. Preference of cabbage white butterflies and  
honey bees for nectar that contains amino acids. – *Oecologia*  
84: 53–57.
- Baker, H. G. and Baker, I. 1982. Chemical constituents of nectar  
in relation to pollination mechanisms and phylogeny. – In:  
110 Nitecki, M. (ed.), *Biochemical aspects of evolutionary biology*.  
Univ. of Chicago Press, pp. 131–171.
- Baker, H. G. and Baker, I. 1983. A brief historical review of the  
chemistry of floral nectar. – In: Bentley, B. and Elias, T. (eds),  
115 *The biology of nectaries*. Columbia Univ. Press, pp. 126–152.
- Baker, H. G. and Baker, I. 1990. The predictive value of nectar  
chemistry to the recognition of pollinator types. – *Isr. J. Bot.*  
39: 157–166.
- Baker, H. G. et al. 1998. Sugar composition of nectars and fruits  
consumed by birds and bats in the tropics and subtropics. –  
121 *Biotropica* 30: 559–586.



- 0 Bates, D. et al. 2015. Fitting linear mixed-effects models using lme4. – J. Stat. Softw. 67: 1–48.
- Baude, M. et al. 2011. Litter inputs and plant interactions affect nectar sugar content. – J. Ecol. 99: 828–837.
- 5 Bertazzini, M. et al. 2010. Amino acid content and nectar choice by forager honeybees (*Apis mellifera* L.). – Amino Acids 39: 315–318.
- Bogo, G. et al. 2019. Effects of non-protein amino acids in nectar on bee survival and behavior. – J. Chem. Ecol. 45: 278–285.
- 10 Borghi, M. and Fernie, A. R. 2017. Floral metabolism of sugars and amino acids: implications for pollinators' preferences and seed and fruit set. – Plant Physiol. 175: 1510–1524.
- Brysch-Herzberg, M. 2004. Ecology of yeasts in plant–bumblebee mutualism in central Europe. – FEMS Microbiol. Ecol. 50: 87–100.
- 15 Canto, A. and Herrera, C. M. 2012. Micro-organisms behind the pollination scenes: microbial imprint on floral nectar sugar variation in a tropical plant community. – Ann. Bot. 110: 1173–1183.
- Canto, A. et al. 2008. Pollinator foraging modifies nectar sugar composition in *Helleborus foetidus* (Ranunculaceae): an experimental test. – Am. J. Bot. 95: 315–320.
- 20 Carter, C. et al. 2006. A novel role for proline in plant floral nectars. – Naturwissenschaften 93: 72–79.
- Chalcoff, V. R. et al. 2017. Pollinator type and secondarily climate are related to nectar sugar composition across the angiosperms. – Evol. Ecol. 31: 585–602.
- 25 Chappell, C. R. and Fukami, T. 2018. Nectar yeasts: a natural microcosm for ecology. – Yeast 35: 417–423.
- Chiang, H.-H and Dandekar, A. M. 1995. Regulation of proline accumulation in *Arabidopsis thaliana* (L.) Heynh during development and in response to desiccation. – Plant. Cell Environ. 18: 1280–1290.
- 30 Cohen, S. A. and Michaud, D. P. 1993. Synthesis of a fluorescent derivatizing reagent, 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate, and its application for the analysis of hydrolysate amino acids via high-performance liquid chromatography. – Anal. Biochem. 211: 279–287.
- 35 De Vega, C. and Herrera, C. M. 2013. Microorganisms transported by ants induce changes in floral nectar composition of an ant-pollinated plant. – Am. J. Bot. 100: 792–800.
- 40 Dray, S. and Dufour, A. B. 2007. The ade4 package: implementing the duality diagram for ecologists. – J. Stat. Softw. 22: 1–20.
- Erhardt, A. and Baker, I. 1990. Pollen amino acids – an additional diet for a nectar feeding butterfly? – Plant Syst. Evol. 169: 111–121.
- 45 Erhardt, A. and Rusterholz, H. P. 1998. Do peacock butterflies (*Inachis io* L.) detect and prefer nectar amino acids and other nitrogenous compounds? – Oecologia 117: 536–542.
- Felicioli, A. et al. 2018. Effects of nonprotein amino acids on survival and locomotion of *Osmia bicornis*. – Insect Mol. Biol. 27: 556–563.
- 50 Fenster, C. B. et al. 2004. Pollination syndromes and floral specialization. – Annu. Rev. Ecol. Evol. Syst. 35: 375–403.
- Galetto, L. and Bernardello, L. M. 1992. Nectar secretion pattern and removal effects in six Argentinean Pitcairnioideae (Bromeliaceae). – Bot. Acta 105: 292–299.
- 55 Gottsberger, G. et al. 1990. Variation in floral nectar amino acids with aging of flowers, pollen contamination and flower damage. – Isr. J. Bot. 39: 167–176.
- Hansen, K. et al. 1998. New aspects of chemoreception in flies. – Ann. N. Y. Acad. Sci. 855: 143–147.
- 60 Hausmann, S. L. et al. 2017. Solving the puzzle of yeast survival in ephemeral nectar systems: exponential growth is not enough. – FEMS Microbiol. Ecol. 93: 1–11.
- Heil, M. 2011. Nectar: generation, regulation and ecological functions. – Trends Plant Sci. 16: 191–200. 65
- Herrera, C. M. et al. 2008. Invisible floral larcenies: microbial communities degrade floral nectar of bumble bee-pollinated plants. – Ecology 89: 2369–2376.
- Herrera, C. M. et al. 2009. Yeasts in floral nectar: a quantitative survey. – Ann. Bot. 103: 1415–1423. 70
- Hong-qi, Z. et al. 1982. Protein synthesis in germinating pollen of *Petunia*: role of proline. – Planta 154: 199–203.
- Inouye, D. W. and Waller, G. D. 1984. Responses of honey bees (*Apis mellifera*) to amino acid solutions mimicking floral nectars. – Ecology 65: 618–625. 75
- Jost, L. 2006. Entropy and diversity. – Oikos 113: 363–375.
- Kress, W. J. 1985. Bat pollination of an old world *Heliconia*. – Biotropica 17: 302.
- Lasso, E. and Naranjo, M. E. 2003. Effect of pollinators and nectar robbers on nectar production and pollen deposition in *Hamelia patens* (rubiceae). – Biotropica 35: 57–66. 80
- Lenth, R. et al. 2020. Estimated marginal means, aka least-squares means. – R Packag. ver. 1.5.1.
- Micheu, S. et al. 2000. Importance of proline and other amino acids during honeybee flight (*Apis mellifera carnica* Pollmann). – Amino Acids 18: 157–175. 85
- Mittelbach, M. et al. 2015. Nectar sugars and bird visitation define a floral niche for basidiomycetous yeast on the Canary Islands. – BMC Ecol. 15: 1–13.
- Mustard, J. A. 2020. Neuroactive nectar: compounds in nectar that interact with neurons. – Arthropod. Plant. Interact. 14: 151–159. 90
- Nakamura, N. and Suzuki, H. 1981. Sugar composition of pollen grain and pollen tube cell walls. – Phytochemistry 20: 981–984.
- Nepi, M. 2014. Beyond nectar sweetness: the hidden ecological role of non-protein amino acids in nectar. – J. Ecol. 102: 108–115. 95
- Nepi, M. and Stpiczynska, M. 2008. The complexity of nectar: secretion and resorption dynamically regulate nectar features. – Naturwissenschaften 95: 177–184.
- Nepi, M. et al. 2010. Variation in nectar-sugar profile of *Anchusa* and allied genera (Boraginaceae). – Bot. J. Linn. Soc. 162: 616–627. 100
- Nepi, M. et al. 2018. Nectar in plant–insect mutualistic relationships: from food reward to partner manipulation. – Front. Plant Sci. 9: 1063.
- Nicolson, S. W. 2007. Amino acid concentrations in the nectars of southern African bird-pollinated flowers, especially *Aloe* and *Erythrina*. – J. Chem. Ecol. 33: 1707–1720. 105
- Nicolson, S. W. and Nepi, M. 2005. Dilute nectar in dry atmospheres: nectar secretion patterns in *Aloe castanea* (Asphodelaceae). – Int. J. Plant Sci. 166: 227–233. 110
- Nicolson, S. W. and Thornburg, R. W. 2007. Nectar chemistry. – In: Nicolson, S. W. et al. (eds), Nectaries and nectar. Springer, pp. 215–264.
- Nocentini, D. et al. 2012. Flower morphology, nectar traits and pollinators of *Cerinthe major* (Boraginaceae–Lithospermeae). – Flora 207: 186–196. 115
- O'Brien, D. M. et al. 2003. Pollen feeding in the butterfly *Heliconius charitonia*: isotopic evidence for essential amino acid transfer from pollen to eggs. – Proc. R. Soc. B 270: 2631–2636.
- Pacini, E. 1996. Types and meaning of pollen carbohydrate reserves. – Sex. Plant Reprod. 9: 362–366. 121

0	Pacini, E. and Nepi, M. 2007. Nectar production and presentation. – In: Nicolson, S. W. et al. (eds), Nectaries and nectar. Springer, pp. 167–214.	61
	Pacini, E. and Nicolson, S. W. 2007. Introduction. – In: Nicolson, S. W. et al. (eds), Nectaries and nectar. Springer, pp. 1–18.	
5	Pálfi, G. et al. 1987. Concentration of proline in pollens: a new taxonomical index. – Acta Biol. 33: 25–34.	65
	Parachnowitsch, A. L. et al. 2019. Evolutionary ecology of nectar. – Ann. Bot. 123: 247–261.	
	Parthasarathy, A. et al. 2019. The synthesis and role of $\beta$ -alanine in plants. – Front. Plant Sci. 10: 921.	70
10	Petanidou, T. et al. 1996. Change in floral nectar components from fresh to senescent flowers of <i>Capparis spinosa</i> (Capparidaceae), a nocturnally flowering Mediterranean shrub. – Plant Syst. Evol. 199: 79–92.	
15	Potter, C. F. and Bertin, R. I. 1988. Amino acids in artificial nectar: feeding preferences of the flesh fly <i>Sarcophaga bullata</i> . – Am. Midl. Nat. 120: 156.	75
	Pozo, M. I. and Jacquemyn, H. 2019. Addition of pollen increases growth of nectar-living yeasts. – FEMS Microbiol. Lett. 366: fnz191.	
20	Pozo, M. I. et al. 2015. Impact of microorganisms on nectar chemistry, pollinator attraction and plant fitness. – In: Peck, R. L. (ed.), Nectar: production, chemical composition and benefits to animals and plants. Nova Science Publishers, pp. 1–40.	
25	Pyke, G. H. 2016. Floral nectar: pollinator attraction or manipulation? – Trends Ecol. Evol. 31: 339–341.	85
	Rivera, G. L. et al. 1996. Nectar secretion pattern, removal effects and breeding system of <i>Ligaria cuneifolia</i> (Loranthaceae). – Can. J. Bot. 74: 1996–2001.	
30	Rossi, M. 2012. Taxonomy, phylogeny and reproductive ecology of <i>Gentiana lutea</i> L. – PhD thesis, Univ. di Bologna, Italy.	90
	Rossi, M. et al. 2014. Bouncy versus idles: on the different role of pollinators in the generalist <i>Gentiana lutea</i> L. – Flora 209: 164–171.	95
35		
	Rossi, M. et al. 2016. The effect of pollination mode on seed performance of <i>Gentiana lutea</i> : a laboratory evaluation of seed germinability. – Nord. J. Bot. 34: 761–768.	100
	Ruhlmann, J. M. et al. 2010. Cell wall invertase 4 is required for nectar production in <i>Arabidopsis</i> . – J. Exp. Bot. 61: 395–404.	105
	Schaeffer, R. N. et al. 2017. Consequences of a nectar yeast for pollinator preference and performance. – Funct. Ecol. 31: 613–621.	
	Stevenson, P. C. et al. 2017. Plant secondary metabolites in nectar: impacts on pollinators and ecological functions. – Funct. Ecol. 31: 65–75.	110
	Teulier, L. et al. 2016. Proline as a fuel for insect flight: enhancing carbohydrate oxidation in hymenopterans. – Proc. R. Soc. B 283: 20160333.	115
	Tutin, T. G. et al. 1972. Flora Europea, Vol. 3. – Cambridge Univ. Press.	
	Vannette, R. L. and Fukami, T. 2018. Contrasting effects of yeasts and bacteria on floral nectar traits. – Ann. Bot. 121: 1343–1349.	
	Vannette, R. L. et al. 2013. Nectar bacteria, but not yeast, weaken a plant–pollinator mutualism. – Proc. R. Soc. B in press.	
	Venables, W. N. and Ripley, B. D. 2002. Modern applied statistics with S, 4th edn. – Springer.	
	Wacht, S. et al. 2000. Chemosensory control of pollen ingestion in the hoverfly <i>Eristalis tenax</i> by labellar taste hairs. – J. Comp. Physiol. A Sens. Neural Behav. Physiol. 186: 193–203.	
	Wickham, H. 2016. ggplot2: elegant graphics for data analysis. – Springer.	
	Willmer, P. G. 1980. The effects of insect visitors on nectar constituents in temperate plants. – Oecologia 47: 270–277.	
	Yang, M. et al. 2019. Nectar yeasts enhance the interaction between <i>Clematis akebioides</i> and its bumblebee pollinator. – Plant Biol. 21: 732–737.	
40		
45		
50		
55		
60		

