

# Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Health Risk Assessment of Potentially Toxic Elements, Persistence of NDL-PCB, PAHs, and Microplastics in the Translocated Edible Freshwater Sinotaia quadrata (Gasteropoda, Viviparidae): A Case Study from the Arno River Basin (Central Italy)

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Pastorino, P., Nocita, A., Ciccotelli, V., Zaccaroni, A., Anselmi, S., Giugliano, R., et al. (2021). Health Risk Assessment of Potentially Toxic Elements, Persistence of NDL-PCB, PAHs, and Microplastics in the Translocated Edible Freshwater Sinotaia quadrata (Gasteropoda, Viviparidae): A Case Study from the Arno River Basin (Central Italy). EXPOSURE AND HEALTH, 13(4), 583-596 [10.1007/s12403-021-00404-w].

*Availability:*

[This version is available at: https://hdl.handle.net/11585/840451 since: 2021-12-02](https://hdl.handle.net/11585/840451)

*Published:*

[DOI: http://doi.org/10.1007/s12403-021-00404-w](http://doi.org/10.1007/s12403-021-00404-w)

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

> This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

# **Health Risk Assessment of Potentially Toxic Elements, Persistence of NDL‑PCB, PAHs, and Microplastics in the Translocated Edible Freshwater** *Sinotaia quadrata* **(Gasteropoda, Viviparidae): A Case Study from the Arno River Basin (Central Italy)**

Paolo Pastorino<sup>1\*</sup>, Annamaria Nocita<sup>2</sup>, Valentina Ciccotelli<sup>1</sup>, Annalisa Zaccaroni<sup>3</sup>, Serena Anselmi<sup>4</sup>, Roberta Giugliano<sup>1</sup>, Mattia Tomasoni<sup>1</sup>, Marina Silvi<sup>3</sup>, Vasco Menconi<sup>1</sup>, Barbara Vivaldi<sup>1</sup>, Elisabetta Pizzul<sup>5</sup>, Monia Renzi<sup>5</sup>, Marino Prearo<sup>1</sup>

<sup>1</sup> The Veterinary Medical Research Institute for Piemonte, Liguria and Valle D'Aosta, via Bologna 148, 10154 Torino, Italy

- <sup>2</sup> Department of Biology, University of Firenze, Via Madonna del Piano 6, 50019 Sesto Fiorentino FI, Italy
- <sup>3</sup> Department of Veterinary Medical Science, University of Bologna, viale Vespucci 2, 47042 Cesenatico, FC, Italy
- <sup>4</sup> Bioscience Research Center, via Aurelia Vecchia 32, 58015 Orbetello, GR, Italy

<sup>5</sup> Department of Life Sciences, University of Trieste, via L. Giorgieri 10, 34127 Trieste, Italy

\* Paolo Pastorino: paolo.pastorino@izsto.it

# **Abstract**

With this study we investigated the accumulation of potentially toxic elements (As, Cd, Cr, Cu, Hg, Pb, Zn), six indicators (28, 52, 101, 138, 153, 180) of non-dioxin-like polychlorinated biphenyls ( $\Sigma_6$  NDL-PCBs), polycyclic aromatic hydrocarbons (PAHs), and microplastics in *S. quadrata* (edible part) collected from two sampling sites (1 and 2) from the Arno River Basin (Central Italy). A risk assessment of the implications for human health was also performed. Levels of potentially toxic elements in gastropods from site 2 were slightly higher and the  $\Sigma_6$  NDL-PCB concentration was significantly higher (7.32 ng g−1 *vs*. 3.07 ng g −1) compared to site 1 due to higher anthropogenic pressures. The concentration of chrysene, benzo[b]fluoranthene, and benzo(a)pyrene was below the limit of quantification (0.5 μg kg<sup>-1</sup>). Benzo[a]anthracene was detected in gastropods from both sites ( $0.5 \pm 0.02 \,\mu$ g kg<sup>-1</sup> and  $0.7 \pm 0.02$  $\mu$ g kg<sup>-1</sup> from site 1 and 2, respectively). The microplastics frequency (mainly polyethylene terephthalate) differed significantly between the sites (site  $1, 0.8 \pm 1.30$ ; site  $2, 1 \pm 0.37$  items/specimen). All contaminant levels were compliant with international regulatory limits and guidelines. Incremental lifetime cancer risk (ILCR) values for As, Cd, Cr, and Pb were far below the safety values of  $1 \times 10^{-4}$ . Similarly, the ILCR values from the Monte Carlo simulation model were all within the safety region of  $1 \times 10^{-4}$  and  $1 \times 10^{-6}$ . Findings from the health risk assessment indicated no adverse effects for human health from any of the contaminants analysed here, except for microplastics for which no limits or legislation are currently in force.

**Keywords** River Arno, Benzo[a]anthracene, Environmental monitoring, Polyethylene terephthalate, Risk assessment

#### **Introduction**

Evaluation of polluting compounds and knowledge of the ecological condition of an ecosystem are requisite for management and prevention of health risks to humans and for environmental protection (Cunningham et al. 2009). Biomonitoring involves the use of organisms to evaluate changes in water or air quality and measure the bioavailable fraction of the compound and its fate along the food web (McIntyre and Beauchamp 2007; Needham et al. 2007).

Among freshwater macroinvertebrates, gastropods are benthic organisms that colonize a wide variety of aquatic environments (Strong et al. 2007). Found in typically lotic and lentic environments, many species have been translocated and become an invasive species in non-native environments (Cianfanelli et al. 2017). The geographical area comprising Florence, Prato, and Pistoia (Tuscany, central Italy) is home to a large Asian community of more than 50,000 inhabitants (over 8% of which are Chinese), ranking second to the Milan metropolitan area (north Italy) population size (ISTAT 2019). In keeping with their traditional cuisine, they have introduced animal and plant species to urban non-native environments outside China. The freshwater gastropod *Sinotaia quadrata* (Caenogastropoda, Viviparidae) is widely used in Chinese cuisine. Its recent discovery in the Arno river basin is the first recorded occurrence in Europe (Cianfanelli et al. 2017). The potential creation of an extensive mollusk farm using the Arno as a natural habitat for *S. quadrata* for human consumption raises concern about the threat to biodiversity and to human health from consumption of this alien species (Cianfanelli et al. 2017).

Cianfanelli et al. (2017) reported evidence for a wellstructured population of *S. quadrata* in sites on the Arno river (downstream of Florence), where the water is rich in organic substances and nutrients that facilitate the action of tolerant suspension feeders like *S. quadrata*. The main anthropogenic inputs to the Arno occur downstream of Florence and derive from direct civil and industrial wastewater discharges. The River Arno is the longest in Tuscany (about 245 km length, catchment area about 8228  $km<sup>2</sup>$ ) and is ranked fifth among the largest basins in Italy. It rises in the northern Apennines and empties into the Tyrrhenian Sea near Pisa. It is polluted by industrial and untreated civil wastewaters from Florence and other cities along the river (Arezzo, upper stretch of the river; Pistoia and Prato north of Florence). Adding to the potential threat to the environment are the wastewaters from industrial zones and nurseries around Pistoia and Prato carried by its tributaries, the Bisenzio and the Ombrone.

The bioaccumulation of contaminants by *S. quadrata* may pose a potential health risk to humans: potentially toxic elements, persistent and accumulative substances such as polychlorinated biphenyls (PCBs) and polycyclic aromatic hydrocarbons (PAHs), plus emerging contaminants such as microplastics can accumulate in aquatic organisms, including gastropods.

Metal contamination is a major environmental concern in aquatic ecosystems because the elements tend to accumulate in the body of organisms at concentrations higher than those found in abiotic compartments (Jia et al. 2017; Pastorino et al. 2019). Some have nutritional functions and are essential for life (copper and zinc) and for maintenance of animal (Esposito et al. 2018), even if become toxic at higher concentrations (Yuan et al. 2017; Pastorino et al. 2020a). Others still, like lead, cadmium, mercury, and arsenic, have no biological function and are extremely toxic even at low concentration (Halder et al. 2020).

PCBs comprise a series of 209 aromatic compounds made up of variously chlorinated biphenyl molecules synthesized at the beginning of the last century and produced commercially since 1930 (Howell et al. 2008); they are currently largely banned due to their toxicity and tendency to bioaccumulate (Oziolor et al. 2018). Only 12 of the 209 PCBs congeners have chemical-physical and toxicological characteristics comparable to dioxins and furans: these are defined as dioxin-like PCBs (DL-PCBs). The sum of the six congeners termed indicators (28, 52, 101, 138, 153, 180) includes about half of all non-dioxin-like PCBs (NDLPCBs) present in feed and food products (Malisch and Kotz 2014). The European Food Safety Authority (EFSA) stated that the sum of these PCBs constitutes an adequate indicator of the occurrence and human exposure to NDL-PCBs since they are analytically predominant in environmental matrices (EFSA 2005; Squadrone et al. 2013).

Polycyclic aromatic hydrocarbons are organic compounds formed by two or more aromatic rings that derive mainly from the incomplete combustion of organic matter (e.g., coal, petroleum derivatives, oil or biomass) (Marzooghi and Di Toro 2017; Wang and Wang 2018; Molsen et al. 2019). Because of their hydrophobicity and long-range transportation, PAHs are widely distributed and can accumulate in diverse environmental compartments, including living organisms (Honda and Suzuki 2020). Generally, PAHs are substances of toxicological interest as some are considered probable or possible carcinogens. The most extensively studied component is benzo(a)pyrene (B[a]P) widespread in the environment at significant concentrations and extremely toxic (Zhang et al. 2019); the PAH4 index (the sum of benzo[a]anthracene-B[a]A, chrysene-Chr, benzo[b]fluoranthene-B[b]FL, and B[a]P) was recognized as an indicator of cancer risk due to dietary exposure to PAHs (EFSA 2008).

Another major environmental issue is plastic pollution because ubiquitous in marine and freshwater ecosystems worldwide (Borrelle et al. 2020). An emergent threat to wildlife and human health is the formation of microplastics from the breakdown of large plastic pieces into tiny particles (< 5 mm) (Mitrano and Wohlleben 2020). Human health and well-being are intimately linked to environmental quality (USEPA, 2021). Supra-optimal levels of pollutants can induce or increase the risk of developing diseases of the respiratory, reproductive, and immune system and cancer in humans (Shahid et al. 2020).

To our best knowledge, there are no investigations into the occurrence of contaminants in *S. quadrata* from the Arno River basin. Furthermore, a risk assessment of implications for human health by their consumption is warranted. The aims of the present study were to: (a) measure the accumulation of potentially toxic elements (As, Cd, Cr, Cu, Hg, Pb, Zn), PAHs (B[a]A, Chr, B[b]FL, and B[a]P), NDL-PCBs (28, 52, 101, 138, 153, 180), and microplastics in tissue samples of *S. quadrata* collected from two sites on the Arno; (b) determine whether contaminant levels are compliant with regulatory limits and guidelines; (c) assess the possible human health risks associated with gastropod consumption.

#### **Material and Methods**

#### **Study Area**

Two sampling sites were selected according to their accessibility and the presence/abundance of *Sinotaia quadrata* (Cianfanelli et al. 2017). Site 1 (Samminiatello; 1663666X 4845701Y) and site 2 (Fibbiana; 1660695X 4844688Y) are both located in the Municipality of Montelupo Fiorentino

(Florence Province) (Fig. 1) where the riverbed is largely disconnected from the perifluvial belt, as it is intensely urbanized. The sites are located near the River Arno Valdarno Inferiore Capraia e Limite (code MAS-108) stations, which is monitored by the Regional Environmental Protection Agency of Tuscany (ARPAT) for assessment of the ecological status sensu Water Framework Directive. At the station site, the River Arno receives the Ombrone Pistoiese and Bisenzio tributaries on the right. The river water quality begins to decrease due to inflow of wastewater from the Baciacavallo treatment plant (Prato Province), which serves the towns of this textile manufacturing area. This stretch of the Arno is noted for its low quality: the ecological status of MAS-108 is "poor" according to the last report (ARPAT 2021).

# **Sampling and Processing of** *Sinotaia quadrata*

Fifty specimens homogeneous in size  $(33.9 \pm 0.12 \text{ mm}$  in height and  $19.4 \pm 0.21 \text{ mm}$  in diameter) were collected by hand at each site directly from stones, pebbles, and riverbed mud (30–100 cm in depth). Samples were placed in precleaned glass beakers to prevent contamination, covered with aluminum foil, and transported refrigerated  $(+ 4 \degree C)$  to the laboratory within a few hours after collection and immediately frozen (−20 °C) whole (in their shell) until chemical analyses.

# **Detection of Potentially Toxic Elements**

Determination of As, Cd, Cr, Cu, Hg, Pb, and Zn was performed on the whole soft body of fifteen *S. quadrata* per site by inductively coupled plasma-optic emission spectrometry (ICP-OES) on a Perkin Elmer Optima 2100 DV instrument (PerkinElmer, Inc., Shelton, CT, USA), coupled with a CETAC U5000AT + ultrasound nebulizer (Cetac Technologies, Inc., Omaha, NE, USA) for mercury, as previously reported by Pastorino et al. (2020b). Briefly, samples were homogenized and microwavedigested in a Milestone ETHOS ONE oven using 4 mL H NO<sub>3</sub> and 1 mL H  $_2O_2$ . All reagents were from Merck (Darmstadt, Germany); acids were of Suprapur grade. Results are presented as µg g<sup>-1</sup>wet weight (w.w.). Analytical performance was determined by processing certified reference materials (HISS-1) in six replicates for corresponding elements, along with blank reagents in each analytical run. All reference material values were within certified limits. Table S1 presents the limit of detection (LOD), the reference material values, and the percentages of recovery.

# **PAH and PCBs Analyses**

Fifteen specimens of *S. quadrata* per site underwent PAHs analysis. Determination of benzo[a]anthracene-B[a]A, chrysene-Chr, benzo[b]fluoranthene-B[b]FL, and B[a]P) was performed using HPLC-FLD. Fifteen specimens of *S. quadrata* per site underwent NDL-PCBs (28, 52, 101, 138, 153, 180) analysis by GC–MS.

The HPLC system consisted of a 1100 Series quaternary pump (Agilent Technologies, Santa Clara, CA, USA). Data acquisition and analysis were performed using ChemStation software. The  $\lambda_{ex}/\lambda_{em}$  for the fluorescence was  $294/404$  nm; the gain was  $12x$ ; the injection volume 1 mL; the stop time at 36 min; and the column temperature 25 °C. Chromatographic separation was performed using a reversed-phase column Envirosep pp (125 m  $\times$  4.6 mm, i.d. 5 µm). The eluents were: water (A) and acetonitrile (B). The flow rate was 1 mL min<sup>-1</sup> and the program was 80% B for 15 min, then 100% B for 10.0 min, and then held for 10 min.

GC–MS analyses were performed on a gas-chromatograph coupled with single quadrupole GC DSQ 70 FOCUS (Thermo Fisher Scientific, Waltham, MA, USA), equipped with an AS 3000 autosampler. Xalibur software was for mass spectrometer control, data acquisition and analysis. The GC column was a DB-5MS (30 m  $\times$  0.25 mm, 0.25 µm), the working conditions were: drying gas He (purity  $> 99.9\%$ ) at 1.2 mL min−1; EI voltage 70 eV; injector temperature 250 °C; splitless mode; split flow 50 mL min−1; gas saver flow 10 mL min−1 (5 min); injection volume 1 µL. The oven temperature program was started at 100 °C for 1 min, then increased by 20 °C min−1 to 190 °C, and held for 2 min, increased by 3 °C min−1 to 250 °C and by 50 °C min−1 to 300 °C, and held for 20 min. The mass spectrometer operated in EI ionization in positive mode, the MS transfer line was 270 °C. Table S2 presents the transitions used for detection.

*PAHs analysis*: 2 g of sample were saponified with 10 mL of a solution of KOH 2 N in EtOH for 2 h and extracted three times with 20 mL of cyclohexane. The organic phase was dried in a rotavapor at 60 °C and added with 4 mL of acetonitrile. The samples were purified with SPE columns and eluted with 12 mL of acetonitrile. The organic phase was dried at  $60 \pm 5$  °C, reconstituted in 1 mL of acetonitrile, and injected into the HPLC system. Analytical blanks and two spiked samples at a concentration of 2  $\mu$ g kg<sup>-1</sup> for all analytes were run in the same way as the samples; concentrations were determined using a calibration curve prepared in acetonitrile using PAH-Mix 9 standard (Dr. Ehrenstorfer GmbH, Germany) at 1 ng mL<sup>-1</sup>, 2 ng mL<sup>-1</sup>, 5 ng mL<sup>-1</sup>, 10 ng mL<sup>-1</sup>, 20 ng mL<sup>-1</sup>, 50 ng mL<sup>-1</sup>, 100 ng mL<sup>-1</sup>, 200 ng mL<sup>-1</sup>. The recovery sample results ranged between 81% and 90%. The LOQ for all analytes was 0.5  $μ$ g Kg $^{-1}$ .

*NDL-PCBs analysis*: 5 g of each sample was added with 20 µL of internal standard (PCB No.198 and PCB No.155, Dr. Ehrenstorfer GmbH). The PCBs were extracted from fat tissue using a hexane/acetone mixture (50/50 v/v) and shaken overnight. For the clean-up, an Extrelut-SPE cartage system was built by putting Extrelut between a glass flask and a SPE cartage; 3 mL of sulfuric acid was maintained in contact with the fat for 10 min in the Extrelut cartage. The sample was eluted with 13 mL of n-hexane. The organic phase was dried at  $60 \pm 5$  °C, added with 0.1 mL of iso-octane, and injected into the GC system. Quantification was based on peak areas and relative response factors derived from calibration curves constructed using PCB-Mix, PCB No.198 and PCB No.155 (Dr. Ehrenstorfer GmbH) at 5 ng  $mL^{-1}$ , 10 ng mL<sup>-1</sup>, 50 ng mL<sup>-1</sup>, 100 ng mL<sup>-1</sup>, 200 ng mL<sup>-1</sup> in isooctane. For quality assurance, an analytical blank and two spiked samples at a concentration of 12  $\mu$ g kg<sup>-1</sup> for all determinations were used, together with each batch of samples, to check for the absence of matrix interferent and the recovery, which was found to range between 94% and 101%. World Health Organization toxic equivalents (WHO-TEQs) are expressed as lower bound concentrations, assuming that all values of the specific NDL-PCB congeners below the LOQ are equal to zero. The LOQ for all analytes was 1.2 ng  $g^{-1}$ .

#### **Microplastics Analysis**

Frozen specimens of *S. quadrata* (*n* = 5 per site) were pooled and analyzed, as reported by Pastorino et al. (2021a). Briefly, analysis was performed under air-controlled conditions to prevent airborne microplastic pollution in a dedicated clean chamber equipped with double HEPA filtration to minimize microplastic pollution. Accurately rinsed glassware was used at each stage and blanks and spiked samples were analyzed to evaluate the performance of the process. Blanks were performed using

extraction solutions as samples ( $n = 5$ ), because of results obtained on blanks (mean  $0.004 \pm 0.009$ ) items/L, recovered items was a white filament of polyethylene terephthalate-PET), data reported in this study were not corrected by microplastics recorded in blanks because it was negligible. Spiked samples were extracted to evaluate recovery. Pooled gastropods tissues were added with 0.14 items g<sup>-1</sup> of marked microplastics (PP 23–60  $\mu$ m blue and PET 35–100  $\mu$ m white; *n* = 5) to evaluate recovery (PP 97.1  $\pm$  7.8%; PET 94.3  $\pm$  6.4%). Pooled samples were extracted by ultrasonic cavitation (35 °C, 50 min, frequency 40 kHz) using an extraction solution of H  $_2O_2$  (15% w/v). The extract was collected by vacuum filtration on an Anodisc™ membrane (Merck) for chemical analysis. The samples were analyzed by microscopy associated with Fourier transform infrared spectroscopy (μFT-IR Nicolet iN 10MX, Thermo Fischer Scientific) using a liquid nitrogencooled MCT-A detector that operates in the spectral range of 7800–650 cm<sup>-1</sup> by transmission technique. Spectral matches acquired on unknown particles were compared with the reference library (OMNIC™ Picta™ software libraries integrated with libraries collected by the Bioscience Research Center on reference materials); only spectral matches >90% were considered; matches between 70% and 90% of correspondence were evaluated by skilled researchers before being excluded. Limit of detection of the chemical analysis was a particle size of 10 µm. Detailed quality assurance/ quality control activities to ensure the general quality of the method were the same as those recently published elsewhere (Pastorino et al. 2021a).

### **Statistical Analysis and Risk Assessment**

### **Statistical Analysis**

The Shapiro–Wilk test was performed to determine whether our dataset was well-modeled by normal distribution. Differences in contaminant (trace elements, NDL-PCBs, PAHs, microplastics) concentration between the two sites were analyzed using the non-parametric Mann–Whitney U test since the null hypothesis for normal distribution could not be rejected. Differences in microplastics shape, type, and color between the samples from the two sites were tested using the chi-square test. Statistical significance was set at *p-*value < 0.05. Statistical analyses were performed using R software (RStudio, Inc., version 1.1.463). A probabilistic approach using Monte Carlo simulation (MCS) was employed for a more realistic assessment of cancer risk related to toxic metals (Orosun et al. 2020, 2021). MCS was performed using Oracle Crystal Ball software version 11.1.2.4.900.

#### **Risk Assessment: Trace Elements**

Exposure parameters for human health risk are presented in Table S3. To evaluate potential hazardous exposure to potentially toxic elements from gastropod consumption by the local Chinese population, the estimated daily intake (EDI;  $\mu$ g kg<sup>-1</sup> day<sup>-1</sup>) and the target hazard quotient (THQ) were calculated for each element using the following formulas, respectively (USEPA 2000a):

$$
EDI = \frac{Cm * CR}{ABW}
$$

where Cm is the average element concentration between the two sites ( $\mu$ g g<sup>-1</sup> w.w.); CR is the consumption rate obtained from Cheng and Yap (2015) that estimated the potential human health risk from metals via mangrove snail consumption (17.86 g day<sup>-1</sup> and 35.70 g day<sup>-1</sup> for average and highlevel mollusk consumers, respectively); ABW is the average human body weight (70 kg for adults);

$$
\text{THQ} = \frac{EF * ED * CR * Cm}{RfD * ABW * AET} * 10 - 3
$$

where EF is exposure frequency (365 days year<sup>-1</sup>); ED is the exposure duration (70 years), equivalent to an average lifetime; CR is the consumption rate (17.86 g day<sup>-1</sup> and 35.7 g days<sup>-1</sup> [edible part] for average and high-level mollusk consumers, respectively); RfD is the oral reference dose for Cr, Cu, Pb, Zn, Cd, Hg, and As (1500 μg kg<sup>-1</sup> day<sup>-1</sup>, 40 μg kg<sup>-1</sup> day<sup>-1</sup>, 3.5 μg kg<sup>-1</sup> day<sup>-1</sup>, 300 μg kg<sup>-1</sup> day<sup>-1</sup>, 1.0 μg kg<sup>-1</sup> day<sup>-1</sup>, 0.7 µg kg<sup>-1</sup> day<sup>-1</sup>, 50 µg kg<sup>-1</sup> day<sup>-1</sup>, respectively) as reported by literature (Hang et al. 2009; Cheng and Yap 2015); ABW is the average body weight (70 kg for adults); AET is the average exposure time for non-carcinogens (365 days year<sup>-1</sup> \* ED); and  $10^{-3}$  is the unit conversion factor. The total target hazard quotient (TTHQ) of trace elements for *S. quadrata* is the sum of each THQ obtained from each element. Values of THQ or TTHQ > 1 mean that the exposed population is likely to experience obvious deleterious effects after the consumption of contaminated gastropods.

The carcinogenic risk assessment was calculated to estimate the possibility of developing cancer after exposure to toxic metals (Orosun et al., 2020, 2021). To do this, we calculated the average daily intake (ADI) for cancer risk (CR). The ADI was calculated (Zeng et al. 2015):

$$
ADI = \frac{Cm * CR * ED * EF}{ABW * AET}
$$

where Cm is the average element concentration between the two sites (mg kg<sup>-1</sup> w.w.), CR is the consumption rate (0.01786 g days<sup>-1</sup> and 0.0357 g days<sup>-1</sup> for average and high-level mollusk consumers, respectively), ED is the exposure duration (70 years), EF is the exposure frequency (365 days year<sup>-1</sup>), ABW is the average body weight (70 kg for adults), and AET is the average exposure time for noncarcinogens (365 days year<sup>-1</sup> \* ED).

The incremental lifetime cancer risk (ILCR) was calculated for each metal by multiplying the ADI (mg/kg/day) by the carcinogenic slope factor (SF; As: 1.5, Cd: 0.38, Cr: 0.5, Pb:  $8.5 \times 10^{-3}$  [mg/kg/day]<sup>-1</sup>) (Orosun et al. 2020):

# $ILCR = ADI * SF$

For example, an ILCR of 10−4 indicates a probability of 1 in 10,000 individuals developing cancer. The risk was then categorized in seven risk levels (level I: extremely low risk,  $\leq 10^{-6}$ ; level VII: extremely high risk,  $>10^{-3}$ ) according to Li et al. (2017) and Orosun et al. (2020). The acceptable range lies between  $1 \times 10^{-4}$  and  $1 \times 10^{-6}$ .

Finally, we used a probabilistic approach with Monte Carlo simulation (MCS) for a more realistic risk assessment related to potentially toxic metals (As, Cd, Cr, Pb). While overestimation of risk can lead to wastage of limited means on needless mitigatory actions, underestimation can cause preventable health consequences to the dwellers (Orosun et al. 2020, 2021). The MCS method allows for creating probability distributions of risk that describe the uncertainty surrounding specific input variables (i.e., body weight, daily intake, concentration of contaminant, and cancer potency factor). Rather than onepoint values, the MCS method uses several values to repeatedly calculate and obtain results with different assurance levels that range from 1 to 99%. The MCS model was run for a total of 10,000 trials. The mean, 5th and 95th percentiles of ILCR distribution were determined.

# **Risk Assessment: NDL‑PCB**

Dietary daily exposure to NDL-PCBs was calculated as follows (Lee et al. 2021):

$$
ADD = \frac{C * CR}{ABW}
$$

where ADD (ng k g<sup>-1</sup> b.w. d ay<sup>-1</sup>) is the average daily dose exposure to NDL-PCBs; C (ng g<sup>-1</sup> w.w.) is the average concentration for the two sites of each NDL-PCBs, considering that the values of the congeners below the LOQ were considered equal to the LOQ (1.2 ng  $g^{-1}$ ); CR is the consumption rate (17.86 g days<sup>-1</sup> and 35.7 g days<sup>-1</sup> for average and high-level mollusk consumers, respectively); and ABW is the average body weight (70 kg).

The non-cancer hazard quotient (HQ) for each of the six congeners was calculated as follows (Lee et al. 2021):

$$
HQ = \frac{ADD}{TDI}
$$

where tolerable daily intake (TDI) is set at 10 ng kg<sup>-1</sup> b.w. day<sup>-1</sup> as proposed by the World Health Organization (AFFSA 2007), assuming that the six indicators of PCB account for about 50% of total dietary NDL-PCB intake (Johnson 2007).

Finally, the hazard index (HI) was calculated according to the formula (Lee et al. 2021):

HI =Σ*HQ*

where  $\Sigma$ HQ is the sum of the individual HQs of each of the six congeners. HQ or HI < 1 indicates a daily exposure dose that might not cause adverse health effects.

# **Risk Assessment: PAHs**

Dietary daily exposure to PAHs was calculated for each of the four congeners as follows (Tongo et al. 2017):

$$
DDI = \frac{C \cdot CR}{ABW}
$$

where DDI is the dietary daily intake ( $\mu$ g kg<sup>-1</sup>b.w. day<sup>-1</sup>), C ( $\mu$ g kg<sup>-1</sup> w.w.) is the average concentration for the two sites of each PAH; CR is the consumption rate  $(0.0175 \text{ kg days}^{-1}$  and  $0.0357 \text{ kg days}^{-1}$  for average and high-level mollusk consumers, respectively); and ABW is the average body weight (70 kg).

Carcinogenic risk assessment of PAHs was assessed using the PAH4 index, individual PAH carcinogenic potencies, and carcinogenic toxic equivalents (TEQs). The PAH4 index was calculated as follows (Tongo et al. 2017):

PAH4 Index (PAH4) =  $\sum (B[a]A + Chr + B[b]FL + B[a]P)$  where  $\sum (B[a]A + Chr + B[b]FL + B[a]P)$  is the sum of the concentration (μg kg<sup>-1</sup> w.w.) of benzo[a]anthracene (B[a] A), chrysene (Chr), benzo[b]fluoranthene (B[b]FL), and benzo[a]pyrene (B[a]P).

The carcinogenic potency of each PAH congener was calculated by multiplying the PAH concentration by the individual toxicity equivalency factor (TEF) (Tongo et al. 2017):

Carcinogenic potencies of individual PAHs (B(A)Pteq) = Ci  $*$  TEFi where Ci (μg kg<sup>-1</sup> w.w.) is the measured concentration of the PAH congener (i) in *S. quadrata* (mean of site 1 and site 2) and TEF is the individual toxicity equivalency factor of the PAH congener (i). TEF are (Nisbet and LaGoy 1992):  $B[a] A = 0.1$ ; Chr = 0.01;  $B[b]FL = 1$ ;  $B[a]P = 0.1$ .

Carcinogenic toxic equivalents (TEQs) were calculated for the PAH4 index as follows (Ding et al. 2012):

 $TEQ = \sum B(A)Pteq$ 

The TEQ was compared with a screening value (SV) to determine the health risks of PAH4 to humans from gastropod consumption. The SV indicates the threshold of chemicals in edible tissue which is of potential public health concern (Cheung et al. 2007):

$$
SV = \frac{\left(\frac{RL}{SF}\right) * ABW}{CR}
$$

where RL is the maximum acceptable risk level dimensionless =  $10^{-5}$  (USEPA 2000b), SF is the oral slope factor (SF): 1 mg kg<sup>-1</sup> day<sup>-1</sup> (USEPA 1993), CR is the consumption rate (0.0175 kg days<sup>-1</sup> and 0.0357 kg days−1 for average and high-level mollusk consumers, respectively), and ABW is the average body weight (70 kg).

# **Results**

# **Potentially Toxic Element Accumulation**

Table 1 presents the mean toxic element concentration in *S. quadrata* from the two sampling sites. The mean concentration of elements ( $\mu$ g g<sup>-1</sup>) at site 1 was: Zn (3.55) > Cu (1.89) > Pb (0.113) > As (0.112)  $>$  Hg (0.045)  $>$  Cr (0.009)  $>$  Cd (0.001); the mean concentration (µg g<sup>-1</sup>) at site 2 was: Zn (3.68)  $>$  Cu  $(2.39)$  > As  $(0.685)$  > Pb  $(0.141)$  > Hg  $(0.060)$  > Cr  $(0.021)$  > Cd  $(0.002)$ . The concentrations of all elements were slightly higher at site 2; however, the difference in element concentration between the two sites was not statistically significant (Mann–Whitney U test,  $p > 0.05$ ).

# **NDL‑PCB Accumulation**

The concentration of  $\Sigma_6$  PCB was significantly higher in the samples from site 2 (7.32 ng g<sup>-1</sup> vs. 3.07 ng  $g^{-1}$ ) (Mann–Whitney U test,  $p = 0.001$ ) (Table 2). Among the single congeners, the concentration of PCB-138 and PCB-153 was analytically predominant: PCB-138 accounted for 53.4% and 29.6% of the  $\Sigma_6$  PCB for site 1 and site 2, respectively; PCB-153 accounted for 46.9% and 24.2% of the  $\Sigma_6$  PCB for

site 1 and site 2, respectively; PCB-180 was detected only in samples from site 2 and accounted for 22% of the  $\Sigma_6$  PCB.

## **PAHs Accumulation**

Among the single congeners, the concentration of Chr, B[b]FL and B[a]P was below the limit of quantification (0.5  $\mu$ g kg<sup>-1</sup>). B[a]A was detected in the samples from both sites (mean values: 0.5 and 0.7 μg kg−1 for site 1 and site 2, respectively) (Table 3).

# **Microplastics**

The number of microplastics frequency was  $0.8 \pm 1.30$  item/ specimen for site 1, and  $1 \pm 0.37$ item/specimen for site 2 (Table 4). There was a significant difference in the items/ specimens between the two sites (Mann–Whitney U test,  $p = 0.008$ ). Fibers ( $n = 6$ ) were found in samples from site 1 ( $n =$ 2) and site 2 ( $n = 4$ ); fragments ( $n = 3$ ) were detected in samples from site 1 ( $n = 1$ ) and site 2 ( $n = 2$ ). There were no differences in shape frequency between the two sites ( $\chi^2$  test,  $p > 0.05$ ). The predominant colors were green (33.33%) and tan (33.33%), followed by blue (22.22%) and black (11.12%). There were no differences in color frequency between the two sites ( $\chi^2$  test,  $p > 0.05$ ).

The chemical composition of microplastic was predominantly PET polyethylene terephthalate (66.67%), followed by nylon (22.22%) and PP polypropylene (11.11%). There were no differences in chemical composition frequencies between the two sites ( $\chi^2$  test,  $p > 0.05$ ).

#### **Risk Assessment**

#### **Potentially Toxic Elements**

Table 5 reports the estimated weekly intake (EDI) and the target hazard quotient (THQ) for potentially toxic elements by the consumption of *S. quadrata* in an average and a high-level mollusk consumer. In an average mollusk consumer, the EDI ranged from 0.0003 μg kg<sup>-1</sup> day<sup>-1</sup> for Cd to 0.92 μg kg<sup>-1</sup> day<sup>-1</sup> for Zn. A similar scenario was observed for a high-level mollusk consumer: EDI ranged from 0.0007  $\mu$ g kg<sup>-1</sup> day<sup>-1</sup> for Cd to 1.84  $\mu$ g kg<sup>-1</sup> day<sup>-1</sup> for Zn. The target hazard quotient (THQ) for all trace elements  $was < 1$ , as was the total target hazard quotient (TTHQ) for an average and a high-level mollusk consumer: 0.0026 and 0.0051, respectively. Average daily intake (ADI) (Table 6), calculated to obtain the ILCR, ranged from 0.00000013 for As to 0.0000039 for Cr and from 0.00000025 for As to 0.0000078 for Cr, for an average and a high consumption level, respectively. The ILCR was also estimated; the mean (for average and high consumption, respectively) was: As (1.90E−7; 3.81E−7), Cd (1.26E−7 ; 2.51E−7), Cr (1.94E−6; 3.88E−6), and Pb (2.76E−7; 5.51E−7), with Cr contributing the most to cancer risk, followed by Pb, As, and Cd (Table 6).

The MCS results are presented in Fig. 2 and Tables S4 and S5. According to the 95% cumulative probability, Cr (1.24E − 5) posed the highest risk, followed by As (1.85E − 6), Pb (1.49E − 6), and Cd (7.01E − 7). The sensitivity chart from the MCS model (Table S5) revealed that carcinogenic slope factor ranks highest, followed by the daily intake of gastropods, concentration of toxic element in *S. quadrata*, and then average adult body weight, which was negative for all potentially toxic elements.

### **NDL‑PCB**

Table 7 presents the average daily dose exposure (ADD) to NDL-PCBs for an average and a high-level mollusk consumer. The ADD ranged from 0.31 ng kg<sup>-1</sup> day<sup>-1</sup> for PCB28 and PCB-101 to 0.49 ng kg<sup>-1</sup> day<sup>-1</sup> for PCB-138 for an average mollusk consumer and from 0.61 ng kg<sup>-1</sup> day<sup>-1</sup> for PCB-28 and PCB-101 to 0.97 ng kg<sup>-1</sup> day<sup>-1</sup> for PCB138 for a high-level mollusk consumer. The ADD values for the  $\Sigma_6$ PCB ranged from 2.24 ng kg<sup>-1</sup> day<sup>-1</sup> to 4.49 ng kg<sup>-1</sup> day<sup>-1</sup> for an average and a high-level mollusk consumer, respectively. All the hazard quotient (HQ) values were  $\leq 1$ . The same results were obtained by calculating the Hazard Index (HI) (sum of the individual HQs), which was < 1 for an average (0.22) and a high-level (0.45) mollusk consumer alike.3.5.3 PAHs.

Table 8 presents the dietary daily intake (DDI) ( $\mu$ g kg<sup>-1</sup>b.w. day<sup>-1</sup>) for an average and a high-level mollusk consumer. The DDI ranged from 0.0001  $\mu$ g kg<sup>-1</sup> day<sup>-1</sup> for B[a]P, Chr and B[b]FL to 0.0002  $\mu$ g  $kg^{-1}$  day<sup>-1</sup> for B[a]A for an average mollusk consumer and from 0.0003 µg kg<sup>-1</sup> day<sup>-1</sup> for B[a]P, Chr and B[b]FL to

0.0004 μg  $kg^{-1}$  day<sup>-1</sup> for B[a]A for a high-level mollusk consumer.

The carcinogenic potencies of PAHs (B(A)Pteq) were: B[a]P:  $0.05 \mu$ g kg<sup>-1</sup>; B[a]A:  $0.06 \mu$ g kg<sup>-1</sup>; Chr: 0.005 μg kg<sup>-1</sup>; B[b]FL: 0.5 μg kg<sup>-1</sup>. The TEQ obtained from PAH4 was 0.615 μg kg<sup>-1</sup>(0.000615 mg  $kg^{-1}$ ). The SV was 0.04 and 0.02 mg kg<sup>-1</sup> for an average and a high-level consumer, respectively.

# **Discussion**

### **Contaminants Accumulation**

The literature documents the use of mollusks as bioindicators to assess water quality in compliance with ecological and human safety legislation (Tietze and De Francesco 2010; Kumar et al. 2017). Generally, freshwater gastropods are considered good indicators of water quality owing to their relatively long-life cycle and low mobility (de Freitas Tallarico 2015). They are highly sensitive to chemical pollution; high levels can cause their disappearance or inhibit their reproduction (Krupnova et al. 2018; Caixeta et al. 2020). They are especially sensitive to pollution by heavy metals (i.e., cadmium, mercury, lead, zinc, copper) (Krupnova et al. 2018). Since contaminants can have a negative impact on freshwater mollusks at environmental concentrations lower than for other invertebrates or vertebrates, mollusks are studied as part of an ecological early warning system (de Freitas Tallarico 2015). In the present study the two sampling sites are located downstream of wastewater treatment plants and civil and industrial discharges, but the levels of contaminants were not of particular concern for human safety.

Although the difference was not statistically significant, the trace element concentration was higher in samples from site 2. This is easily explained by the fact that site 2 is located downstream of site 1 and receives the waters from the Pesa River near Montelupo Fiorentino (Florence Province). The Pesa River is under anthropogenic pressures that deteriorate the water quality at site 2. Despite this, the concentration was low. Cortecci et al. (2009) measured the main toxic trace elements (As, Cu, Zn, Ni, Pb, Cd) in water samples from the River Arno and found that concentrations were below the acceptable limits of current European regulations for the quality of waters for potable use and fish life. In a later study, Mecatti et al. (2017) measured the concentration of Hg, Cd, As, Pb, and Cr in the muscle of the European catfish (*Silurus glanis*) from the Arno and found very low levels of these toxic elements.

Unfortunately, there are no studies on trace element accumulation in *S. quadrata* or other mollusk species for comparison with our study. However, drawing on data for other members of the same gastropod family (Viviparide), the concentrations we measured were much lower than those reported by Gundacker (2000) who detected Cd, Cu, Pb, and Zn in *Viviparus* sp. soft bodies at five sampling sites in Vienna (Austria). Of note is that *S. quadrata* is a filter-feeder organism, which is why the trace elements accumulating in its soft body are more closely related to ingestion of suspended matter and filtrate from river water. Metals concentration in water recorded by ARPAT at the MAS-108 station (located near the two sampling sites) and during the same month (July 2020) as our sampling revealed low values for all elements (As 1 μg L<sup>-1</sup>; Cd < 0.2 μg L<sup>-1</sup>; Cr < 1 μg L<sup>-1</sup>; Pb < 1 μg L<sup>-1</sup>; Cu 5 μg L<sup>-1</sup>; Zn 35 μg L<sup>-1</sup>; Hg 0.035 μg L<sup>-1</sup>) (ARPAT 2021).

Among the NDL-PCBs, the six congeners (28, 52, 101, 138, 153, 180) were selected for this study because they are easily quantified compared to other PCBs (Squadrone et al. 2013). Moreover, they account for all relevant levels of chlorination. The EFSA stated that the sum of these six congeners is an adequate indicator of the occurrence of NDLPCBs and human exposure to them (EFSA 2005). As for the trace elements, the  $\Sigma_6$  NDL-PCB was also higher at site 2 compared to site 1. Congeners PCB-153 and PCB-138 were analytically predominant compared to the other congeners. PCB-52 and PCB-180 were also detected in samples from site 2. The non-planar PCB-138, PCB-153, and PCB-180 are the most predominant and persistent congeners reported in humans (Humphrey et al. 2000; Leijs et al. 2019). Squadrone et al. (2013) found that congeners PCB-153 and PCB-138 were analytically predominant (40% and 30%, respectively) in the muscle of *Silurus glanis* from north Italian rivers. The concentrations we measured are in line with those detected by Nhan et al. (2001) in the soft tissue of the freshwater mollusk (*Angulyagra* sp.; Viviparidae) from water canals in the region of Hanoi City (Vietnam).

The PAHs concentration was below the LOQ

(0.5  $\mu$ g kg<sup>-1</sup>) for the majority of the PAHs congeners (B[a]

P, Chr and B[b]FL), except for B[a]A which was near the LOQ for both sites. Primary exposure to benz[a]anthracene occurs mainly from smoking or second-hand smoke, air polluted with combustion products or food and water polluted with combustion products (Librando et al. 2014). Generally, B[a]A do not accumulate in fish (due to biotransformation) (Bleeker and Verbruggen 2009). Its accumulation in *Sinotaia quadrata* may be related to the absence of an enzyme for metabolic activation. Moreover, aquatic mollusks in general are limited in their ability to excrete pollutants directly through their excretory organs and tissues compared to other invertebrates such as arthropods or vertebrates, which is why gastropods can be used in ecological early warning systems (Oehlmann and Schulte-Oehlmann 2003).

The number of microplastic items/specimen was similar to those reported by Akindele et al. (2019): 1.70 ± 0.42 items/specimen in the freshwater gastropod *Melanoides tuberculate* (Thiaridae) from the Osun River system (Nigeria). PET (polyethylene terephthalate) was the main chemical type we found, as reported in other freshwater ecosystems (Li et al. 2020; Pastorino et al. 2021a, 2021b). PET is a synthetic material in the polyester family made from oil, natural gas or vegetable raw materials. Thanks to its properties, PET is used in diverse areas: from the production of containers (plastic bottles) to textiles (Piccardo et al. 2020). The dominant microplastics shape in the samples from both sites was fibers. This finding is shared by previous studies that reported fibers as the most dominant shape in freshwater environments (McCormick et al. 2014; Baldwin et al. 2016; Pastorino et al. 2021a). The presence of fibers can be attributed to their diverse origins: washing, domestic wastewater, and other anthropogenic activities. Their occurrence in our samples could be related to the wastewater from textile factories in Prato that reaches the River Arno through its Bisenzio tributary.

# **Compliance with National and International Regulatory Bodies and Health Risk Assessment**

Generally, analysis of the results for accumulation highlights low concentrations for all contaminates analyzed. Since no arsenic limit has been established by any regulatory body, its concentration in *S. quadrata* could not be assessed for potential risk by direct safety guideline comparison. Cd concentration was within the limit of 1.0  $\mu$ g g<sup>-1</sup> imposed by Regulation (EC) 1881/2006 (EC 2006). Cr concentration in samples from both sampling sites were well below the maximum allowable limit of 13  $\mu$ g g<sup>-1</sup> for shellfish as established by the US Food and Drug Administration (USFDA 2011). The Cu levels were below the limits recommended by the US Environmental Protection Agency (USEPA 1983) (120  $\mu$ g g<sup>-1</sup>), the UK Ministry of Agriculture, Fisheries and Food (MAFF 2000) (20  $\mu$ g g<sup>-1</sup>), and the WHO (1996) (30  $\mu$ g g<sup>-1</sup>). The Hg levels were below the limit of 0.5  $\mu$ g g<sup>-1</sup> imposed by Regulation (EC) 1881/2006 (EC 2006). Pb levels were below the limit of 4.0  $\mu$ g g<sup>-1</sup> recommended by the USEPA (USEPA 1983), 2.0  $\mu$ g g<sup>-1</sup> by the UK Ministry of Agriculture, Fisheries and Food (MAFF 2000), 2.0  $\mu$ g  $g^{-1}$  by the WHO (WHO 1996) 0.5 µg  $g^{-1}$  by the UN Food and Agriculture Organization (FAO 1983) and 1.5  $\mu$ g g<sup>-1</sup> imposed by Regulation (EC) 1881/2006 (EC 2006). Zn levels were also within the limits (480  $\mu$ g g<sup>-1</sup>) established by the USEPA (1983), the UK Ministry of Agriculture, Fisheries and Food (MAFF 2000) (50  $\mu$ g g<sup>-1</sup>), and those (30  $\mu$ g g<sup>-1</sup>) determined by FAO (1983).

The sum of the six indicators NDL-PCBs 28, 52, 101, 138, 153, 180 ( $\Sigma_6$  NDL-PCBs) was well below the maximum tolerable levels (MLs) of 75 ng  $g^{-1}$  established by Regulation 1259/2011 (EC 2011b).

Knowledge of the dietary intake of *S. quadrata* is an important factor in assessing risk to human health from potentially harmful chemicals in food. For both an average and a high-level consumer the trace elements EDI was lower than the oral reference dose (Cheng and Yap 2015). Generally, the EDI we calculated was lower than that reported in a study by Cheng and Yap (2015) that estimated the health risks from toxic metals via mangrove snail consumption from the Malaysian Peninsula.

Target hazard quotients (THQ) were also calculated since they are widely used in the risk assessment of heavy metals in contaminated foods (Storelli 2008). Our findings revealed that all THQs were < 1 for all elements, indicating that the health risks associated with As, Cd, Cr, Cu, Hg, Pb, Zn exposure for average and high-level mollusk consumers are insignificant. The TTHQ was also < 1, suggesting that the consumption of *S. quadrata* does not carry health risks.

The ILCR was far below the safety values of  $1 \times 10^{-4}$  as recommended by the USEPA (USEPA 2021) for the toxic elements studied here. The ILCR calculated by the MCS model was within the safety region of  $1 \times 10^{-4}$  and  $1 \times 10^{-6}$ , as recommended by the USEPA (USEPA 2021). Based on the Delphi method (Li et al. 2017), the risk ranged from level I to level II, from completely acceptable to not willing to care about the risk.

The average daily dose (ADD) of the  $\Sigma_6$  NDL-PCB was 2.24 and 4.49 ng kg<sup>-1</sup> day<sup>-1</sup> b.w. for an average and a highlevel mollusk consumer, respectively. The two ADD were much lower than the reference dose of the  $\Sigma_6$  NDL-PCB (10 ng kg<sup>-1</sup> day<sup>-1</sup> b.w.) (Lee et al. 2021). The average daily intake of total NDL-PCBs for adults in Europe is estimated to be between 10 ng kg<sup>-1</sup> day<sup>-1</sup> and 45 ng kg<sup>-1</sup>  $day^{-1}$  b.w. (EFSA 2005) and lower than the ADD we calculated.

The non-cancer risk (HQ) and the hazard index (HI) associated with NDL-PCBs exposure from an average and a high consumption of gastropods were all  $\leq 1$ , indicating no adverse effects for human health. Unfortunately, the lack of similar studies on freshwater gastropods precludes comparison with the literature.

The dietary daily intake (DDI) for the PAH4 index was much lower than the range (0.0195–0.0345 μg kg day<sup>-1</sup> b.w.) estimated by the EFSA (EFSA 2008) for each congener. Also, B[a]P daily intake was below the acceptable daily intake of 0.01  $\mu$ g kg<sup>-1</sup> day<sup>-1</sup> suggested by the Joint FAO/ WHO Expert Committee on Food Additives (JFCFA) and for the range 0.039–0.065  $\mu$ g kg<sup>-1</sup> day<sup>-1</sup> reported by the EFSA (EFSA 2008). Concentrations of B[a]P and PAH4 index were lower than the legal limit (fishery products) of 2  $\mu$ g kg<sup>-1</sup> and 12  $\mu$ g kg<sup>-1</sup>, respectively, imposed by Regulation (EC) 835/2011 (EC 2011a).

The SV for average and high consumption was compared to the TEQ to determine the health risk to humans via gastropod consumption. The TEQ (0.000615 mg  $kg^{-1}$ ) was far below the SV (0.04 and 0.02 mg kg−1 , respectively), indicating no risk to human health from the consumption of *S. quadrata.*

We were unable to assess the health risk from exposure to microplastics, since too little data are available to fully understand their implications for human health, as declared by the EFSA (German Federal Institute for Risk Assessment et al. 2020). Currently, there is no legislation concerning microplastics as contaminants in food. Nonetheless, negative effects may be expected due to their physical (size, and shape) and chemical proprieties, concentration, occurrence of contaminants adsorbed or microbial biofilm growth (Campanale et al. 2020).

#### **Conclusions**

Contaminant levels in specimens of *S. quadrata* collected from two sites on the Arno were compliant with regulatory limits and guidelines. The findings from health indices indicated no adverse effects for human health by exposure to trace elements, NDL-PCB, and PAHs from gastropod consumption. However, the ingestion of microplastics could expose humans to chemicals known to be harmful. Substantial data gaps with respect to exposure as well as toxicity of such particles impede the risk assessment. Finally, regular monitoring of the River Arno for organic and inorganic pollutants is warranted to detect eventual changes in contaminant levels.

**Acknowledgements** We would like to thank Mr. Floriano Pratelli for his help during sampling.

**Author Contributions** Conceptualization: Paolo Pastorino and Marino Prearo; data curation, Paolo Pastorino; investigation: Paolo Pastorino, Annamaria Nocita, Valentina Ciccotelli, Annalisa Zaccaroni, Serena Anselmi, Roberta Giugliano, Mattia Tomasoni, Marina Silvi, Vasco Menconi, Barbara Vivaldi, Elisabetta Pizzul, Monia Renzi and Marino Prearo; methodology: Paolo Pastorino, Valentina Ciccotelli, Annalisa Zaccaroni, Serena Anselmi, Roberta Giugliano, Mattia Tomasoni, Marina Silvi, Vasco Menconi, Barbara Vivaldi, Elisabetta Pizzul, Monia Renzi, Marino Prearo; supervision: Marino Prearo; writingoriginal draft: Paolo Pastorino; writing-review and editing: Paolo Pastorino, Annamaria Nocita, Valentina Ciccotelli, Annalisa Zaccaroni, Serena Anselmi, Roberta Giugliano, Mattia Tomasoni, Marina Silvi, Vasco Menconi, Barbara Vivaldi, Elisabetta Pizzul, Monia Renzi, Marino Prearo.

**Data Availability** The data supporting the findings of this study are available on request from the corresponding author.

# **Declarations**

**Conflict of Interest** The authors declare no conflict of interest.

**Ethical approval** Not applicable.

**Consent to participate** Not applicable.

**Consent for publication** All authors have read and agreed to the published version of the manuscript.

# **References**

- AFFSA (2007). Opinion of the French food safety agency on the establishment of relevant maximum levels for non dioxin-like polychlorobiphenyls (NDL-PCB) in some foodstuffs afssa – Request No. 2006-SA-0305. [https://w ww.anses.f r](https://www.anses.fr/fr/system/files/RCCP2006sa0305EN.pdf) [/fr/s ystem](https://www.anses.fr/fr/system/files/RCCP2006sa0305EN.pdf) [/files/R CCP](https://www.anses.fr/fr/system/files/RCCP2006sa0305EN.pdf) [2006](https://www.anses.fr/fr/system/files/RCCP2006sa0305EN.pdf) [sa](https://www.anses.fr/fr/system/files/RCCP2006sa0305EN.pdf) [0305EN. pdf.](https://www.anses.fr/fr/system/files/RCCP2006sa0305EN.pdf) Accessed 20 Feb 2021
- Akindele EO, Ehlers SM, Koop JH (2019) First empirical study of freshwater microplastics in West Africa using gastropods from

Nigeria as bioindicators. Limnologica 78:125708

- ARPAT, 2021. Dati e mappe. [http://w ww.arpat.t osc](http://www.arpat.toscana.it/datiemappe#c9=banche.dati&c9=bollettini&c9=dati&c9=mappe&c0=5&b_start=0&c1=acque.interne) [ana.i t](http://www.arpat.toscana.it/datiemappe#c9=banche.dati&c9=bollettini&c9=dati&c9=mappe&c0=5&b_start=0&c1=acque.interne) [/datiem appe](http://www.arpat.toscana.it/datiemappe#c9=banche.dati&c9=bollettini&c9=dati&c9=mappe&c0=5&b_start=0&c1=acque.interne) [#](http://www.arpat.toscana.it/datiemappe#c9=banche.dati&c9=bollettini&c9=dati&c9=mappe&c0=5&b_start=0&c1=acque.interne) [c9=b anche.dati&c 9](http://www.arpat.toscana.it/datiemappe#c9=banche.dati&c9=bollettini&c9=dati&c9=mappe&c0=5&b_start=0&c1=acque.interne) [=bollet](http://www.arpat.toscana.it/datiemappe#c9=banche.dati&c9=bollettini&c9=dati&c9=mappe&c0=5&b_start=0&c1=acque.interne)  [tin](http://www.arpat.toscana.it/datiemappe#c9=banche.dati&c9=bollettini&c9=dati&c9=mappe&c0=5&b_start=0&c1=acque.interne) [i&c 9](http://www.arpat.toscana.it/datiemappe#c9=banche.dati&c9=bollettini&c9=dati&c9=mappe&c0=5&b_start=0&c1=acque.interne) [=dati&c 9](http://www.arpat.toscana.it/datiemappe#c9=banche.dati&c9=bollettini&c9=dati&c9=mappe&c0=5&b_start=0&c1=acque.interne) [=mappe&](http://www.arpat.toscana.it/datiemappe#c9=banche.dati&c9=bollettini&c9=dati&c9=mappe&c0=5&b_start=0&c1=acque.interne) [c0=5 &b](http://www.arpat.toscana.it/datiemappe#c9=banche.dati&c9=bollettini&c9=dati&c9=mappe&c0=5&b_start=0&c1=acque.interne) start=  $0& \text{c1}$  = acque. inter ne Accessed 20 Feb 2021
- Baldwin AK, Corsi SR, Mason SA (2016) Plastic debris in 29 Great Lakes tributaries: relations to watershed attributes and hydrology.

Environ Sci Technol 50:10377–10385

- Bleeker J, Verbruggen J (2009) Bioaccumulation of polycyclic aromatic hydrocarbons in aquatic organisms Bioaccumulation of polycyclic aromatic hydrocarbons in aquatic organisms. Natl Inst Public Heal Environ 56.
- Borrelle SB, Ringma J, Law KL, Monnahan CC, Lebreton L, McGivern A et al (2020) Predicted growth in plastic waste exceeds efforts to mitigate plastic pollution. Science 369(6510):1515–1518
- Caixeta MB, Araújo PS, Gonçalves BB, Silva LD, Grano-Maldonado MI, Rocha TL (2020) Toxicity of engineered nanomaterials to aquatic and land snails: A scientometric and systematic review. Chemosphere 260:127654
- Campanale C, Massarelli C, Savino I, Locaputo V, Uricchio VF (2020) A detailed review study on potential effects of microplastics and additives of concern on human health. Int J Environ Res Public Health 17(4):1212
- Cheng WH, Yap CK (2015) Potential human health risks from toxic metals via mangrove snail consumption and their ecological risk assessments in the habitat sediment from Peninsular Malaysia. Chemosphere 135:156–165
- Cheung KC, Leung HM, Kong KY, Wong MH (2007) Residual levels of DDTs and PAHs in freshwater and marine fish from Hong Kong markets and their health risk assessment. Chemosphere 66(3):460– 468
- Cianfanelli S, Stasolla G, Inghilesi AF, Tricarico E, Goti E, Strangi A, Bodon M (2017) First European record of *Sinotaia quadrata* (Benson, 1842), an alien invasive freshwater species: accidental or voluntary introduction?: (Caenogastropoda: Viviparidae). Bollettino Malacologico 53:150–160
- Cortecci G, Boschetti T, Dinelli E, Cidu R, Podda F, Doveri M (2009) Geochemistry of trace elements in surface waters of the Arno River Basin, northern Tuscany. Italy Appl Geochem 24(5):1005–1022
- Cunningham VL, Binks SP, Olson MJ (2009) Human health risk assessment from the presence of human pharmaceuticals in the aquatic environment. Regul Toxicol Pharmacol 53(1):39–45
- de Freitas TL (2015) Freshwater gastropods as a tool for ecotoxicology assessments in Latin America. Am Malacol Bull 33(2):330–336
- Ding C, Ni HG, Zeng H (2012) Parent and halogenated polycyclic aromatic hydrocarbons in rice and implications for human health in China. Environ Pollut 168:80–86
- EC (2006) Commission Regulation N. 1881/2006 of 19 December 2006. Setting maximum levels of certain contaminants in foodstuff. Off J Eur Union 65:5–24
- EC (2011) Commission Regulation (EU) No 1259/2011 of 2 December 2011 amending Regulation (EC) No 1881/2006 as regards maximum levels for dioxins, dioxin-like PCBs and non dioxinlike PCBs in foodstuffs. Off J Eur Union 320:18–23
- EC (2011) Commission Regulation (EU) No 835/2011 of 19 August 2011 amending Regulation (EC) No 1881/2006 as regards maximum levels for polycyclic aromatic hydrocarbons in foodstuffs. Off J Eur Union 215(4):1–5
- EFSA (2005) Opinion of the scientific panel on contaminants in the food chain on a request from the commission related to the presence of the non-dioxin-like polychlorinated biphenyls (PCB) in feed and food (Adopted on 8 November 2005) (Question No. EFSA-Q-2003-114). EFSA J 284:1–137
- EFSA (2008) Scientific opinion of the panel on contaminants in the food chain on a request from the European Commission on polycyclic aromatic hydrocarbons in food. EFSA J 724:1–114
- Esposito G, Meloni D, Abete MC, Colombero G, Mantia M, Pastorino P, Prearo M, Pais A, Antuofermo E, Squadrone S (2018) The bivalve *Ruditapes decussatus*: A biomonitor of trace elements pollution in Sardinian coastal lagoons (Italy). Environ Pollut 242:1720–1728
- FAO (1983) Compilation of legal limits for hazardous substances in fish and fishery products. FAO Fisheries Circular. 764. FAO, Rome, pp. 102, http:// www. fao. org/ fi/ oldsi te/ eims\_ search/ 1\_ dett. asp? calli ng= simple s resul t& lang= fr&p ub  $id= 65155$ .
- German Federal Institute for Risk Assessment (BfR), Department of Food Safety, Unit Effect‐based Analytics and Toxicogenomics Unit and Nanotoxicology Junior Research Group, Berlin, Germany, Shopova S, Sieg H, Braeuning A (2020) Risk assessment and toxicological research on micro-and nanoplastics after oral exposure via food products. EFSA J. 18:e181102
- Gundacker C (2000) Comparison of heavy metal bioaccumulation in freshwater molluscs of urban river habitats in Vienna. Environ Pollut 110(1):61–71
- Halder D, Saha JK, Biswas A (2020) Accumulation of essential and non-essential trace elements in rice grain: possible health impacts on rice consumers in West Bengal India. Sci. Total Environ. 706:135944
- Hang X, Wang H, Zhou J, Ma C, Du C, Chen X (2009) Risk assessment of potentially toxic element pollution in soils and rice (*Oryza sativa*) in a typical area of the Yangtze River Delta. Environ Pollut 157(8–9):2542–2549
- Honda M, Suzuki N (2020) Toxicities of polycyclic aromatic hydrocarbons for aquatic animals. Int J Environ Res Public Health 17(4):1363
- Howell NL, Suarez MP, Rifai HS, Koenig L (2008) Concentrations of polychlorinated biphenyls (PCBs) in water, sediment, and aquatic biota in the Houston Ship Channel. Texas Chemosphere 70(4):593– 606
- Humphrey HE, Gardiner JC, Pandya JR, Sweeney AM, Gasior DM, McCaffrey RJ, Schantz SL (2000) PCB congener profile in the serum of humans consuming Great Lakes fish. Environ Health Perspect 108(2):167–172
- ISTAT (2019) Il Censimento Permanente Della Popolazione: Dati
- Regionali. [https:// www. istat. it/ it/ archi vio/ stran ieri](https://www.istat.it/it/archivio/stranieri) [A](https://www.istat.it/it/archivio/stranieri)ccessed 20 Feb 2021
- Jia Y, Wang L, Qu Z, Wang C, Yang Z (2017) Effects on heavy metal accumulation in freshwater fishes: species, tissues, and sizes. Environ Sci Pollut Res Int 24(10):9379–9386
- Johnson JW (2007) Final report on the safety assessment of capsicum annuum extract, capsicum annuum fruit extract, capsicum annuum resin, capsicum annuum fruit powder, capsicum frutescens fruit, capsicum frutescens fruit extract, capsicum frutescens resin, and capsaicin Int. J Toxicol 26(1):3– 106
- Krupnova TG, Mashkova IV, Kostryukova AM, Schelkanova EE, Gavrilkina SV (2018) Gastropods as potential biomonitors of contamination caused by heavy metals in South Ural lakes. Russia Ecol Indic 95:1001–1007
- Kumar N, Krishnani KK, Singh NP (2017) Oxidative and cellular stress as bioindicators for metal contamination in freshwater mollusk *Lamellidens marginalis*. Environ Sci Pollut Res Int 24(19):16137–16147
- Lee CC, Chang WH, Hung CF, Chen HL (2021) Fish consumption is an indicator of exposure to nondioxin like polychlorinated biphenyls in cumulative risk assessments based on a probabilistic and sensitive approach. Environ Pollut 268:115732
- Leijs MM, Gan L, De Boever P, Esser A, Amann PM, Ziegler P, Fietkau K, Schettgen T, Kraus T, Merk HF, Baron JM (2019) Altered gene expression in dioxin-like and non-dioxin-like PCB exposed peripheral blood mononuclear cells. Int J Environ Res Public Health 16(12):2090
- Li C, Busquets R, Campos LC (2020) Assessment of microplastics in freshwater systems: a review. Sci Total Environ 707:135578
- Li F, Qiu Z, Zhang J, Liu C, Cai Y, Xiao M (2017) Spatial distribution and fuzzy health risk assessment of trace elements in surface water from Honghu Lake. Int J Environ Res Public Health 14(9):1011
- Librando V, Bracchitta G, de Guidi G, Minniti Z, Perrini G, Catalfo A (2014) Photodegradation of anthracene and benzo[a]anthracene in polar and apolar media: new pathways of photodegradation. Polycyclic Aromat Compd 34(3):263–279
- MAFF (2000) Monitoring and surveillance of non-radioactive contaminants in the aquatic environment and activities regulating the disposal of wastes at sea, 1997. Aquatic environment monitoring report number 52.
- Malisch R, Kotz A (2014) Dioxins and PCBs in feed and food—review from European perspective. Sci Total Environ 491:2–10
- Marzooghi S, Di Toro DM (2017) A critical review of polycyclic aromatic hydrocarbon phototoxicity models. Environ Toxicol Chem 36(5):1138–1148
- McCormick A, Hoellein TJ, Mason SA, Schluep J, Kelly JJ (2014) Microplastic is an abundant and distinct microbial habitat in an urban river. Environ Sci Technol 48:11863–11871
- McIntyre JK, Beauchamp DA (2007) Age and trophic position dominate bioaccumulation of mercury and organochlorines in the food web of Lake Washington. Sci Total Environ 372:571–584

Mecatti M, Gualtieri M, Cecchi G, Gattai K, Mariotto L, Bossù T

(2017) Biological and nutritional characteristics of wild European catfish (*Silurus glanis*): a case study from Italy. Italian J Freshwater Ichthyol 2:33–42

- Mitrano DM, Wohlleben W (2020) Microplastic regulation should be more precise to incentivize both innovation and environmental safety. Nat Commun 11(1):1–12
- Moslen M, Miebaka CA, Boisa N (2019) Bioaccumulation of Polycyclic Aromatic Hydrocarbon (PAH) in a bivalve (Arca senilis-blood cockles) and health risk assessment. Toxicol Rep 6:990–997
- Needham LL, Calafat AM, Barr DB (2007) Uses and issues of biomonitoring. Int J Hyg Environ Health 210(3–4):229–238
- Nhan DD, Carvalho FP, Am NM, Tuan NQ, Yen NTH, Villeneuve JP, Cattini C (2001) Chlorinated pesticides and PCBs in sediments and molluscs from freshwater canals in the Hanoi region. Environ Pollut 112(3):311–320
- Nisbet IC, Lagoy PK (1992) Toxic equivalency factors (TEFs) for polycyclic aromatic hydrocarbons (PAHs). Regul Toxicol Pharmacol 16(3):290–300
- Oehlmann J, Schulte-Oehlmann U (2003) Molluscs as bioindicators. In: Markert BA, Breure AM, Zechmeister HG (eds) Bioindicators and biomonitors—principles, concepts and applications. Elsevier, Amsterdam, Lausanne, New York, pp 577–635
- Orosun MM, Adewuyi AD, Salawu NB, Isinkaye MO, Orosun OR, Oniku AS (2020) Monte Carlo approach to risk assessment of heavy metals at automobile spare part and recycling market in Ilorin. Nigeria Sci Rep 1:22084
- Orosun MM (2021) Assessment of arsenic and its associated health risks due to mining activities in parts of North-Central Nigeria: probabilistic approach using Monte Carlo. J Hazard Mater 412:125262
- Oziolor EM, Apell JN, Winfield ZC, Back JA, Usenko S, Matson CW (2018) Polychlorinated biphenyl (PCB) contamination in Galveston Bay, Texas: Comparing concentrations and profiles in sediments, passive samplers, and fish. Environ Pollut 236:609–661
- Pastorino P, Bertoli M, Squadrone S, Brizio P, Piazza G, Oss Noser AG, Prearo M, Abete MC, Pizzul E (2019) Detection of trace elements in freshwater macrobenthic invertebrates of different functional feeding guilds: A case study in Northeast Italy. Ecohydrol Hydrobiol 19(3):428–440
- Pastorino P, Pizzul E, Bertoli M, Anselmi S, Kušće M, Menconi V, Prearo M, Renzi M (2021) First insights into plastic and microplastic occurrence in biotic and abiotic compartments, and snow from a high-mountain lake (Carnic Alps). Chemosphere 265:129121
- Pastorino P, Prearo M, Anselmi S, Menconi V, Bertoli M, Dondo A, Pizzul E, Renzi M (2021) Use of the Zebra Mussel *Dreissena polymorpha* (Mollusca, Bivalvia) as a bioindicator of microplastics pollution in freshwater ecosystems: a case study from Lake Iseo (North Italy). Water 13(4):434
- Pastorino P, Prearo M, Bertoli M, Abete MC, Dondo A, Salvi G, Zaccaroni A, Elia AC, Pizzul E (2020) Accumulation of As, Cd, Pb, and Zn in sediment, chironomids and fish from a high-mountain lake: First insights from the Carnic Alps. Sci Total Environ. 729:139007
- Pastorino P, Zaccaroni A, Doretto A, Falasco E, Silvi M, Dondo A, Elia AC, Prearo M, Bona F (2020) Functional feeding groups of aquatic insects influence trace element accumulation: findings for filterers, scrapers and predators from the Po basin. Biology 9(9):288
- Piccardo M, Provenza F, Grazioli E, Cavallo A, Terlizzi A, Renzi M (2020) PET microplastics toxicity on marine key species is influenced by pH, particle size and food variations. Sci Total Environ 715:136947
- Shahid M, Nadeem M, Bakhat HF (2020) Environmental toxicology and associated human health risks. Environ Sci Pollut Res 27:39671–39675
- Squadrone S, Favaro L, Prearo M, Vivaldi B, Brizio P, Abete MC (2013) NDL-PCBs in muscle of the European catfish (*Silurus glanis*): an alert from Italian rivers. Chemosphere 93(3):521–525
- Storelli MM (2008) Potential human health risks from metals (Hg, Cd, and Pb) and polychlorinated biphenyls (PCBs) via seafood consumption: estimation of target hazard quotients (THQs) and toxic equivalents (TEQs). Food Chem Toxicol 46(8):2782–2788
- Strong EE, Gargominy O, Ponder WF, Bouchet P (2007) Global diversity of gastropods (Gastropoda; Mollusca) in freshwater. Freshwater animal diversity assessment. Springer, Dordrecht, pp 149–166
- Tietze E, De Francesco CG (2010) Environmental significance of freshwater mollusks in the Southern Pampas, Argentina: to what detail can local environments be inferred from mollusk composition? Hydrobiologia 641(1):133–143
- Tongo I, Ogbeide O, Ezemonye L (2017) Human health risk assessment of polycyclic aromatic hydrocarbons (PAHs) in smoked fish species from markets in Southern Nigeria. Toxicol Rep 4:55– 61

USEPA (1983) Methods for chemical analysis of water and waste. EPA Report 600/4-79-020. Cincinnati: Office of Water, USEPA.

- USEPA (1993) Provisional Guidance for Quantitative Risk Assessment of Polycyclic Aromatic Hydrocarbons. EPA/600/R-93/089, U.S. Environmental Protection Agency Office of Research and Development, Washington, DC
- USEPA (2000a) Risk-Based Concentration Table. Philadelphia PA: USEPA, Washington, DC.
- USEPA (2000b) Guidance for Assessing Chemical Contaminant. Data for Use in Fish Advisories. Fish Sampling and Analysis (3rd ed.), Office of Water, Washington DC.
- USEPA (2021) U.S. Environmental Protection Agency. Integrated

Risk Information System. pp. 14–19. [https:// www. epa. gov/ iris/.](https://www.epa.gov/iris/) Accessed 20 April 2021

- USFDA (2011) Fish and fisheries products hazards and controls guidance, third ed. Center for food safety and applied nutrition. US Food and Drug Administration.
- Wang W, Wang J (2018) Different partition of polycyclic aromatic hydrocarbon on environmental particulates in freshwater: microplastics in comparison to natural sediment. Ecotoxicol Environ Saf 147:648–655
- WHO (1996) Health criteria other supporting information. Guidelines for drinking water quality, 2nd edn. WHO, Geneva, pp 31–388
- Yuan Z, Yao J, Wang F, Guo Z, Dong Z, Chen F, Hu Y, Sunahara G (2017) Potentially toxic trace element contamination, sources, and pollution assessment in farmlands, Bijie City, southwestern China. Environ Monit Assess 189(1):1–10
- Zeng F, Wei W, Li M, Huang R, Yang F, Duan Y (2015) Heavy metal contamination in rice-producing soils of Hunan province, China and potential health risks. Int J Environ Res Public Health 12(12):15584–15593
- Zhang QL, Guo J, Deng XY, Wang F, Chen JY, Lin LB (2019) Comparative transcriptomic analysis provides insights into the response to the benzo(a)pyrene stress in aquatic firefly (*Luciola leii*). Sci Total Environ 661:226–234



**Fig. 1** Location of sampling sites (site 1: Samminiatello; site 2: Fibbiana), Tuscany, central Italy. The study sites are shown in the insert



**Fig. 2** Probability and frequency plot of cancer risk due to *Sinotaia quadrata* consumption: **a** arsenic (As), **b** cadmium (Cd), **c** chromium (Cr), **d** lead (Pb)



**Table 1** Potentially toxic element concentrations (µg g−1) in *Sinotaia quadrata* from

Plus-minus values are the mean  $\pm$  standard deviation

**Table 2** Concentration of the six indicators of NDL-PCBs (28, 52, 101, 138, 153, 180) in *Sinotaia quadrata* from Samminiatello (site 1) and<br>Fibbiano (site 2)  $\int$ aita 2)

$F1001$ ana (site $\angle$ )								
			Site Lipid (%) PCB-28 PCB-52 PCB-101 PCB-138 PCB-153 PCB-180					$\Sigma_6$
								PCB.
	32		$\langle$ LOQ $\langle$ LOQ $\langle$ LOQ $1.64 \pm 0.08$ 1.43 $\pm 0.10$ $\langle$ LOQ					3.07
	32		$\leq$ LOQ 1.77 $\pm$ 0.05 $\leq$ LOQ 2.17 $\pm$ 0.07 1.77 $\pm$ 0.12 1.61 $\pm$ 0.07 7.32					

Data are reported as the mean of three pools  $(n = 10)$ ;  $\lt$  LOQ = concentration below limit of quantification (1.2 ng g<sup>-1</sup> w.w.). Plus-minus values are the mean  $\pm$  standard deviation

**Table 3** Concentration (μg kg−1 w.w.) of benzo[a]pyrene (B[a]P), benzo[a]anthracene (B[a]A), chrysene (Chr), and benzo[b]fluoranthene (B[b]FL) detected in *Sinotaia quadrata* from Samminiatello (site 1) and Fibbiana (site 2);  $\leq$  LOQ = concentration below the limit of quantification (0.5 μg k g<sup>-1</sup> w.w.)

Site	B[a]P	B[a]A	Chr	B[b]FL
		$\langle$ (LOQ) $0.5 \pm 0.02$ $\langle$ LOQ		
				LOO
$\mathcal{D}_{\mathcal{L}}$		$\langle$ (LOQ) $0.7 \pm 0.02$ $\langle$ LOQ		
				LOO

Plus-minus values are the mean  $\pm$  standard deviation



**Table 4** Microplastics detected in *Sinotaia quadrata* at the two sites (n = 5 per site) are reported as total items, items/ animal  $\pm$  standard deviation (SD), mean size ( $\mu$ m)  $\pm$  SD, shape, color, and chemical type

*PET* polyethylene terephthalate, *PP* polypropylene

Table 5 Estimated weekly intake (EDI; μg kg<sup>-1</sup> day<sup>-1</sup>), target hazard quotient (THQ), and total target hazard quotient (TTHQ; sum of each THQ obtained from each element) by the consumption of *Sinotaia quadrata* in an average and a high-level mollusk consumer

	Index Consumption As		C <sub>d</sub>	Cr	Cu	Hg	Pb	Zn	<b>TTHQ</b>
<b>EDI</b>	Average	0.10	0.0003	0.004	0.54	0.01	0.03	0.92	
	High	0.20	0.0007	0.008	1.09	0.03	0.06	1.84	
<b>THQ</b>	Average High	0.004		$0.002$ $0.000001$ $0.000000003$ $0.00001$ $0.00002$ $0.000009$ $0.000003$ $0.0026$ $0.0000003$ $0.000000005$ $0.00003$ $0.00004$ $0.00002$ $0.000006$ $0.0051$					

**Table 6** Average daily intake (ADI; mg k  $g^{-1}$  day<sup>-1</sup>) and incremental lifetime cancer risk (ILCR) associated with the consumption of *Sinotaia quadrata* in an average and a high-level mollusk consumer



Table 7 Average daily dose exposure (ADD; ng kg<sup>-1</sup> day<sup>-1</sup>) and hazard quotient (HQ) of NDL-PCBs (PCB-28, PCB-52, PCB-101, PCB-138, PCB-153, PCB-180) for an average and a high-level mollusk consumer

Index	Consumption PCB-28 PCB-52 PCB-101 PCB-138 PCB-153 PCB-180 $\Sigma$ 6 PCB HI								
<b>ADD</b>	Average High	0.31 0.61	0.38 0.76	0.31 0.61	0.49 0.97	0.41 0.82	0.36 0.72	2.24 4.49	
HQ	Average High	0.03 0.06	0.04 0.08	0.03 0.06	0.05 0.09	0.04 0.08	0.04 0.07	—	0.22 0.45

**Table 8** Dietary daily intake (DDI) (μg kg−1 day−1 b.w.) of benzo[a]pyrene (B[a]P), benzo[a]anthracene (B[a]A), chrysene (Chr), benzo[b] fluoranthene (B[b]FL), and PAH4 index (sum of B[a]P, B[a]A, Chr, B[b]FL) for an average and a high-level mollusk consumer



# Supplementary files



**Table S1.** Limit of Detection (LOD) ( $\mu$ g g<sup>-1</sup>), reference material values and percentages (%) of recovery for each element.

<b>PCB</b>	Quantification Qualification			
	$\text{Ion} \left( \text{m/z} \right)$	Ions (m/z)		
28	256	186		
		258		
52	292	220		
		290		
155	360	290		
(IS)		362		
101	326	254		
		328		
153	360	290		
		325		
138		290		
	360	325		
180	394	324		
		396		
198		358		
(IS)	430	393		

**Table S2**. MS transitions used for PCBs detection.

**Table S3.** Exposure parameters used to calculate the human health risk by the consumption of *Sinotaia* 



*quadrata*.

**Table S4.** Summary of the Monte Carlo simulation.



