

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Annihilation operators for exponential spaces in subdivision

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Conti C., Lopez-Urena S., Romani L. (2022). Annihilation operators for exponential spaces in subdivision. APPLIED MATHEMATICS AND COMPUTATION, 418, 1-7 [10.1016/j.amc.2021.126796].

Availability:

This version is available at: <https://hdl.handle.net/11585/840671> since: 2021-12-05

Published:

DOI: <http://doi.org/10.1016/j.amc.2021.126796>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Costanza Conti, Sergio López-Ureña and Lucia Romani, Annihilation operators for exponential spaces in subdivision, Applied Mathematics and Computation 418, 126796 (2022)

The final published version is available online at
<https://dx.doi.org/10.1016/j.amc.2021.126796>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Annihilation Operators for Exponential Spaces in Subdivision

Costanza Conti^{a,*}, Sergio López-Ureña^b, Lucia Romani^c

^a*Dip. di Ingegneria Industriale, Università di Firenze, Viale Morgagni 40/44, 50134 Firenze, Italy*

^b*Dep. de Matemàtiques, Universitat de València, Doctor Moliner Street 50, 46100, Burjassot, Valencia, Spain*

^c*Dip. di Matematica, Alma Mater Studiorum Università di Bologna, Piazza P.S. Donato 5, 40126 Bologna, Italy*

Abstract

We investigate properties of differential and difference operators annihilating certain finite-dimensional spaces of exponential functions in two variables that are connected to the representation of real-valued trigonometric and hyperbolic functions. Although exponential functions appear in a variety of contexts, the motivation behind this technical note comes from considering subdivision schemes where annihilation operators play an important role. Indeed, subdivision schemes with the capability of preserving exponential functions can be used to obtain an exact description of surfaces parametrized in terms of trigonometric and hyperbolic functions, and annihilation operators are useful to automatically detect the frequencies of such functions.

Keywords: Subdivision scheme; Exponential function preservation; Difference operator annihilating exponentials.

1. Introduction

In this technical note we investigate properties of differential and difference operators annihilating a particular type of spaces of exponential functions (for short exponential spaces). The motivation to study these operators comes from considering non-stationary subdivision schemes preserving exponential polynomials (see, e.g., [5, 11, 12, 19]), which are efficient iterative algorithms for the definition of curves and surfaces. Non-stationary subdivision schemes reproducing exponential polynomials, first proposed in [8], have been the core subject of numerous papers in the last two decades, and they have already proved to be very useful in applications like geometric modelling (see, e.g., [10, 16, 23]) as well as curve and surface design from gradient fields (see, e.g., [3, 6]). However, there are still several application contexts (like, e.g., isogeometric analysis, multigrid, image analysis or wavelets construction) that could further benefit from their use since, as yet, the mostly exploited subdivision schemes have been mainly the stationary ones (see, e.g., [15, 18] and [4] for the applications of subdivision in isogeometric analysis and multigrid, whereas see, e.g., [1, 13, 17] and [2] as to the use of subdivision in image analysis and wavelets construction, respectively).

Up to now, in all subdivision schemes reproducing exponential polynomials, the exponential frequencies and their multiplicities are explicitly involved in the refinement rules. This is a major drawback since, usually, there is no a priori knowledge of them from the initial data. Since a common machinery in designing linear, non-stationary subdivision schemes is the identification of both a difference operator that annihilates the function space of interest, and a subdivision operator that reproduces it [21], the aim of this technical note is twofold. On the one hand, we investigate properties of differential and difference operators annihilating the type of exponential spaces of major interest in subdivision; on the other hand, following the seminal

*Corresponding author

Email addresses: costanza.conti@unifi.it (Costanza Conti), sergio.lopez-urena@uv.es (Sergio López-Ureña), lucia.romani@unibo.it (Lucia Romani)

work [8], we show how to use them for the automatic detection of the unknown exponential frequencies. These ideas are already used in the univariate non-linear case (see [7]), where the derivation of subdivision rules that guarantee the preservation of exponential polynomials rely on the definition of an *annihilation operator* (also called *annihilator*) whose kernel consists of them. In [7], the construction of an interpolatory, non-linear, stationary subdivision scheme capable of reproducing functions in

$$\text{span}\{1, \exp(\gamma z), \exp(-\gamma z)\}, \quad \text{where } \gamma \in \mathbb{G} \setminus \{0\}, \quad \text{and } \mathbb{G} := \mathbb{R}_{\geq 0} \cup i(0, \pi),$$

is indeed based on an annihilation operator. For $f : \mathbb{R} \rightarrow \mathbb{C}$, this annihilator is obtained by the repeated application of the differential operator $D^\gamma f(z) = f'(z) - \gamma f(z)$, $z \in \mathbb{R}$. For the considered space, the annihilator is $D^0 D^\gamma D^{-\gamma} f(z) = f'''(z) - \gamma^2 f'(z)$ since

$$D^0 D^\gamma D^{-\gamma} f(z) = 0, \quad \forall z \in \mathbb{R} \quad \Leftrightarrow \quad f \in \text{span}\{1, \exp(\gamma z), \exp(-\gamma z)\} \quad \text{with } \gamma \in \mathbb{G} \setminus \{0\}.$$

Replacing the differential operator D^γ with the difference operator Δ_t^γ defined as $\Delta_t^\gamma f(z) = f(z+t) - \exp(\gamma t)f(z)$, $t, z \in \mathbb{R}$, the discrete operator $\Delta_t^0 \Delta_t^\gamma \Delta_t^{-\gamma}$ turns out to be a discrete version of the annihilator $D^0 D^\gamma D^{-\gamma}$ in the sense that

$$\Delta_t^0 \Delta_t^\gamma \Delta_t^{-\gamma} f(z) = 0, \quad \forall t, z \in \mathbb{R} \quad \Leftrightarrow \quad f \in \text{span}\{1, \exp(\gamma z), \exp(-\gamma z)\}, \quad \gamma \in \mathbb{G} \setminus \{0\}. \quad (1)$$

If applied to a *discrete* function $\mathbf{f}^k := \{f(2^{-k}\alpha), \alpha \in \mathbb{Z}\}$, $k \in \mathbb{N}$, basic ingredient of any subdivision scheme, the identity in (1), for $t = 2^{-k}$ and $z = 2^{-k}(\alpha - 1)$, yields an equation satisfied for all $\alpha \in \mathbb{Z}$ and $f \in \text{span}\{1, \exp(\gamma z), \exp(-\gamma z)\}$, $\gamma \in \mathbb{G} \setminus \{0\}$, that reads as

$$f(2^{-k}(\alpha - 1)) - (2 \cosh(2^{-k}\gamma) + 1)f(2^{-k}\alpha) + (2 \cosh(2^{-k}\gamma) + 1)f(2^{-k}(\alpha + 1)) - f(2^{-k}(\alpha + 2)) = 0. \quad (2)$$

Equation (2) can be used to compute $\cosh(2^{-k}\gamma)$, that is to identify γ , and to set the subdivision refinement rules able to reproduce the above-mentioned space of exponential functions (see [7] for all details).

The goal of this work is to investigate a similar idea for the bivariate case. We remark that we are aware that differential/difference operators associated with exponential functions are of interest in several domains of mathematics (other than subdivision schemes) and that we are omitting many related works. For example, [20] provides a complete characterization of exponential polynomials using finite difference operators in the very general but certainly more involved context of exponential Box-splines. But, here, our aim is to provide a simple and focused analysis of these operators helpful to understand the reproduction capabilities of bivariate subdivision schemes and related topics. Indeed, the results we here present on bivariate exponential spaces are complete and easy-to-follow, and make their understanding and use simple.

The rest of this paper consists of two main sections. Section 2 characterizes spaces of exponential functions as the kernel of an annihilation operator based on the repeated application of a particular differential operator. In Section 2 we also show how this differential operator can be replaced by a discrete operator, which is the generalization to the bivariate setting of the finite difference operator in (1), proposed for univariate functions in [7]. Section 3 shows how to apply annihilation operators in the context of bivariate subdivision schemes that reproduce an important class of exponential functions. Conclusions are drawn in Section 4.

2. Characterization of bivariate exponential functions via differential and difference operators

Let us start by defining the space of exponential functions we want to work with, relying on the standard column vector notation.

Definition 2.1. Let $n \in \mathbb{N}$, $\mathbb{D} := \mathbb{R} \cup i(-\pi, \pi)$,

$$\gamma^\ell \in \mathbb{D}^2, \quad \ell = 1, \dots, n \quad \text{with } \gamma^\ell \neq \gamma^j \quad \text{if } \ell \neq j, \quad \text{and } \Gamma := \{\gamma^1, \dots, \gamma^n\}. \quad (3)$$

The space of exponential functions associated to the set of n distinct frequencies Γ is

$$E_\Gamma := \text{span} \left\{ \exp((\gamma^1)^T \mathbf{z}), \exp((\gamma^2)^T \mathbf{z}), \dots, \exp((\gamma^n)^T \mathbf{z}) : \gamma^\ell \in \Gamma, \ell = 1, 2, \dots, n \right\},$$

where $\mathbf{z} \in \mathbb{R}^2$ is the function variable.

Definition 2.2. Let $F : \mathbb{R}^2 \rightarrow \mathbb{C}$ be a differentiable function and let $\nabla F(\mathbf{z}) = \left(\frac{\partial F}{\partial z_1}, \frac{\partial F}{\partial z_2} \right)^T$. Given $\mathbf{v} \in \mathbb{R}^2 \setminus \{\mathbf{0}\}$ and $\gamma \in \mathbb{D}^2$, we define the differential operator

$$\mathcal{D}_{\mathbf{v}}^{\gamma} F(\mathbf{z}) := (\nabla F(\mathbf{z}) - \gamma F(\mathbf{z}))^T \mathbf{v}, \quad \mathbf{z} \in \mathbb{R}^2. \quad (4)$$

Our aim is first to show that the exponential functions in E_{Γ} are in the kernel of the differential operator $\mathcal{D}_{\mathbf{v}^1}^{\gamma_1} \mathcal{D}_{\mathbf{v}^2}^{\gamma_2} \dots \mathcal{D}_{\mathbf{v}^n}^{\gamma_n}$, where $\mathbf{v}^1, \mathbf{v}^2, \dots, \mathbf{v}^n \in \mathbb{R}^2 \setminus \{\mathbf{0}\}$ is an arbitrary set of n non-zero directions. Second, to show that such exponential functions are also in the kernel of the discrete version of $\mathcal{D}_{\mathbf{v}^1}^{\gamma_1} \mathcal{D}_{\mathbf{v}^2}^{\gamma_2} \dots \mathcal{D}_{\mathbf{v}^n}^{\gamma_n}$.

Remark 2.1. Let $\mathbf{v} = (v_1, v_2)^T \in \mathbb{R}^2 \setminus \{\mathbf{0}\}$ and $|\mathbf{v}| := \sqrt{\mathbf{v}^T \mathbf{v}} = \sqrt{v_1^2 + v_2^2}$. It is easy to show that

$$\mathcal{D}_{\mathbf{v}}^{\gamma} F(\mathbf{z}) = 0 \Leftrightarrow \mathcal{D}_{\mathbf{v}/|\mathbf{v}|}^{\gamma} F(\mathbf{z}) = 0, \quad \text{for all } \mathbf{z} \in \mathbb{R}^2.$$

Thus, from now on, we continue by assuming \mathbf{v} to be a unit vector of \mathbb{R}^2 and by writing $\mathbf{v} \in \mathbb{U} := \{\mathbf{u} \in \mathbb{R}^2 : |\mathbf{u}| = 1\}$.

We go on with two preliminary results needed as basic steps of an induction argument used to prove Theorem 2.4. The latter discusses solutions to differential equations of the type $\mathcal{D}_{\mathbf{v}^1}^{\gamma_1} \mathcal{D}_{\mathbf{v}^2}^{\gamma_2} \dots \mathcal{D}_{\mathbf{v}^n}^{\gamma_n} F = 0$ where $\mathbf{v}^1, \mathbf{v}^2, \dots, \mathbf{v}^n \in \mathbb{U}$ is an arbitrary set of n unit vectors, and shows that such solutions are independent of the vectors $\mathbf{v}^1, \mathbf{v}^2, \dots, \mathbf{v}^n$.

The first preliminary result considers the homogeneous case.

Proposition 2.2. Let $\mathbf{v} \in \mathbb{U}$ and $\gamma \in \mathbb{D}^2$ be such that $\gamma^T \mathbf{v} \neq 0$ and $\gamma^T \mathbf{v}^{\perp} \neq 0$ where $\mathbf{v}^{\perp} := (-v_2, v_1)^T$. Then,

$$\mathcal{D}_{\mathbf{v}}^{\gamma} F = 0, \quad \mathbf{w} \in \{\mathbf{v}, \mathbf{v}^{\perp}\} \quad \Leftrightarrow \quad F(\mathbf{z}) = c \exp(\gamma^T \mathbf{z}) \quad \text{with } c \in \mathbb{C}, \quad \mathbf{z} \in \mathbb{R}^2.$$

Proof. The implication \Leftarrow is easily verified. To prove \Rightarrow we first show that, for $\mathbf{z} = (z_1, z_2)^T$,

$$F(\mathbf{z}) = \kappa_1 (\mathbf{z}^T \mathbf{v}^{\perp}) \exp((\gamma^T \mathbf{v})(\mathbf{z}^T \mathbf{v})) = \kappa_2 (\mathbf{z}^T \mathbf{v}) \exp((\gamma^T \mathbf{v}^{\perp})(\mathbf{z}^T \mathbf{v}^{\perp})), \quad (5)$$

where $\kappa_1, \kappa_2 : \mathbb{R} \rightarrow \mathbb{C}$. Indeed, from the assumption $\mathcal{D}_{\mathbf{v}}^{\gamma} F = 0$ we deduce that the univariate function $G_{t_2}(t_1) := F(t_1 \mathbf{v} + t_2 \mathbf{v}^{\perp})$ satisfies the differential equation

$$G'_{t_2}(t_1) = (\gamma^T \mathbf{v}) G_{t_2}(t_1). \quad (6)$$

Similarly, from the assumption $\mathcal{D}_{\mathbf{v}^{\perp}}^{\gamma} F = 0$ we deduce that the univariate function $H_{t_1}(t_2) := F(t_1 \mathbf{v} + t_2 \mathbf{v}^{\perp})$ satisfies the differential equation

$$H'_{t_1}(t_2) = (\gamma^T \mathbf{v}^{\perp}) H_{t_1}(t_2). \quad (7)$$

Now, as well known, the set of solutions of the previous differential equations (6) and (7) are $\{\kappa_1(t_2) \exp(\gamma^T \mathbf{v} t_1) : \kappa_1(t_2) \in \mathbb{C}\}$ and $\{\kappa_2(t_1) \exp(\gamma^T \mathbf{v}^{\perp} t_2) : \kappa_2(t_1) \in \mathbb{C}\}$, respectively. Hence, by selecting $t_1 = \mathbf{z}^T \mathbf{v}$ and $t_2 = \mathbf{z}^T \mathbf{v}^{\perp}$, we arrive at

$$G_{\mathbf{z}^T \mathbf{v}^{\perp}}(\mathbf{z}^T \mathbf{v}) = \kappa_1(\mathbf{z}^T \mathbf{v}^{\perp}) \exp((\gamma^T \mathbf{v})(\mathbf{z}^T \mathbf{v})) \quad \text{and} \quad H_{\mathbf{z}^T \mathbf{v}}(\mathbf{z}^T \mathbf{v}^{\perp}) = \kappa_2(\mathbf{z}^T \mathbf{v}) \exp((\gamma^T \mathbf{v}^{\perp})(\mathbf{z}^T \mathbf{v}^{\perp})).$$

Taking into consideration that $F(\mathbf{z}) = G_{\mathbf{z}^T \mathbf{v}^{\perp}}(\mathbf{z}^T \mathbf{v}) = H_{\mathbf{z}^T \mathbf{v}}(\mathbf{z}^T \mathbf{v}^{\perp})$, (5) is shown. Next, taking $\mathbf{z} = t\mathbf{v}$, $t \in \mathbb{R}$, we obtain

$$\kappa_1(0) \exp(\gamma^T \mathbf{v} t) = \kappa_2(t),$$

where we used that $\mathbf{v} \in \mathbb{U}$. Denoting $c = \kappa_1(0)$, we can write (from the expression of $F(\mathbf{z})$ involving κ_2) that

$$F(\mathbf{z}) = c \exp((\gamma^T \mathbf{v})(\mathbf{z}^T \mathbf{v})) \exp((\gamma^T \mathbf{v}^{\perp})(\mathbf{z}^T \mathbf{v}^{\perp})) = c \exp((\gamma^T \mathbf{v})(\mathbf{v}^T \mathbf{z})) \exp((\gamma^T \mathbf{v}^{\perp})((\mathbf{v}^{\perp})^T \mathbf{z})).$$

Next we apply the additive property of exponentials and the distributivity of matrix operations, so obtaining

$$F(\mathbf{z}) = c \exp\left(\boldsymbol{\gamma}^T (\mathbf{v}\mathbf{v}^T + \mathbf{v}^\perp(\mathbf{v}^\perp)^T) \mathbf{z}\right).$$

Since $\mathbf{v}\mathbf{v}^T + \mathbf{v}^\perp(\mathbf{v}^\perp)^T = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, we arrive at $F(\mathbf{z}) = c \exp(\boldsymbol{\gamma}^T \mathbf{z})$, so concluding the proof. \square

The second preliminary result considers the non-homogeneous case and can be proven by standard arguments (look at [22, chapter 3], for example).

Lemma 2.3. *Let $\mathbf{v} \in \mathbb{U}$, $\boldsymbol{\gamma}^1, \boldsymbol{\gamma}^2, \dots, \boldsymbol{\gamma}^n$ be defined as in (3) and $(\boldsymbol{\gamma}^\ell - \boldsymbol{\gamma}^n)^T \mathbf{v} \neq 0$, $\ell = 1, \dots, n-1$. Then, for $c_\ell \in \mathbb{C}$, $\ell = 1, \dots, n-1$ there exist $d_\ell(\mathbf{v}) \in \mathbb{C}$, $\ell = 1, \dots, n-1$ such that*

$$\mathcal{D}_{\mathbf{v}}^{\boldsymbol{\gamma}^n} F(\mathbf{z}) = \sum_{\ell=1}^{n-1} c_\ell \exp((\boldsymbol{\gamma}^\ell)^T \mathbf{z}) \Rightarrow F(\mathbf{z}) = \sum_{\ell=1}^{n-1} d_\ell(\mathbf{v}) \exp((\boldsymbol{\gamma}^\ell)^T \mathbf{z}) + \kappa(\mathbf{z}^T \mathbf{v}^\perp) \exp(((\boldsymbol{\gamma}^n)^T \mathbf{v})(\mathbf{z}^T \mathbf{v}))$$

where $\kappa : \mathbb{R} \rightarrow \mathbb{C}$ is a particular function.

Exploiting Lemma 2.3, with an induction argument, we arrive at the sought result.

Theorem 2.4. *Given $\boldsymbol{\gamma}^1, \boldsymbol{\gamma}^2, \dots, \boldsymbol{\gamma}^n$ as in (3) we have that,*

$$\mathcal{D}_{\mathbf{v}^1}^{\boldsymbol{\gamma}^1} \mathcal{D}_{\mathbf{v}^2}^{\boldsymbol{\gamma}^2} \dots \mathcal{D}_{\mathbf{v}^n}^{\boldsymbol{\gamma}^n} F = 0, \quad \forall \mathbf{v}^1, \mathbf{v}^2, \dots, \mathbf{v}^n \in \mathbb{U} \Leftrightarrow F(\mathbf{z}) = \sum_{\ell=1}^n c_\ell \exp((\boldsymbol{\gamma}^\ell)^T \mathbf{z}) \text{ with } c_\ell \in \mathbb{C}, \mathbf{z} \in \mathbb{R}^2.$$

Proof. To show \Leftarrow we first exploit the linearity and then the commutativity of the differential operator, so obtaining

$$\mathcal{D}_{\mathbf{v}^1}^{\boldsymbol{\gamma}^1} \mathcal{D}_{\mathbf{v}^2}^{\boldsymbol{\gamma}^2} \dots \mathcal{D}_{\mathbf{v}^n}^{\boldsymbol{\gamma}^n} F(\mathbf{z}) = \sum_{\ell=1}^n c_\ell \mathcal{D}_{\mathbf{v}^1}^{\boldsymbol{\gamma}^1} \dots \mathcal{D}_{\mathbf{v}^\ell}^{\boldsymbol{\gamma}^\ell} \dots \mathcal{D}_{\mathbf{v}^n}^{\boldsymbol{\gamma}^n} \exp((\boldsymbol{\gamma}^\ell)^T \mathbf{z}) = \sum_{\ell=1}^n c_\ell \mathcal{D}_{\mathbf{v}^1}^{\boldsymbol{\gamma}^1} \dots \mathcal{D}_{\mathbf{v}^n}^{\boldsymbol{\gamma}^n} \mathcal{D}_{\mathbf{v}^\ell}^{\boldsymbol{\gamma}^\ell} \exp((\boldsymbol{\gamma}^\ell)^T \mathbf{z}) = 0.$$

To prove \Rightarrow we use induction on n . Proposition 2.2 proves the case $n = 1$. For a general n , let \mathbf{v} be a unit vector such that $\mathbf{v}^T(\boldsymbol{\gamma}^\ell - \boldsymbol{\gamma}^n) \neq 0$, $\ell = 1, 2, \dots, n-1$. Applying the induction hypothesis, we have

$$\mathcal{D}_{\mathbf{v}^1}^{\boldsymbol{\gamma}^1} \mathcal{D}_{\mathbf{v}^2}^{\boldsymbol{\gamma}^2} \dots \mathcal{D}_{\mathbf{v}^{n-1}}^{\boldsymbol{\gamma}^{n-1}} \mathcal{D}_{\mathbf{v}}^{\boldsymbol{\gamma}^n} F(\mathbf{z}) = 0, \quad \forall \mathbf{v}^1, \mathbf{v}^2, \dots, \mathbf{v}^{n-1} \in \mathbb{U} \Rightarrow \mathcal{D}_{\mathbf{v}}^{\boldsymbol{\gamma}^n} F(\mathbf{z}) = \sum_{\ell=1}^{n-1} c_\ell(\mathbf{v}) \exp((\boldsymbol{\gamma}^\ell)^T \mathbf{z}),$$

where the dependence on \mathbf{v} of the coefficients $c_\ell(\mathbf{v})$ is due to the fact that, for each value of \mathbf{v} , $\mathcal{D}_{\mathbf{v}}^{\boldsymbol{\gamma}^n} F$ is a different function. By Lemma 2.3 there exists $\kappa : \mathbb{R} \rightarrow \mathbb{C}$ such that

$$F(\mathbf{z}) = \sum_{\ell=1}^{n-1} d_\ell(\mathbf{v}) \exp((\boldsymbol{\gamma}^\ell)^T \mathbf{z}) + \kappa(\mathbf{z}^T \mathbf{v}^\perp) \exp(((\boldsymbol{\gamma}^n)^T \mathbf{v})(\mathbf{z}^T \mathbf{v})). \quad (8)$$

Setting $\mathbf{v}^1 = \mathbf{v}^2 = \dots = \mathbf{v}^{n-1} = \mathbf{v}$, we have

$$\mathcal{D}_{\mathbf{v}}^{\boldsymbol{\gamma}^1} \mathcal{D}_{\mathbf{v}}^{\boldsymbol{\gamma}^2} \dots \mathcal{D}_{\mathbf{v}}^{\boldsymbol{\gamma}^{n-1}} F(\mathbf{z}) = \kappa(\mathbf{z}^T \mathbf{v}^\perp) \exp(((\boldsymbol{\gamma}^n)^T \mathbf{v})(\mathbf{z}^T \mathbf{v})) \prod_{\ell=1}^{n-1} (\boldsymbol{\gamma}^\ell - \boldsymbol{\gamma}^n)^T \mathbf{v}. \quad (9)$$

Exploiting the commutativity property and using the induction for $n = 1$, we obtain

$$\mathcal{D}_{\mathbf{w}}^{\boldsymbol{\gamma}^n} \left(\mathcal{D}_{\mathbf{v}}^{\boldsymbol{\gamma}^1} \mathcal{D}_{\mathbf{v}}^{\boldsymbol{\gamma}^2} \dots \mathcal{D}_{\mathbf{v}}^{\boldsymbol{\gamma}^{n-1}} F(\mathbf{z}) \right) = 0, \quad \forall \mathbf{w} \in \mathbb{U} \implies \mathcal{D}_{\mathbf{v}}^{\boldsymbol{\gamma}^1} \mathcal{D}_{\mathbf{v}}^{\boldsymbol{\gamma}^2} \dots \mathcal{D}_{\mathbf{v}}^{\boldsymbol{\gamma}^{n-1}} F(\mathbf{z}) = C(\mathbf{v}) \exp((\boldsymbol{\gamma}^n)^T \mathbf{z}),$$

where $C(\mathbf{v}) \in \mathbb{C}$ is again dependent on \mathbf{v} since we apply the induction for each choice of a preliminary fixed \mathbf{v} . Comparing (9) with the last expression of $\mathcal{D}_{\mathbf{v}}^{\gamma^1} \mathcal{D}_{\mathbf{v}}^{\gamma^2} \dots \mathcal{D}_{\mathbf{v}}^{\gamma^{n-1}} F(\mathbf{z})$ we find that,

$$\kappa(\mathbf{z}^T \mathbf{v}^\perp) \exp((\gamma^n)^T \mathbf{z}) = c_n(\mathbf{v}) \exp((\gamma^n)^T \mathbf{z}), \quad \text{for some } c_n(\mathbf{v}) \in \mathbb{C}.$$

Thus, replacing the latter in (8), we conclude that $F(\mathbf{z}) = \sum_{\ell=1}^n c_\ell(\mathbf{v}) \exp((\gamma^\ell)^T \mathbf{z})$, with $c_\ell(\mathbf{v}) = d_\ell(\mathbf{v})$, $\ell = 1, \dots, n-1$. To deduce that c_ℓ , $\ell = 1, \dots, n$, do not depend on \mathbf{v} , we take an arbitrary \mathbf{v} inside the open set $\{\mathbf{w} \in \mathbb{U} : \mathbf{w}^T(\gamma^\ell - \gamma^n) \neq 0, \ell = 1, \dots, n-1\}$, and differentiate F with respect to \mathbf{v} , providing

$$0 = \sum_{\ell=1}^n (\nabla c_\ell(\mathbf{v})) \exp((\gamma^\ell)^T \mathbf{z}).$$

From the latter and the linear independence of $\{\exp((\gamma^\ell)^T \mathbf{z})\}_{\ell=1}^n$, we deduce that c_ℓ , $\ell = 1, \dots, n$, are indeed independent of \mathbf{v} . \square

The annihilation operator obtained via the differential operator in (4) admits a discrete analogue based on the repeated application of the difference operator defined next. Note that the presence of the scalar t is to deal with non-unit vectors.

Definition 2.3. Given $F : \mathbb{R}^2 \rightarrow \mathbb{C}$, $\mathbf{v} \in \mathbb{U}$, $\gamma \in \mathbb{D}^2$, we define the difference operator

$$\Delta_{t\mathbf{v}}^\gamma F(\mathbf{z}) := F(\mathbf{z} + t\mathbf{v}) - \exp(\gamma^T \mathbf{v} t) F(\mathbf{z}), \quad \mathbf{z} \in \mathbb{R}^2, \quad t \in \mathbb{R}_+.$$

In the next theorem we show that $\mathcal{D}_{\mathbf{v}}^\gamma$ can be replaced by $\Delta_{t\mathbf{v}}^\gamma$, $t \in \mathbb{R}_+$, to characterize functions in E_Γ .

Theorem 2.5. Given $\gamma^1, \gamma^2, \dots, \gamma^n$ as in (3) we have that

$$\Delta_{t_1 \mathbf{v}^1}^{\gamma^1} \Delta_{t_2 \mathbf{v}^2}^{\gamma^2} \dots \Delta_{t_n \mathbf{v}^n}^{\gamma^n} F = 0, \quad \forall t_i \mathbf{v}^i \in \mathbb{R}^2 \setminus \{\mathbf{0}\}, \quad i = 1, \dots, n \quad \Leftrightarrow \quad F(\mathbf{z}) = \sum_{\ell=1}^n c_\ell \exp((\gamma^\ell)^T \mathbf{z}) \quad \text{with } c_\ell \in \mathbb{C}, \mathbf{z} \in \mathbb{R}^2.$$

Proof. We will prove that

$$\Delta_{t_1 \mathbf{v}^1}^{\gamma^1} \Delta_{t_2 \mathbf{v}^2}^{\gamma^2} \dots \Delta_{t_n \mathbf{v}^n}^{\gamma^n} F = 0 \quad \text{if and only if} \quad \mathcal{D}_{\mathbf{v}^1}^{\gamma^1} \mathcal{D}_{\mathbf{v}^2}^{\gamma^2} \dots \mathcal{D}_{\mathbf{v}^n}^{\gamma^n} F(\mathbf{z}) = 0, \quad \forall t_i \mathbf{v}^i \in \mathbb{R}^2 \setminus \{\mathbf{0}\}, \quad i = 1, \dots, n. \quad (10)$$

The left implication can be easily proven with the help of Theorem 2.4, while for the right implication we use an induction argument. For $n = 1$, we know that $\Delta_{t\mathbf{v}}^\gamma F = 0$ means $F(\mathbf{z} + t\mathbf{v}) = \exp(\gamma^T \mathbf{v} t) F(\mathbf{z})$ from which we get

$$\nabla F(\mathbf{z})^T \mathbf{v} = \lim_{t \rightarrow 0} t^{-1} (F(\mathbf{z} + t\mathbf{v}) - F(\mathbf{z})) = F(\mathbf{z}) \lim_{t \rightarrow 0} t^{-1} (\exp(\gamma^T \mathbf{v} t) - 1) = F(\mathbf{z}) \gamma^T \mathbf{v},$$

and therefore $\mathcal{D}_{\mathbf{v}}^\gamma F(\mathbf{z}) = (\nabla F(\mathbf{z}) - \gamma F(\mathbf{z}))^T \mathbf{v} = 0$. For $n > 1$, we use the easy-to-check commutativity of the differential and difference operators, i.e.,

$$\mathcal{D}_{\mathbf{v}}^\gamma \Delta_{t\mathbf{w}}^\mu = \Delta_{t\mathbf{w}}^\mu \mathcal{D}_{\mathbf{v}}^\gamma, \quad \forall \gamma, \mu \in \mathbb{D}^2, \quad \mathbf{v}, \mathbf{w} \in \mathbb{U}, \quad t \in \mathbb{R}_+.$$

From above we see by induction that for $\mathbf{v}^1, \mathbf{v}^2, \dots, \mathbf{v}^n \in \mathbb{U}$ and $\gamma^1, \gamma^2, \dots, \gamma^n$ as in (3),

$$\mathcal{D}_{\mathbf{v}^1}^{\gamma^1} \mathcal{D}_{\mathbf{v}^2}^{\gamma^2} \dots \mathcal{D}_{\mathbf{v}^n}^{\gamma^n} F = 0 \quad \Leftrightarrow \quad \Delta_{t_1 \mathbf{v}^1}^{\gamma^1} \Delta_{t_2 \mathbf{v}^2}^{\gamma^2} \dots \Delta_{t_n \mathbf{v}^n}^{\gamma^n} F = 0, \quad \forall t_i \in \mathbb{R}_+, \quad i = 1, \dots, n, \quad (11)$$

which concludes the proof. \square

Remark 2.6. Note that, when $t\mathbf{v} \in \mathbb{Z}^2$, the operator $\Delta_{t\mathbf{v}}^\gamma$ can also be applied to discrete data, as needed when we deal with subdivision schemes. Indeed, for the sequence $\mathbf{F} = \{F(\boldsymbol{\alpha}), \boldsymbol{\alpha} \in \mathbb{Z}^2\}$ seen as a function from \mathbb{Z}^2 to \mathbb{C} , the action of $\Delta_{t\mathbf{v}}^\gamma$ reads as

$$\Delta_{t\mathbf{v}}^\gamma F(\boldsymbol{\alpha}) := F(\boldsymbol{\alpha} + t\mathbf{v}) - \exp(\gamma^T \mathbf{v} t) F(\boldsymbol{\alpha}), \quad \boldsymbol{\alpha} \in \mathbb{Z}^2, \quad t\mathbf{v} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}.$$

Note also that, for a fixed $\boldsymbol{\alpha}$, the repeated application of the operator $\Delta_{t\mathbf{v}}^\gamma$ with $t\mathbf{v} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}$, namely $\Delta_{t_1 \mathbf{v}^1}^{\gamma^1} \dots \Delta_{t_n \mathbf{v}^n}^{\gamma^n} F(\boldsymbol{\alpha})$, can be made local, i.e. involving only points of \mathbb{Z}^2 around the point $\boldsymbol{\alpha}$.

3. Applying annihilation operators in bivariate subdivision

We continue this technical note with a section discussing the use of bivariate, discrete annihilation operators that provide the natural generalization of the univariate, discrete annihilation operator in (1). Such operators are applied to discrete data and used to identify a special space of exponential functions of interest in subdivision. This is spanned by real-valued exponential functions obtained by assuming n odd and considering the symmetric set of frequencies

$$\Gamma = \{\mathbf{0}\} \cup \{\gamma^\ell, -\gamma^\ell\}_{\ell=1, \dots, \frac{n-1}{2}} \quad \text{with} \quad \gamma^\ell \in \mathbb{G}^2 \setminus \{\mathbf{0}\} \quad \text{and} \quad \mathbb{G} := \mathbb{R}_{\geq 0} \cup i(0, \pi).$$

Specifically, in the following we focus our attention on $\Gamma = \{\mathbf{0}, \gamma, -\gamma, \tilde{\gamma}, -\tilde{\gamma}\}$ with $\gamma := (\gamma_1, \gamma_2)^T$, $\tilde{\gamma} := (\gamma_1, -\gamma_2)^T$ and $\gamma, \tilde{\gamma} \in \mathbb{G}^2 \setminus \{\mathbf{0}\}$, since this set of frequencies is the one needed to reproduce spheres, hyperbolic paraboloids and other quadric surfaces. In fact, the coordinate components of the parametric representations of such surfaces are bivariate functions in

$$E_\Gamma = \text{span}\{1, \exp(\gamma^T \mathbf{z}), \exp(-\gamma^T \mathbf{z}), \exp(\tilde{\gamma}^T \mathbf{z}), \exp(-\tilde{\gamma}^T \mathbf{z})\}, \quad \gamma, \tilde{\gamma} \in \mathbb{G}^2 \setminus \{\mathbf{0}\}. \quad (12)$$

Remark 3.1. For later use we observe that, if E_Γ is the space of exponential functions in (12), and a bivariate function $F \in E_\Gamma$ assumes a constant value at 5 distinct points of \mathbb{Z}^2 that correspond to the vertices of the pair of grey triangles displayed in each illustration of Figure 1, then F is a constant function that can be associated to the frequency $\gamma = \mathbf{0}$.




Figure 1: Pairs of triangles with vertices in \mathbb{Z}^2 used to check whether the function $F \in E_\Gamma$ is constant.

We have already mentioned in the Introduction that, to model parametric surfaces like spheres, hyperbolic paraboloids and quadric surfaces, we can conveniently use subdivision schemes. Roughly speaking, subdivision schemes are efficient iterative methods capable of generating surfaces from samples of a function on \mathbb{Z}^2 by means of the repeated application of simple and local refinement rules. At level k , such rules use values on the grid $2^{-k}\mathbb{Z}^2$ to generate values on the grid $2^{-k-1}\mathbb{Z}^2$, and so on (see for example the recent survey [5]). As it is well known, the refinement rules of subdivision schemes reproducing exponential polynomials are not only level-dependent, but also explicitly involving the exponential frequencies γ (see, e.g., [7, 8, 11, 16]). This is a major drawback since, usually, there is no a priori knowledge of γ from the initial data. Hence, the development of techniques to automatically estimate γ from the given data are very important. The latter observation motivates a discussion on how discrete difference operators can be used as *annihilation operators* for the automatic detection of exponential frequencies. In the context of subdivision schemes it is also important to use annihilators that are *local* and involve the same sets of points -generally called *stencils*- used by the refinement rules of the scheme. To keep the exposition simple, we stay focused both on the space (12) (where $\Gamma = \{\mathbf{0}, \gamma, -\gamma, \tilde{\gamma}, -\tilde{\gamma}\}$ and whose importance is clear from the above discussion), and on the stencils of the *extended butterfly* subdivision scheme (see [9]). The refinement rules of the extended butterfly subdivision scheme involve points distributed as in Figure 2.

From Theorem 2.5 we deduce that, for $F \in E_\Gamma$ and $\alpha \in \mathbb{Z}^2$,

$$\Delta_{t_0 \mathbf{v}^0}^0 \Delta_{t_1 \mathbf{v}^1}^\gamma \Delta_{t_2 \mathbf{v}^2}^{-\gamma} \Delta_{t_3 \mathbf{v}^3}^{\tilde{\gamma}} \Delta_{t_4 \mathbf{v}^4}^{-\tilde{\gamma}} F(\alpha) = 0, \quad \forall \mathbf{v}^i \in \mathbb{U}, t_i \in \mathbb{R}_+ \text{ such that } t_i \mathbf{v}^i \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}, i = 0, \dots, 4. \quad (13)$$




Figure 2: From left to right: the horizontal, vertical and diagonal stencils of the extended butterfly subdivision scheme and the union of their points. The thicker edges correspond to locations where the new points will be inserted by the refinement rules.

Therefore, the operator $\mathcal{N}_{\alpha, t_0 \mathbf{v}^0, t_1 \mathbf{v}^1, t_2 \mathbf{v}^2, t_3 \mathbf{v}^3, t_4 \mathbf{v}^4}(F, \gamma) = \Delta_{t_0 \mathbf{v}^0}^0 \Delta_{t_1 \mathbf{v}^1}^\gamma \Delta_{t_2 \mathbf{v}^2}^{-\gamma} \Delta_{t_3 \mathbf{v}^3}^{\tilde{\gamma}} \Delta_{t_4 \mathbf{v}^4}^{-\tilde{\gamma}} F(\alpha)$ is an annihilation operator for E_Γ . Now, observe that for specific choices of $t_0 \mathbf{v}^0, t_1 \mathbf{v}^1, t_2 \mathbf{v}^2, t_3 \mathbf{v}^3, t_4 \mathbf{v}^4$, some finite differences coincide. For instance,

$$\Delta_{(1,0)^T}^\gamma = \Delta_{(1,0)^T}^{\tilde{\gamma}}, \quad \Delta_{(0,1)^T}^\gamma = \Delta_{(0,1)^T}^{-\tilde{\gamma}}, \quad \Delta_{(1,0)^T}^{-\gamma} = \Delta_{(1,0)^T}^{-\tilde{\gamma}} \quad \text{and} \quad \Delta_{(0,1)^T}^{-\gamma} = \Delta_{(0,1)^T}^{\tilde{\gamma}}. \quad (14)$$

Hence, it turns out that to annihilate E_Γ it is sufficient to consider just 3 finite difference operators, that is the annihilators

$$\mathcal{N}_{\alpha, t\mathbf{v}, \mathbf{e}, \mathbf{e}}(F, \gamma) = \Delta_{t\mathbf{v}}^0 \Delta_{\mathbf{e}}^\gamma \Delta_{-\mathbf{e}}^{-\gamma} F(\alpha), \quad \mathbf{e} \in \{(1,0)^T, (0,1)^T\}, \quad \alpha \in \mathbb{Z}^2, \quad t\mathbf{v} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}. \quad (15)$$

Note that $\mathcal{N}_{\alpha, t\mathbf{v}, \mathbf{e}, \mathbf{e}}(F, \gamma)$ involves function evaluations at $\alpha + \lambda t\mathbf{v} + \mu \mathbf{e}$ with $\lambda \in \{0, 1\}$, $\mu \in \{0, 1, 2\}$, i.e. at 6 points of \mathbb{Z}^2 at most, and this is certainly a great advantage compared with the 14 function evaluations that $\mathcal{N}_{\alpha, t_0 \mathbf{v}^0, t_1 \mathbf{v}^1, t_2 \mathbf{v}^2, t_3 \mathbf{v}^3, t_4 \mathbf{v}^4}(F, \gamma)$ requires. For the application of annihilators in the context of subdivision schemes this is a crucial feature. Now, expansion of $\mathcal{N}_{\alpha, t\mathbf{v}, \mathbf{e}, \mathbf{e}}(F, \gamma) = 0$ provides the equation

$$-2 \cosh(\gamma^T \mathbf{e}) \Delta_{t\mathbf{v}}^0 F(\alpha + \mathbf{e}) + \Delta_{t\mathbf{v}}^0 F(\alpha + 2\mathbf{e}) + \Delta_{t\mathbf{v}}^0 F(\alpha) = 0, \quad \alpha \in \mathbb{Z}^2, \quad (16)$$

from which, whenever $F \in E_\Gamma$ and $\Delta_{t\mathbf{v}}^0 F(\alpha + \mathbf{e}) \neq 0$, we can identify the unknown parameter γ . As a matter of fact, from (16) we can write

$$\cosh(\gamma^T \mathbf{e}) = \frac{\Delta_{t\mathbf{v}}^0 F(\alpha + 2\mathbf{e}) + \Delta_{t\mathbf{v}}^0 F(\alpha)}{2 \Delta_{t\mathbf{v}}^0 F(\alpha + \mathbf{e})}, \quad \forall \alpha \in \mathbb{Z}^2, \quad (17)$$

and, with the choices $\mathbf{e} = (1,0)^T$ and $\mathbf{e} = (0,1)^T$, we indeed univocally identify $\cosh(\gamma_1)$ and $\cosh(\gamma_2)$, respectively, and hence γ . Note that, in case $\Delta_{t\mathbf{v}}^0 F(\alpha + \mathbf{e}) = 0$, to make it non-zero we change $t\mathbf{v}$ with the restriction of involving only points belonging to the union of the three subdivision stencils of the extended butterfly scheme shown in the rightmost picture of Figure 2. According to Remark 3.1, for $\mathbf{e} = (1,0)^T$ and a fixed selection of α , there are two different sets of choices for $t\mathbf{v}$ that are enough to be considered (see Figure 1 where the vertex shared by the two triangles represents α). Precisely, as to the leftmost picture, the vectors are $t\mathbf{v} \in \{(0,1)^T, (1,1)^T, (0,-1)^T, (-1,-1)^T\}$ whereas, as to the rightmost picture, they are $t\mathbf{v} \in \{(1,0)^T, (1,1)^T, (-1,0)^T, (-1,-1)^T\}$, respectively. Therefore, if for a fixed selection of α , $\Delta_{t\mathbf{v}}^0 F(\alpha + \mathbf{e}) = 0$ for all the choices of $t\mathbf{v}$ in one of the two sets mentioned above, this means that F is constant, which anyway implies the identification of γ with $\mathbf{0}$. The same obviously holds for $\mathbf{e} = (0,1)^T$.

4. Closing remarks

This technical note investigates properties of differential and difference operators annihilating certain finite-dimensional spaces of bivariate exponential functions connected to the representation of real-valued

trigonometric and hyperbolic functions used in the context of subdivision schemes. The example illustrated in Section 3 has shown the benefits of such annihilation operators to identify the frequency γ involved in the definition of subdivision rules. For further details concerning the use of annihilators in subdivision schemes, we refer the interested reader to the follow-up of this theoretical work recently appeared in [14]. In that paper, annihilation operators as those in (15) are shown in action, not only to detect frequencies, but also to locally identify the correct subdivision rule to be applied. This demonstrates the important role that annihilators can play.

Acknowledgement

The first and the last author are members of INdAM-GNCS, Italy, which has partially supported this work. The last author is also member of the Alma Mater research center on Applied Mathematics (AM^2). The second author has been supported by grant MTM2017-83942 funded by Spanish MINEC, by grant PID2020-117211GB-I00 funded by MCIN/AEI/10.13039/501100011033 and by the scholarships FPU14/02216 and EST16/00133 by Spanish MEC.

References

- [1] Badoual, A., Schmitter, D., Uhlmann, V., Unser, M.: Multiresolution subdivision snakes. *IEEE Transactions on Image Processing* 26 (3), 1188-1201 (2016).
- [2] Charina, M., Conti, C., Cotronei, M., Sauer, T.: Bivariate two-band wavelets demystified. *Linear Algebra and Its Applications* 608, 13–36 (2021).
- [3] Charina, M., Conti, C., Mejsirik, T., Merrien, J.L.: Joint Spectral Radius and Ternary Hermite Subdivision. *Adv. Comput. Math.* 47, art. 25 (2021).
- [4] Charina, M., Donatelli, M., Romani, L., Turati, V.: Anisotropic bivariate subdivision with applications to multigrid. *Appl. Numer. Math.* 135, 333–366 (2019).
- [5] Conti, C., Dyn, N.: Non-stationary Subdivision Schemes: State of the Art and Perspectives. In: Fasshauer G.E., Neamtu M., Schumaker L.L. (eds) *Approximation Theory XVI. AT 2019. Springer Proceedings in Mathematics & Statistics*, vol 336. Springer, Cham. (2021).
- [6] Cotronei, M., Moosmüller, C., Sauer, T., Sissouno, N.: Level-Dependent Interpolatory Hermite Subdivision Schemes and Wavelets. *Constructive Approximation* 50(2), 341–366 (2019).
- [7] Donat, R., López-Ureña, S.: Nonlinear stationary subdivision schemes reproducing hyperbolic and trigonometric functions. *Adv. Comput. Math.* 45, 3137–3172 (2019).
- [8] Dyn, N., Levin, D., Luzzatto, A.: Exponentials reproducing subdivision schemes. *Found. Comput. Math.* 3(2), 187–206 (2003).
- [9] Hed, S.: Analysis of subdivision schemes for surface generation. M.Sc. thesis, Tel Aviv University (1992).
- [10] Jena, H., Jena, M.K.: A hybrid non-stationary subdivision scheme based on triangulation. *International Journal of Applied and Computational Mathematics* 7(4), art. 169 (2021).
- [11] Jeong, B., Kim, H.O., Lee, Y.J., Yoon, J.: Exponential polynomial reproducing property of non-stationary symmetric subdivision schemes and normalized exponential B-splines. *Adv. Comput. Math.* 38(3), 647–666 (2013).
- [12] Jeong, B., Lee, Y.J., Yoon, J.: A family of non-stationary subdivision schemes reproducing exponential polynomials. *J. Math. Anal. Appl.* 402(1), 207–219 (2013).
- [13] Kitslaar, P.H., van't Klooster, R., Staring, M., Lelieveldt, B.P.F. and van der Geest, R.J.: Segmentation of branching vascular structures using adaptive subdivision surface fitting. In: *Proceedings of the SPIE International Symposium on Medical Imaging: Image Processing (MI'15)*, vol. 9413, Orlando, FL, USA, February 21-26, 2015, art. 94133Z.
- [14] López-Ureña, S., Viscardi, A.: An annihilator-based strategy for the automatic detection of exponential polynomial spaces in subdivision. *Comput. Aided Geom. Design* 85, art. 101976 (2021).
- [15] Ma, Y., Ma, W.: A subdivision scheme for unstructured quadrilateral meshes with improved convergence rate for isogeometric analysis. *Graphical Models* 106, art. 101043 (2019).
- [16] Novara, P., Romani, L.: Building blocks for designing arbitrarily smooth subdivision schemes with conic precision. *J. Comput. Appl. Math.* 279, 67–79 (2015).
- [17] Orderud, F., Rabben, S.I.: Real-time 3D segmentation of the left ventricle using deformable subdivision surfaces. In: *Proceedings of the IEEE Conference of Computer Vision and Pattern Recognition (CVPR'08)*, Anchorage AK, USA, June 23-28, 2008, pp.1–8.
- [18] Pan, Q., Rabczuk, T., Yang, X.: Subdivision-based isogeometric analysis for second order partial differential equations on surfaces. *Comput Mech* 68, 1205—1221 (2021).
- [19] Romani, L.: Creating a bridge between cardinal Br-spline fundamental functions for interpolation and subdivision. *Appl. Math. Comput.* 401, art. 126071 (2021)
- [20] Ron, A.: Exponential box splines. *Constr. Approx.* 4(1), 357–378 (1988).

- [21] Sauer, T.: Kernels of discrete convolutions and application to stationary subdivision operators. *Acta Appl. Math.* 145, 115–131 (2016).
- [22] Sneddon, I.N.: *Elements of partial differential equations*, Dover Publications Inc., 2013.
- [23] Warren, J., Weimer, H.: *Subdivision Methods for Geometric Design*. Morgan Kaufmann, San Francisco, 2002.