

# Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Genetic characterization of Italian and Spanish wild and domesticated chestnut trees

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Alessandri S., Cabrer A.M.R., Martin M.A., Mattioni C., Pereira-Lorenzo S., Dondini L. (2022). Genetic characterization of Italian and Spanish wild and domesticated chestnut trees. SCIENTIA HORTICULTURAE, 295(March 2022), 1-8 [10.1016/j.scienta.2022.110882].

Availability:

This version is available at: https://hdl.handle.net/11585/849447 since: 2022-01-31

Published:

DOI: http://doi.org/10.1016/j.scienta.2022.110882

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

1	Genetic characterization	of Italian	and	Spanish	wild	and	domesticated	chestnut
2	trees							

- 3 Sara Alessandri<sup>1</sup>, Ana Maria Ramos Cabrer<sup>2</sup>, M. Angela Martìn<sup>3</sup>, Claudia Mattioni<sup>4</sup>, Santia-
- 4 go Pereira-Lorenzo<sup>2</sup>, Luca Dondini<sup>1</sup>
- <sup>5</sup> <sup>1</sup>DISTAL <sup>-</sup> Department of Agricultural and Food Sciences, University of Bologna, Bologna, Italy
- 6 <sup>2</sup>USC <sup>-</sup> Department of Crop Production and Projects of Engineering, University of Santiago de
- 7 Compostela, Lugo, Spain
- 8 <sup>3</sup>UCO Department of Genetics, University of Cordoba, Cordoba, Spain
- 9<sup>4</sup> IRET Research Institute on Terrestrial Ecosystems, National Research Council, Porano, Italy
- 10 Corresponding author: Sara Alessandri sara.alessandri4@unibo.it
- 11

### 12 Abstract

13 The only native species of the Castanea genus in Europe is Castanea sativa Mill., a widespread and 14 important multipurpose tree species in the Mediterranean area that provides fruit, wood and shelter 15 for hives. With the aim of expanding the knowledge of the genetic variability of the chestnut spe-16 cies (wild trees and varieties) and promoting the traceability of local products, an analysis based on 17 16 SSRs was carried out on 630 single trees from Italy and Spain. 319 unique genotypes were iden-18 tified. A Bayesian approach combined with the Markov Chain Monte Carlo (MCMC) simulation 19 method revealed the existence of two genetically distinct groups of chestnuts: Cluster 1 (Spain) and 20 Cluster 2 (Italy), with a clear separation between the cultivars from (northern and southern) Spain 21 and from Italy. The results also confirmed a common genetic structure between chestnut popula-22 tions from southern Spain and southern Italy, which is the result of historical events and long-term

human impact. The results showed no genetic differentiation between chestnut cultivars (grafted
trees) and wild chestnut trees, probably as a consequence of the proximity of orchards and natural
populations, which resulted in a gene flow between them.

26

Keyword: *Castanea sativa* Mill., microsatellite markers (SSRs), Genetic diversity, Germplasm
conservation, Structure analysis

29

#### 30 1 Introduction

The *Castanea sativa* Mill. species belongs to the *Fagaceae* family and is the only native species of the genus *Castanea* in Europe. Nowadays, its widespread distribution is the result of both natural and anthropogenic factors that affected the species over time (Conedera et al., 2004). Climatic conditions have influenced chestnut distribution since the Pleistocene; during this glacial period, the species was confined to limited and climatically stable areas called refugia (Krebs et al., 2004; Mattioni et al., 2013). Subsequently, during the postglacial period, the natural recolonization of chestnuts started from these areas.

However, chestnut distribution was also influenced by human colonization and migration. In southern Italy, chestnut trees were introduced by the ancient Greeks about 5,000 years ago. Later, the
Romans spread this species in Europe (Huntley and Birks, 1983; Bernetti, 1995; Krebs et al., 2004
and 2019; Roces-Diaz et al., 2018).

In the last 20 years, several studies have described in detail the fundamental role of the historical and natural processes that resulted through time in the establishment of the varietal genetic diversity of chestnuts that exists to this day (Martín et al., 2009; Pereira-Lorenzo et al., 2010, 2011; Marinoni et al., 2013; Lusini et al., 2014, Villani et al., 1999; Martín et al., 2012; Fernández-Cruz and Fernández-López, 2016; Mattioni et al., 2013 and 2017). While the genetic pool of the domesticated

47 trees was influenced by human selection, propagation and hybridization (to improve fruit or wood 48 quality traits, as well as resistance to abiotic and biotic stress), wild trees were subjected to natural 49 selection to adapt to different environmental and geographical conditions (Barrett and Schluter, 50 2008; Nishio et al., 2021). In particular, the selective pressure to which wild populations were sub-51 jected favored different allele pools involved in the trees' adaptation to different regional condi-52 tions, such as adaptation to drought (Soto et al., 2019; Alcaide et al., 2019; Castellana et al., 2021). 53 Furthermore, wild chestnut genetic variability has also been affected by the proximity of orchards, 54 mainly due to pollen diffusion and male-sterility, which have contributed to the inclusion of local 55 and nonlocal new alleles (Lopez et al., 2021). It is also worth mentioning the importance of non-56 grafted giant chestnuts as a reservoir of genetic diversity, which represent the basis from which the selection and cultivation process commenced (Pereira-Lorenzo et al., 2019). 57 58 Despite the different selection process, domesticated chestnuts have preserved most of the diversity 59 found in the oldest wild tree populations (non-grafted giant chestnuts; Pereira-Lorenzo et al., 2019). 60 However, the hybridization conducted by humans improved chestnut genetic variability. For in-61 stance, the crosses between C. sativa and the Asiatic species allowed the creation of new hybrid cul-62 tivars that are resistant to several pathogens (e.g., to the pathogenic fungus Gnomoniopsis pascoe 63 and the Chinese wasp Dryocosmus kuriphilus) and tolerant to biotic stress (Sartor et al., 2009 and 64 2015; Dini et al., 2012; Botta et al., 2012; Alcaide et al., 2020 and 2021). 65 Low differences in genetic variability have been observed between chestnut cultivars and wild trees 66 (Pereira-Lorenzo et al., 2019; Bouffartigue et al., 2019 and 2020). 67 Therefore, in this study, we assessed a genetic characterization among chestnut cultivars and wild 68 chestnut trees from the Iberian Peninsula and Italy. The main aims of this research were: a) to inves-69 tigate the gene pools of chestnut trees from Italy and the Iberian Peninsula based on the reference 70 SSR; b) to estimate the genetic diversity between wild chestnut trees and chestnut varieties.

#### 72 2 Materials and methods

73 2.1 Plant material

74 A total of 630 wild chestnut trees and chestnut varieties were analyzed: in particular, 244 were new-

75 ly collected samples and 386 were derived from previous studies (Table S1). More in detail, 520

samples were varieties and the remaining 110 derived from wild chestnut trees, as single and isolat-ed trees.

13 representative samples of *Castanea pumila, C. crenata, C. mollissima* and the 'Volos' cultivar

79 were added to the analysis in order to collect information on interspecific hybrids versus Italian and

80 Spanish chestnut samples.

81 The 630 genotypes were standardized with the SSR profiles of the unique accessions available in

82 the European Chestnut Database (Pereira-Lorenzo et al., 2017).

83

84 2.2 DNA extraction and PCR amplification of microsatellites (SSRs)

85 Young leaves used for DNA extraction were frozen in liquid nitrogen and stored at -80 ° C or ly-

86 ophilized. The extraction was performed on samples of 0.1-0.5 grams of fresh leaves previously

87 ground in liquid nitrogen, or on 5 mg of ground lyophilized leaves. The DNA was extracted follow-

88 ing the CTAB protocol developed by Maguire et al. (1994). dsDNA was quantified using a

89 Nanodrop<sup>TM</sup> ND-1000 Spectrophotometer (Thermo Scientific, Wilmington, DE, USA) and diluted

90 to 10 ng/μl.

91 The microsatellites chosen were selected from the CsCAT and EMCs series, and OAL (Marinoni et

92 al., 2003; Buck et al., 2003 and Gobbin et al., 2007) and QrZAG (Kampfer et al., 1998) series. The

93 primers were used in multiplex sets according to Pereira-Lorenzo et al. (2017).

94 The PCR reaction was performed in 10  $\mu$ l final volume containing 6.45  $\mu$ l of sterile H<sub>2</sub>O, 1  $\mu$ l of

95 GeneAmp<sup>®</sup> 10X reaction buffer, 0.8 μl of MgCl<sub>2</sub>, 0.25 of dNTPs and 0.1 units AmpliTaq Gold<sup>TM</sup> 4

DNA polymerase. Amplification products were sequenced on a 3130 Genetic Analyzer capillary
sequencer (Applied Biosystems, USA). The internal GeneScan TM size standard 500 LIZ (-250)
was included in each sample. The allele sizes were detected using Peak Scanner TM software (Applied Biosystems). The samples collected were suitably standardized with the alleles found in the
European dataset for the 16 SSR being assessed.

101

102 2.3 Cluster analysis

103 The genetic distance between the 630 samples was calculated by using the R software. The con-

104 struction of the genetic distance dendrogram was elaborated using the Unweighted Pair-Group

105 Method (UPGMA) by R software with the function 'hclust, method = 'average'', packages 'adege-

106 nett".

107

108 2.4 Genetic diversity and gene pools structure

The genetic diversity indices were assessed by the Cervus software, version 3.0.3 (Kalinowski et al., 2007): the number of alleles per locus (k), the expected and the observed heterozygosity ( $H_e$  and  $H_o$ ), polymorphism information content (PIC) and the probability of the allele null (F-null) were estimated. A PIC threshold of 0.7 was considered for defining loci as highly polymorphic and informative. The frequency of the null alleles (F-null) for each locus was calculated using the maximum likelihood (ML) estimator of Kalinowski (2007) implemented in Cervus. The Marker index (MI) was calculated with iMEC: Online Marker Efficiency Calculator (Amirvousefi et al., 2018).

116 The MI allowed the measurement of polymorphism information for the individual markers used.

117 The STRUCTURE 2.3.3 (Pritchard et al., 2000) software was used to evaluate the genetic pool sub-

118 division of 319 genotypes (varieties and wild trees) and to calculate the estimated membership coef-

119 ficient (Q-value) that indicates the membership of each individual in each cluster. This analysis was

120 conducted with a Bayesian approach combined with the Markov Chain Monte Carlo (MCMC) sim-

121 ulation method and was performed using an "admixture model" and correlated allele frequencies.

122 Following the protocol of Pereira-Lorenzo et al. (2019), 30 replicate runs of STRUCTURE were 123 performed by setting the number of clusters (K) from 1 to 15. Each run consisted of a burning peri-124 od of 200,000 steps followed by 200,000 MCMC replicates, with the usage options: locprior=0, 125 popinfo = 0, popflag = 0 (Pereira-Lorenzo et al., 2019; Porras-Hurtado et al., 2013). 126 In order to check if the inclusion of hybrid samples affected the STRUCTURE analysis, the hybrid 127 samples were removed, and a second analysis was performed on a total of 306 samples with the 128 same conditions described above (Figure S3). A Q threshold of 0.8 was used to infer an accession to 129 a specific cluster. The  $\Delta K$  value (defined as the most probable number of clusters in the population) 130 was calculated through Structure Harvester v.09.93 (Earl, 2012) by testing the change of the log-131 likelihood between K values ( $\Delta$ K) as described by Evanno (2005). If a sample has a Q-value < 0.8, 132 it is considered an admixed sample.

133

134 2.5 Genetic differentiation

135 To validate the genetic structure revealed by the Bayesian model-based clustering, a multivariate 136 Principal Coordinate Analysis (PCoA) was elaborated with GenAlEx version 6.502 (Peakall and 137 Smouse, 2006). The PCoA representation was determined on the genetic distance measured by Jac-138 card coefficient, based on the estimates of  $\Delta K$  from STRUCTURE for 306 samples (hybrid samples 139 were not considered). A set of analysis to estimate the population differentiation was conducted un-140 der four scenarios: a) the two main groups (Cluster 1 vs Cluster 2) resulting from the Structure 141 analysis; b) the sub-groups (K=3, K=4), c) chestnut varieties versus wild trees (200 vs 106, respec-142 tively) and d) chestnut varieties and the wild trees separated in the two main clusters (Cluster 1 and 143 Cluster 2). Pairwise F<sub>ST</sub> values and private alleles (Np) were estimated for the different partitioning 144 levels considered using GeneAlEx version 6.502; missing data were coded as 0. The F<sub>ST</sub> value 145 ranges between -1 (absent inbreeding, excess of heterozygous) and 1 (non-random reproduction, 146 excess homozygous).

The gene flow (Nm) was estimated for the different partitioning levels considered (K=2, 3 and 4) using GenAlEx version 6.502 (Peakall and Smouse, 2012). Hierarchical analysis of molecular variance (AMOVA) was implemented in the GeneAlEx version 6.502 (Peakall and Smouse, 2006) in order to evaluate the genetic variation among and within Clusters. Tests of significance were performed using 9999 permutations within the total dataset of 306 samples. The 13 hybrids were removed from this analysis.

#### 153 3 Results

154 3.1 Genetic variability of the 16 microsatellites

155 The 16 SSRs showed high levels of polymorphism and discriminating power and revealed a total of 156 212 alleles, with an average number of 13.25 alleles per locus (Table 1). The average PIC was 157 0.735, ranging between 0.879 for CsCAT3 and 0.593 for EMCs2. Furthermore, expected 158 heterozygosity varied between 0.889 for CsCAT3 and 0.619 for CsCAT1, with a mean value of 0.763 (Table 1). CsCAT41 was known to amplify two different genomic sites (A and B); for this 159 160 reason, the CsCAT41A locus was removed from the dataset before the analyses (Pereira-Lorenzo et 161 al., 2010). CsCAT2 and EMCs38 showed a high frequency value of null alleles (0.209 and 0.118, 162 respectively); consequently, these two loci were removed from the subsequent analysis.

163

- 164
- 165

166 Table 1: Number of alleles (k), observed (Ho) and expected (He) heterozygosity, polymorphic information content

(PIC), null alleles frequencies (F[null]) and marker index (MI) for 319 unique genotypes of *C. sativa* accessions evaluated with 16 SSRs.

Locus	k	Но	Не	PIC	F(Null)	MI
CsCAT41B	12	0.690	0.812	0.793	0.071	0.270

CsCAT16	12	0.777	0.812	0.785	0.018	0.262
CsCAT6	19	0.846	0.872	0.858	0.013	0.186
CsCAT1	16	0.596	0.619	0.601	0.022	0.197
CsCAT3	26	0.803	0.889	0.879	0.050	0.165
QrZAG96	12	0.596	0.718	0.694	0.082	0.311
EMCs15	7	0.599	0.661	0.605	0.053	0.385
EMCs38	19	0.693	0.883	0.871	0.118	0.204
EMCs2	6	0.624	0.661	0.593	0.030	0.427
EMCs22	10	0.652	0.682	0.654	0.020	0.301
CsCAT2	16	0.555	0.855	0.841	0.209	0.240
CsCAT17	13	0.771	0.844	0.827	0.042	0.284
CsCAT14	10	0.721	0.753	0.711	0.015	0.313
CsCAT15	11	0.665	0.666	0.605	-0.003	0.266
CsCAT8	11	0.727	0.843	0.821	0.068	0.341
OAL	12	0.586	0.651	0.629	0.048	0.268
Mean	13.25	0.681	0.763	0.735	0.005	0.257

- 169
- 170 3.2 Cluster analysis
- A dendrogram was constructed using the UPGMA method with R software to evaluate the genetic
  diversity and relatedness between the 630 wild and grafted chestnut trees.
- 173 Clustering according to the UPGMA method allowed the authentication of accessions (true-to-type
- 174 cultivars), as well as the indication of possible homonyms, synonyms and incorrect denominations.

175 The 630 samples corresponded to 319 unique chestnut genotypes (Figure S1).

- 176
- 177 3.3 Gene pools

178 The genetic structure of the 319 unique genotypes (including varieties and wild chestnut trees) was

179 evaluated using the 14 loci that did not show the presence of null alleles. In the first set of STRUC-

180 TURE analysis (Figure S2), the  $\Delta K$  statistics gave a maximum value of K=2 ( $\Delta K$  = 78.61), alt-

181 hough a small peak of  $\Delta K$  was also observed for K=3 ( $\Delta K$ =18.42), K=6 ( $\Delta K$ =6.21) and for K=8

182 (ΔK=6.05).

- 183 For K=2, genotypes were grouped into two main clusters with a clear distinction between Spanish
- 184 genotypes, represented by Cluster 1 (with 102 accessions), and Italian genotypes, represented by
- 185 Cluster 2 (with 163 accessions). The threshold for membership determination was Q > 80%. Addi-

tionally, 54 admixed samples were found (Q < 80%) (Figure 1A).

- 187 For K=3, a further separation of hybrid samples was observed. Cluster 3 included the Euro-Chinese
- 188 hybrids (Figure 1B) and some significant Italian cultivars ('Marrone Fiorentino'; 'Madonna', 'Lu-
- 189 cente' and 'Inserta') as well as two varieties from the Andalucía region ('Tomasa' and 'Capilla').



191 Figure 1A: Representation of 319 samples for K=2 by STRUCTURE Software. Each individual is represented by a

192 vertical line and each cluster by different colors: Cluster 1 in red, Cluster 2 in green.





Figure 1B: Representation of 319 samples for K=3 by STRUCTURE Software. Each individual is represented by a vertical line and each cluster by different colors: Cluster 1 in red, Cluster 2 in green and Cluster 3 in blue.

198 3.4 Genetic structure of chestnut accessions of the Iberian and Italian Peninsulas (varieties and wild199 trees)

200

201 The results confirmed the clear separation between the two main clusters (Cluster 1 represented by 202 167 Spanish samples; Cluster 2 represented by 104 Italian and 35 admixed samples). The compari-203 son of chestnut varieties and wild chestnut trees within each cluster did not highlight genetic struc-204 tural differences (Table S3). The separation for K=3 and K=4, with  $\Delta$ K=27.34 and 25.72 respectively, was also considered (Figure S3, Table S3). Accordingly, the separation between the Italian and 205 206 Spanish varieties was also maintained in further subdivisions. The Italian cluster was represented as 207 follows: Cluster 1 for subdivision K=3 included 89 samples; Cluster 4 for subdivision K=4 had 86 208 samples. The Italian cluster included both wild chestnut trees (such as samples named 'Matildico',

- 209 'Legno') and varieties from the Tuscan-Emilian Apennines (such as samples named 'Pastanese',
- 210 'Ceppa', 'Pistolese', 'Piusela' and 'Lisanese').

211 An important group of chestnut varieties was Cluster 2 for subdivision K=3 with different relevant

212 varieties: (e.g. 'Marrone Fiorentino') from northern Italy (Emilia-Romagna, Trentino Alto Adige

- and Piedmont); (e.g., 'Riggiola') from southern Italy (Calabria and Campania); (e.g., "Temprana")
- 214 from southern Spain (Canary Islands, Extremadura, Andalucía). Varieties from Galicia Asturias and
- 215 Andalucía, such as "Luguesa", "Longal", "Temprana" and "Miguelina", as well as "Martahiña",
- 216 were also represented in Cluster 2 (Table S3).
- 217 Most of the varieties contained in Cluster 2 for subdivision K= 3 were also included in Cluster 1 for
- 218 subdivision K=4. In Cluster 2, only a few wild chestnut trees were present (Table S3).
- 219 Finally, Cluster 3 for subdivision K=3 included the main varieties from northern Spain: Galicia
- 220 ('Famosa', 'Inxerta'), Castilla-León ('Negral'), Asturias ('Parede', 'Rapuca' and 'Chamberga'),
- 221 Cantabria, some accessions from Extremadura ('Verata') and from Canary Islands ('Mollar', 'Mu-
- 222 lata' and 'Armentina'; Table S3). In addition, most part of the wild chestnut trees from Galicia,
- 223 Castilla-León and Cantabria were grouped in Cluster 3 (such as 'Pesaguero', 'Alcobilla',
- 224 'Peixeroos').
- 225
- 226 3.5 Genetic differentiation
- 227 The PCoA analysis performed on subdivision K=2 corroborated the results of the STRUCTURE
- 228 analysis by showing a clear separation between Italian (green) and Spanish (red) accessions (Figure
- 229 2A). Some admixed samples were observed as reported by the STRUCTURE analysis.
- 230 Results obtained for subdivision K=3 separated northern (blue) from southern (red) Spanish varie-
- ties. The northern Italian cluster appeared particularly well separated (green) from the two Spanish
- clusters (Figure 2B).
- 233 Furthermore, a larger number of wild chestnut trees was found in the northern Spanish and Italian
- clusters than in the one from southern Spain, in which most samples were varieties (Figure 2B).

Subdivision K=4 showed the presence of two further (sub) clusters (mainly chestnut varieties) in
southern Spain: the first is a cluster deriving from southern Spain and Italy and the second a cluster

237 deriving from central-southern Spain (represented by Cluster 1 and Cluster 2) (Figure 2C).

238 The genetic differentiation between the two main clusters was  $F_{ST} = 0.077$ , P < 0.001 (Table S4.A),

239 suggesting a genetic structure for the chestnut at Italian and Spanish level, also confirmed by the

AMOVA results (8%) (Table 2A). A higher gene flow (Nm) (9.235) was detected for subdivision

241 K=2. In addition, the number of private alleles detected for subdivision K=2 and the observed val-

242 ues are quite similar (0.8571 and 0.7142 respectively; Table S5).

Similar AMOVA results were found for K=3 and K=4, with a 6% and 7% of variance component among the populations, respectively (Table 2, B and C). In contrast, high gene flows (Nm) (6.118 and 3.302 respectively) were detected for subdivisions K=3 and K=4.

246 The largest differentiation between pairs of groups was found between the northern Italian cluster

247 (Cluster 4), with samples mainly from the Tuscan-Emilian Apennines, and the northern-central

248 Spanish cluster (Cluster 3) for K=4 (F<sub>ST</sub>=0.133, P<0.001), as shown in Table S4.C.

249 A high F<sub>ST</sub> value was observed also between Cluster 1, containing cultivars from Italy and southern

250 Spain, and Cluster 4, represented by the Italian cluster with  $F_{ST}=0.113$ , P<0.001; similarly, between

251 Cluster 1 and Cluster 2, which included central Spain's varieties (F<sub>ST</sub>=0.112, P<0.001).

252 In addition, AMOVA analysis, conducted between wild trees and chestnut varieties, showed no dif-

253 ferences between them, in agreement with the STRUCTURE division into two main clusters (Table

254 2. D and E). The variance components among populations were 1% and 5% respectively, confirmed

also by the F<sub>ST</sub> index (0.0012 with P<0.001, Table S4. D and E). The Nm values between wild

256 chestnut trees and varieties and between the two main clusters subject of the STRUCTURE analysis

were 8.112 and 6.364 respectively. Notably, the number of private alleles (Np) in chestnut varieties

258 (0.9285) was higher than the number of private alleles in wild chestnut trees (0.5714, Table S5).

A) Principal Coordinates (PCoA) K=2











260 Figure 2: Principal Coordinate Analysis (PCoA) based on polymorphism at 14 SSR loci for 306 unique genotypes. A),

261 K=2; B), K=3; C) K=4. Accession color reflects the consistent assignment using Bayesian analysis to the sub-groups

262 defined in Fig. 3. Wild chestnut trees are represented by empty symbols inside the sub-groups.

264 Table 2: Analysis of molecular variance (AMOVA) based on the 14 SSR loci of 306 chestnut accessions defined by

		D	f	Variance components (%)		
	Populations	Among Popu- lations	Within Popu- lations	Among Popu- lations	Within Popu- lations	p-value
۸)	Structure Cluster K=2, No hybrids included (306					
A)	samples)	2	609	8	92	0.077
B)	Structure Cluster K=3, No hybrids included (306 samples)	3	608	6	94	0.060
C)	Structure Cluster K=4, No hybrids included (306 samples)	4	607	7	93	0.070
D)	306 chestnut trees divided in chestnut varieties and wild	1	610	1	99	0.012
E)	306 chestnut trees divided in chestnut varieties and wild in Cluster 1/Cluster 2	4	607	5	95	0.048

265 STRUCTURE analysis. All estimates were highly significant - P<0.001.

### 267 **4 Discussion**

268 Our results confirmed the high degree of variability of *C. sativa* and the selected 16 SSRs markers

as powerful tools to evaluate the genetic diversity of EU chestnut germplasm (Pereira-Lorenzo et

270 al., 2011, 2017; Martín et al., 2017a). In particular, the CsCAT3 (PIC=0.879) and EMCs15

271 (PIC=0.604) loci appeared to be the most and least informative loci respectively, as previously re-

272 ported by Pereira-Lorenzo et al. (2010, 2011) and Martín et al. (2012). The average number of al-

leles per locus in this study was 13.25 for 16 SSRs, as in Pereira-Lorenzo et al. (2017), with 8.92

average using 24 SSRs.

275 Our results highlighted the complex structure and genetic diversity of chestnut trees. Genetic diver-

sity is linked to climatic conditions, mainly temperature and precipitation gradient (Pereira-Lorenzo

et al., 2010) and to the domestication process carried out by humans through the centuries (Pereira-

278 Lorenzo et al., 2011, 2019).

279 Hybridization could also have played an important role in the diversification process, as previously

suggested by Pereira-Lorenzo et al. (2011), and, furthermore, it explains the great diversity found in

281 small geographical areas such as the Tuscan-Emilian Apennines (central-northern Italy) and Galicia

282 (northern Spain), as shown by the STRUCTURE and PCoA analysis. In these regions, wild and

283 domesticated chestnuts were found to be genetically similar and cannot be separated. This is in

284 agreement with previous studies, in which no substantial differences between chestnut varieties and 285 wild chestnut trees were found (Fst=0.007 in Pereira-Lorenzo et al., 2019; Fst=0.008 in Bouffartigue et al., 2020). This evidence is supported by the AMOVA analysis that showed a low F<sub>ST</sub> value 286 287 among wild trees and cultivars ( $F_{ST}$ = 0.012 with P<0.001). The Fst and the Nm values observed in 288 samples from the Tuscan-Emilian Apennines and Galicia can be explained considering these areas 289 as the probable sites where the domestication process started, favored by the high number of au-290 tochthonous chestnut populations. In addition, private alleles were also detected in higher values in 291 chestnut varieties, due probably to selection practices, and they have evolutionary significance (Petit 292 et al., 1998).

Our results therefore highlight a low gene flow between chestnut varieties and wild trees. The gene flow between wild trees and chestnut varieties is maintained by changes in forest use over time and the practices related to how the forests were used (Pereira-Lorenzo et al., 2019).

296 At least three major results were obtained by the STRUCTURE analysis.

297 Firstly, the STRUCTURE analysis evidenced a clear separation between the Spanish and the Italian

298 chestnut trees with admixed samples. Similar results, differentiating Spanish and Italian varieties,

were previously reported for adaptive markers (Martín et al., 2017b).

300 The genetic differentiation between the two clusters and the admixed samples was low ( $F_{ST}=0.019$ 

and 0.032, P<0.001 respectively). In addition, the AMOVA analysis which compared varieties and

302 wild trees within the two clusters did not show genetic structural differences ( $F_{ST} = 0.012$ , P<0.001).

303 The large number of admixed samples can indicate a hybridization between the two clusters as

304 suggested also by Pereira-Lorenzo et al. (2012). Furthermore, the aforesaid author evidenced that

305 the hybridization process occurred before the 15<sup>th</sup> century by considering the oldest giant trees from

306 Andalucía with Italian genetic background (Pereira-Lorenzo et al., 2019).

307 Secondly, the STRUCTURE analysis evidenced a separation among samples belonging to the

308 northern and central/southern Spanish Clusters. In particular, the Northern Cluster contained most

309 of the wild chestnut trees while the Central/Southern Cluster included far more chestnut varieties,

310 probably due to human selection (and vegetative propagation of the selected genotypes) related to 311 the use of new seedlings from local cultivars selected for the superior traits of their nuts (Auge and 312 Brandl, 1997; Forneck, 2005; Pereira-Lorenzo et al., 2010). In this respect, Pereira-Lorenzo et al. 313 (2010), reported that the distribution of seedlings from the main cultivar groups from the northern 314 and central Iberian Peninsula was used to create new orchards in southern Spain, in Andalucía and 315 the Canary Islands (e.g., the "Longal", "Reborda" and "Dieguina" cultivars). Our results confirmed 316 the relationship between "Longal" and the main different varieties from southern Spain, such as 317 "Laga", "Temprana" and "Pelona", and from Extremadura, with "Injerta". This was also in line 318 with previous studies (Pereira-Lorenzo et al., 2006, and Costa et al., 2008), which identified "Lon-319 gal" as a cultivar used for genetic contribution to create new cultivars in different regions of Spain. This may also explain the huge number of admixed samples between the sub-clusters being as-320 321 sessed.

Finally, the STRUCTURE analysis evidenced that part of the varieties from southern Spain (Andalucía) shared a higher number of alleles with both varieties from southern Italy (Calabria and Campania regions) and with "Marrone Fiorentino", the most important northern Italian variety.

325 As shown in Figure 3, northern Italy produced a high introgression (30% of the genotypes) in cen-

326 tral-southern Spain and an even higher introgression in central-southern Italy (50%). Moreover, in-

327 trogressions from both the north and the south of Italy were also noticed in northern Spain (16%),

328 with 5% from northern and 11% from southern Italy.

The above results are in line with Pereira-Lorenzo et al. (2019), who supports an early introduction of chestnut cultivations from Italy into Spain. In particular, the introduction of chestnut cultivations started in the Andalucía and Extremadura regions, with contacts also in Castilla-León and Galicia (see the results related to the "Luguesa" cultivar, which was included in Cluster 2 for K=3 and Cluster 1 for K=4 with the main southern Italian varieties).

- 334
- 335





Figure 3: Gene pool distribution of chestnut cultivars for K=3 between the north and the south, both of Italy and Spain.
Green – Mainly northern Italian and France (Cluster 1); Red – Italy, central and southern Spain (Cluster 2); Blue –
northern Spain (Cluster 3).

## 341 5 Conclusions

342 This study contributes to improving knowledge on the genetic relationships between chestnut varie-343 ties and wild chestnut trees in the Iberian and Italian Peninsula and to expanding the Chestnut Euro-344 pean Genetic Dataset. The genetic variations between and within Italian and Spanish clusters, in-345 cluding both chestnut varieties and wild trees, reflect a combination of historical migrations and se-346 lection processes. This is highlighted by the high number of admixed in relation to the chestnut 347 populations of central-southern Spain and central-southern Italy. Adaptation to different environ-348 ments and hybridization led to a wide genetic variability in limited areas such as the Tuscan-349 Emilian Apennines and the Galicia region. In these areas, no genetic structure differences between 350 wild chestnut trees and chestnut varieties was found.

#### 351 Acknowledgments

352 Field and laboratory work were funded by Sara Alessandri's fellowship PhD program, Agricultural, 353 Environmental, and Food Science and Technology (STAAA) offered by the Department of Agriculture and 354 Food Sciences (DISTAL, University of Bologna). The Italian accessions were analyzed with the contribution 355 of the PSR "BIODIVERSAMENTE CASTAGNO" regional project and of the National Academy of 356 Agriculture (ANA). The authors would like to thank USC - Department of Crop Production and Projects of 357 Engineering, in particular, Prof. S. Pereira-Lorenzo and A.M. Ramos Cabrer for the hospitality and efforts to 358 align samples and for the genetic diversity analysis. SSRs from the EU Database were obtained with funds 359 from the Spanish Government for the project "An integrated approach to sustainable management of 360 chestnut in Spain" (AGL2013-48017).

#### 361 **Bibliography**

362 Alcaide, F.; Solla, A.; Cuenca, B.; Martín, M. Á. (2021). Molecular evidence of introgression of

Asian germplasm into a natural *Castanea sativa* forest in Spain. Forestry: An International Journal
 of Forest Research.

365

Alcaide, F.; Solla, A.; Cherubini, M.; Mattioni, C.; Cuenca, B.; Camisón, Á.; Martín, M. Á. (2020).
Adaptive evolution of chestnut forests to the impact of ink disease in Spain. Journal of Systematics
and Evolution, 58(4), 504-516.

369

370 Alcaide, F.; Solla, A.; Mattioni, C.; Castellana, S.; Martín, MA (2019) Adaptive diversity and

371 drought tolerance in Castanea sativa assessed through EST-SSR genic markers. Forestry 92:287–

372 296. https:// doi.org/10.1093/forestry/cpz007

373

- 374 Amiryousefi, A.; Hyvönen, J.; Poczai, P. (2018). iMEC: online marker efficiency calcula-
- tor. Applications in plant sciences, 6(6), e01159.

377	Auge, H.; Brandl, R. (1997). Seedling recruitment in the invasive clonal shrub, Mahonia aquifolium
378	Pursh (Nutt.). Oecologia (Berl.), 110(2): 205–211. doi: 10.1007/s004420050151.
379	
380	Bernetti, G. (1995). Selvicoltura speciale. Unione tipografico-editrice Torinese.
381	
382	Bouffartigue, C.; Debille, S.; Fabreguette, O.; Ramos Cabrer, A.; Pereira-Lorenzo, S.; Flutre, T.;
383	Harvengt, L. (2019). Genetic diversity and structure of sweet chestnut (Castanea sativa Mill.) in
384	France: At the intersection between Spain and Italy. bioRxiv 792259; doi:
385	https://doi.org/10.1101/792259
386	
387	Bouffartigue, C.; Debille, S.; Fabreguettes, O.; Cabrer, A. R.; Pereira-Lorenzo, S.; Flutre, T.;
388	Harvengt, L. (2020). High admixture between forest and cultivated chestnut (Castanea sativa Mill.)
389	in France. bioRxiv, 792259.
390	
391	Buck, E.J.; Hadonou, M.; James, C.J.; Blakesley, D.; Russell, K. (2003) Isolation and characteriza-
392	tion of polymorphic microsatellites in European chestnut (Castanea sativa Mill.) Mol Ecol Notes
393	3:239–241. https://doi.org/10.1046/j.1471-8286.2003.00410.x
394	
395	Castellana, S.; Martin, M. Á.; Solla, A; Alcaide, F.; Villani, F.; Cherubini, M.; Mattioni, C. (2021).
396	Signatures of local adaptation to climate in natural populations of sweet chestnut (Castanea sativa
397	mill.) from southern Europe. Annals of Forest Science, 78(2), 1-21.
398	
399	Conedera, M. (1996). Die Kastanie: der Brotbaum. Vergangenheit, Gegenwart und Zukunft der
400	"Waldfrucht par excellence"". Bu"ndnerwald, 49: 28–46.
401	
	20

402	Conedera, M.; Krebs, P.; Tinner, W.; Pradella, M.; Torriani, D. (2004). The cultivation of Castanea
403	sativa (Mill.) in Europe, from its origin to its diffusion on a continental scale. Vegetation History
404	and Archaeobotany, 13(3), 161-179. https://doi.org/10.1007/s00334-004-0038-7
405	

- 406 Costa, R.; Ribeiro, C.; Valdiviesso, T.; Afonso, S.; Borges, O.; Soeiro, J.; Costa, H.; Fonseca, L.;
- 407 Augusta, C.; Cruz, M.H.; Salazar, M.; Matos Soares, F.; Sequeira, J.; Assunção, A.; Correia, P.;
- 408 Lima, M.J. (2008). Variedades de Castanha das Regiões Centro e Norte de Portugal. INRB.I.P.409
- 410 Earl, D.A.; Vonholdt, B.M. (2012) STRUCTURE HARVESTER: a website and program for visual-
- 411 izing STRUCTURE output and implementing the Evanno method. Conserv Genet Resour 4:359–
- 412 361. https://doi.org/10.1007/s12686-011-9548-7
- 413
- 414 Evanno, G.; Regnaut, S.; Goudet, J. (2005). Detecting the number of clusters of individuals using
- 415 the software STRUCTURE: A simulation study. Molecular Ecology, 14(8), 2611–2620.
- 416 https://doi.org/10.1111/j.1365-294X.2005.02553.x
- 417
- 418 Fernández Cruz, J.; Fernández López, J. (2016) Genetic structure of wild sweet chestnut (Castanea
- 419 sativa Mill.) populations in northwest of Spain and their differences with other European stands.

420 Conserv Genet 17:949–967. https://doi.org/10.1007/s10592-016-0835-4

- 421
- 422 Fernández-López, J.; Fernández-Cruz, J.; Míguez-Soto, B. (2021). The demographic history of *Cas-*423 *tanea sativa* Mill. in southwest Europe: a natural population structure modified by translocations.
- 424 Molecular Ecology.
- 425
- 426 Fideghelli, C. (2017). Atlante dei fruttiferi autoctoni italiani. Crea Centro di frutticoltura, Roma.
- 427

428	Forneck, A. (2005). Plant breeding: clonality — a concept for stability and variability during vege-
429	tative propagation. Prog. Bot. 66: 164-183. doi: 10.1007/3-540-27043-4_8
430	
431	Gobbin, D.; Hohl, L.; Conza, L.; Jermini, M.; Gessler, C.; Conedera, M. (2007) Microsatellite-
432	based characterization of the Castanea sativa cultivar heritage of southern Switzerland. Genome

- 433 50:1089–1103. doi: 10.1139/G07-086
- 434
- Hummer, K.E.; Dempewolf, H.; Bramel, P.; Markham, R.; Stover, E. (2015) Status of global strategies for horticultural fruit crops. Acta horticulturae

438 Huntley, B. and Birks, H. J. B. (1983). Atlas of past and present pollen maps for Europe, 0-13,000

439 years ago. Cambridge University Press.

440

441 iMEC: Online Marker Efficiency Calculator, Ali Amiryousefi, Jaakko Hyvönen, Péter Poczai Appl
442 Plant Sci. (2018); 6(6): e01159.

443

- 444 Jarman, R.; Mattioni, C.; Russell, K.; Chambers, F. M.; Bartlett, D.; Martin, M. A.; Webb, J.
- 445 (2019). DNA analysis of Castanea sativa (sweet chestnut) in Britain and Ireland: Elucidating Euro-

446 pean origins and genepool diversity. PloS one, 14(9), e0222936.

- 447
- 448 Kalinowski, S. T.; Taper, M. L.; Marshall, T. C. (2007). Revising how the computer program
- 449 CERVUS accommodates genotyping error increases success in paternity assignment. Molecular
- 450 Ecology, 16(5), 1099–1106. https://doi.org/10.1111/j.1365294X.2007.03089.x
- 451

452	Kampfer, S.; Lexer, C.; Glossl, J.; Steinkellner, H. (1998). Characterization of (GA)(n) microsatel-
453	lite loci from Quercus robur. Hereditas, 129(2), 183–186. https://doi.org/10.1111/j.1601-
454	5223.1998.00183.x

456 Krebs, P.; Conedera, M.; Pradella, M.; Torriani, D.; Felber, M.; Tinner W. (2004) Quaternary refu-

457 gia of the sweet chestnut (Castanea sativa Miller): an extended palynological approach. Vegetation

458 History and Archaeobotany, 13, 145–160.

459

460 Krebs, P.; Pezzatti, G. B.; Beffa, G.; Tinner, W.; Conedera, M. (2019). Revising the sweet chestnut

461 (Castanea sativa Mill.) refugia history of the last glacial period with extended pollen and macrofos-

462 sil evidence. Quaternary Science Reviews, 206, 111-128.

463

Lusini, I.; Velichkov, I.; Pollegioni, P.; Chiocchini, F.; Hinkov, G.; Zlatanov, T. et al. (2014). Estimating the genetic diversity and spatial structure of Bulgarian *Castanea sativa* populations by
SSRs: implications for conservation. Conservation genetics, 15(2), 283-293.

467

468 Maguire, T. L.; Collins, G. G.; Sedgley, M. (1994). A modified CTAB DNA extraction procedure

469 for plants belonging to the family proteaceae. Plant Molecular Biology Reporter, 12(2), 106–109.

470 https://doi.org/10.1007/BF02668371

471

472 Marinoni, D. T.; Akkak, A.; Beltramo, C.; Guaraldo, P.; Boccacci, P.; Bounous, G.; Botta, R.

473 (2013). Genetic and morphological characterization of chestnut (Castanea sativa Mill.) germplasm

474 in Piedmont (north-western Italy). Tree genetics & genomes, 9(4), 1017-1030. DOI

475 10.1007/s11295-013-0613-0

476

- 477 Marinoni, D.; Akkak, A.; Bounous, G.; Edwards, K.J.; Botta, R. (2003). Development and charac-
- 478 terization of microsatellite markers in Castanea sativa (Mill.). Mol. Breed. 11 (2), 127–136

479 https://doi.org/10.1023/ A:1022456013692.

- 480
- 481 Martín, M.A.; Alvarez, J.B.; Mattioni, C.; Cherubini, M.; Villani, F.; Martín, L.M. (2009) Identifi-
- 482 cation and characterization of traditional chestnut varieties of southern Spain using morphological
- 483 and simple sequence repeat (SSRs) markers. Ann Appl Biol 154:389–398.
- 484 https://doi.org/10.1111/j.1744-7348.2008.00309.x
- 485
- 486 Martín, M.A.; Mattioni, C.; Molina, J.R.; Alvarez, J.A.; Cherubini, M.; Herrera, M.A. et al. (2012)
- 487 Landscape genetics structure of chestnut (Castanea sativa Mill.) in Spain. Tree Genet Genomes

488 8:127–136. https://doi.org/10.1007/s11295-011-0427-x

- 489
- 490 Martín, M. A.; Monedero, E.; Martín, L. M. (2017a). Genetic monitoring of traditional chestnut or-
- 491 chards reveals a complex genetic structure. Annals of Forest Science, 74(1).
- 492 https://doi.org/10.1007/s13595-016-0610-1
- 493
- 494 Martín, M. A.; Mattioni, C.; Cherubini, M.; Villani, F.; Martín, L. M. (2017b). A comparative study
- 495 of European chestnut varieties in relation to adaptive markers. Agroforestry systems, 91(1), 97-109.496
- 497 Mattioni, C.; Martin, A.M.; Pollegioni, P.; Cherubini, M.; Villani, F. (2013). Microsatellite markers
- 498 reveal a strong geographical structure in European populations of Castanea sativa (Fagaceae): Evi-
- 499 dence for multiple glacial refugia. American Journal of Botany, 100(5), 951–961.
- 500 https://doi.org/10.3732/ajb.1200194
- 501

502	Mattioni, C.; Martin, M. A.; Chiocchini, F.; Cherubini, M.; Gaudet, M.; Pollegioni, P. et al. (2017).
503	Landscape genetics structure of European sweet chestnut (Castanea sativa Mill): indications for
504	conservation priorities. Tree Genetics and Genomes, 13(2). https://doi.org/10.1007/s11295-017-
505	1123-2
506	
507	Mellano, M.G.; Torello-Marinoni, D.; Boccacci, P. (2018) Ex situ conservation and characterization
508	of the geneticdiversity of Castanea spp. Acta Horticulturae 1-6. doi:
509	10.17660/ActaHortic.2018.1220.1
510	
511	Míguez-Soto, B.; Fernández-Cruz, J.; Fernández-López, J. (2019). Mediterranean and Northern Ibe-
512	rian gene pools of wild Castanea sativa Mill. are two differentiated ecotypes originated under natu-
513	ral divergent selection. PloS one, 14(2), e0211315.
514	
515	Neri, L.; Dimitri, G.; & Sacchetti, G. (2010). Chemical composition and antioxidant activity of
516	cured chestnuts from three sweet chestnut (Castanea sativa Mill.) ecotypes from Italy. Journal of
517	Food Composition and Analysis, 23(1), 23–29. https://doi.org/10.1016/j.jfca.2009.03.002
518	
519	Nishio, S.; Takada, N.; Terakami, S.; Takeuchi, Y.; Kimura, M. K.; Isoda, K.; Saito, T.; Iketani, H.
520	(2021). Genetic structure analysis of cultivated and wild chestnut populations reveals gene flow
521	from cultivars to natural stands. Scientific Reports, 11(1), 1-13.
522	
523	Peakall, R.; Smouse, P.E. (2006) Genalex 6: genetic analysis in Excel. Population genetic software
524	for teaching and research. Mol Ecol Notes 6:288-295. https://doi.org/10.1111/j.1471-
525	8286.2005.01155.x
526	

- 527 Peakall, R., Smouse, P.E., (2012). GenAlEx 6.5: genetic analysis in Excel. Population genetic soft-
- 528 ware for teaching and research-an update. Bioinformatics 28, 2537e2539.
- 529 https://doi.org/10.1093/bioinformatics/bts460.
- 530
- 531 Petit, R.J., El Mousadik, A., Pons, O., (1998). Identifying populations for conservation on the basis
- of genetic markers. Conserv. Biol. 12, 844e855. https://doi.org/10.1111/j.1523-1739.1998.96489.x.
- 534 Pereira-Lorenzo, S.; Costa, R.; Ramos-Cabrer, A.M.; Ribeiro, C.; Serra da Silva, C.; Manzano, G.;
- 535 Barreneche, T. (2010) Variation in grafted European chestnut and hybrids by microsatellites reveals
- two main origins in the Iberian Peninsula. Tree Genet Genomes 6:701–715.
- 537 https://doi.org/10.1007/s11295-010-0285-y
- 538
- 539 Pereira-Lorenzo, S.; Lourenço Costa, R.M.; Ramos-Cabrer, A.M.; Ciordia-Ara, M.; Marques Ribei-
- 540 ro, C.A.; Borges, O.; Barreneche, T. (2011) Chestnut cultivar diversification process in the Iberian
- 541 Peninsula, Canary Islands, and Azores. Genome 54:301–315. https://doi.org/10.1139/g10-122
- 542
- 543 Pereira-Lorenzo, S.; Ramos-Cabrer, A. M.; Barreneche, T.; Mattioni, C.; Villani, F.; Díaz-
- 544 Hernández, M.B. et al. (2017). Database of European chestnut cultivars and definition of a core col-
- 545 lection using simple sequence repeats. Tree Genetics and Genomes, 13(5).
- 546 https://doi.org/10.1007/s11295-017-1197-x
- 547
- 548 Pereira-Lorenzo, S.; Ramos-Cabrer, A. M.; Barreneche, T.; Mattioni, C.; Villani, F.; Díaz-
- 549 Hernández, B.; Martín, A. (2019). Instant domestication process of European chestnut cultivars.
- 550 Annals of Applied Biology, 174(1), 74–85. https://doi.org/10.1111/aab.12474
- 551

- 552 Pereira-Lorenzo, S.; Bischofberger, Y.; Conedera, M.; Piattini, P.; Crovadore, J.; Chablais, R. et al.
- 553 (2020) Reservoir of the European chestnut diversity in Switzerland. Biodivers Conserv (2020).

554 https://doi.org/10.1007/s10531-020-01970-2

- 556 Pitte, J.R. (1986). Terres de castanide. Hommes et paysages du chataignier de l'Antiquite'a nos
  557 jours. Librairie Artheme Fayard.
- 558
- 559 Porras-Hurtado, L.; Ruiz, Y.; Santos, C.; Phillips, C.; Carracedo, A.; Lareu, M. (2013) An overview
- of STRUCTURE: applications, parameter settings, and supporting software. Front Genet 4:98.
- 561 https://doi.org/10. 3389/fgene.2013.00098
- 562
- 563 Pritchard, J.K.; Stephens, M.; Donnelly, P. (2000) Inference of population structure using multi564 locus genotype data. Genetics 155:945–959
- 565
- 566 Roces-Díaz, J. V.; Jiménez-Alfaro, B.; Chytrý, M.; Díaz-Varela, E. R.; Álvarez-Álvarez, P. (2018).
- 567 Glacial refugia and mid-Holocene expansion delineate the current distribution of *Castanea sativa* in
- 568 Europe. Palaeogeography, Palaeoclimatology, Palaeoecology, 491, 152-160.
- 569
- 570 Silvanini, A.; Marinoni, D. T.; Beccaro, G. L.; Ganino, T. (2011). La caratterizzazione varietale del
- 571 germoplasma di *Castanea sativa* Mill. Italus Hortus, 18, 47-61.
- 572
- 573 Villani, F.; Sansotta, A.; Cherubini, M.; Cesaroni, D.; Sbordoni, V. (1999). Genetic structure of
- 574 natural populations of Castanea sativa in Turkey: evidence of a hybrid zone. JOURNAL OF EVO-
- 575 LUTIONARY BIOLOGY, 12(2), 233-244.
- 576
- 577
- 27

578	Support	ting int	format	tion
570	Suppor		i vi ma	101

579 **Table S1:** unique genotypes analyzed in this study (319 samples).

580

581 Table S2: list of the different varieties collected with their countries of origin, the STRUCTURE 582 subdivision for K=2 and K=3 for 319 samples. A Q threshold of 0.8 was used to infer an accession to a specific cluster. 583 584 585 Table S3: list of the different varieties collected with their countries of origin, the STRUCTURE 586 subdivision for K=2, K=3 and K=4 for 306 samples (no hybrids included) and varieties/wild subdi-587 vision. A Q threshold of 0.8 was used to infer an accession to a specific cluster. 588 589 Table S4: pairwise estimate of Fst value based on the 14 SSR loci of 306 chestnut accessions de-590 fined by STRUCTURE analysis: A), K=2; B), K=3; C) K=4; D) the total population (n= 306) di-591 vided into wild trees and chestnut varieties; E) wild trees between chestnut varieties in K=2. All es-592 timates were highly significant - P<0.001. 593 594 Figure S1: UPGMA tree indicating the relationships among 630 chestnut samples. 595

596 Figure S2: Estimates of  $\triangle$  k calculated based on Evanno et al., (2005), based on k-subdivision for 597 319 samples.

- 599 Figure S3: Estimates of  $\triangle$  k calculated as described by Evanno et al., (2005), based on k-
- 600 subdivision for 306 samples.