



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Corrigendum and Addendum to “Polarized Parallel Transport and Uniruled Divisors on Generalized Kummer Varieties”

This is the final peer-reviewed author’s accepted manuscript (postprint) of the following publication:

Published Version:

Corrigendum and Addendum to “Polarized Parallel Transport and Uniruled Divisors on Generalized Kummer Varieties” / Mongardi G.; Pacienza G.. - In: INTERNATIONAL MATHEMATICS RESEARCH NOTICES. - ISSN 1073-7928. - STAMPA. - 2021:18(2021), pp. 14398-14404. [10.1093/imrn/rnz163]

Availability:

This version is available at: <https://hdl.handle.net/11585/850332> since: 2023-08-09

Published:

DOI: <http://doi.org/10.1093/imrn/rnz163>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Giovanni Mongardi , Gianluca Pacienza, Corrigendum and Addendum to “Polarized Parallel Transport and Uniruled Divisors on Generalized Kummer Varieties”, *International Mathematics Research Notices*, Volume 2021, Issue 18, September 2021, Pages 14398–14404

The final published version is available online at <https://dx.doi.org/10.1093/imrn/rnz163>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

CORRIGENDUM AND ADDENDUM TO “POLARIZED PARALLEL TRANSPORT AND UNIRULED DIVISORS ON GENERALIZED KUMMER VARIETIES”

GIOVANNI MONGARDI AND GIANLUCA PACIENZA

ABSTRACT. We correct the statement of the main result of [MP] and provide some further precisions.

The goal of this short note is to state correctly the main result of [MP]. For the definitions, the notations and the motivations we refer the reader to [MP]. The correct statement is the following:

Theorem 0.1. *Let $n \geq 1$ be an integer. Let $\mathfrak{M} = \cup_{d>0} \mathfrak{M}_{2d}$ be the union of the moduli spaces \mathfrak{M}_{2d} of projective irreducible holomorphic symplectic varieties of $K_n(A)$ -type polarized by a line bundle of degree $2d$. For all $(X, H) \in \mathfrak{M}$, outside at most a finite number of connected components determined by the monodromy orbit of H , the linear system $|mH|$, for some m , contains a uniruled divisor covered by rational curves of primitive class.*

Let q be the Beauville-Bogomolov quadratic form on $H^2(X, \mathbb{Z})$. This induces an embedding $H^2(X, \mathbb{Z}) \hookrightarrow H_2(X, \mathbb{Z}), H \mapsto H^\vee$. By abuse of notation we denote again by q the quadratic form on $H_2(X, \mathbb{Z})$.

Remark 0.2. The statement above insures precisely existence of uniruled divisors covered by primitive rational curves if there exist integers p, g and ϵ such that $p \geq g$ and $\epsilon = 0$ or 1 with

- (i) the class $\alpha := \frac{H^\vee}{\text{div}(H)} \in H_2(X, \mathbb{Z})$ can be written as $\gamma + (2g - \epsilon)\eta$ with η in the monodromy orbit of the class of the exceptional curve on a $K_n(A)$;
- (ii) $\gamma \in \eta^\perp$, $q(\gamma) = 2p - 2$ (hence $q(\alpha) = 2p - 2 - \frac{(2g-\epsilon)^2}{2n+2}$).

Remark 0.3.

- (i) It follows from Proposition 2.1 that if $q(\alpha) > n + 1$, then a multiple of H is uniruled by primitive rational curves of class α .
- (ii) If $\rho(X) \geq 2$ then X always contains an ample uniruled divisor covered by primitive rational curves (cf. Corollary 2.3).
- (iii) If $n \leq 5$ then the conclusion of the theorem holds for *all* the connected components of \mathfrak{M} (cf. Remark 2.4).
- (iv) If $n + 1$ is a power of a prime number, then by [Mark, Mon2], the monodromy group is maximal. Therefore it suffices to check that the square $q(\alpha)$ is of the form $2p - 2 - \frac{(2g-\epsilon)^2}{2n+2}$, with $p \geq g$.

The original proof was based on 3 ingredients: the first was a deformation theoretic statement, saying that rational curves whose deformations cover a divisor in irreducible holomorphic symplectic manifolds are non-obstructed (see [CP, Corollary 3.5]). The second is the characterization of polarized parallel transport operators on polarized irreducible holomorphic symplectic varieties (X, H) of $K_n(A)$ -type (see [MP, Theorem 1.1]) which allows to obtain an explicit description of

Date: June 19, 2019.

the polarized deformation equivalence (see [MP, Theorem 4.2]). These two ingredients are true. The third argument consists in the construction of explicit examples of uniruled divisors on the generalized Kummer variety associated to a polarised abelian surface (A, H_A) with $NS(A) = \mathbb{Z}H_A$ such that $p_a(H_A) \geq g \geq 2$. The construction is also correct, but the examples that we provided cannot yield all the possible primitive polarizations, as we tacitly and erroneously assumed in [MP]. Even without taking the monodromy orbit into account, this is simply because it may happen that the number $2p - 2 - \frac{(2g-\epsilon)^2}{2n+2}, \epsilon = 0, 1$, is positive even with $p < g$, which obviously renders our geometric argument empty. Indeed the rational curves are constructed as \mathfrak{g}_n^1 on the normalization of a nodal curve of geometric genus g lying in the hyperplane linear system $|H_A|$, which is supposed to have $p_a(H_A) = p$. We also take the occasion of this note to provide the full proof (see Proposition 1.1) of a technical point which we claimed in [MP, Section 4.2] to follow from a dimension count as in [Voi15, Example 4.1, 3]). The statement is correct, but the argument cannot be the same as in [Voi15, Example 4.1, 3]) because we deal here with a locally closed subset (the Severi variety) of a complete linear system, and not with the full complete linear system.

The $K3^{[n]}$ -type case, initially treated in [CP], is subject to the same considerations and will be treated in [CMP].

We realized our mistake after the appearance of [OSY], which provides counterexamples in the $K3^{[n]}$ -case which apply exactly in all the cases not covered by the similar geometric constructions for the Hilbert scheme of points on a general projective $K3$. Contrary to the $K3^{[n]}$ -type case as far as we know there are no known counterexamples to the existence of uniruled divisors ruled by a primitive curve class in the $K_n(A)$ -type case. Nevertheless we have no reasons to believe that the $K_n(A)$ -type case could be exempt from this type of sporadic pathologies.

ACKNOWLEDGEMENTS

We would like to thank Q. Yin for useful discussions and F. Charles for reading a preliminary version. GM is supported by “Progetto di ricerca INdAM per giovani ricercatori: Pursuit of IHS” and would like to thank the Université de Lorraine for hosting him while this work was finished. GP is partially supported by the Projet ANR-16-CE40-0008 “Foliage”.

1. EXISTENCE OF UNIRULED DIVISORS ON $K_n(A)$

In [MP, Section 4.2 “Examples”] we claimed that “the natural map from $\tilde{\mathcal{C}}_{g+1}^1 \rightarrow A^{[g+1]}$ is finite onto its image” invoking a dimension count made in [Voi15, Example 4.1, 3]). However the same argument cannot work because we do not work with the full continuous system, but with a locally closed subset (the Severi variety). Hence we take the occasion to provide a full proof of that statement in the following.

Proposition 1.1. *Let g be an integer ≥ 2 and (A, H_A) be a general polarized abelian surface with $p_a(H_A) =: p \geq g$. Then $A^{[g+1]}$ contains a uniruled divisor covered by the \mathfrak{g}_{g+1}^1 on nodal genus g curves in the continuous system $\{H_A\}$.*

Proof. To prove the statement we can actually work over a very general polarized abelian surface, so let us suppose that $NS(A) = \mathbb{Z}H_A$. We will prove this statement by induction on g . It is sufficient to show it on the symmetric product of A .

Observe that, by [KLM, Thm. 1.1], for all $2 \leq g \leq p_a(H_A)$, that the Severi variety parametrizing nodal genus g curves inside $\{H_A\}$ is non-empty of the expected dimension g .

It is sufficient to show the claim on the symmetric product $A^{(g+1)}$ of A . More precisely, we will prove the following statement: there exists an irreducible component V of the (Zariski closure of the) Severi variety parametrizing nodal genus g curves inside $\{H_A\}$ such that, if $\mathcal{C}_V \rightarrow V$ denotes the universal curve and $\mathcal{C}_V^{(g+1)} \rightarrow V$ the relative symmetric product, the natural morphism

$$\mathcal{C}_V^{(g+1)} \rightarrow A^{(g+1)}$$

is generically finite onto its image. Note that this is equivalent to saying that $(g+1)$ generic points on a generic curve of the family lie only on a finite number of curves of the family.

Indeed as

$$\dim \mathcal{C}_V^{(g+1)} = \text{reldim}(\mathcal{C}_V^{(g+1)}) + \dim V = (g+1) + g = 2g+1$$

it follows that the image is a divisor inside $A^{(g+1)}$. Since the k -th symmetric product of a curve is uniruled for k greater than the genus of the curve, as a by-product we have that such divisor is uniruled.

Note also that positive dimensional fibers of the morphism $\mathcal{C}_V^{(g+1)} \rightarrow A^{(g+1)}$ cannot lie in a fiber of $\mathcal{C}_V^{(g+1)} \rightarrow V$, as $C_t^{(g+1)}$ injects into $A^{(g+1)}$ for every $t \in V$.

We start with the case $g=2$. Let C be one of the (finitely many) nodal curves of geometric genus 2 inside the linear system $|H_A|$. In this case the points of the component V of the Severi variety containing C are given by all the translates of C . The third symmetric product $C^{(3)}$ injects as a 3-dimensional subvariety inside $A^{(3)}$. The action of A on $C^{(3)}$ by translation has no positive-dimensional stabilizer (as A is general, hence simple). Therefore the orbit of $C^{(3)}$ under this action is a divisor. Using the notation above such divisor is the image of $\mathcal{C}_V^{(2+1)}$ in $A^{(3)}$.

By inductive hypothesis, there exists an irreducible component W of the (Zariski closure of the) Severi variety parametrizing nodal genus $g-1$ curves inside $\{H_A\}$ such that, if $\mathcal{C}_W \rightarrow W$ denotes the universal curve and $\mathcal{C}_W^{(g)} \rightarrow W$ the relative symmetric product, the natural morphism

$$\mathcal{C}_W^{(g)} \rightarrow A^{(g)}$$

is generically finite onto its image.

Now let V be (the Zariski closure of) an irreducible component of the Severi variety of nodal genus g curves in $\{H_A\}$ obtained by smoothing one node of the curves in W (which can be done by the regularity of the Severi variety, [CS, Example 1.3]). By construction $W \subset V$. Let $\mathcal{C}_V \rightarrow V$ be the universal curve. Its restriction over W yields a map $\mathcal{C}_W \rightarrow W$. Let D be the image of the morphism

$$\mathcal{C}_V^{(g+1)} \rightarrow A^{(g+1)}.$$

Observe that D contains the image D_W of

$$\mathcal{C}_W^{(g+1)} \rightarrow A^{(g+1)}.$$

We claim that by the inductive hypothesis D_W has codimension 2, or, equivalently, that the morphism $\mathcal{C}_W^{(g+1)} \rightarrow A^{(g+1)}$ is generically finite onto its image. Indeed if $\xi = x_1 + \dots + x_{g+1}$ is a generic point of the image, then, say, $x_1 + \dots + x_g$ is a generic point of the image of the morphism $\mathcal{C}_W^{(g)} \rightarrow A^{(g)}$. By the inductive hypothesis the points x_1, \dots, x_g lie on finitely many curves of the family W , *a fortiori* that will be true for x_1, \dots, x_g, x_{g+1} and the claim follows.

We want to prove that D contains D_W strictly. If this were not the case, by irreducibility, we would have $D = D_W$. Let $U \subset D$ be an open subset over which the morphisms $\mathcal{C}_W^{(g+1)} \rightarrow A^{(g+1)}$ and $\mathcal{C}_V^{(g+1)} \rightarrow A^{(g+1)}$ are smooth and let $p_1 + p_2 + \dots + p_{g+1}$ be a point in U . Let C be a nodal genus g curve in V containing these points. Let us fix the first g points p_1, \dots, p_g . By induction these points are contained inside a finite number of curves of genus $g-1$ belonging to W . Let

B_1, \dots, B_m be all such curves. Let $U_C \subset C$ be an open subset such that for all $q \in U_C$ we have $p_1 + \dots + p_g + q \in U$. As we have seen above p_1, \dots, p_g, q lie on finitely many curves of genus $g - 1$ belonging to W , and these curves must be B_1, \dots, B_m . Therefore, as q varies in U_C , we deduce that U_C is a subset of a finite union of genus $(g - 1)$ curves. As C is irreducible, there is an i such that $C = B_i$, which is clearly a contradiction. Therefore D must strictly contain D_W and be a divisor, which is necessarily uniruled. □

The rest of the proof remains the same and we refer the reader to [MP] for the details.

2. WHERE IT DOES NOT WORK

In this section we prove that, for every dimension, there is at most a finite number of components of the moduli space of polarized manifolds (X, H) of $K_n(A)$ -type where the strategy of the previous section does not work.

The uniruled divisors we constructed have a cohomology class which is a multiple of $H_A - (2g)\tau$ (or $H_A - (2g - 1)\tau$) where $2p - 2 = H_A^2$ and H_A is the primitive polarization on the abelian surface. We have the following:

Proposition 2.1. *Let X be a projective irreducible holomorphic symplectic variety of $K_n(A)$ -type. Let $C \in H_2(X, \mathbb{Z}) \cap N_1(X)$ be a primitive class such that its square $q(C)$ with respect to the Beauville-Bogomolov form is $> n + 1$. Then, the class C is deformation equivalent to the class of one of the curves constructed in the previous section.*

Proof. We know by [MP, Theorem 4.2] that C is deformation equivalent to either $H_A - 2g\tau$ or $H_A - (2g - 1)\tau$, with $g \leq n + 1$. If $q(C) > n + 1$, the square of $H_A - (2n + 2)\tau$ is positive, that is $H_A^2 = 2p - 2$ with $p > n + 1$. Thus, $p > n + 1 \geq g$ which means that $H_A - 2g\tau$ can be represented by the class of a \mathfrak{g}_{g+1}^1 on a nodal curve in $\{H_A\}$. □

Corollary 2.2. *Let \mathcal{M}_n be the moduli space of all polarized manifolds of $K_n(A)$ -type with n fixed. Then, the number of components of \mathcal{M}_n whose general points (X, H) do not have a uniruled divisor ruled by a rational curve of primitive class is at most finite.*

Proof. The components of \mathcal{M}_n are in bijective correspondence with the monodromy orbits of a given class of positive square in $L_n := U^3 \oplus (-2n - 2) \cong H^2(X, \mathbb{Z})$, see [Ono, Thm. 2.8].

For a fixed square of H , there is a finite number of orbits (computed again in [Ono, Thm. 2.8]), so it follows that if X has a uniruled divisor when $q(H)$ is big enough, our claim will hold. The dual curve to H is given by $H/\text{div}(H)$, where $\text{div}(H)$ is the divisibility of H which is the positive generator of the ideal $q(H, H^2(X, \mathbb{Z}))$. The divisibility is at most $2n + 2$, therefore if $q(H) \geq (2n + 2)^2(n + 1)$ the dual curve has square at least $n + 1$, so that Proposition 2.1 applies and our claim follows. □

Corollary 2.3. *Let X be a projective manifold of $K_n(A)$ -type with Picard rank at least two. Then X has an ample divisor ruled by primitive rational curves.*

Proof. Since X is projective and has Picard rank at least two, its Picard lattice is indefinite and contains primitive elements of positive arbitrary Beauville-Bogomolov square, and so does the ample cone. Let H be an ample divisor such that $q(H) \geq (2n + 2)^2(n + 1)$. Let C be its dual curve in $H_2(X, \mathbb{Z})$. As the divisibility of H is at most $2n + 2$ it follows that $q(C) \geq n + 1$ and Proposition 2.1 yields our claim. □

Remark 2.4. The estimate of Proposition 2.1 is definitely not sharp, indeed all primitive curves of positive square on manifolds of $K_n(A)$ -type with $n \leq 5$ are deformation equivalent to the curves we construct in Proposition 1.1. Indeed, by [MP, Theorem 4.2] we can suppose that our pair (X, C) with $q(C) > 0$ is $(K_n(A), H_A - \mu\tau)$ with $0 \leq \mu \leq n + 1$ and A is an abelian surface of genus p . The class $H_A - \mu\tau$ is given by the class of the rational curves constructed in Proposition 1.1, which have class $H_A - 2g\tau$, with the eventual addition of a tail of class τ , so that $2g \leq n + 2$. By contradiction let us suppose that $g > p$ and $n \leq 5$. We have $q(H_A - 2g\tau) = 2p - 2 - 2\frac{g^2}{n+1} \leq 2p - 2 - 2\frac{(p+1)^2}{n+1} \leq 2p - 2 - 2\frac{(p+1)^2}{6}$. However, the last value is never positive, hence $q(H_A - 2g\tau)$ cannot be positive and we reach a contradiction. Analogously, for $C = H_A - (2g - 1)\tau$, we have $q(C) \leq \frac{20p - 25 - 4p^2}{12}$ with $g \geq p + 1$ and $2g \leq n + 2$, which is again not positive.

REFERENCES

- [CP] F. Charles, G. Pacienza, Families of rational curves on holomorphic symplectic varieties, [arXiv:1401.4071v2](#) [math.AG].
- [CMP] F. Charles, G. Mongardi, G. Pacienza, Families of rational curves on holomorphic symplectic varieties of $K3^{[n]}$ -type, forthcoming.
- [CS] L. Chiantini, E. Sernesi, Nodal curves on surfaces of general type, *Math. Annalen* 307 (1997), 41-56.
- [KLM] A. L. Knutsen, M. Lelli-Chiesa, and G. Mongardi, Severi varieties and Brill-Noether theory of curves on abelian surfaces, *Journal für die reine und angewandte Mathematik (Crelles Journal)*, Vol. 749 (2019), Pages 161–200.
- [KLM2] A.-L. Knutsen, M. Lelli-Chiesa, G. Mongardi, Wall divisors and algebraically coisotropic subvarieties of irreducible holomorphic symplectic manifolds, *Trans. Amer. Math. Soc.* vol. 371 (2019) no. 2, 1403–1438.
- [LS] H. Lange, E. Sernesi, Severi varieties and branch curves of abelian surfaces of type (1,3), *International J. of Math.* 13 (2002), 227-244.
- [Mon2] G. Mongardi, On the monodromy of irreducible symplectic manifolds, *Alg. Geom.* (2016) no. 3, 385–391.
- [Mark] E. Markman, The monodromy of generalized Kummer varieties and algebraic cycles on their intermediate Jacobians, [arXiv:1805.11574](#).
- [MP] G. Mongardi, G. Pacienza, Polarized parallel transport and uniruled divisors on deformations of generalized Kummer varieties, *International Mathematics Research Notices Volume 2018* (2018), no. 11, Pages 3606–3620.
- [OSY] G. Oberdieck, J. Shen and Q. Yin, Rational curves in the Fano varieties of cubic 4-folds and Gromov-Witten invariants, [arXiv:1805.07001v1](#).
- [Ono] C. Onorati, Connected components of moduli spaces of generalized Kummer varieties, [arXiv:1608.06465](#)
- [Voi15] C. Voisin, Remarks and questions on coisotropic subvarieties and 0-cycles of hyper-Kähler varieties, “K3 Surfaces and Their Moduli”, *Progress in Math*, Proceedings of the Schiermonnikoog conference 2014, Birkhäuser.

ALMA MATER STUDIORUM UNIVERSITÀ DI BOLOGNA DIPARTIMENTO DI MATEMATICA, PIAZZA DI PORTA SAN DONATO 5, BOLOGNA, 40126 ITALIA

Email address: giovanni.mongardi2@unibo.it

INSTITUT ELIE CARTAN DE LORRAINE, UNIVERSITÉ DE LORRAINE, B.P. 70239, F-54506 VANDOEUVRE-LÉS-NANCY CEDEX FRANCE

Email address: gianluca.pacienza@univ-lorraine.fr