

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Rescue tracheostomy for patients with unresectable large growing neck masses

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Puma F., Ceccarelli S., Potenza R., Italiani A., Melis A., Cagini L., et al. (2021). Rescue tracheostomy for patients with unresectable large growing neck masses. ANNALS OF THORACIC SURGERY, 112(5), e383-e386 [10.1016/j.athoracsur.2021.02.087].

Availability:

This version is available at: <https://hdl.handle.net/11585/853354> since: 2022-02-05

Published:

DOI: <http://doi.org/10.1016/j.athoracsur.2021.02.087>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1 **Rescue Tracheostomy for Patients with Unresectable Large Growing Neck**
2 **Masses**

3
4 **Running Head: Tracheostomy for Unresectable Masses**
5
6

7 **Francesco Puma, MD; Silvia Ceccarelli, MD, PhD; Rossella Potenza, MD; Alberto**
8 **Italiani, MD; Alberto Melis, MD; Lucio Cagini, MD; Massimo Monacelli, MD.**

9
10
11 **Department of Thoracic Surgery, University of Perugia Medical School, Perugia, Italy**
12
13
14
15
16

17
18 **Word count: 1500**
19
20

21 **Corresponding author: Silvia Ceccarelli MD,PhD Department of Thoracic**
22 **Surgery, S.Maria della Misericordia Hospital, 06156 Perugia, Italy; email:**
23 **s.ceccarelli1983@gmail.com**

24 **Abstract**

25

26 Most patients with undifferentiated thyroid cancer have an unresectable disease with

27 very high rate of airway compromise. Tracheostomy typically entails technical issues in

28 these cases. In fact, it can be very difficult to expose or simply locate the trachea beneath

29 the mass, and the extensive soft tissue involvement can force the surgeon to cut the

30 tumor to place the tracheostomy tube.

31

32

33 The combined use of rigid bronchoscopy and percutaneous tracheostomy techniques,

34 applied in an open surgical procedure, can greatly simplify the procedure. Furthermore,

35 by this method, the airways are quickly secured and the risk of intraoperative bleeding

36 reduced.

37 Patients with undifferentiated thyroid carcinomas frequently have serious airway-related
38 issues throughout the natural history of the disease, because of the high rate of tracheal
39 invasion and bilateral vocal cord paralysis [1]. Although the ethical dilemma of whether or
40 not to perform a tracheostomy in such a lethal form of cancer has not resolved, the surgeon
41 is often involved in managing difficult airways in this clinical setting [2,3]. Although the
42 current guidelines recommend that the tracheostomy must be avoided as long as possible in
43 patients with Anaplastic Thyroid Cancer [4], the issue of impending asphyxia frequently is
44 addressed under extremely difficult technical conditions. Tracheostomy can be a real
45 challenge in these patients mainly because the trachea is displaced and/or infiltrated by the
46 tumor and its exact position beneath the mass is not easy detectable from the operative field
47 without precise anatomical landmarks.

48
49
50
51
52

A simple technique, developed for such situation, is herein presented.

53 **Technique**

54
55
56

Clinical setting: Patients with large infiltrating neck masses, requiring urgent
tracheostomy (Fig 1-2).

57
58

The procedure must be performed in an operating room, under general anesthesia.

The steps of the technique are the following:

59
60

-The patient is placed in a supine position with his/her head slightly extended
(sniffing/intubation position).

61
62

-The techniques of pre-oxygenation and induction of anesthesia have been described
elsewhere [5].

63
64
65
66

-Tracheal intubation is performed with an 8.5 rigid bronchoscope (Storz Medical AG,
Tagerwilen, Switzerland). The rigid bronchoscope is used to reopen the airway in case
of endoluminal tumor growth; otherwise the tip of the bronchoscope is simply advanced
up until the distal trachea.

67 -Spontaneous assisted ventilation is ensured through the channel of the rigid
68 bronchoscope.

69 - Standard skin prepping and draping is performed.
70
71 -A collar cervical incision is made and the tumor obstructing tracheal access is exposed.
72
73 -Both the operating lamp and the operating room lights have to be turned off. The rigid
74 bronchoscope is withdrawn to the cricoid level. The oblique telescope at 30° of the
75 bronchoscope is faced forward and the intensity of xenon light is set at the maximum
76 level so that the airways are transilluminated and the position of the trachea can be
77 identified from the operating field.

78 -A partial debulking of the tumor can be required whenever the pretracheal mass is
79 remarkably thick. The Harmonic Scalpel (Ethicon: Somerville, New Jersey, USA) can be
80 used for this purpose. The tumor debulking is conducted until the position of the trachea
81 is detectable by the transillumination maneuver.

82 -From the operating field a 23 Gauges needle is inserted through the tumor mass into the
83 trachea, on the transillumination guide (Fig. 3A).

84 -When the endoscopic control shows that the needle is positioned in the midline of the
85 trachea at the chosen level, the needle is removed and the puncture cannula of the
86 PercuTwist set (Rüsch GmbH, Kernen, Germany) is inserted into the trachea.

87 -The internal metal guide of the puncture cannula is removed and the guide wire with the
88 flexible J-tip is pushed forward into the distal trachea through the indwelling plastic
89 cannula (Fig. 3B).

90 -The wet PercuTwist introducing dilator is advanced over the guide wire.
91
92 -The PercuTwist dilator is then screwed until the required dilation level is reached, while
93 maintaining the dilator at an obtuse angle (about 130°) with respect to the carina (Fig.
94 3C). During this maneuver, the tip of the rigid bronchoscope is positioned proximally to
95 the tracheostomy site, so that it can exert countertraction on the dilator.

96 -At this point the dilator is carefully removed, rotating it in a counter-clockwise direction
97 while the guide wire remains in the trachea.

98 -A long tracheostomy cannula with the completely deflated tube cuff is loaded onto the
99 introduction stylet (Fig. 3D) and placed into the trachea through the guide wire, using
100 Seldinger's technique.

101 -Finally, there is the removal of the introduction stylet and guide wire, the inflation of the
102 tracheostomy tube cuff, and the connection of the tracheostomy tube to a ventilator.

103

104

105

106

107

Comment

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

Patients with rapidly growing anterior neck masses may have a highly malignant thyroid cancer [5] and many of them develop severe airway problems. The aim of palliation in this setting needs to avoid death by suffocation, which is rarely achievable by endotracheal stents or laser vaporization of intraluminal growths [1].

In acute airway distress, unless the patient denies the consent, urgent tracheostomy should be considered, entailing two major technical problems related to intubation and surgical management. In our experience, the combined use of a rigid bronchoscopy and percutaneous tracheostomy techniques, but used in an open surgical procedure, has resulted in an easier and safer management, even in extremely challenging situations. Rigid bronchoscopy is to be considered the most efficient method for controlling the airway, even in critical situations [5,6]. A rigid bronchoscope allows for several advantages: the airways are under control during the entire procedure; in the case of an endotracheal tumor, the airways can be reopened; the tracheal dilation is performed under direct sight; blood and secretions are adequately suctioned and the positioning of the tracheostomy tube is carried out under direct monitoring. Furthermore, the tip of the rigid bronchoscope exerts an effective countertraction on the tracheal dilator, thus preventing any risk of a tear regarding the posterior part of the trachea during the dilation maneuver. Another significant advantage of this method is its simplicity. The transillumination allows to identify the position of the trachea underneath the tumor with relative ease and, once the needle has been introduced into the selected location of the

127 trachea, the main technical problems can be resolved, thanks to the use of the
128 percutaneous tracheostomy devices.

129 Dr. Majid et al. have already highlighted the advantages of the rigid bronchoscopy for
130 successful percutaneous tracheostomies, in patients where the latter are relatively
131 contraindicated [7]. Among the different percutaneous tracheostomy methods currently
132 available, we prefer to use the PercuTwist technique in these circumstances. This is due
133 to the fact that we found the screw dilator with its sharp threads to be highly reliable for
134 creating the stoma opening through the neoplastic tissue. Although the procedure is
135 theoretically associated with a high risk of bleeding, the PercuTwist dilator is able to
136 produce an effective mechanical haemostasis and no procedure-related bleeding has been
137 observed in our cases.

138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157

References

1. Shaha AR, Ferlito A, Owen RP, et al. Airway issues in anaplastic thyroid carcinoma. *Eur Arch Otorhinolaryngol.* 2013;270:2579-83.
2. Avenia N, Vannucci J, Monacelli M, et al. Thyroid cancer invading the airway: diagnosis and management. *Int J Surg.* 2016;28:S75-8.
3. Mani N, McNamara K, Lowe N, Loughran S, Yap BK. Management of the compromised airway and role of tracheotomy in anaplastic thyroid carcinoma. *Head Neck.* 2016;38:85-8.
4. Smallridge RC, Ain KB, Asa SL, et al. American Thyroid Association Anaplastic Thyroid Cancer Guidelines Taskforce. American Thyroid Association guidelines for management of patients with anaplastic thyroid cancer. *Thyroid.* 2012;22:1104-39.
5. Puma F, Meattelli M, Kolodziejek M, et al. An Alternative Method for Airway Management With Combined Tracheal Intubation and Rigid Bronchoscope. *Ann Thorac Surg.* 2019;107:e435-e436
6. Nicastrì, DG, Weiser, TS. Rigid bronchoscopy: indications and techniques. *Oper Tech Thorac Cardiovasc Surg.* 2012;17:44-51.
7. Majid A, Cheng GZ, Kent MS, Gangadharan SP, Whyte R, Folch E. Evaluation of rigid bronchoscopy-guided percutaneous dilational tracheostomy. A pilot study. *Ann Am Thorac Soc.* 2014;11:789-94.

158 **FIGURE LEGENDS**

159
160


161 Fig 1: Computed tomographic images of the neck: 87-year-old patient with acute
162 respiratory distress due to Anaplastic Thyroid Cancer infiltrating the airway.

163
164
165


166 Fig 2: Computed tomographic images of the neck: 84-year-old patient, with bilateral
167 vocal cord paralysis due to necrotic undifferentiated thyroid carcinoma, marginally
168 infiltrating the trachea.

168
169
170


171 Fig 3: The main steps of the technique from the view of the rigid bronchoscope: (A) the
172 needle is inserted into the trachea from the operating field; (B) the guide wire is pushed
173 forward into the distal trachea through the indwelling plastic cannula; (C) the tip of the
174 PercuTwist introducing dilator is screwed inside the trachea; (D) the introduction stylet
has been inserted and the tracheostomy tube is ready to be positioned into the trachea.






A




B




C




177




A



B



C



D

178