

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

AtPng1 knockout mutant of Arabidopsis thaliana shows a juvenile phenotype, morpho-functional changes, altered stress response and cell wall modifications

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

AtPng1 knockout mutant of Arabidopsis thaliana shows a juvenile phenotype, morpho-functional changes, altered stress response and cell wall modifications / Serafini-Fracassini D.; Della Mea M.; Parrotta L.; Faleri C.; Cai G.; Del Duca S.; Aloisi I.. - In: PLANT PHYSIOLOGY AND BIOCHEMISTRY. - ISSN 0981-9428. - ELETTRONICO. - 167:(2021), pp. 11-21. [10.1016/j.plaphy.2021.07.024]

Availability:

This version is available at: <https://hdl.handle.net/11585/855173> since: 2022-02-10

Published:

DOI: <http://doi.org/10.1016/j.plaphy.2021.07.024>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Serafini-Fracassini D.; Della Mea M.; Parrotta L.; Faleri C.; Cai G.; Del Duca S.; Aloisi I.: *AtPng1 knockout mutant of Arabidopsis thaliana shows a juvenile phenotype, morpho-functional changes, altered stress response and cell wall modifications*

PLANT PHYSIOLOGY AND BIOCHEMISTRY VOL. 167. ISSN: 0981-9428

DOI: 10.1016/j.plaphy.2021.07.024

The final published version is available online at:

<https://dx.doi.org/10.1016/j.plaphy.2021.07.024>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

1 ***AtPng1* knockout mutant of *Arabidopsis thaliana* shows a juvenile phenotype,**
2 **morpho-functional changes, altered stress response and cell wall modifications**

3

4 Serafini-Fracassini D.^{1§}, Della Mea M.^{1§}, Parrotta L.¹, , Faleri C.², Cai G.², Del Duca S.^{1*}, Aloisi I.¹

5

6 ¹ Dipartimento di Scienze Biologiche, Geologiche e Ambientali, Università degli Studi di Bologna,
7 via Irnerio, Bologna 40126, Italy;

8 ² Dipartimento di Scienze della Vita, Università degli Studi di Siena, via Mattioli 4, Siena 53100,
9 Italy;

10

11 § These authors contributed equally to the work

12 * Author to whom correspondence should be addressed; E-Mail: stefano.delduca@unibo.it; Tel. +39
13 0512091283; Fax: +39 051242576.

14

1 **Abstract**

2 In order to ascertain the role of plant transglutaminases (TGase) in growth and abiotic stress response,
3 the *AtPng1* knock out (KO) line of *A. thaliana* has been analyzed during plant development and under
4 heat and wound stress. Comparing wild type (WT) and KO lines a 58-kDa band was immunodetected
5 by anti-AtPng1p antibody in the cell wall and chloroplasts only in the WT line. A residual TGase
6 activity, not showing correlation with development nor stress response, was still present in the KO
7 line. The KO line was less developed, with a juvenile phenotype characterized by fewer, smaller and
8 less differentiated cells. Chloroplast TGase activity was insensitive to mutation. Data on stressed
9 plants showed that (i) KO plants under heat stress were more juvenile compared to WT, (ii) different
10 responses between WT and KO lines after wounding took place. TGase activity was not completely
11 absent in the KO line, presenting high activity in the plastidial fraction. In general, the mutation
12 affected *A. thaliana* growth and development, causing less differentiated cytological and anatomical
13 features.

14

15 **Key words:** *A. thaliana*, Transglutaminase, AtPng1p, Differentiation, Phenotype, Abiotic stress, Cell
16 Wall, Polyamines.

17

18 **List of Abbreviations:**

19 AtPng1p: *A. thaliana* peptide N-glycanase 1 protein

20 PAs: Polyamines

21 KO: knock out

22 TGase: Transglutaminase

23 WT: Wild Type.

24

1 **1. Introduction**

2 Plant growth is a process regulated by a cohort of environmental and internal factors, which tune a
3 coordinated morphogenesis. Several recent reviews summarize the increasing relevance of the main
4 aliphatic polyamines (PAs), in cell differentiation and wound stress response (Handa et al. 2018;
5 Aloisi et al. 2016a; Chen et al. 2019). Due to their polycationic nature, PAs easily react with other
6 biological molecules, *i.e.* DNA/RNA and proteins (Iacomino et al. 2012; Antognoni et al. 1999). In
7 plants, the formation of cross-links with hydroxycinnamic acids might also occur in cell walls
8 (Bassard et al. 2010) and in addition, by forming covalent linkages, PAs make cross-links between
9 proteins, forming supramolecular nets (Serafini-Fracassini and Del Duca 2008).

10 Transglutaminases (TGase: E.C. 2.3.2.13), a family of enzymes present in all living
11 organisms, catalyze the covalent linkages of PAs with proteins (Griffin et al. 2002; Beninati et al.
12 2013; Lorand and Graham 2003). Plants TGases are Ca^{2+} -dependent enzymes that catalyze an acyl-
13 transfer reaction between primary amino groups and specific protein-bound Gln residues. A variety
14 of primary amino groups may act as amine donors, including the ϵ -amino group of protein-bound Lys
15 residues, which result in the formation of inter- or intra-molecular protein cross-links, or one or two
16 primary amino groups of PAs to specific protein-bound Gln residues giving rise to N-(γ -glutamyl)-
17 polyamine bonds. In case of PAs, the linkage with one of the two terminal amino groups increases
18 the positive charge on the protein surface (cationization), while two linkages with both terminal amino
19 groups result in covalent crosslinking between the proteins.

20 In plants the enzyme is also present with the same mechanism of action (Aloisi et al. 2016b; Del
21 Duca and Serafini-Fracassini 2005). Several isoforms of TGases with a broad range of molecular
22 mass have been found in different organisms (Del Duca and Serafini-Fracassini 2005) and among
23 angiosperms in different organs, such as seeds, pollen, meristems and vegetative organs (Del Duca et
24 al. 2013; Campos et al. 2013; Falcone et al. 1993). TGases localize in distinct subcellular
25 compartments, *e.g.* cytoplasm, chloroplast, microsome fraction, nuclei and cell wall, suggesting either
26 different roles or different regulatory mechanisms of enzyme activity (Campos et al. 2013; Campos
27 et al. 2010; Ioannidis et al. 2012; Della Mea et al. 2007; Mandrone et al. 2019). Cytosolic enzyme is
28 translocated to the cell wall by probably two different mechanisms, both involving and secretion (Del
29 Duca et al. 2013). In mature plants, the enzyme activity is extremely low, while it is enhanced during
30 growth and development, and as response to external stimuli such stresses (Aloisi et al. 2020;
31 Sobieszczuk-Nowicka et al. 2015; Del Duca et al. 2014; Shu et al. 2020; Zhong et al. 2019).

32 There is no evidence whether these enzymes result from posttranslational modifications of a
33 single gene or whether multiple genes are present, as reported for mammals in which 9 genes are

1 known. In *A. thaliana* a single *AtPng1* gene was found, containing the catalytic domain with the Cys–
2 His–Asp triad typical of the TGases superfamily (Suzuki et al. 2000; Della Mea et al. 2004a). The
3 encoded protein, the first TGase identified in plants, was shown to have a Ca²⁺- and GTP-dependent
4 transamidase activity and to catalyze the formation of glutamyl-PA derivatives in several organs of
5 *A. thaliana* (Della Mea et al. 2004a). As some of the TGase substrates are, in addition to plastid
6 proteins, also structural proteins, like cytoskeleton and cell wall components, the activity of the
7 enzyme could be relevant for the cell structure and consequently for the tissues and the general plant
8 structure.

9 In an attempt to clarify the role of TGase on plant growth by identifying the cell component
10 eventually affected, a TGase knock-out (KO) *A. thaliana* line was phenotypically and physiologically
11 analyzed, compared to WT. Thus, the TGase 5' UTR loss-of-function line was analyzed, verifying
12 the expression, localization and activity of TGase by phenotypic observations, cyto- and histological
13 analysis of plant tissues, and quantitation of various parameters. To highlight differences in KO and
14 WT lines, TGase activity has also been stimulated by abiotic stresses, *i.e.* heat and wound treatments.
15 To our knowledge, this is the first paper in which a *AtPng1* mutant was phenotypically and
16 physiologically analyzed.

17

18

1 **2. Materials and methods**

2 2.1. Chemicals, plant material and stress conditions

3 All chemicals (unless otherwise indicated) were obtained from Sigma–Aldrich (Milan, Italy). Mutant
4 (KO) and WT lines were in the Columbia (Col-0) background. T-DNA insertional mutants
5 SALK_076538 was identified from the SALK collection (Alonso et al. 2003). All phenotypic
6 characterizations were performed with homozygous plants (<http://natural.salk.edu/geno/sum.txt>). A
7 map of the gene structure of the three T-DNA insertion mutant lines is reported in Supplementary
8 material.

9 All plants were grown in greenhouse conditions (23°C) with 16 h light /8 h dark (light
10 intensity, 55 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$). Heat stress treatment was performed on surface-sterilized seeds
11 plated on half-strength MS (Murashige and Skoog 1962) medium (2.15g/L), sown at 30 °C and
12 allowed to develop for 5 and 10 days.

13 With a razor blade, wound stress was performed by a longitudinal radial cut of about 2 mm
14 deep until the medullary parenchyma on stems of two-months old plants. Transverse slices cut and
15 stained as reported below, were microscopically observed at different days (day 0, namely
16 immediately -1-5-9-14 until day 21) in the region between the basal rosette leaves and the first node.
17 Leaves were injured with 5 cuts perpendicular to the midrib and analysed at different times. The other
18 leaf half symmetrically opposite to the wounded half was left unscathed. Controls are represented by
19 not-wounded leaves of the same age and taken from the same plant.

20

21 2.2. mRNA extraction and TGase expression analysis

22 mRNA was isolated by the kit Nucleospin RNA plant (Macherey-Nagel) starting from 100 mg of leaf
23 tissue and following manufacturer's instruction. mRNA quality was checked spectrophotometrically
24 and by gel electrophoresis. Ten micrograms of RNA were used to synthesize cDNAs using oligo(dT)
25 primers according to manufacturer's instructions (Invitrogen, San Giuliano Milanese, Italy). TGase
26 and actin (housekeeping gene) amplification was performed as previously described (Della Mea et
27 al. 2004a). Amplification products were checked on 1% agarose gel and the density of the bands was
28 analysed by the AIDA software (Fuji Inc., Tokyo, Japan). Primers used are listed below:

29 AtPng1 FW: TTGTCGCTGCACGAGATAACG; AtPng1 REV: CTTCTGTACAGATCGATG
30 CTCCC; Actin FW: GACTCTGGAGATGGTGTG; Actin REV: ATCTGCTGGAAGGTACTGAG

31

32 2.3. Preparation of cellular fractions, protein extraction and Western-blot analyses

1 Cell wall, microsomal, soluble and plastidial fractions were obtained from 21-d-old *A. thaliana* leaves
2 according to previous literature with some modifications (Bregoli et al. 1997). Nuclei enriched
3 fraction was prepared as previously described (Zhao et al. 2001).

4 Total proteins were extracted as previously described (Parrotta et al. 2020), from 21-d-old
5 plants. In detail, entire plants were grinded in liquid nitrogen and proteins were extracted in agitation
6 for 30 min at 4 °C in extraction buffer (60 mg/mL) containing 100 mM Tris-HCl pH 8.5, 10 mM 2-
7 Mercaptoethanol, 0.2% Triton X-100 and protease inhibitor cocktail. Large cell debris were removed
8 from the total homogenate by centrifugation at 12,000 rpm for 10 min at 4 °C. Protein concentration
9 was estimated on the supernatant by the Bradford method with bovine serum albumin (BSA) as the
10 standard protein. Extraction was repeated in triplicate. 30 µg of protein from each enriched fraction
11 were loaded onto a denaturing 10% (w/v) SDS-PAGE gel and migrated using a Mini-protean II
12 apparatus (Bio-Rad, Italy). Protein were blotted to a nitrocellulose membrane (Amersham
13 Biosciences, Buckinghamshire, UK) using a semidry Trans-Blot system (Bio-Rad, Italy). Western
14 blotting was performed as previously described, using an anti AtPNG1 antibody as primary antibody
15 (1:2000 dilution) (Della Mea et al. 2007). An anti-chicken antibody conjugated to alkaline
16 phosphatase was used as a secondary antibody (1:3000 dilution).

17

18 2.4. Nε-(γ glutamyl)-lysine quantification

19 The presence of the TGase activity product, the Nε-(γ glutamyl)-lysine was analysed by
20 immunodetection with 81D4 antibody (Covalab, Bron, France) in an ELISA assay. ELISA was
21 carried out as described previously (Mandrone et al. 2019). As positive control X-linked casein was
22 tested in the same conditions. Casein (final concentration 2 mg/mL) X-linking was induced for 2
23 hours in 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 2.5 mM CaCl₂, 10 mM DTT in presence of 5 µg of
24 Guinea pig liver (gpl) TGase. The final volume of the reaction was 250 µL. For negative control,
25 casein was incubated in the same conditions with the omission of gpl TGase.

26

27 2.5. Analysis of TGase activity: microplate-based and radioactivity assays

28 The *in vitro* TGase activity was measured by the conjugation of biotinylated cadaverine to N, N'-
29 dimethylcasein as previously described (Paris et al. 2017; Del Duca et al. 2018). Specific activity was
30 determined as a change in A450 of 0.1 per hour per mg of protein after subtraction of the value of the
31 controls treated with 20mM EGTA. Alternatively, TGase activity was radioactively determined (Del
32 Duca et al. 2018; Della Mea et al. 2004a).

33

1 2.6. Chlorophyll and carotenoids determination

2 Chlorophyll *a* and *b*, as well as total carotenoids (xanthophylls and carotenes) were extracted with
3 cold 80% acetone then centrifuged at 4000 g for 10 min at 4 °C. Their levels were determined
4 according to the literature (Porra et al. 1989).

5

6 2.7. Microscopy analysis and immunolocalization of TGase

7 Fresh inflorescence stems were transversely cut between the basal rosette leaves and the first node
8 and observed unfixed. Slices were observed by optical microscopy either by white light or by
9 fluorescence microscopy at 380 or at 450 nm. Observations were made using a Zeiss Axio Imager
10 microscope. Images were acquired with an AxioCam MRm camera using the software AxioVision.
11 Specific staining for lignin and suberin were performed by acid floroglucine and Sudan III,
12 respectively (Angelini et al. 2008). The measures of the cells were performed by NIS Element
13 program (Nikon). Measures of the major cell diameters or of the wall thickness were repeated ten
14 times and the mean calculated and reported as frequency classes.

15 For TGase immunolocalization, inflorescence was directly thawed in a buffer solution (100
16 mM Pipes pH 6.8, 10 mM EGTA, 10 mM MgCl₂, 0.1% NaN₃) plus detergent and fixative (0.05%
17 Triton X-100, 1.5% paraformaldehyde, 0.05% glutaraldehyde) for 30 min on ice and then at 4 °C for
18 an additional 30 min. For the localization of TGase, samples were cut along their length and placed
19 in the buffer solution containing 0.75% cellulysin and 0.75% pectinase for 7 min. For
20 immunofluorescence microscopy, samples were washed in the above buffer and incubated with the
21 anti AtPNG1 antibody diluted 1:20 in the buffer; incubation was 1 h at 37 °C according to previous
22 literature (Del Duca et al. 2013). After washing with buffer, samples were incubated with The
23 secondary antibody was Alexa Fluor 488-conjugated goat anti-chicken (Thermo Fisher Scientific)
24 secondary antibody diluted 1:50 in the buffer solution, for 45 min at 37 °C in the dark. Samples were
25 observed with a Zeiss Axiophot fluorescence microscope (Ex-Max 490 nm/Em-Max 525 nm)
26 equipped with an MRm video camera and a 63× oil immersion objective.

27

28 2.8. Data and statistics

29 Each experiment was repeated at least three times. Differences between samples sets were determined
30 by analysis of variance (one-way ANOVA, with a threshold P-value of 0.05) using GraphPad Prism.

31

1 3. Results

2 3.1. The enzyme TGase is present in different cellular compartments in *A. thaliana*.

3 To investigate the presence and distribution of TGase in the *A. thaliana* WT line, TGase activity was
4 investigated in several cellular compartments, *i.e.* cell wall, nuclei, chloroplasts as well as in the
5 microsomal and soluble fractions (Fig. 1A). Apart from the soluble fraction, where enzymatic activity
6 was under the detection limit, all other subcellular compartments showed TGase activity, with the
7 cell wall fraction having about 5 times more activity in comparison to nuclei, and the microsomal
8 fraction and 3 times in comparison to chloroplasts. The higher activities in cell walls and chloroplasts
9 could be due to a more abundant accumulation of TGase in these fractions, as shown by western
10 blotting with the anti-AtPNG1 polyclonal antibody, which proved a cross-reacting band of 58 kDa
11 (Fig.1B). No immuno-signal was detected in the nuclei compartment, nor in the microsomal and
12 soluble fractions (data not shown).

13
14 3.2. The *AtPng1* KO line shows altered morphological differentiation

15 KO line was compared with that of the WT line in an attempt to verify possible morphological
16 differences caused by its loss of function. As shown in Fig. 2A, the WT line was characterized by
17 many basal rosette leaves with several cauline inflorescences, relatively large and whose stems were
18 pink-pigmented (Fig. 2A, arrow). The cauline leaves were lost during plant growth and numerous
19 siliques were produced. Compared to WT, KO plants showed a reduced number of rosette leaves, as
20 well as numerous larger green cauline leaves (insert in Fig. 2A) with inflorescences characterized by
21 fewer siliques. Cauline leaves in the KO line were characterized by a higher percentage of fresh
22 weight: 92.6 ± 1 with respect to 90.1 ± 0.6 in WT and by a higher content in chlorophylls *a/b* (96.4
23 $\mu\text{g/g fw} \pm 4.8$ in KO and $90.5 \mu\text{g/g fw} \pm 0.9$ in WT). Carotenoid content was not significantly different
24 ($9.5 \mu\text{g/g fw} \pm 1$ in KO and $10.5 \mu\text{g/g fw} \pm 0.8$ in WT).

25 The KO and WT lines differed in stem diameter, due to a reduced development mainly of the
26 sclerenchyma and medullar tissues in the KO (Fig. 2C) compared to the WT line (Fig. 2B). The
27 photosynthetic cortex of WT (Fig. 2B, red fluorescence) surrounded a thick layer of sclerenchyma
28 cells (green fluorescence). In the WT line, the sclerenchyma developed as a thick continuous layer
29 deepen into the medulla which incorporated several large vascular bundles. This sclerenchyma layer
30 was less developed in the KO line (Fig 2C). In the inserts of Fig. 2B-C epidermal, cortical and
31 sclerenchyma tissues are shown at high magnification. The multiple nonhomogeneous cell layers
32 formed a continuous sclerenchyma ring in the WT line (insert in Fig. 2B, yellow line), while the

1 sclerenchyma in KO line was formed by only 3 or 4 regular cell layers separated in short arches which
2 did not include vascular bundles (insert in Fig. 2C, yellow line).

3 Details of the basal stem sections of WT and KO lines at the same age are shown in Fig. 3. In
4 particular, the sclerenchyma layer (blue arrows) and the vascular bundles (yellow arrows) were
5 considerably larger in the WT stem (Fig. 3A) compared to KO stem (Fig. 3B). In the WT stem (Fig.
6 3C) the cortex layer (red arrow) showed a pink color in the subepidermal region (white arrow) whose
7 color was visible also macroscopically, as observed in Fig. 2A, whereas these pigments were absent
8 in the green photosynthetic cortex of KO stem (Fig. 3D).

9 Histological observations of the stem tissues of both lines are comparatively shown in Fig. 4
10 and in Fig. 5. In WT, the epidermal cells were larger, with a thicker cell wall and distributed in several
11 size classes (ranging from 6 to 30 μm) compared to the KO (Fig. 4A1 and 4A3). In the KO line, cells
12 were smaller, missing the largest categories (maximum 18-21 μm) (Fig. 4A2 and 4A3). Xylem vessels
13 showed well-developed cell walls in the WT line (Fig. 4B1 and 4B2 respectively). In the KO line, the
14 diameter range of the highest percentage of vessels was smaller when compared to the highest
15 percentage diameter range of WT vessels (Fig. 4B3).

16 Sclerenchyma cells showed variable cell diameters in WT (Fig. 5A1 and 5A2) whereas KO
17 was characterized by a prevalent frequency of small cells, large ones being sporadic (Fig. 5A3). Also
18 sclerenchyma cell wall thickness was reduced in the KO line, where about 70% of cells showed a cell
19 wall thickness of 0.5-1 μm (Fig. 5B1, 5B2 and 5B3).. In summary, epidermis, vessels and
20 sclerenchyma all showed that in KO cells were homogeneous, smaller with the cell walls apparently
21 thinner.

22 Summarizing the morphological and histological evidence, KO plants showed a reduced
23 differentiation and altered development, presenting a pronounced juvenile phenotype, characterized
24 by a lower number of floral stems and rosette leaves, by the longer permanence of cauline leaves on
25 the stem, by the lower production of siliques and by cytological features.

26 3.3. TGase enzymatic activity persist in the plastid fraction of the *AtPng1* KO line

27 The WT line showed a TGase of 58 kDa, band that was not detectable in the KO line (Fig. 6A).
28 Concomitantly, the isopeptide N ϵ -(γ glutamyl)-lysine, the product of *in vitro* TGase reaction, was
29 significantly higher in the WT line compared to the KO mutant (Fig. 6B) .

30 To detect the enzymatic activity *in vitro*, whole plants of WT and KO were analyzed along
31 with their life span during vegetative and reproductive phases. In the WT line, TGase activity
32 significantly increased during plant growth, decreasing after 60-70 days when plants developed the
33

1 inflorescence, and reaching the lowest value during senescence. On contrary, TGase activity in KO
2 line was halved compared to the WT line and did not significantly change during growth. In the KO
3 line, TGase activity decreased after inflorescence development (90-100 days), to values not
4 significantly different from those of the WT line (Fig. 6C).

5 As a critical point *in vivo* is the onset of flowering, TGase activity was determined in the two
6 life stages, vegetative and reproductive, in the organelles of rosette leaves and of inflorescence (Fig.
7 7). The activity was prevalent in the chloroplast fraction and it was not significantly different in WT
8 or KO, either when measured in the inflorescence or rosette. Soluble and microsomal fraction showed
9 a low activity in both lines.

10 Together, these data showed that the KO line did not present (or its level is too low to be
11 detected) the 58 kDa protein immuno-recognized in the WT line by the antibody against AtPNG1P;
12 nevertheless, a low level of TGase activity was present in the plastid fraction of the KO line.

13

14 3.4. KO and WT lines differently respond to abiotic stresses

15 As TGase activity was shown to be at basal level in physiological conditions, to further emphasize
16 the differences between KO and WT, plants of both lines were subjected to heat and wound stresses.
17 Seeds of both lines were sown on sterile medium at 30 °C and plantlets were allowed to develop for
18 5 and 10 days. Both roots and cotyledons were affected by heat stress, as shown in Fig. 8. Compared
19 to WT (Fig. 8A1, B1), the KO line showed a reduced primary roots development , followed by
20 impaired development of lateral roots after 5 days (Fig. 8C1) and after 10 days (Fig. 8D1). WT
21 cotyledons were green after 5 days (Fig. 8A2) and developed trichomes after 10 days (Fig. 8B2),
22 whereas those of the KO line were smaller, and accumulated red pigments (Fig. 8C2); cotyledons
23 neither grew significantly nor differentiated trichomes until day 10 (Fig. 8D2).

24 Leaves and inflorescence stems were studied also after wound stress. The change in TGase
25 activity in WT leaves at the wound was rapid but transient (Fig. 9A). TGase activity increased within
26 15 min of about three-fold in respect to the basal activity observed at time 0 (Fig. 9B), remained
27 constant until 30 min and decreased further on. On the contrary, in KO leaves the wound was
28 ineffective, being the activity constantly at low level, until 24 h.

29 Wound induced *AtPng1* transcription within the first 5 min and reached minimum
30 transcription levels after 15-30 min in the WT line (Fig. 9C). Subsequently, TGase expression
31 increased after 1 hour and it remained visible for several hours. In the KO line, wound stress in leaves
32 did not induce any stimulation of *AtPng1* transcription (Fig. 9C).

1 Wound stress was induced also on the cauline inflorescence, and the progression of wound
2 repair was studied until day 21. The non-wounded unstained stems (day 0, Fig. 10A) showed a
3 chlorophyll rich subepidermal tissue (characterized by red autofluorescence at 450 nm), a multilayer
4 of peripheral sclerenchyma (green under fluorescence) and a medullary parenchyma (dark under
5 fluorescence and colorless under white light). In the WT line, at day 1 (Fig. 10B, d1) no signs of
6 repair were visible, whereas on day 5 (Fig. 10C, d5) a lignin- and suberin-enriched tissue (data not
7 shown), impermeabilized the wound edges (Fig. 10C, red arrows). The underlying cells of the
8 medullary parenchyma enlarged and divided disorderly, forming a small mass of meristematic tissue
9 (Fig. 10C, d5, green arrow). At day 9 the protecting process continued (Fig. 10D2, red arrows and
10 details in D1 and D3). At day 14 the shape of the wound assumed a “V” form, forming an angle of
11 about 30 degrees (Fig. 10E1). Under some dead cells, the wound edges were straight and formed by
12 a bi- three-layer of modified cells, especially in their external cell walls (Fig. 10E2, E3). Wound edges
13 appeared protected by protruding tissue (Fig. 10E1, red arrows). At day 21 the edges of the wound
14 were straight (red parenthesis in Fig. 10F1 and F3), formed by cells with a modified cell walls
15 (fluorescent towards the wound, but not modified towards medullary parenchyma, Fig. 10F2). Some
16 parenchyma cells were still dividing, showing two brother nuclei on either side of the newly formed
17 wall (Fig. 10F4, green arrow).

18 The most significant difference between the WT and KO lines was the shape of wound, as
19 shown in Fig. 11. Whereas in WT (Fig. 11A) the wound had a “V” shape with straight sides and
20 protruding edges, in the KO line the wound assumed a “U” shape (Fig. 11B). The layer of repair cells
21 was thinner and the underlying parenchyma was formed by large spherical cells (details in Fig. 11C
22 and D).

23 As TGase was identified in a highly active form in the cell walls (Fig. 1A), its presence in
24 repairing cells was investigated by fluorescence microscopy with AtPNG1P antibody. At time 0 after
25 wound induction, TGase was hardly immunodetected (Fig. 12A), while at day 21 wound edges clearly
26 showed TGase labeling (Fig. 12B, white arrows). In the KO mutant, TGase was not immunodetected
27 at time 0 nor after 21 days (Fig. 12C and D).

28

29 **4. Discussion**

30 The comparison between KO mutant and the WT line clarify TGase role during *A. thaliana*
31 development. The data here reported allow concluding that AtPNG1P and its TGase activity are
32 involved in plant cell development and in stress response. Macroscopically, the KO line showed
33 delayed and reduced development, reduced dimension of the inflorescence, bigger cauline leaves and

1 a decreased production of smaller siliques. Anatomically, in the KO, vascular bundles were less
2 numerous and smaller and the sclerenchyma layer in the stem was less developed. Also, the cortical
3 layer was less developed and did not present the pink color typical of WT stem. Epidermis, vessel,
4 and sclerenchyma cells were significantly smaller in the KO line and the thickness of the
5 sclerenchyma cell wall was significantly reduced. All those aspects were related to a less
6 differentiated secondary cell walls, and to a reduced number of cells, generally of smaller dimension
7 in KO than WT.

8 The macro- and microscopical differences between the two lines could be related to the
9 absence of a 58 kDa TGase in the KO line, which, on the contrary, was clearly present in chloroplasts
10 and in cell walls of WT. Nevertheless, a low residual TGase activity was detectable also in the KO
11 line, suggesting the possible existence of other active forms, mainly in the chloroplasts. The
12 hypothesis of the existence of other TGase forms is supported by data in literature (Della Mea et al.
13 2004b). The 58 kDa TGase form is probably the most common one, also in other plants, as shown in
14 the plastid, microsome and cell wall fractions of the *Nicotiana* flower corolla (Della Mea et al. 2007).
15 Other TGase isoforms have been identified in different plant organelles, for example, a 52 kDa
16 isoform was found in the soluble fraction, while 39 and 58 kDa isoforms were studied in the
17 chloroplast of higher plants (Dondini et al. 2003). In the apical meristem of *Helianthus tuberosus*,
18 three isoforms of 58, 75 and 85 kDa have been identified, while a 70 kDa TGase was found in apple
19 pollen (Beninati et al. 2013; Di Sandro et al. 2010; Del Duca et al. 2009; Serafini-Fracassini and Del
20 Duca 2008). These forms could derive by post-translational modification of a single protein of 86
21 kDa immunodetected in *Arabidopsis* enriched microsomal fraction, identified as the first plant TGase
22 by Della Mea et al. (2004).

23 The residual activity in KO could thus be tentatively attributed, at least in part, to TGase
24 isoforms presents in the chloroplasts. In fact, plastidial TGase activity was not significantly different
25 in WT or KO, either when measured in the inflorescence or in the basal leaf, probably due to a
26 chloroplast-encoded isoform.

27 Here it was found that KO residual TGase activity poorly correlated to the development
28 events; in WT, on the contrary, the activity increased with stem growth and decreased during the
29 flowering stages, namely approaching senescence.

30 The effects of the KO mutation, could be explained at least in part, by highlighting the
31 involvement of PAs, substrates of TGase, in the deposition of secondary wall, in cell differentiation
32 and cell death in early metaxylem (Tisi et al. 2011; Aloisi et al. 2017). In addition, the amount of PAs
33 is known to be severely modified after different stresses (Minocha et al. 2014; Tiburcio et al. 2014;

1 Liu et al. 2015; Aloisi et al. 2016a), and induced stress conditions enhanced the differences between
2 WT and KO. Nevertheless, TGase might also be directly affected by stress, as shown, for example,
3 by salt and light stresses in *Dunaliella salina* (Dondini et al. 2001). Finally, wound stress has been
4 reported to affect the neo-synthesis and activity of two TGase forms, one of which of 58-kDa, in
5 dormant tissues of *Helianthus tuberosus* when activated *in vitro* (Del Duca et al. 2000).

6 In the present work, plantlets of the TGaseKO line showed a more juvenile aspect, both under
7 standard and heat stress conditions. These evidences could be explained, at least in part, by the
8 involvement of PAs, in particular spermine, in *A. thaliana* stress tolerance (Sagor et al. 2013). Finally
9 higher levels of free and bound PAs were reported in heat tolerant rice (Pang et al. 2007). In stressed
10 KO plantlets, as suggested by their more juvenile aspect, a lower catalysis of conjugated PAs can be
11 hypothesized, due to the lack of 58 kDa TGase. By transcriptomic assays, among thousand genes
12 involved in tomato heat-stressed seedlings, as unique report, at least to our knowledge,
13 overexpression of TGase was found to enhance, within few hours, heat tolerance by maintaining
14 membrane integrity and holding higher net photosynthetic rate. Numerous genes were closely
15 associated with the protein processing in the endoplasmic reticulum, carbon fixation, and
16 photosynthetic metabolism (Jahan et al. 2021).

17 The possibility that TGase can participate in healing processes is rather unknown for plants
18 but already demonstrated for some animals (Greenberg et al. 1991; Telci and Griffin 2006; Qin et al.
19 2013). Present data confirmed that wound stress affected TGase, thereby causing clear differences in
20 reactivity between WT and KO lines. In WT, the transcription of TGase occurred very rapidly within
21 5 minutes and allowed the enzyme activity to reach a three-fold peak already within 15 minutes, while
22 in the KO line the enzyme transcription was not detectable.

23 The evaluation of wound stress in the stem allowed to observe the recovery from wounding
24 for several days and the formation of a scarring tissue that covered the entire wound. TGase
25 accumulated in the stem of the WT line, in the 2-3 cell layers underneath some superficial dead cells.
26 This distribution pattern suggests that TGase could be involved in the injury response, probably by
27 exerting its glue function and strengthening cell walls, as previously observed during senescence in
28 *Nicotiana* petals (Della Mea et al. 2007). The lack of TGase activity in the cell walls of the KO line,
29 might be related to the weaker anatomical structure observed, characterized by large intercellular
30 spaces in the parenchyma, and large, spherical parenchyma cells, which is probably due to reduced
31 stiffness of the KO cell wall, failing to counteract the internal turgor pressure (Carter et al. 2017). The
32 different cell features of the wounded stem tissues cause differences in the shape of the wound,
33 exhibiting a “V” shape in WT and a “U” shape in KO line (Waffenschmidt et al. 1999; Iranzo et al.

1 2002). The cytological comparison between WT and *AtPng1* KO plants suggested that many of the
2 differences between KO and WT lines were related to a different distribution of TGase in the cell
3 walls of the scarring layer, probably leading to a less differentiated secondary cell walls in the former.

4 Present data allow to assume the important role of cell wall TGase in cell differentiation,
5 organization, and plant development. In fact, the *AtPng1* KO was characterized by lower development
6 and by a reduced reactivity to abiotic stresses when compared to WT. In conclusion, our data highlight
7 the involvement of TGase, and consequently probably its reaction products, in the normal growth and
8 differentiation of *A. thaliana*, as shown by its phenotypic and physiological profiling analysis.

9 **Fundings**

10 This research did not receive any specific grant from funding agencies in the public, commercial, or
11 not-for-profit sectors.

13 **Declaration of competing interest**

14 The authors declare that they have no known competing financial interests or personal relationships
15 that could have appeared to influence the work reported in this paper. The research involves neither
16 human nor animals.

18 **Credit authorship contribution statement**

19 **I. Aloisi** and **L. Parrotta**: Investigation, Methodology, Formal analysis, Writing - original draft.

20 **C. Faleri** and **M. Della Mea**: Investigation, Methodology, Formal analysis.

21 **G. Cai**, **S. Del Duca** and **D. Serafini-Fracassini**: Investigation, Writing - review and editing.

22
23 **Acknowledgements:** We are very grateful to Joan Rigau, Inma Claparols and David Caparròs-Ruiz
24 of the Consorci CSIC-IRTA, Laboratory de Genètica Molecular vegetal, Barcelona, Spain, for their
25 help in growth of *A. thaliana* plants mutants. We are grateful to Saïd El Alaoui (Covalab, Bron,
26 France) for useful indications regarding the 81D4 antibody. Thanks also for their collaboration during
27 their undergraduate thesis to Dr. E. Turco, S. Di Rubbo, F. Costa, and F. De Filippis. We thank the
28 Salk Institute Genomic Analysis Laboratory for providing the sequence-indexed Arabidopsis TDNA
29 insertion mutants and NASC for providing our laboratory with the plant material.

1
2
3

1 **Figure legends**

2 **Fig. 1. TGase in WT.** TGase activity in subcellular enriched fractions, *i.e.* Cell Wall (CW), Nuclei,
3 microsomal (MICRO), soluble (SOL) and chloroplast (CHL). Means were compared by Dunnett's
4 Multiple Comparison Test. Bars with the same letter are not significantly different ($p \geq 0.01$) (**A**).
5 Immunorecognition in Western blot analysis by the *A. thaliana* TGase antibody (AtPNG1P) of the
6 cell walls (CW), and chloroplast (CHL) fractions. Non-immunoreacted fractions (*i.e.* Microsomal,
7 Nuclei, Soluble) are not shown (**B**).

8

9 **Fig. 2. Comparison of WT and KO lines.** Morphological comparison of 3 months-old plants. Arrow
10 highlights pink color pigmentation in WT stems (**A**). Particular of the caulinar leaves is reported in
11 the squared detail. Stem transverse section observed by fluorescence microscopy in WT line (**B**) and
12 *AtPng1* KO line (**C**). Green fluorescence: lignified sclerenchyma and xylem elements. Red
13 fluorescence: chlorenchyma. The yellow lines in the magnifications reported in squares label the
14 thickness of the sclerenchyma layers. Bars in figure B and C 250 μm ; bars in inserts 50 μm .

15

16 **Fig. 3. Histological comparison of WT and KO lines.** Basal stem transverse section in WT line,
17 stained with acid fluoroglucine (**A**) and unstained KO (**B**). Blue arrows indicate the thickness of the
18 sclerenchyma layer; yellow arrows indicate the radial diameter of vascular collateral bundles. Detail
19 of the unstained cortex of WT (**C**) and stained KO (**D**) inflorescence. Red arrows mark the cortical
20 parenchyma in both lines; in WT the external layers showed a natural pink color (C, white arrow).
21 Bars: 50 μm .

22

23 **Fig. 4. Comparison of cell dimension of WT and KO line epidermis and vessels.** Transverse
24 section of the epidermis of the basal stem in WT (**A1**) and KO (**A2**) lines. Comparison of epidermis
25 cell diameter, recorded by dimension classes, is reported in **A3**. Transverse section of the vascular
26 bundles of the basal stem in WT (**B1**) and KO (**B2**) lines. Comparison of vessel diameter, recorded
27 by dimension classes, is reported in **B3**. Bars: 50 μm .

28

29 **Fig. 5. Comparison of cell dimension and cell wall thickness of WT and KO line sclerenchyma.**
30 Transverse section of the basal stem in WT (**A1**) and KO (**A2**) lines. Comparison of sclerenchyma
31 cell diameter, recorded by dimension classes, is reported in **A3**. Transverse section of the basal stem
32 in WT (**B1**) and KO (**B2**) lines. Comparison of thickness of sclerenchyma cell wall, recorded by

1 dimension classes, is reported in **B3**. Bars in figure A1 and A2 50 μm ; bars in figure B1 and B2 10
2 μm .

3

4 **Fig. 6. TGase presence and activity in WT and KO lines.** Immunodetection by Western blotting
5 of TGase in WT and KO. Total lysate was probed (A). TGase products detected in ELISA assay by
6 the 81D4 antibody in WT and KO lines. Casein was used as internal control. C+ = TGase-treated
7 casein, C- = untreated casein (B). Total TGase activity in WT and KO lines during *in vivo* plant
8 growth (from 45 to 126 days) (C). Means of samples were compared by Dunnett's Multiple
9 Comparison Test. *** = $p \leq 0.001$

10

11 **Fig. 7. Comparison of the subcellular activity of TGase of the WT and KO lines.** The activity
12 was measured in soluble (SOL), microsomal (MICRO) and chloroplasts (CHL) fractions of leaves
13 and inflorescence. Means were compared by Dunnett's Multiple Comparison Test. Bars with the same
14 letter are not significantly different ($p \geq 0.01$).

15

16 **Fig. 8. Comparison of WT and KO lines germinating seeds under heat stress.** Seeds were exposed
17 to 30°C for 5 and 10 days. Root development of WT (A1) and KO (C1) and cotyledon development
18 of WT (A2) and KO (C2) at 5 days. Root development of WT (B1) and KO (D1) and cotyledon
19 development of WT (B2) and KO (D2) at 10 days.

20

21 **Fig. 9. TGase expression and activity after leaf wounding in WT and KO lines.** Leaves from the
22 basal rosette were wounded by longitudinal cutting along the mid-rib (A). Time-course TGase activity
23 in leaves until 24 hours after wounding. Means of samples were compared by Dunnett's Multiple
24 Comparison Test. *** = $p \leq 0.001$ (B). Time-course expression of AtPng1 revealed by PCR after
25 wounding (C). Actin was used as housekeeping gene.

26

27 **Fig. 10. Time-course histological observations after stem wounding in WT line.** Transversal
28 sections of the basal inflorescence observed by white light or fluorescence microscopy before
29 longitudinal radial wounding; part of the collateral vascular bundles and its phloem (Phl) are in
30 evidence (A). At day 1 the radial wound (W) enlarged laterally (B). At day 5, a new irregular
31 protection cell layer surrounded the wound edges (red arrows) covering the wounded epidermis,
32 photosynthetic cortex, phloem and connected with sclerenchyma (green fluorescence in squares). The
33 medullary parenchyma formed new cells, some of which very large (green arrow) (C). At day 9, the

1 wound surface was completely covered by neo-formed protective tissue (blue fluorescence, red
2 arrows) (**D2**). Phl was completely covered (details in **D1** and **D3**). At day 14, the wound enlarged in
3 a “V” shape. The thickness of the new protective layer increased covering the wound margin (**E1**, red
4 arrows), and magnification of the wound bottom in **E2**) and their cells show lignified cell walls (**E3**).
5 At day 21, the protective tissue (red arrows, **F1**) and a lignified cell layers still increased (**F2**), as
6 highlighted by autofluorescence (red parenthesis, **F3**). Dividing cells were visible below the
7 protecting layer (green arrow, **F4**). Bars in figure A, B, C, D1, D3, E2, E3, F2 and F3 50 µm; bars in
8 figure E1 and F4 20 µm. Bar in figure D2: 250 µm. Bar in figure F1: 200 µm.

9

10 **Fig. 11. Comparative stem anatomy after wounding in WT and KO lines.** Sections of the
11 inflorescence stained by acid floroglucine and observed after 21 days of WT line (**A**) and of KO line
12 (**B**) and relative magnifications (**C-D**). Bars: 50 µm.

13

14 **Fig. 12. Comparative immunofluorescence of TGase distribution after wounding of stem in WT**
15 **and KO line. (A-B) WT line. (A)** At time 0, anti-TGase (AtPNG1p and secondary Alexa-Fluor 488
16 antibodies) labelling was weakly diffused. (**B**) At day 21, TGase labelling was located along the
17 wound edge (white arrows). (**C-D**) KO mutant line. (**C**) Anti-TGase labelling at day 0 revealed a
18 pattern similar to WT. (**D**) At day 21, the wound edge exhibits weak or no labelling. The red dotted
19 lines indicate the wound position. Bars: 50 µm.

20

21

1 **References**

- 2 Aloisi I, Cai G, Faleri C, Navazio L, Serafini-Fracassini D, Del Duca S (2017) Spermine Regulates
3 Pollen Tube Growth by Modulating Ca²⁺-Dependent Actin Organization and Cell Wall
4 Structure. *Front Plant Sci* 8. doi:10.3389/fpls.2017.01701
- 5 Aloisi I, Cai G, Serafini-Fracassini D, Del Duca S (2016a) Polyamines in Pollen: From
6 Microsporogenesis to Fertilization. *Front Plant Sci* 7:155. doi:10.3389/fpls.2016.00155
- 7 Aloisi I, Cai G, Serafini-Fracassini D, Del Duca S (2016b) Transglutaminase as polyamine mediator
8 in plant growth and differentiation. *Amino Acids* 48 (10):2467-2478. doi:10.1007/s00726-
9 016-2235-y
- 10 Aloisi I, Distefano G, Antognoni F, Potente G, Parrotta L, Faleri C, Gentile A, Bennici S, Mareri L,
11 Cai G (2020) Temperature-dependent compatible and incompatible pollen-style interactions
12 in *Citrus clementina* Hort. ex Tan. show different transglutaminase features and polyamine
13 pattern. *Front Plant Sci* 11:1018. doi:10.3389/fpls.2020.01018
- 14 Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman J,
15 Barajas P, Cheuk R (2003) Genome-wide insertional mutagenesis of *Arabidopsis thaliana*.
16 *Science* 301 (5633):653-657. doi:10.1126/science.1086391
- 17 Antognoni F, Del Duca S, Kuraishi A, Kawabe E, Fukuchi-Shimogori T, Kashiwagi K, Igarashi K
18 (1999) Transcriptional inhibition of the operon for the spermidine uptake system by the
19 substrate-binding protein PotD. *J Biol Chem* 274 (4):1942-1948. doi:10.1074/jbc.274.4.1942
- 20 Bassard JE, Ullmann P, Bernier F, Werck-Reichhart D (2010) Phenolamides: bridging polyamines to
21 the phenolic metabolism. *Phytochemistry* 71 (16):1808-1824.
22 doi:10.1016/j.phytochem.2010.08.003
- 23 Beninati S, Iorio RA, Tasco G, Serafini-Fracassini D, Casadio R, Del Duca S (2013) Expression of
24 different forms of transglutaminases by immature cells of *Helianthus tuberosus* sprout apices.
25 *Amino Acids* 44 (1):271-283. doi:10.1007/s00726-012-1411-y
- 26 Bregoli A, Crosti P, Cavallini A, Cionini G, Del Duca S, Malerba M, Natali L, Serafini-Fracassini D,
27 D'amato F (1997) Nuclear DNA distribution and amitotic processes in activated *Helianthus*
28 *tuberosus* tuber parenchyma. *Plant Biosystems* 131 (1):3-12.
29 doi:10.1080/11263504.1997.10654161
- 30 Campos A, Carvajal-Vallejos PK, Villalobos E, Franco CF, Almeida AM, Coelho AV, Torné JM,
31 Santos M (2010) Characterization of *Zea mays* L. plastidial transglutaminase: interactions
32 with thylakoid membrane proteins. *Plant Biol* 12:712-716. doi:10.1111/j.1438-
33 8677.2009.00280.x
- 34 Campos N, Castañón S, Urreta I, Santos M, Torné JM (2013) Rice transglutaminase gene:
35 identification, protein expression, functionality, light dependence and specific cell location.
36 *Plant Sci* 206:97-110. doi:10.1016/j.plantsci.2013.01.014
- 37 Carter R, Woolfenden H, Baillie A, Amsbury S, Carroll S, Healicon E, Sovatzoglou S, Braybrook S,
38 Gray JE, Hobbs J (2017) Stomatal opening involves polar, not radial, stiffening of guard cells.
39 *Curr Biol* 27 (19):2974-2983. e2972. doi:10.1016/j.cub.2017.08.006
- 40 Chen D, Shao Q, Yin L, Younis A, Zheng B (2019) Polyamine function in plants: metabolism,
41 regulation on development, and roles in abiotic stress responses. *Front Plant Sci* 9:1945.
42 doi:10.3389/fpls.2018.01945
- 43 Del Duca S, Allué Creus J, D'Orazi D, Dondini L, Bregoli AM, Serafini-Fracassini D (2000) Tuber
44 vegetative stages and cell cycle in *Helianthus tuberosus*: protein pattern and their modification
45 by spermidine. *J Plant Physiol* 156:17-25. doi:10.1016/S0176-1617(00)80267-9
- 46 Del Duca S, Bonner P, Aloisi I, Serafini-Fracassini D, Cai G (2018) Determination of
47 transglutaminase activity in plants. In: *Polyamines*. Springer, pp 173-200. doi: 10.1007/978-
48 1-4939-7398-9_18

- 1 Del Duca S, Faleri C, Iorio RA, Cresti M, Serafini-Fracassini D, Cai G (2013) Distribution of
2 transglutaminase in pear pollen tubes in relation to cytoskeleton and membrane dynamics.
3 Plant Physiol 161 (4):1706-1721. doi:10.1104/pp.112.212225
- 4 Del Duca S, Serafini-Fracassini D (2005) Transglutaminases of higher, lower plants and fungi. Prog
5 Exp Tumor Res 38:223-247
- 6 Del Duca S, Serafini-Fracassini D, Bonner P, Cresti M, Cai G (2009) Effects of post-translational
7 modifications catalysed by pollen transglutaminase on the functional properties of
8 microtubules and actin filaments. Biochem J 418 (3):651-664. doi:10.1042/BJ20081781
- 9 Del Duca S, Serafini-Fracassini D, Cai G (2014) Senescence and programmed cell death in plants:
10 polyamine action mediated by transglutaminase. Front Plant Sci 5:120.
11 doi:10.3389/fpls.2014.00120
- 12 Della Mea M, Caparros-Ruiz D, Claparols I, Serafini-Fracassini D, Rigau J (2004a) AtPng1p. The
13 first plant transglutaminase. Plant Physiol 135 (4):2046-2054. doi:10.1104/pp.104.042549
- 14 Della Mea M, De Filippis F, Genovesi V, Serafini-Fracassini D, Del Duca S (2007) The acropetal
15 wave of developmental cell death of tobacco corolla is preceded by activation of
16 transglutaminase in different cell compartments. Plant Physiol 144 (2):1211-1222.
17 doi:10.1104/pp.106.092072
- 18 Della Mea M, Di Sandro A, Dondini L, Del Duca S, Vantini F, Bergamini C, Bassi R, Serafini-
19 Fracassini D (2004b) A Zea mays 39-kDa thylakoid transglutaminase catalyses the
20 modification by polyamines of light-harvesting complex II in a light-dependent way. Planta
21 219 (5):754-764. doi:10.1007/s00425-004-1278-6
- 22 Di Sandro A, Del Duca S, Verderio E, Hargreaves AJ, Scarpellini A, Cai G, Cresti M, Faleri C, Iorio
23 RA, Hirose S, Furutani Y, Coutts IG, Griffin M, Bonner PL, Serafini-Fracassini D (2010) An
24 extracellular transglutaminase is required for apple pollen tube growth. Biochem J 429
25 (2):261-271. doi:10.1042/BJ20100291
- 26 Dondini L, Bonazzi S, Del Duca S, Bregoli AM, Serafini-Fracassini D (2001) Acclimation of
27 chloroplast transglutaminase to high NaCl concentration in a polyamine-deficient variant
28 strain of *Dunaliella salina* and in its wild type. J Plant Physiol 158:185-197.
29 doi:10.1078/0176-1617-00099
- 30 Dondini L, Del Duca S, Dall'Agata L, Bassi R, Gastaldelli M, Della Mea M, Di Sandro A, Claparols
31 I, Serafini-Fracassini D (2003) Suborganellar localisation and effect of light on *Helianthus*
32 *tuberosus* chloroplast transglutaminases and their substrates. Planta 217 (1):84-95.
33 doi:10.1007/s00425-003-0998-3
- 34 Falcone P, Serafini-Fracassini D, Del Duca S (1993) Comparative studies of transglutaminase-like
35 activity and substrates in different organs of *Helianthus tuberosus*. J Plant Physiol 142:265-
36 273
- 37 Greenberg CS, Birckbichler PJ, Rice RH (1991) Transglutaminases: multifunctional cross-linking
38 enzymes that stabilize tissues. The FASEB Journal 5 (15):3071-3077.
39 doi:10.1096/fasebj.5.15.1683845
- 40 Griffin M, Casadio R, Bergamini CM (2002) Transglutaminases: nature's biological glues. Biochem
41 J 368 (Pt 2):377-396. doi:10.1042/BJ20021234
- 42 Handa AK, Fatima T, Mattoo AK (2018) Polyamines: bio-molecules with diverse functions in plant
43 and human health and disease. Front Chem 6:10. doi:10.1096/fasebj.5.15.1683845
- 44 Iacomino G, Picariello G, D'Agostino L (2012) DNA and nuclear aggregates of polyamines. Biochim
45 Biophys Acta 1823 (10):1745-1755. doi:10.1016/j.bbamcr.2012.05.033
- 46 Ioannidis NE, Lopera O, Santos M, Torne JM, Kotzabasis K (2012) Role of plastid transglutaminase
47 in LHCII polyamination and thylakoid electron and proton flow. PLoS One 7 (7):e41979.
48 doi:10.1371/journal.pone.0041979

- 1 Iranzo Ma, Aguado C, Pallotti C, Cañizares JV, Mormeneo S (2002) Transglutaminase activity is
2 involved in *Saccharomyces cerevisiae* wall construction. *Microbiology* 148 (5):1329-1334.
3 doi:10.1099/00221287-148-5-1329
- 4 Jahan MS, Shi ZR, Zhong M, Zhang YM, Zhou RR, El-Mogy MM, Sun J, Shu S, Guo SR, Wang Y
5 (2021) Comparative transcriptome analysis reveals gene network regulation of TGase-
6 induced thermotolerance in tomato. *Not Bot Horti Agrobo* 49 (1).
7 doi:10.15835/nbha49112208
- 8 Liu J-H, Wang W, Wu H, Gong X, Moriguchi T (2015) Polyamines function in stress tolerance: from
9 synthesis to regulation. *Front Plant Sci* 6:827. doi:10.3389/fpls.2015.00827
- 10 Lorand L, Graham RM (2003) Transglutaminases: crosslinking enzymes with pleiotropic functions.
11 *Nat Rev Mol Cell Biol* 4 (2):140-156. doi:10.1038/nrm1014
- 12 Mandrone M, Antognoni F, Aloisi I, Potente G, Poli F, Cai G, Faleri C, Parrotta L, Del Duca S (2019)
13 Compatible and incompatible pollen-styles interaction in *Pyrus communis* L. show different
14 transglutaminase features, polyamine pattern and metabolomics profiles. *Front Plant Sci*
15 10:741. doi:10.3389/fpls.2019.00741
- 16 Minocha R, Majumdar R, Minocha SC (2014) Polyamines and abiotic stress in plants: a complex
17 relationship. *Front Plant Sci* 5:175. doi:10.3389/fpls.2014.00175
- 18 Murashige T, Skoog F (1962) A revised medium for rapid growth and bio assays with tobacco tissue
19 cultures. *Physiol Plant* 15 (3):473-497. doi:10.1111/j.1399-3054.1962.tb08052.x
- 20 Pang X-M, Zhang Z-Y, Wen X-P, Ban Y, Moriguchi T (2007) Polyamines, all-purpose players in
21 response to environment stresses in plants. *Plant Stress* 1 (2):173-188
- 22 Paris R, Pagliarani G, Savazzini F, Aloisi I, Iorio RA, Tartarini S, Ricci G, Del Duca S (2017)
23 Comparative analysis of allergen genes and pro-inflammatory factors in pollen and fruit of
24 apple varieties. *Plant Sci* 264:57-68. doi:10.1016/j.plantsci.2017.08.006
- 25 Parrotta L, Aloisi I, Faleri C, Romi M, Del Duca S, Cai G (2020) Chronic heat stress affects the
26 photosynthetic apparatus of *Solanum lycopersicum* L. cv Micro-Tom. *Plant Physiology and*
27 *Biochemistry* 154:463-475. doi:10.1016/j.plaphy.2020.06.047
- 28 Porra R, Thompson W, Kriedemann P (1989) Determination of accurate extinction coefficients and
29 simultaneous equations for assaying chlorophylls a and b extracted with four different
30 solvents: verification of the concentration of chlorophyll standards by atomic absorption
31 spectroscopy. *Biochimica et Biophysica Acta (BBA)-Bioenergetics* 975 (3):384-394
- 32 Qin Y, Guo XW, Li L, Wang HW, Kim W (2013) The antioxidant property of chitosan green tea
33 polyphenols complex induces transglutaminase activation in wound healing. *J Med Food* 16
34 (6):487-498. doi: 10.1089/jmf.2012.2623
- 35 Sagor G, Liu T, Takahashi H, Niitsu M, Berberich T, Kusano T (2013) Longer uncommon
36 polyamines have a stronger defense gene-induction activity and a higher suppressing activity
37 of Cucumber mosaic virus multiplication compared to that of spermine in *Arabidopsis*
38 *thaliana*. *Plant cell reports* 32 (9):1477-1488. doi: 10.1007/s00299-013-1459-5
- 39 Serafini-Fracassini D, Del Duca S (2008) Transglutaminases: widespread cross-linking enzymes in
40 plants. *Ann Bot* 102 (2):145-152. doi:10.1093/aob/mcn075
- 41 Shu S, Tang Y, Zhou X, Jahan MS, Sun J, Wang Y, Guo S (2020) Physiological Mechanism of
42 Transglutaminase-mediated Improvement in Salt Tolerance of Cucumber Seedlings. *Plant*
43 *Physiology and Biochemistry*
- 44 Sobieszczuk-Nowicka E, Zmienko A, Samelak-Czajka A, Luczak M, Pietrowska-Borek M, Iorio R,
45 Del Duca S, Figlerowicz M, Legocka J (2015) Dark-induced senescence of barley leaves
46 involves activation of plastid transglutaminases. *Amino Acids* 47 (4):825-838.
47 doi:10.1007/s00726-014-1912-y
- 48 Suzuki T, Park H, Hollingsworth NM, Sternglanz R, Lennarz WJ (2000) PNG1, a yeast gene
49 encoding a highly conserved peptide: N-glycanase. *J Cell Biol* 149 (5):1039-1052.
50 doi:10.1083/jcb.149.5.1039

- 1 Telci D, Griffin M (2006) Tissue transglutaminase (TG2)--a wound response enzyme. *Front Biosci*
2 11:867-882. doi:10.2741/1843
- 3 Tiburcio AF, Altabella T, Bitrian M, Alcazar R (2014) The roles of polyamines during the lifespan
4 of plants: from development to stress. *Planta* 240 (1):1-18. doi:10.1007/s00425-014-2055-9
- 5 Tisi A, Federico R, Moreno S, Lucretti S, Moschou PN, Roubelakis-Angelakis KA, Angelini R, Cona
6 A (2011) Perturbation of Polyamine Catabolism Can Strongly Affect Root Development and
7 Xylem Differentiation. *Plant Physiol* 157 (1):200-215. doi:10.1104/pp.111.173153
- 8 Waffenschmidt S, Kusch T, Woessner JP (1999) A transglutaminase immunologically related to
9 tissue transglutaminase catalyzes cross-linking of cell wall proteins in *Chlamydomonas*
10 *reinhardtii*. *Plant Physiol* 121 (3):1003-1015. doi:10.1104/pp.121.3.1003
- 11 Zhao J, Morozova N, Williams L, Libs L, Avivi Y, Grafi G (2001) Two phases of chromatin
12 decondensation during dedifferentiation of plant cells distinction between competence for cell
13 fate switch and a commitment for S phase. *J Biol Chem* 276 (25):22772-22778.
14 doi:10.1074/jbc.M101756200
- 15 Zhong M, Wang Y, Zhang Y, Shu S, Sun J, Guo S (2019) Overexpression of transglutaminase from
16 cucumber in tobacco increases salt tolerance through regulation of photosynthesis. *Int J Mol*
17 *Sci* 20 (4):894. doi:10.3390/ijms20040894

18
19