

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Effect of plasma activated water (PAW) on rocket leaves decontamination and nutritional value

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Laurita, R., Gozzi, G., Tappi, S., Capelli, F., Bisag, A., Laghi, G., et al. (2021). Effect of plasma activated water (PAW) on rocket leaves decontamination and nutritional value. INNOVATIVE FOOD SCIENCE & EMERGING TECHNOLOGIES, 73(October 2021), 1-8 [10.1016/j.ifset.2021.102805].

Availability:

This version is available at: <https://hdl.handle.net/11585/865756> since: 2024-04-11

Published:

DOI: <http://doi.org/10.1016/j.ifset.2021.102805>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the peer-reviewed accepted manuscript of:

Laurita R.; Gozzi G.; Tappi S.; Capelli F.; Bisag A.; Laghi G.; Gherardi M.; Cellini B.;
Abouelenein D.; Vittori S.; Colombo V.; Rocculi P.; Dalla Rosa M.; Vannini L.

Effect of plasma activated water (PAW) on rocket leaves decontamination and
nutritional value

which has been published in final form in INNOVATIVE FOOD SCIENCE & EMERGING
TECHNOLOGIES Volume 73, October 2021, n. 102805

The final published version is available online at:

<http://dx.doi.org/10.1016/j.ifset.2021.102805>

© 2021 Elsevier. This manuscript version is made available under the Creative Commons
Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) 4.0 International License
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

1 **Effect of plasma activated water (PAW) on rocket leaves decontamination and** 2 **nutritional value**

3 Romolo Laurita ^{a,b*}, Giorgia Gozzi ^{c*}, Silvia Tappi ^{c,d**}, Filippo Capelli ^a, Alina Bisag ^a, Giulia
4 Laghi ^a, Matteo Gherardi ^{a,b}, Beatrice Cellini ^c, Doaa Abouelenein ^{e,f}, Sauro Vittori ^e, Vittorio
5 Colombo ^{a,b,d}, Pietro Rocculi ^{c,d}, Marco Dalla Rosa ^{c,d}, Lucia Vannini ^{c,d}

6 ^a *Department of Industrial Engineering (DIN), University of Bologna, Via Saragozza 8, Bologna, Italy*

7 ^b *Interdepartmental Center for Industrial Research for Advanced Mechanics and Materials, University of*
8 *Bologna, Via Saragozza 8 Bologna Italy*

9 ^c *Department of Agricultural and Food Sciences, University of Bologna, Piazza Goidanich, 60, Cesena FC,*
10 *Italy*

11 ^d *Interdepartmental Centre for Agri-Food Industrial Research, University of Bologna, via Q. Bucci 336,*
12 *Cesena, FC, Italy*

13 ^e *School of Pharmacy, University of Camerino, Italy*

14 ^f *Department of Pharmacognosy, Faculty of Pharmacy, Zagazig University, 44519 Zagazig, Egypt.*

15

16 * These two authors contributed in equal measure to this research

17 **Corresponding author. Mail: silvia.tappi2@unibo.it

18

19

20 **Abstract**

21 Plasma Activated Water (PAW) obtained by exposing water to cold atmospheric pressure plasma,
22 has recently emerged as a promising alternative for food decontamination, compared to the use of
23 traditional chemical sanitizers. The aim of the study was to evaluate the efficacy of PAW treatments
24 for rocket salad decontamination. Washing with PAW for 2, 5, 10 and 20 min was assessed against
25 different endogenous spoilage microorganisms and compared to **untreated water and** hypochlorite
26 solution. The chemical composition of PAW as a function of treatment and delay time was
27 characterized and the effect on product quality and nutritional parameters was evaluated.

28 Results showed that PAW allowed an average reduction of **1.7-3 Log** CFU/g for total mesophilic and
29 psychrotrophic bacteria and *Enterobacteriaceae* following 2-5 min washing with minimal variation
30 of qualitative and nutritional parameters. Overall, experimental results highlighted the potentiality of
31 PAW treatments as a promising alternative to chlorine having the advantage of a minor adverse
32 impact on environment and consumers' health.

33

34 **Industrial relevance**

To meet consumers demand, the minimally processed fruit and vegetable industry needs to find sustainable solutions as alternative to the use of traditional chemical sanitizers that allow to increase product shelf-life and preserve safety, qualitative and nutritional characteristics.

Plasma activated water represents a promising strategy for food decontamination, but its effects on foods have been only limitedly investigated. The present research is the first study on the use of plasma activated water on fresh rocket leaves, providing new and important information on microbial inactivation and quality of the fresh cut product.

Keywords: emerging technologies; plasma activated water; microbial decontamination; food quality; food spoilage

1. Introduction

The market of fresh and minimally processed vegetables has rapidly grown in the last decades as these products are very attractive to consumers looking for healthy and convenient meals. However, they are characterized by a very short shelf-life that limit their market opportunity and that leads to high amount of wasted product. The main limiting factor for these products shelf-life is surely microbial spoilage. Mesophilic aerobic bacteria, coliforms, *Pseudomonas* spp., *Bacillus* spp., lactic acid bacteria, as well as yeast and moulds are considered to be the main microorganisms found in fresh and ready-to-eat salads with loads up to 10^7 Log CFU/g. Decontamination of fresh products from pathogenic and spoilage microorganisms is usually performed through washing with chemical sanitizers, being sodium hypochlorite the most commonly used one. However, there are several concerns related to human health and environmental pollution that have led to the search for alternative methods. Moreover, wide variability in disinfection efficacy has been reported in numerous studies due to several factors, e.g. microbial protection by internalization within the vegetable tissue or biofilm formation and depletion of free-chlorine during washing (Yoon & Lee, 2018).

On one side, microorganisms are becoming more resistant to existent traditional technologies, and they are able to survive under various stress conditions, resulting in potential risk for human health (Zhao, Patange, Sun, & Tiwari, 2020). On the other side, the necessity to preserve the fresh-like characteristics and the content of bioactive compounds requires a mild technology in the decontamination process. Hence, the combination of effective sanitizing measures, while preserving quality and nutritional features, is the main issues for producers to improve the fresh products' shelf-life. Moreover, the environmental sustainability of the overall process has to be taken into account.

68 The use of cold atmospheric pressure plasma (CAP) represents a novel approach for sanitation of
69 food products that involves the application of an ionised gas mixture composed by charged particles,
70 electric fields, ultraviolet (UV) photons, and reactive species exploiting its strong oxidative power.
71 Previous research has demonstrated that CAP can efficiently inactivate a wide range of
72 microorganisms, including bacteria, fungi, viruses, bacterial spores and biofilms (Ikawa, Tani,
73 Nakashima, & Kitano, 2016; Kolb et al., 2008; Machala et al., 2013; Pan et al., 2013). Reactive
74 oxygen and nitrogen species (RONS) are the major bactericidal agents, which cause damage to DNA
75 and proteins in microbial cells (Joshi et al., 2011; Laurita, Barbieri, Gherardi, Colombo, & Lukes,
76 2015).

77 Plasma-activated water (PAW) is generated by exposing water to a plasma discharge above or
78 underneath water surface. The plasma-generated reactive species interact with water molecules and
79 trigger several chemical reactions, which produce a unique mixture of biochemical reactive
80 chemistries. In recent years, PAW has also been confirmed to possess outstanding biological activity
81 in biomedical and agricultural sectors (Kaushik et al., 2018; Sajib et al., 2020).

82 PAW can represent a promising alternative to washing in chlorinated water for fresh fruit and
83 vegetables, since it does not leave any residual chemicals in the products and it is considered more
84 environmentally friendly compared with some traditional chemical sanitizers (Zhao et al., 2020).

85 There are few studies (Baier et al., 2013; Giannoglou et al., 2020; Hertrich, Boyd, Sites, & Niemira,
86 2017; Silvetti et al., 2021) that report the efficacy of cold plasma on the microbial inactivation of
87 fresh cut salads, assessing endogenous microbiota or inoculated pathogens load reduction. However,
88 in all of these studies plasma in the gaseous form was employed, while, to our knowledge, no reports
89 are available on the use of PAW.

90 On the contrary, several authors have reported on the efficiency of PAW in inactivating natural
91 microbiota and pathogens deliberately contaminated on other types of fresh fruits and vegetables,
92 such as strawberries, button mushrooms, Chinese bayberries, and fresh-cut apples (Liu et al., 2020;
93 Ma et al., 2016; Ma et al., 2015; Xu, Tian, Ma, Liu, & Zhang, 2016), but no study on the effect of
94 PAW on background spoilage microbiota of rocket leaves is available in literature. Moreover, the
95 effect on microbial load has been investigated by many authors, while the effect on products quality
96 and nutritional value is often scarcely considered.

97 Fresh fruit and vegetables are rich in bioactive compounds that exert a pronounced effect on human
98 health, due to their antioxidant properties. Hence, the preservation of these compounds is fundamental
99 for product quality. A previous research showed that exposure to RONS produced by CAP treatment
100 reactive species could lead to the oxidation of some phenolic compounds in leafy vegetables such as
101 lams lettuce, but mainly on model systems, while the matrix protected these compounds

(Grzegorzewski, Ehlbeck, Schlüter, Kroh, & Rohn, 2011). Nevertheless, no differences in the antioxidant activity of fresh cut apples (Liu et al., 2020), pears (Chen et al., 2019) and button mushrooms (Xu et al., 2016) were observed after washing with PAW.

The aim of the present research was to investigate the decontamination of endogenous spoilage microbiota of rocket leaves induced by PAW washing and compare it to the effect of hypochlorite solution washing, used as traditional reference method. The effect on product quality was evaluated by measuring pH, color parameters and some bioactive compounds content.

2. Materials and Methods

2.1 Raw materials

Fresh rocket (*Rucola sativa*) leaves were purchased at the local market in Cesena, Italy and stored in a refrigerated cell at 2 ± 1 °C for maximum 24 h before processing.

2.2 Corona discharge for the production of PAW: electrical analysis and chemical characterization of RONS in PAW.

PAW was produced treating distilled water with a corona discharge, as reported in Laurita et al. (2021). Briefly, the plasma source consists of a stainless-steel pin-electrode connected to a **high voltage** microsecond pulsed generator (AlmaPulse, AlmaPlasma s.r.l.) at a distance of 5 mm from the liquid surface. A volume of 450 ml of distilled water, in a borosilicate Erlenmeyer flask on a stirrer (IKA Magnetic Stirrers RCT basic), was grounded. **The CAP is generated in the air gap between the tip of the high voltage electrode and the liquid surface.** The generator was operated at a peak voltage of 9 kV, with a fixed frequency of 5 kHz. The voltage (V) and the current (i) were measured using a high voltage probe (Tektronix P6015A) and a current probe (Pearson 6585) connected to a digital oscilloscope (Tektronix DPO4034, 350 MHz, 2.5 GSa s⁻¹). The average power (P) dissipated in the discharge was determined applying the following formula:

$$P = \frac{1}{T} \int_0^T i(t) V(t) dt, \quad (1)$$

where T is the applied voltage period.

The concentrations of H₂O₂, NO₂⁻ and dissolved O₃ produced in PAW by the CAP treatment were measured. The Amplex® Red Hydrogen Peroxide Assay Kit (Thermo Fisher Scientific, Waltham, MA, USA) and the Nitrate/Nitrite colorimetric assay (ROCHE, Baesel, Switzerland) were used for the measurement of H₂O₂ and NO₂⁻, respectively. The analysis of their concentrations was performed photometrically according to the manufacturer's protocols using a microplate reader (Rayto, P.R.

China). The O₃ concentration produced in the liquid phase was evaluated according to the manufacturer's protocols using Chematest 20 (SWAN Analytics). Moreover, the pH and the conductivity were evaluated by the means of inoLab® pH 7110 and Oakton Instrument: Con 6+ Meter, respectively. The RONS concentrations were assessed for four different treatment times (1, 2, 3 and 4 min) and four different delay times (2, 5, 10 and 20 min). All measurements were performed at least 3 times.

2.3 PAW treatment

PAW was produced exposing distilled water the aforementioned corona discharge for 4 min. Immediately after the treatment, rocket samples were immersed in PAW for 2, 5, 10 and 20 min in the ratio product:liquid of 1:20 (w:v) at room temperature. 20 g of rocket leaves were treated each time. During immersion, samples were constantly agitated in an orbital agitator.

After dipping, rocket leaves were removed from PAW and blotted with adsorbent paper to remove excess liquid. For each treatment time, two independent treatments were carried out.

Untreated rocket leaves were considered as control. Moreover, in order to evaluate the effect of PAW treatment, washing in untreated water was tested and a dipping in sodium hypochlorite (NaClO) 100 ppm for 2, 5, 10 and 20 min was used as common reference washing procedure.

For analysis of bioactive compounds, after each treatment time, rocket samples from both washing replicates were immediately freeze dried after treatment, the obtained samples were stored at -20°C until analysis.

2.4 Analytical determinations

2.4.1 Microbiological analysis

Immediately after each washing treatment, rocket samples were aseptically transferred to sterile saline solution (0.9 %, w/v, 1:10 w/v) and homogenized in a Stomacher 400 apparatus (Seward Ltd., Worthing, West Sussex, UK) for 2 min. Ten-fold serial dilutions of the homogenates were then performed and inoculated on suitable media for counting viable cells of various microbial groups. Total aerobic mesophilic and psychrotrophic microorganisms were evaluated onto Standard Plate Count Agar (Oxoid, Basingstoke, UK) plates incubated at 30 °C for 48 h and at 4 °C for 10 d, respectively. *Enterobacteriaceae* were counted on Violet Red Bile Glucose agar (Oxoid, Basingstoke, UK) incubated at 37 °C for 24 h. For each washing replicate (n=2), microbiological analyses were carried out at least in duplicate and the data were expressed as the mean of log colony forming units (CFU) per gram ±standard deviation.

170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203

2.4.2 Physico-chemical parameters

pH was measured in triplicate with a pH-meter (Crison, Barcellona, Spain) on the juice obtained by 10 g of treated or untreated sample. Water content was determined in triplicate by gravimetric method, drying samples in an oven at 70°C until constant weight was achieved. Results were expressed as g/100g.

Image analysis was used to determine samples color. Digital images were acquired placing the samples inside a black box and illuminated using four parallel lamps (mod. TL-D deluxe, natural daylight, 18W/965, Philips, NY, USA) with a colour temperature of 6500 K. The Colour Digital Camera (mod. D7000, Nikon, Shinjuku, Japan) used was equipped with a 60 mm lens (mod. AF-S micro, Nikkor) and located vertically over the sample at a fixed distance of 23 cm and at an angle with the lightning source of approximately 45°. The pre-processing of RGB images, segmentation and colour quantification were performed with ImageJ analysis software (NIH, USA). The average value of the segmented pixels was registered as L, R, G and B values. For each sample, images of 10 leaves were acquired and analysed.

2.4.3 Nutritional quality

Total Phenolic Content (TPC) was determined spectrophotometrically according to Mustafa et al. (2016) method with some modifications. Briefly, 0.5 mL of extracts solution (100 mg freeze dried sample extracted with 10 mL ethanol 70% with Formic acid (1.5%, v/v) was introduced into test tubes, then 2.5 mL of Folin-Ciocalteu reagent solution and 7 mL of Na₂CO₃ (7.5% w/w in water) solution were added. The reaction mixture allowed to stand at room temperature in the dark for 2 h and absorption was measured at 765 nm. The TPC was expressed as mg of gallic acid equivalents (GAE) per 100 g of dry weight of the rocket samples using gallic acid calibration curve.

Total Flavonoid Content (TFC) was determined following a method described by (Chen, Chen, Xiao, & Fu, 2018) with slight variations. Briefly, 0.5 mL of extract solution (10 mg/mL), 0.15 mL of NaNO₂ (0.5 M), 3.2 mL of methanol (30% v/v) and 0.15 mL of AlCl₃·6H₂O (0.3 M) were mixed. After 5 min, 1 mL of NaOH (1 M) was added. The solution was mixed well, and the absorbance was measured against the blank reagent at 506 nm. The standard calibration curve for TFC was made using quercetin standard solution (0 to 100 mg/L) under the same procedure as described above. TFC was expressed as mg of quercetin equivalents (QE) per 100g of dried extract.

Analytical determinations of TPC and TFC were carried out in triplicate for each freeze-dried sample.

2.5 Statistical analysis

To investigate statistical significance of observed differences in microbial, qualitative and nutritional data, a one-way analysis of variance (ANOVA) followed by the Tukey's HSD test (multiple range test) was used ($p < 0.05$). Analysis was conducted using Microsoft Excel 365 and Minitab ver. 19.0.

3. Results and Discussion

3.1 Electrical analysis and chemical characterization of RONS in PAW

Figure 1 shows current and voltage waveforms for the corona discharge operated at a peak voltage of 9 kV and a frequency of 5 kHz, referring to two applied voltage periods representative of the discharge electrical behavior during the treatment of the distilled water. Each applied voltage period (200 μ s) exhibited four voltage peaks, whose intensities reached maximum values up to 9.24 kV. The average discharge dissipated power, which can be calculated directly from the measured current and voltage reported in **Figure 1**, is 480.95 ± 33.42 W.

pH and conductivity of PAW as a function of treatment time are shown in **Figure 2**. As previously reported, the plasma treatment induced a reduction of pH (Laurita et al., 2015) up to 3.3 (dotted black line) after 4 min of plasma treatment, and an increase of conductivity (dotted purple line) up to 200 mS/cm.

Figures 3 and **4** report the behaviors of the concentration of the investigated RONS as a function of the treatment time and of the delay time. H_2O_2 , NO_2^- and O_3 concentrations increased with the increasing of the treatment times up to 4.5 ± 0.1 mg/L, 30.4 ± 0.9 mg/L and 0.3 ± 0.1 mg/L respectively. Focusing on the post-discharge kinetic of reactive species reported in **Figure 3b**, the decrease in concentration of both H_2O_2 and NO_2^- is related to the post-discharge processes in acid environment mediated by peroxynitrite (ONOOH) (Laurita et al., 2015; Lukes, Dolezalova, Sisrova, & Clupek, 2014). Finally, it is worth to note that the O_3 decay reported in **Figure 4b** is in accordance with the half-life values reported by Gardoni, Vailati, & Canziani (2012).

3.2 Effect of PAW on microbial load

The effect of PAW on the microbial loads of the target microbial groups of the rocket leaves was evaluated after washing for 2, 5, 10, and 20 min and compared to **untreated water and** hypochlorite solution (**Figure 5**).

Initial contamination levels of the untreated samples were about 5.74 ± 0.49 , 6.17 ± 0.70 and 3.67 ± 0.49 Log CFU/g for total mesophilic and psychrotrophic bacteria and *Enterobacteriaceae*,

238 respectively. These data are consistent with literature reporting rather high contamination levels for
 239 rocket leaves with counts for the aerobic mesophilic and psychrotrophic microorganisms up to 7.5
 240 Log CFU/g, yeasts and moulds about 5.8 Log CFU/g, and the range for *Enterobacteriaceae* 4.3–5.9
 241 Log CFU/g (Abadias, M., Usall, Anguera, Solsona, & Viñas, 2008).

242 The results presented in **Figure 5** clearly show that untreated water cannot be used to effectively
 243 decrease surface contamination of fresh rocket leaves since mean reductions for the target spoilage
 244 bacteria did not exceed 0.6 Log units regardless washing time. By contrast, all the considered
 245 microbial populations were significantly ($p < 0.05$) affected by PAW treatments depending on
 246 washing time. Psychrotrophic bacterial population was the most sensible one and it was reduced by
 247 2 Log CFU/g after a 2-min washing. By extending treatments up to 5 min, the decontaminating effect
 248 of PAW was enhanced resulting in a 3 Log CFU/g reduction. However, no further significant
 249 improvement was observed following the longest treatments. On the other hand, a 5-min washing
 250 was effective in decreasing counts of mesophilic bacteria and *Enterobacteriaceae* by 1.7-2.5 Log
 251 CFU/g. Similarly to the psychrotrophic population, PAW decontamination activity against
 252 mesophiles and enterobacteria did not significantly increase by prolonging the treatment time.

253 Overall, PAW had a stronger effect on the spoilage-related microorganisms with respect to the
 254 reference method (NaClO, 100 ppm) indicating that it can be an alternative to conventional washing
 255 with a chlorine-based sanitiser. More precisely, no significant influence on the presence of total
 256 mesophilic bacteria was detected over 20 min treatment with NaClO. Moreover, longer treatments
 257 were necessary with NaClO than PAW to have significant differences compared to untreated samples
 258 for enterobacteria and total psychrophilic population. In particular, this was observed only after 10
 259 min with maximum reductions of about 1.4-1.7 Log CFU/g. However, it was less effective than PAW
 260 since resulted in lower population decreases for both microbial groups which did not further improve
 261 by lengthening the treatment to 20 min.

262 It is not surprising that the chlorine treatments achieved rather limited reductions and its use for
 263 sanitizing fresh produce is controversial due to several issues which limit its effectiveness (Yoon &
 264 Lee, 2018). Besides the possibility for indigenous microbiota to internalize within tissues of fresh
 265 produce and to produce biofilm, depletion of free chlorine due to the presence of organic matters is
 266 well known to affect the disinfection capacity of the treatment, thus making processing time little
 267 relevant. On the other hand, the bactericidal action of PAW relies on the RONS content and its low
 268 pH. An important feature of PAW used in the present study is that only limited changes occurred in
 269 the content of nitrite anion by increasing washing time being its content about 25 ppm even after 20
 270 min (**Figure 3**).

Several studies showed that PAW can efficiently inactivate a wide variety of microorganisms from different fresh fruit and vegetables, but no study on rocket has been performed yet. It should however be considered that effectiveness of washing with PAW depends on several aspects related to the mode and conditions of PAW generation, which affect its chemico-physical properties, as well as on the different sizes, shapes, surface-to-volume ratio, and initial microbial contamination level of various fruits and vegetables. Despite differences in these factors, our data are in agreement with previous studies. Choi et al. (2019) reported reductions of 2.0, 2.2, 1.8, and 0.9 Log CFU/g in mesophilic aerobic bacteria, lactic acid bacteria, yeast and moulds, and coliforms, respectively of shredded salted Chinese cabbages. While such values are consistent with those obtained in this study, it is important to point out that the PAW they used was generated following a 120 min discharge resulting in NO_2^- and NO_3^- concentrations up to 88 and 394 ppm, respectively, while H_2O_2 was negligible. Xiang et al. (2019) showed significant reductions in populations of total aerobic bacteria and yeasts and moulds (2.32- and 2.84-Log CFU/g, respectively) of mung bean sprouts after washing for 30 min with PAW produced by 30 sec (APPJ; 5 kV, 40 kHz) and characterized for a content in H_2O_2 , NO_3^- and NO_2^- of about 17, 571 and 531 $\mu\text{mol/L}$, respectively. Reductions up to 1.7 Log CFU/g for mung bean seeds inoculated with *Escherichia coli* O154 were achieved by Machado-Moreira, Tiwari, Richards, Abram, & Burgess (2021) but only after 3h of treatment with PAW. A shorter washing time of 5 min, which is consistent with our study, was demonstrated to have a significant effect in decreasing initial counts of aerobic bacteria and mould and yeast of fresh-cut ‘Fuji’ apples (Liu et al., 2020). Not only PAW treatment reduced by approximately 1-2 Log CFU/g the indigenous spoilage microbiota, but also limited their proliferation over refrigerated storage while reducing superficial browning and maintaining quality features thus confirming its efficacy in prolonging the product shelf-life. Compared to Liu et al. (2020), our results indicate that PAW can be a suitable treatment to reduce bacterial population in rocket salad being active even after the shortest treatment. To successfully extend the produce shelf-life it would be however important to test also the fate of the surviving cells during storage to check whether bacteria were lethally or sub-lethally damaged by PAW and hence could recover or develop stress resistance responses.

3.3. Effect on quality and nutritional parameters

Water content and pH of rocket leaves did not show any significant variation during washing in PAW compared to the untreated sample (data not reported). Although the pH of PAW is significantly lower compared to rocket leaves one (6.17 ± 0.05), washing up to 20 min did not promote any change, in agreement with Xu et al. (2016).

Table 1 reports the colorimetric parameters measured by the image analysis. A significant decrease was observed for the parameters of L (luminosity), R (red components) and G (green parameter) in all samples, but variations do not appear to be proportional to treatment time. According to literature, the effect of plasma exposure on colour of fresh produce is generally quite minimal but depends on the specific matrix (Baier, Ehlbeck, Knorr, Herppich, & Schlüter, 2015). In a previous study, Ramazzina et al. (2015) observed an immediate reduction of chlorophyll content in kiwifruit subjected to gaseous plasma treatment that was attributed to the Type II chlorophylls degradation induced by reactive species. However, the author also observed a better retaining of colour during the following storage, that was probably due to degradative enzymes inactivation. Changes in colour of fresh products could be due to pigment degradation but also to changes in the tissue microstructure. However, to our knowledge there are no reports on the use of PAW on rocket leaves, so the observed effect on colour should be further clarified. Moreover, it is important to determine if these changes are perceptible by consumers and how they evolve during the storage of the product.

Consumption of bioactive compounds, especially polyphenols, plays a central role in the protection of human organs against inflammatory and neurodegenerative disorders and oxidative stress. However, the preservation of natural bioactive compounds in food products is a major challenge for the food industry. In this sense, cold plasma has been proposed as one of the most promising solutions to maximize the retention of polyphenol in food products.

Table 2 shows the time dependent changes in the total phenolic content (TPC) and total flavonoid content (TFC) of PAW treated rocket samples. According to the obtained results, all PAW treated samples showed non-significant differences in the TPC ($p > 0.05$) compared to the control samples, except for the 5 min treatment which induced a slight reduction. These results are in good accordance with the previous findings reported by Xiang et al. (2019) in which the PAW treatment did not induce any significant change in the total phenolic content of mung bean sprouts.

A significant variation in total flavonoid content was observed for PAW treated rocket samples with higher values for 10 and 20 min (3098.85 ± 53.20 and 3046.88 ± 38.58 mg gallic acid/ 100g, respectively) compared to the control samples (2524.88 ± 39.53 mg quercetin/ 100g).

The increase in the content of some specific phenolic compounds was observed after a short exposition to plasma in apples (Tappi et al., 2018) and blueberries (Sarangapani, O'Toole, Cullen, & Bourke, 2017) and after PAW treatment on mung bean sprouts (Fan, Liu, Ma, & Xiang, 2020). This effect was attributed to a physiological response of the tissue to the stress promoted by the reactive species. However, the increase of exposure time generally promoted oxidative reactions and the progressive decrease in the phenolic content and antioxidant activity. In the present research, TPC

339 was not affected even at the longer treatment time, while flavonoids accumulation occurred. A deeper
340 analysis of the single phenolic compounds could help to clarify the response of this specific matrix
341 to the exposure to plasma reactive species.

342

343 **4. Conclusions**

344

345 In summary, this study demonstrates PAW has the potential to decontaminate rocket salad with
346 limited changes in quality parameters. Compared to hypochlorite, which was chosen as a reference
347 sanitizer due to its widespread use in the food industry, PAW treatments resulted to be more effective
348 against the target background microbiota. In fact, shorter treatments were necessary to significantly
349 reduce *Enterobacteriaceae* and psychrotrophic bacterial populations, and greater inactivations were
350 achieved for all the spoilage microbial groups after 2 min.

351 However, the observed slight changes in color and bioactive compounds content should be further
352 clarified. Moreover, the behavior of the surviving microbiota and quality parameters should be
353 evaluated during storage of the packed product in order to determine the effect of PAW on the shelf-
354 life in real distribution and storage conditions.

355

356 **Acknowledgment**

357 The present work is part of the research activities developed within the project “PLASMAFOOD -
358 Study and optimization of cold atmospheric plasma treatment for food safety and quality
359 improvement” founded by MIUR - Ministero dell’Istruzione dell’Università e della Ricerca - PRIN:
360 Progetti di Ricerca di Rilevante Interesse Nazionale, Bando 2017.

361

362 **References**

- 363 Abadias, M., Usall, J., Anguera, M., Solsona, C., & Viñas, I. (2008). Microbiological quality of fresh,
364 minimally-processed fruit and vegetables, and sprouts from retail establishments. *International*
365 *Journal of Food Microbiology*, 123, 121–129.
- 366 Baier, M., Foerster, J., Schnabel, U., Knorr, D., Ehlbeck, J., Herppich, W. B., & Schlüter, O. (2013).
367 Direct non-thermal plasma treatment for the sanitation of fresh corn salad leaves: Evaluation of
368 physical and physiological effects and antimicrobial efficacy. *Postharvest Biology and*
369 *Technology*, 84, 81–87. <https://doi.org/10.1016/j.postharvbio.2013.03.022>
- 370 Baier, Matthias, Ehlbeck, J., Knorr, D., Herppich, W. B., & Schlüter, O. (2015). Impact of plasma
371 processed air (PPA) on quality parameters of fresh produce. *Postharvest Biology and*
372 *Technology*, 100, 120–126. <https://doi.org/10.1016/j.postharvbio.2014.09.015>

373 Chen, C., Liu, C., Jiang, A., Guan, Q., Sun, X., Liu, S., ... Hu, W. (2019). The effects of cold plasma-
 374 activated water treatment on the microbial growth and antioxidant properties of fresh-cut pears.
 375 *Food and Bioprocess Technology*, 12(11), 1842–1851.

376 Chen, G. L., Chen, S. G., Xiao, Y., & Fu, N. L. (2018). Antioxidant capacities and total phenolic
 377 contents of 30 flowers. *Industrial Crops and Products*, 111, 430–445.

378 Choi, E. J., Park, H. W., Kim, S. B., Ryu, S., Lim, J., Hong, E. J., ... Chun, H. H. (2019). Sequential
 379 application of plasma-activated water and mild heating improves microbiological quality of
 380 ready-to-use shredded salted kimchi cabbage (*Brassica pekinensis* L.)., *Food Control*, 98, 501-
 381 509.

382 Fan, L., Liu, X., Ma, Y., & Xiang, Q. (2020). Effects of plasma-activated water treatment on seed
 383 germination and growth of mung bean sprouts. *Journal of Taibah University for Science*, 14(1),
 384 823–830.

385 Gardoni, D., Vailati, A., & Canziani, R. (2012). Decay of ozone in water: a review. *Ozone: Science*
 386 *& Engineering*, 34(4), 233–242.

387 Giannoglou, M., Stergiou, P., Dimitrakellis, P., Gogolides, E., Stoforos, N. G., & Katsaros, G. (2020).
 388 Effect of Cold Atmospheric Plasma processing on quality and shelf-life of ready-to-eat rocket
 389 leafy salad. *Innovative Food Science and Emerging Technologies*, 66(August), 102502.
 390 <https://doi.org/10.1016/j.ifset.2020.102502>

391 Grzegorzewski, F., Ehlbeck, J., Schlüter, O., Kroh, L. W., & Rohn, S. (2011). Treating lamb's lettuce
 392 with a cold plasma - Influence of atmospheric pressure Ar plasma immanent species on the
 393 phenolic profile of *Valerianella locusta*. *LWT - Food Science and Technology*.
 394 <https://doi.org/10.1016/j.lwt.2011.05.004>

395 Hertrich, S. M., Boyd, G., Sites, J., & Niemira, B. A. (2017). Cold plasma inactivation of *Salmonella*
 396 in prepackaged, mixed salads is influenced by cross-contamination sequence. *Journal of Food*
 397 *Protection*, 80(12), 2132–2136.

398 Ikawa, S., Tani, A., Nakashima, Y., & Kitano, K. (2016). Physicochemical properties of bactericidal
 399 plasma-treated water. *Journal of Physics D: Applied Physics*, 49(42), 425401.

400 Joshi, S. G., Cooper, M., Yost, A., Paff, M., Ercan, U. K., Fridman, G., ... Brooks, A. D. (2011).
 401 Nonthermal dielectric-barrier discharge plasma-induced inactivation involves oxidative DNA
 402 damage and membrane lipid peroxidation in *Escherichia coli*. *Antimicrobial Agents and*
 403 *Chemotherapy*, 55(3), 1053–1062.

404 Kaushik, N. K., Ghimire, B., Li, Y., Adhikari, M., Veerana, M., Kaushik, N., ... Choi, E. H. (2018).
 405 Biological and medical applications of plasma-activated media, water and solutions. *Biological*
 406 *Chemistry*, 400(1), 39–62.

- 407 Kolb, J. F., Mohamed, A. A. H., Price, R. O., Swanson, R. J., Bowman, A., Chiavarini, R. L., ...
 408 Schoenbach, K. H. (2008). Cold atmospheric pressure air plasma jet for medical applications.
 409 *Applied Physics Letters*, 92(24), 241501.
- 410 Laurita, R., Barbieri, D., Gherardi, M., Colombo, V., & Lukes, P. (2015). Chemical analysis of
 411 reactive species and antimicrobial activity of water treated by nanosecond pulsed DBD air
 412 plasma. *Clinical Plasma Medicine*, 3(2), 53.
- 413 Laurita, R., Contaldo, N., Zambon, Y., Bisag, A., Canel, A., Gherardi, M., & Colombo, V. (2021).
 414 Laurita, R., Contaldo, N., Zambon, Y., Bisag, A., Canel, A., Gherardi, M., ... & Colombo, V.
 415 (2021). The use of plasma-activated water in viticulture: Induction of resistance and agronomic
 416 performance in greenhouse and open field. *Plasma Processes and Polymers*, 18(1), 2000206.
- 417 Liu, C., Chen, C., Jiang, A., Sun, X., Guan, Q., & Hu, W. (2020). Effects of plasma-activated water
 418 on microbial growth and storage quality of fresh-cut apple. *Innovative Food Science and*
 419 *Emerging Technologies*, 59, 102256.
- 420 Lukes, P., Dolezalova, E., Sisrova, I., & Clupek, M. (2014). Aqueous-phase chemistry and
 421 bactericidal effects from an air discharge plasma in contact with water: evidence for the
 422 formation of peroxynitrite through a pseudo-second-order post-discharge reaction of H₂O₂ and
 423 HNO₂. *Plasma Sources Science and Technology*, 23(1), 015019.
- 424 Ma, R., Yu, S., Tian, Y., Wang, K., Sun, C., Li, X., ... Fang, J. (2016). Effect of non-thermal plasma-
 425 activated water on fruit decay and quality in postharvest Chinese bayberries. *Food and*
 426 *Bioprocess Technology*, 9(11), 1825–1834.
- 427 Ma, Ruonan, Wang, G., Tian, Y., Wang, K., Zhang, J., & Fang, J. (2015). Non-thermal plasma-
 428 activated water inactivation of food-borne pathogen on fresh produce. *Journal of Hazardous*
 429 *Materials*, 300. <https://doi.org/10.1016/j.jhazmat.2015.07.061>
- 430 Machado-Moreira, B., Tiwari, B. K., Richards, K. G., Abram, F., & Burgess, C. M. (2021).
 431 Application of plasma activated water for decontamination of alfalfa and mung bean seeds. *Food*
 432 *Microbiology*, 96, 103708.
- 433 Machala, Z., Tarabova, B., Hensel, K., Spetlikova, E., Sikurova, L., & Lukes, P. (2013). Formation
 434 of ROS and RNS in Water Electro-Sprayed through Transient Spark Discharge in Air and their
 435 Bactericidal Effects. *Plasma Processes and Polymers*, 10(7), 649–659.
- 436 Mustafa, A. M., Maggi, F., Öztürk, N., Öztürk, Y., Sagratini, G., Torregiani, E., ... Caprioli, G.
 437 (2016). Chemical and biological analysis of the by-product obtained by processing *Gentiana*
 438 *lutea* L. and other herbs during production of bitter liqueurs. *Industrial Crops and Products*, 80,
 439 131–140.
- 440 Pan, J., Sun, K., Liang, Y., Sun, P., Yang, X., Wang, J., ... Becker, K. H. (2013). Cold plasma therapy

of a tooth root canal infected with *Enterococcus faecalis* biofilms in vitro. *Journal of Endodontics*, 39(1), 105–110.

Ramazzina, I., Berardinelli, A., Rizzi, F., Tappi, S., Ragni, L., Sacchetti, G., & Rocculi, P. (2015). Effect of cold plasma treatment on physico-chemical parameters and antioxidant activity of minimally processed kiwifruit. *Postharvest Biology and Technology*, 107. <https://doi.org/10.1016/j.postharvbio.2015.04.008>

Sajib, S. A., Billah, M., Mahmud, S., Miah, M., Hossain, F., Omar, F. B., ... Reza, M. A. (2020). Plasma activated water: The next generation eco-friendly stimulant for enhancing plant seed germination, vigor and increased enzyme activity, a study on black gram (*Vigna mungo* L.). *Plasma Chemistry and Plasma Processing*, 40(1), 119–143.

Sarangapani, C., O'Toole, G., Cullen, P. J., & Bourke, P. (2017). Atmospheric cold plasma dissipation efficiency of agrochemicals on blueberries. *Innovative Food Science and Emerging Technologies*, 44. <https://doi.org/10.1016/j.ifset.2017.02.012>

Silvetti, T., Pedroni, M., Brasca, M., Vassallo, E., Cocetta, G., Ferrante, A., ... Morandi, S. (2021). Assessment of possible application of an atmospheric pressure plasma jet for shelf life extension of fresh-cut salad. *Foods*, 10(3), 1–13. <https://doi.org/10.3390/foods10030513>

Tappi, S., Ramazzina, I., Rizzi, F., Sacchetti, G., Ragni, L., & Rocculi, P. (2018). Effect of plasma exposure time on the polyphenolic profile and antioxidant activity of fresh-cut apples. *Applied Sciences (Switzerland)*, 8(10). <https://doi.org/10.3390/app8101939>

Xiang, Q., Liu, X., Liu, S., Ma, Y., Xu, C., & Bai, Y. (2019). Effect of plasma-activated water on microbial quality and physicochemical characteristics of mung bean sprouts. *Innovative Food Science and Emerging Technologies*, 52, 49–56.

Xu, Y., Tian, Y., Ma, R., Liu, Q., & Zhang, J. (2016). Effect of plasma activated water on the postharvest quality of button mushrooms, *Agaricus bisporus*. *Food Chemistry*, 197, 436–444.

Yoon, J.-H., & Lee, S.-H. (2018). Comparison of the effectiveness of decontaminating strategies for fresh fruits and vegetables and related limitations. *Critical Reviews in Food Science and Nutrition*, 58, 3189–3208.

Zhao, Y. M., Patange, A., Sun, D. W., & Tiwari, B. (2020). Plasma-activated water: Physicochemical properties, microbial inactivation mechanisms, factors influencing antimicrobial effectiveness, and applications in the food industry. *Comprehensive Reviews in Food Science and Food Safety*, 19(6), 3951–3979.

473 **Figure captions**

474

475 **Figure 1.** Voltage and current waveforms during distilled water treatment

476 **Figure 2.** pH (black) and conductivity (purple) values of not treated solution (NT) and 1, 2, 3 and 4
477 min of plasma treatment.

478 **Figure 3.** (a) Chemical analysis of the concentrations of H_2O_2 and NO_2^- after 1, 2, 3 and 4 min of
479 plasma treatment and (b) their decay kinetic at room temperature.

480 **Figure 4.** (a) Chemical analysis of the concentrations of O_3 after 1, 2, 3 and 4 min of plasma treatment
481 and (b) its decay kinetic at room temperature.

482 **Figure 5.** Mean counts of (a) total aerobic mesophilic and (b) psychrotrophic bacteria, and (c)
483 *Enterobacteriaceae* on rocket leaves collected before and after the washing with PAW, **untreated**
484 **water** or **sodium** hypochlorite (**NaClO , 100 ppm**) at different times. Mean values with different letters
485 significantly differ among treatment times and washing solutions ($p < 0.05$).

486

487