

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Morpho-functional analyses reveal that changes in the chemical structure of a marine bisindole alkaloid alter the cytotoxic effect of its derivatives

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Sabrina Burattini, M.B. (2022). Morpho-functional analyses reveal that changes in the chemical structure of a marine bisindole alkaloid alter the cytotoxic effect of its derivatives. MICROSCOPY RESEARCH AND TECHNIQUE, 86(7), 1-9 [10.1002/jemt.24092].

Availability:

This version is available at: <https://hdl.handle.net/11585/877804> since: 2022-03-08

Published:

DOI: <http://doi.org/10.1002/jemt.24092>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Changes in the chemical structure of the marine bisindole alkaloid 2,2-bis(6-bromo-1*H*-indol-3-yl)ethanamine alter the cytotoxic effect of its derivatives

Sabrina Burattini^{1a}, Michela Battistelli^{1a}, Michele Verboni¹, Elisabetta Falcieri¹, Simone Lucarini^{1*} and Sara Salucci^{1,2*}

¹Department of Biomolecular Sciences (DISB), University of Urbino Carlo Bo, 61029 Urbino, Italy

²Cellular Signalling Laboratory, Department of Biomedical and Neuro Motor Sciences (DIBINEM), University of Bologna, 40126 Bologna, Italy

^aequally contributed

Corresponding authors:

Dr. Sara Salucci, Department of Biomolecular Sciences, University of Urbino, via Ca' Le Suore 2, 61029 Urbino and Cellular Signalling Laboratory, Department of Biomedical and Neuromotor Sciences, University of Bologna, Bologna, Italy. Email: sara.salucci@uniurb.it; sara.salucci@unibo.it

Dr. Simone Lucarini - Department of Biomolecular Sciences, University of Urbino Carlo Bo, piazza del Rinascimento 6, 61029 Urbino (PU), Italy; orcid.org/0000-0002-3667-1207; Phone: +390722303333; Email: simone.lucarini@uniurb.it

Abstract

2,2-bis(6-bromo-1*H*-indol-3-yl)ethanamine, a marine alkaloid, appeared a potential anticancer agent against several tumor cell models thanks to the presence of a 3,3'-diindolylmethane scaffold. Here, the modifications in its chemical structure into alkaloid-like derivatives, have been evaluated, to investigate changes in its biological activities. Three derivatives have been considered and their potential apoptotic action has been evaluated through morpho-functional analyses in a human cancer cell line.

Apoptosis appears strongly decreased in the derivative compounds without the bromine units (**1**) and in those where the bromine groups have been substituted with fluorine atoms (**2**). On the contrary, the methylation of indole NH (**3**) does not alter the alkaloid apoptotic activity that occurs through the involvement of an increased oxidative stress, leading to peroxidation events and mitochondria dysfunctionality.

This manuscript highlights the alkaloid derivative cytotoxic effect, which is strictly correlated to the maintenance of bisindole structure and bromine atoms. Since molecular therapies, by targeting mitochondria pathways, have shown positive outcomes against several cancer cells, the alkaloid with bisindole scaffold and the two bromine units can be considered a promising candidate to develop new derivatives with strong anticancer property.

Key words: Marine alkaloids; bisindoles; structure-activity relationships, apoptotic activity; mitochondria, anticancer activity, U937 cell line

Introduction

Marine bisindole alkaloids showed significant cytotoxicity, antineoplastic and antimicrobial properties [1,2]. In particular, 2,2-bis(6-bromo-3-indolyl) ethanamine (**BrDIMEA**), isolated from the Californian tunicate *Didemnum candidum* and the New Caledonian sponge *Orina* [3], is a strong cytotoxic agent in various tumor cell lines [3-5]. In our previous paper, its apoptotic mechanism of action, through caspase activation and regulated by Bcl-2 protein family, has been demonstrated in U937 cells. For this potential anticancer activity, **BrDIMEA**, a member of the large 3,3'-diindolylmethane family of alkaloids, has obtained much attention [3].




Figure 1. The marine alkaloid **BrDIMEA** and some natural antitumoral agents sharing the 3,3'-diindolylmethane (**DIM**) scaffold.

Its anticancer properties are possibly due to the common 3,3-diindolylmethane molecular unit (**DIM**), a chemical group (Figure 1) which exhibit itself antiproliferative or apoptotic activities [6-9]. The use of DIM is limited for lipophilic nature and *in vivo* chemical instability and it seems that the presence of the alkylamino side chain improves its chemical and therapeutic activities [4]. In addition, the presence of two bromoindole units could enhance the anti-cancer action. Therefore,

BrDIMEA can be considered a lead molecule for the synthesis of new compounds with a wide range of biological properties, including the antineoplastic action.


In this study, the potential apoptotic effect of three different **BrDIMEA** derivatives (**1-3**), has been evaluated through morpho-functional analyses on U937 human cancer cells by, comparing their action with that of the natural compound.

Materials and Methods

Chemicals

All organic solvents used in this study were purchased from Sigma–Aldrich (St. Louis, MO, USA), Alfa Aesar (Haverhill, MA, USA), or TCI (Tokyo, Japan). Prior to use, acetonitrile, dichloromethane and toluene were dried with molecular sieves with an effective pore diameter of 4 Å. Column chromatography purifications were performed under “flash” conditions using Merck (Darmstadt, Germany) 230–400 mesh silica gel. Analytical thin-layer chromatography (TLC) was carried out on Merck silica gel plates (silica gel 60 F254), which were visualized by exposure to ultraviolet light and an aqueous solution of cerium ammonium molybdate (CAM). ESI-MS spectra were recorded with a Waters (Milford, MA, USA) Micromass ZQ spectrometer. ¹H NMR and ¹³C NMR spectra were recorded on a Bruker (Billerica, MA, USA) AC 400 or 100, respectively, spectrometer and analyzed using the TopSpin 1.3 (2013) software package. Chemical shifts were measured by using the central peak of the solvent. The final compounds were analyzed on a ThermoQuest (Italia) FlashEA 1112 elemental analyzer for C, H, and N. The percentages found were within ±0.4% of the theoretical values. All the tested compounds were >95% pure as determined by elemental analysis.

88 Marine bisindole alkaloid **BrDIMEA** and compounds **1-3** were prepared as described in Scheme
89 1.



92 **Scheme 1.** Reaction conditions: (a) diphenyl phosphate, acetonitrile, 80°C, 24 h; (b) K₂CO₃,
93 methanol, reflux, 2 h.

95 *General procedure for the synthesis of derivatives **BrDIMEA** and **1-3***

96 Diphenyl phosphate (0.02 mmol) was added to a solution of the appropriate indole derivative (0.4
97 mmol) and (trifluoroacetyl amino)acetaldehyde dimethyl acetal (0.2 mmol) in anhydrous
98 acetonitrile (0.2 mL), and the resulting mixture was stirred at 80 °C for 24 h in a sealed tube,
99 monitoring the progress of the reaction by TLC and HPLC-MS. After cooling to room temperature,
100 saturated aqueous NaHCO₃ (30 mL) and dichloromethane (30 mL) were added and the two phases
101 were separated. The aqueous solution was extracted with dichloromethane (3 x 20 mL). After
102 drying over dry Na₂SO₄, the combined organic phases were concentrated in vacuum and the
103 resulting crude product was utilized without further purification. A mixture of that crude
104 trifluoroacetamide derivative and potassium carbonate (1 mmol) in MeOH (1.87 mL) and H₂O
105 (0.13 mL) was stirred and heated at reflux for 2 h. The MeOH was removed under reduced pressure
106 and water was added (30 mL). The aqueous solution was extracted with dichloromethane (3 x 30

mL) and the resulting solution was dried with Na₂SO₄ and concentrated in vacuum. The crude material was purified by flash chromatography on neutral alumina.

The physico-chemical data of **BrDIMEA** [4,10] and **1-3** are in agreement with those reported.

Cell Culture

U937 human myelomonocytic lymphoma cell line, grown in RPMI 1640, supplemented with 10% heat-inactivated fetal bovine serum, 2 mM glutamine, 1% antibiotics, was maintained at 37°C in humidified air with 5% CO₂ [11]. Cell behavior was monitored with the Inverted Microscopy (IM; Eclipse TE2000-S Nikon; objective 10x).

Apoptosis induction

U937 cells have been exposed to **BrDIMEA** and its derivatives **1-3**, at the final concentration in the cell medium of 7.5 µM. This latter is the better apoptotic dose of **BrDIMEA**, already demonstrated by Salucci et al., 2018 [3].

For apoptosis induction, cells (seeded at 1x10⁶ cells/mL) were exposed to three derivatives and to

BrDIMEA.

Trypan Blue (TB) exclusion assay [12] has been used to evaluate living and dead cells in control condition and in treated samples.

Transmission Electron Microscopy (TEM)

U937 pellets were immediately fixed in 2.5% glutaraldehyde in 0.1 M in phosphate buffer, post-fixed in 1% OsO₄ in the same buffer, dehydrated with ethanol and embedded in araldite [13].

Semithin sections have been counterstained with Blue of Toluidine and observed at light microscope (LM). Thin sections, collected on nickel grids and stained with uranyl acetate and lead

citrate, were observed with an electron microscope (Philips CM10, 80KV, FEI Company, Italy).

Confocal Laser Scanning Microscopy (CLSM)

TUNEL

Control and treated cells were fixed with 4% paraformaldehyde in phosphate buffer saline (PBS) pH 7.4 for 30 min., then deposited on poly-lysinated coverslips in Petri dishes overnight at 4 °C. After PBS washing, samples were permeabilized with a 2:1 mixture of ethanol and acetic acid for 5 min at –20 °C. For the TUNEL technique, all reagents were part of the Apoptag Plus kit (D.B.A., Oncor, Dallas, TX, USA) and procedures were carried out according to the manufacturer's instructions and as described in Salucci et al., 2014 [14]. Finally, slides were mounted with an antifading medium. Specimens were observed with a Leica TCS-SP5 confocal laser scanning microscope (CLSM) connected to a DMI 6000 CS inverted microscope (Leica Microsystems CMS GmbH, Mannheim, Germany); excitation was at 488 nm and emission signals were detected at 517 nm.

Mitochondrial behavior

To monitor mitochondrial behavior, fresh cells were treated with 10 nonyl-acridine orange (NAO) for evaluating mitochondria membrane integrity through peroxidation event evaluation. Fresh cells were exposed to 50 nM NAO for 10 min at room temperature and then observed through a Leica TCS-SP5 Confocal connected to a DMI 6000 CS Inverted Microscope (Leica Microsystems CMS GmbH; objectives 40x and 60x).

Excitation was at 488nm (FITC and NAO); emission signals were detected at 525 nm (FITC) and 519 nm (NAO). CLSM images are presented as single-plane images or Z-stack projections [15].

Results

In the previous paper, the best **BrDIMEA** apoptotic action has been demonstrated in U937 cells at the dose of 7.5 μ M.

In this work the effect of its derivatives, which show some modifications in their chemical structures respect to the original alkaloid, has been analyzed. First of all, the ability of these compounds to induce cytotoxicity in the same cell line has been evaluated through TB assay and the number of living cells, expressed as mean value percentage \pm standard deviation, has been calculated. Figure 2 shows micrographs obtained after IM and LM observations which revealed living and dead cells in all experimental conditions together with a graph on cell viability percentage. A good cell preservation confirmed by a high living cell number appears in control condition (Fig. 2A, B). Cells treated with compound **1** or **2** (in which bromide atoms are deleted and in the second compound substituted with the fluorine ones), show, at IM (Fig. 2C, E) or LM (Fig. 2D, F), a morphological behavior similar to that observed in control condition. In both situations, a reduction of cell viability about 10% has been quantified. On the other hand, the compound **3**, which maintains the DID molecular unit and the bromide groups and in which two hydrogen atoms from indole have been deleted, induces a diffuse presence of dead cells (Fig. 2G, H). Therefore, the compound **3** causes a cytotoxic effect comparable to that observed after **BrDIMEA** treatment in our previous work [3].

Morphological observations allow to distinguish the type of death and have been carried out for the compound **3** alone. Compared to control cells, which show a preserved ultrastructure (Fig. 3A, B), those exposed to 7.5 μ M compound **3**, became apoptotic and, in particular, late apoptotic events can be observed (Fig. 3C, D). Therefore, an evident cell shrinkage, cytoplasmic vacuolization, mitochondria damage, chromatin condensation, micronuclei and, sometimes, secondary necrosis appear. Since compound **3** induces an apoptotic phenotype in U937 cells, the *in situ* DNA

fragmentation has been investigated, and the presence of a diffuse number of TUNEL positive nuclei (Fig. 3 inset D), absent in control condition (Fig. 3 inset A) has been observed. In particular, the fluorescent staining is localized inside micronuclei (Fig. 3 inset D).

For what concerns derivative **3**, the mitochondrial behavior has been also investigated. In particular, the presence of cardiolipin peroxidation events have been analyzed with the fluorescent probe NAO at CLSM. Respect to control cells, in which mitochondrial membrane maintains its integrity (Fig. 3E, F), those treated with compound **3** show a decline of fluorescence intensity (Fig. 3G, H) suggesting that the loss of mitochondrial membrane integrity could favor the release of pro-apoptotic proteins from the mitochondria. This latter result confirms the ultrastructural observations, in fact cells treated with compound **3** showed dysfunctional mitochondria which appear empty or with altered cristae, a typical condition induced by an increased oxidative stress.

Discussion

Apoptosis is a programmed cell death which maintains the healthy survival/death balance in cells [16,17]. Misregulation of apoptosis can cause cancer or autoimmunity, while enhanced apoptosis may cause degenerative diseases [11,18]. Researchers are involved in the development of some promising cancer treatment strategies which target apoptotic inhibitors including Bcl-2 family proteins and other substrates. One strong apoptotic agent in various cell models appears a marine bisindole alkaloid [19,20], which induces apoptosis in U937 cells following both extrinsic and intrinsic pathways and by involving the Bcl-2 protein family. In particular, Salucci and collaborators demonstrated the ability of this chemical alkaloid to up-regulate bax and to inactivate Bcl-2 and Bcl-xL [3], the anti-apoptotic factors, which block the release of cytochrome c from mitochondria and thus promote cell survival. Since the 2,2-bis(6-bromo-3-indolyl)-ethanamine targets anti-apoptotic Bcl-2 protein and, consequently, can have success in killing many types of

cancer, its chemical structure can be considered the basic scaffold to obtain synthetic derivatives with similar biological properties but with an improved chemical stability [21] .

For that, in this study, we have evaluated the anti-cancer potential action of three bisindole alkaloid derivatives. To identify and confirm the chemical unit responsible of apoptotic effect, a preliminary structure-activity relationships of 2,2-bis(6-bromo-3-indolyl)-ethanamine have been carried out.

Morpho-functional analyses revealed that, in the same cell model and at the same dosage, only the compound **3** maintains an apoptotic effect similar to that observed after **BrDIMEA**. Moreover, compound **3** is able to induce apoptotic cell death through the oligonucleosomic DNA cleavage and by involving the mitochondrial pathway. In fact, mitochondrial membrane integrity appears loss due to cardiolipin peroxidation, which leads to the formation of reactive aldehydes, able to react with proteins and DNA [22]. Ultrastructural analyses reveal the presence of a high number of empty mitochondria suggestive of oxidative stress increase. This data enhances that oxidative stress production can be considered one of the major causes of apoptosis in cancer cells and it could be recognized into a promising therapeutic approach for the cancer treatment [23].

These findings show that the **BrDIMEA** derivative **3** induces apoptosis via mitochondrial pathway and with a rate similar to that observed for the marine alkaloid lead compound, demonstrating that the bisindole scaffold, even N-methylated, as well as the bromine atoms, are necessary and must be maintained for developing new anticancer drug.

Acknowledgments: This work has been possible thanks to the DISB 2017 Enhancement Project of Urbino University.

Author Contributions:

227 Methodology, S.B.,M.B., S.L.,S.S; Investigation, S.S.; Data Curation, S.S. and S.L.; Writing –
228 Original Draft Preparation, S.S and S.L.; Writing – Review & Editing, S.S and S.L.; Supervision,
229 S.L. and S.S.

References

1. Veale, C.G.; Davies-Coleman, M.T. Chapter one: Marine Bi-, Bis-, and Trisindole Alkaloids. *In: Alkaloids Chem Biol, Elsevier* **2014**, 73, 1- 64.
2. Kamel, M.M.; Abdel-Hameid, M.K.; El-Nassan, H.B.; El-Khouly, E.A. Synthesis and Cytotoxic Activity of Novel Mono- and Bis-Indole Derivatives: Analogues of Marine Alkaloid Nortopsentin. *Med Chem.* **2020**, 17, 779-789.
3. Salucci, S.; Burattini, S.; Buontempo, F.; Orsini, E.; Furiassi, L.; Mari, M.; Lucarini, S.; Martelli, A.M.; Falcieri, E. Marine bisindole alkaloid: A potential apoptotic inducer in human cancer cells. *Eur J Histochem.* **2018**, 62, 2881.
4. Mari, M.; Tassoni, A.; Lucarini, S.; Fanelli, M.; Piersanti, G.; Spadoni, G. Brønsted Acid Catalyzed Bisindolization of α -Amido Acetals: Synthesis and Anticancer Activity of Bis(indolyl)ethanamino Derivatives. *Eur J Org Chem.* **2014**, 2014, 3822-3830.
5. Mantenuto, S.; Lucarini, S.; De Santi, M.; Piersanti, G.; Brandi, G.; Favi, G. One-Pot Synthesis of Biheterocycles Based on Indole and Azole Scaffolds Using Tryptamines and 1,2-Diaza-1,3-dienes as Building Blocks. *Eur J Org Chem.* **2016**, 2016, 3193-3199.
6. Junaid, M.; Dash, R.; Islam, N.; Chowdhury, J.; Alam, M.J.; Nath, S.D.; Shakil, M.A.S.; Azam, A.; Quader, S.M.; Zahid Hosen, S.M. Molecular Simulation Studies of 3,3'-Diindolylmethane as a Potent MicroRNA-21 Antagonist. *J Pharm Bioallied Sci.* **2017**, 9, 259-265.
7. Tian, X.; Liu, K.; Zu, X.; Ma, F.; Li, Z.; Lee, M.; Chen, H.; Li, Y.; Zhao, Y.; Liu, F.; Oi, N.; Bode, A.M.; Dong, Z.; Kim, D.J. 3,3'-Diindolylmethane inhibits patient-derived xenograft colon tumor growth by targeting COX1/2 and ERK1/2. *Cancer Lett.* **2019**, 448, 20-30.
8. Lanza-Jacoby, S.; Cheng, G. 3,3'-Diindolylmethane enhances apoptosis in docetaxel-treated breast cancer cells by generation of reactive oxygen species. *Pharm Biol.* **2018**, 56, 407-414.
9. Condello, M.; Pellegrini, E.; Multari, G.; Gallo, F.R.; Meschini, S. Voacamine: Alkaloid with

its essential dimeric units to reverse tumor multidrug resistance. *Toxicol In Vitro*. **2020**, *65*, 104819.

10. Campana, R.; Favi, G.; Baffone, W.; Lucarini, S. Marine Alkaloid 2,2-Bis(6-bromo-3-indolyl) Ethylamine and Its Synthetic Derivatives Inhibit Microbial Biofilms Formation and Disaggregate Developed Biofilms. *Microorganisms* 2019, *7*, 28.

11. Salucci, S.; Battistelli, M.; Burattini, S.; Sbrana, F.; Falcieri E. Holotomographic microscopy: A new approach to detect apoptotic cell features. *Microsc Res Tech*. **2020**, *83*, 1464-1470.

12. Salucci, S.; Burattini, S.; Giordano, F.M.; Diamantini, G.; Falcieri, E. Further highlighting on the prevention of oxidative damage by polyphenol-rich wine extracts. *J Med Food*, **2017**, *20*, 410-419.

13. Giordano, F.M.; Burattini, S.; Buontempo, F.; Canonico, B.; Martelli, A.M.; Papa, S.; Sampaolesi, M.; Falcieri, E.; Salucci, S. Diet Modulation Restores Autophagic Flux in Damaged Skeletal Muscle Cells. *J Nutr Health Aging*. **2019**, *23*, 739-745.

14. Salucci, S.; Burattini, S.; Battistelli, M.; Baldassarri, V.; Curzi, D.; Valmori, A.; Falcieri, E. Melatonin prevents chemical-induced haemopoietic cell death. *Int J Mol Sci*. **2014a**, *15*, 6625-6640.

15. Salucci, S.; Battistelli, M.; Baldassarri, V.; Burini, D.; Falcieri, E.; Burattini, S. Melatonin prevents mitochondrial dysfunctions and death in differentiated skeletal muscle cells. *Microsc Res Tech*. **2017**, *80*, 1174-1181.

16. Salucci, S.; Burattini, S.; Curzi, D.; Buontempo, F.; Martelli, A.M.; Zappia, G.; Falcieri, E.; Battistelli, M. Antioxidants in the prevention of UVB-induced keratinocyte apoptosis. *J Photochem Photobiol B*. **2014b**, *141*, 1-9.

17. Voss, A.K.; Strasser, A. The essentials of developmental apoptosis. *F1000Res*. **2020**, *9*, F1000 Faculty Rev-148.

- 278 18. Carneiro, B.A.; El-Deiry, W.S. Targeting apoptosis in cancer therapy. *Nat Rev Clin Oncol*.
279 **2020**, *17*, 395-417.
- 280 19. Zhang, Y.; Hu, C. Anticancer activity of bisindole alkaloids derived from natural sources and
281 synthetic bisindole hybrids. *Arch Pharm (Weinheim)*. **2020**, *353*, e2000092.
- 282 20. Lunagariya, J.; Bhadja, P.; Zhong, S.; Vekariya, R.; Xu, S. Marine Natural Product Bis-indole
283 Alkaloid Caulerpin: Chemistry and Biology. *Mini Rev Med Chem*. **2019**, *19*, 751-761.
- 284 21. Imran, S.; Taha, M. Ismail, N.H. A Review of Bisindolylmethane as an Important Scaffold for
285 Drug Discovery. *Curr Med Chem*. **2015**, *22*, 4412-4433.
- 286 22. Pizzimenti, S.; Toaldo, C.; Pettazzoni, P., Dianzani, M.U.; Barrera, G. The "two-faced" effects
287 of reactive oxygen species and the lipid peroxidation product 4-hydroxynonenal in the hallmarks
288 of cancer. *Cancers (Basel)*. **2010**, *2*, 338-363.
- 289 23. Barrera, G.; Gentile, F.; Pizzimenti, S.; Canuto, R.A.; Daga, M.; Arcaro, A.; Cetrangolo, G.P.;
290 Lepore, A.; Ferretti, C.; Dianzani, C.; Muzio, G. Mitochondrial Dysfunction in Cancer and
291 Neurodegenerative Diseases: Spotlight on Fatty Acid Oxidation and Lipoperoxidation Products.
292 *Antioxidants (Basel)*. **2016**, *5*, 7.

293

294

295 **Figure legends**

296




Figure 2. **IM** and LM observations of control condition (A, B) and BrDIMEA derivative 1 (C, D), 2 (E, F) 3 (G, H) treated cells. The graph in I shows TB exclusion assay which reveals a cell viability reduction after drug administration. Data are from three separate experiments and furnished as mean +/- standard deviation. Bars: 10 μ m




Figure 3. TEM (A-D) and CLSM observations (inset A and C, E-H) of control cells (A, inset A, B, E, F) and those exposed to compound 3 (C, D, inset D, G, H). Apoptotic features, altered mitochondria (arrows) and TUNEL positive nuclei, absent in control condition (A, B and inset A) appears after compound 3 administration (C, D, inset D). If control cells maintain the mitochondrial membrane integrity (E, F), treated cells (G, H) show a decrease of fluorescence after NAO staining evidencing the presence of peroxidation events. Bars: 500nm for A-D; 5µm for inset D; 10 µm for inset A, E-H.




Fig. 3