

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Asymptotic growth of Betti numbers of ordered configuration spaces of an elliptic curve

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Asymptotic growth of Betti numbers of ordered configuration spaces of an elliptic curve / Pagaria R.. - In: EUROPEAN JOURNAL OF MATHEMATICS. - ISSN 2199-675X. - STAMPA. - 8:2(2022), pp. 427-445.
[10.1007/s40879-022-00534-8]

Availability:

This version is available at: <https://hdl.handle.net/11585/886798> since: 2022-05-23

Published:

DOI: <http://doi.org/10.1007/s40879-022-00534-8>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Pagaria, R. Asymptotic growth of Betti numbers of ordered configuration spaces of an elliptic curve. *European Journal of Mathematics* 8, 427–445 (2022)

The final published version is available online at <https://dx.doi.org/10.1007/s40879-022-00534-8>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

ASYMPTOTIC GROWTH OF BETTI NUMBERS OF ORDERED CONFIGURATION SPACES ON AN ELLIPTIC CURVE

ROBERTO PAGARIA

ABSTRACT. We construct a dga to computing the cohomology of ordered configuration spaces on an algebraic variety with vanishing Euler characteristic. It follows that the k -th Betti number of $\text{Conf}(C, n)$ (C is an elliptic curve) grows as a polynomial of degree exactly $2k - 2$. We also compute $H^k(\text{Conf}(C, n))$ for $k \leq 5$ and arbitrary n .

The ordered configuration space of n points on a smooth projective variety X is

$$\text{Conf}(X, n) = \{(p_1, \dots, p_n) \in X^n \mid p_i \neq p_j\}.$$

A central problem in the theory of configuration spaces is understand the cohomology of these topological spaces. The main tool is the Križ model, introduced at the same time by Križ [Kri94] and by Totaro [Tot96], it is a differential graded algebra $E(X, n)$ that codifies the rational homotopy type of $\text{Conf}(X, n)$ (see Section 1). Often is useful to study all these configuration spaces $\text{Conf}(X, n)$ for $n \in \mathbb{N}$ all together. In the case of vanishing Euler characteristic $\chi(X) = 0$, we construct a filtration F_\bullet of the Križ model $E(X, n)$ and we prove that the differential is strict with respect to this filtration (Theorem 1.15). Passing to the graded model we obtain a simpler differential graded algebra that is more feasible for computing cohomology. The mixed Hodge numbers of $\text{Conf}(X, n)$ (indeed any polynomial function $\mathbb{N} \rightarrow \mathbb{N}$) can be written as linear combination of binomials with coefficient in \mathbb{Z} . We prove in Proposition 1.18 that these coefficients are nonnegative integers for any algebraic variety with $\chi(X) = 0$, this is not true in a wider generality.

We apply the simpler dga to configuration on an elliptic curve C . It was known from [Chu12] that the k -th Betti number of $\text{Conf}(C, n)$ grows as polynomial of degree at most $2k$. We improve that result by showing that the Betti numbers grow as a polynomial of degree exactly $2k - 2$ (Theorem 2.9). We developed a method to give lower bounds for the mixed Hodge numbers (and so for Betti numbers) involving certain partitions, see Lemma 2.6. The lower bounds are extremely good for understand the asymptotic behaviour of these numbers.

The upper bound for these number is more difficult to determine, so we developed an algorithm for computing the (not so) small cases. The algorithm uses in a essential way the aforementioned filtration of the Križ model. We finally present the mixed Hodge numbers of $\text{Conf}(C, n)$ for $n \leq 7$ (Appendix A). Using the theory development here and the computation of small cases, we also determine the dim $H^k(\text{Conf}(C, n); \mathbb{Q})$ for $k \leq 5$ and arbitrary $n \in \mathbb{N}$ (Theorem 2.9).

A recent tool to deal with sequence of homogeneous objects is the representation stability introduced by Church and Farb [CF13]. We use a variant of that: we substitute the category of finite sets with injections with the category \mathcal{F} of finite

The author is supported by PRIN 2017YRA3LK.

sets with all maps. The representation theory of F is developed by [Wil14], [SS17], and [Ryb18]. Ellenberg and Wiltshire-Gordon [EW15] have shown that the spaces $\text{Conf}(X, n)$ are an F -module if X has two linearly independent vector fields. Their result implies that the cohomology is an F -module, we give a short proof that the Križ model is an F -module. We use this structure in an essential way to define the filtration F and to prove that is strictly compatible with the differential.

1. REPRESENTATION THEORY OF THE KRIŽ MODEL

The Križ model. Let X be a smooth projective variety. For each element $x \in H^\bullet(X)$ we denote by $x_i \in H^\bullet(X^n)$ its image under the map $H^\bullet(X) \hookrightarrow H^\bullet(X^n)$ induced by the projection $X^n \rightarrow X$ on the i -th factor.

The class of the diagonal $\Delta \in H^{2 \dim_{\mathbb{C}} X}(X \times X)$ is the cohomological class of the subvariety $\{(x, x) \mid x \in X\}$ in $X \times X$. If we fix a graded basis $\{b_1, \dots, b_k\}$ of $H^\bullet(X)$ and consider the dual basis $\{b_1^*, \dots, b_k^*\}$ with respect to the cup product, then

$$\Delta = \sum_{j=1}^k (-1)^{\deg b_j^*} b_j \otimes b_j^* \quad (1)$$

Consider also $\Delta_{i,j} \in H^\bullet(X^n)$ as the pullback of the diagonal Δ with respect to the projection $X^n \rightarrow X^2$ on the i -th and j -th factors.

Definition 1.1. Let $E(X, n)$ be the differential bigraded algebra

$$H^\bullet(X^n)[\{G_{i,j}\}_{i < j}] / I,$$

where $\deg(x_i) = (\deg x, 0)$, $\deg(G_{i,j}) = (0, 2 \dim_{\mathbb{C}} X - 1)$, and I is the ideal generated by

$$\begin{aligned} G_{i,j}(x_i - x_j) & \quad \text{for } i < j \text{ and } x \in H^\bullet(X) \\ G_{i,j}G_{j,k} - G_{i,j}G_{i,k} + G_{j,k}G_{i,k} & \quad \text{for } i < j < k. \end{aligned}$$

The differential is defined by $d(x_i) = 0$ for all x and i and $d(G_{i,j}) = \Delta_{i,j}$.

For the sake of notation, we define for $i > j$ $G_{i,j} = G_{j,i}$. The following theorem was proven by Križ [Kri94] and Totaro [Tot96] independently

Theorem 1.2. The dga $E(X, n)$ is a rational model for $\text{Conf}(X, n)$. Therefore $H^\bullet(\text{Conf}(X, n)) \cong H^\bullet(E(X, n), d)$.

The F module structure. We will consider the spaces $\text{Conf}(X, n)$ and their cohomology groups for $n \in \mathbb{N}$ all together. Consider the *category of finite sets* F whose objects are the sets $[n] = \{1, 2, \dots, n\}$ for all $n \in \mathbb{N}$ and with morphisms all the maps of sets between them. Notice that the automorphism of the object $[n]$ $\text{Aut}_F([n]) \cong \mathfrak{S}_n$ is the symmetric group.

An F -representation over \mathcal{V} is a functor V from the category F to the category \mathcal{V} . A F -module V is *finitely generated* if there exists elements $v_i \in V([n_i])$ for i in a finite set I such that no proper submodule of V contains all v_i for $i \in I$. A morphism $\tau: V \rightarrow W$ of F -module is a natural transformation between the functors V and W .

Definition 1.3. The F -module $E(X)$ is the functor defined by $[n] \mapsto E(X, n)$ on the objects and for each map $f: [n] \rightarrow [m]$ associate the morphism $f_*: E(X, n) \rightarrow$

$E(X, m)$ defined by

$$x_i \mapsto x_{f(i)}, \quad G_{i,j} \mapsto \begin{cases} G_{f(i),f(j)} & \text{if } f(i) \neq f(j) \\ 0 & \text{otherwise} \end{cases}$$

The submodules $E^{p,q}(X)$ are finitely generated and $d: E(X) \rightarrow E(X)$ commutes with f_* for all injective maps f .

Proposition 1.4. If the Euler characteristic of X is zero, i.e. $\chi(X) = 0$. Then d is a morphism of F -modules.

Proof. We verify that $f(i) = f(j)$ implies $f_*(d(G_{i,j})) = 0$, the other conditions are trivial. Let $\{b_1, \dots, b_k\}$ be a graded basis of $H^\bullet(X)$ and $\{b_1^*, \dots, b_k^*\}$ be the dual basis, we use Equation (1):

$$\begin{aligned} f_*(d(G_{i,j})) &= f_* \left(\sum_{l=1}^k (-1)^{\deg b_l^*} (b_l)_i (b_l^*)_j \right) \\ &= \sum_{l=1}^k (-1)^{\deg b_l^*} (b_l)_{f(i)} (b_l^*)_{f(i)} \\ &= \sum_{l=1}^k (-1)^{\deg b_l^*} [X]_{f(i)} \\ &= \sum_{r=1}^{2 \dim(X)} (-1)^r h_r(X) [X]_{f(i)} \\ &= \chi(X) [X]_{f(i)}, \end{aligned}$$

where $h_r(X)$ is the r -th Betti number of X . The hypothesis $\chi(X) = 0$ completes the proof. \square

A topological interpretation of the above result can be found in [EW15].

Representation theory of F . The irreducible representations of the symmetric group \mathfrak{S}_n are parametrized by the partitions $\lambda \vdash n$ of the number n , we denote them by V_λ . The category \mathcal{V}^F of the F representation over \mathcal{V} is not semisimple, however has the Jordan Hölder property as shown in [Wil14, Corollary 5.4]. The *Schur projective* representation of type $\lambda \vdash k$ is the functor

$$\mathbb{P}_\lambda([n]) = \mathbb{S}_\lambda(\mathbb{Q}^n),$$

where \mathbb{S}_λ is the Schur functor (see [Wey03, Mac15] for an introductory exposition of Schur functors). The dimension $\dim \mathbb{P}_\lambda([n])$ is an evaluation of the Schur symmetric polynomial, i.e. $s_\lambda(1^n)$. These projective representations have the following property

$$\mathrm{Hom}_{\mathcal{V}^F}(\mathbb{P}_\lambda, V) \cong \mathrm{Hom}_{\mathfrak{S}_k}(V_\lambda, V([k])), \quad (2)$$

where $\lambda \vdash k$.

The simple representations are classified by Wiltshire-Gordon and they are of two kinds: D_k for $k \in \mathbb{N}$ and C_λ for $\lambda \vdash k$ with $\lambda_1 > 1$. The representation C_λ for $\lambda \vdash k$ is defined by $C_\lambda([n]) = 0$ for $n < k$ and by

$$C_\lambda([n]) = \mathrm{Ind}_{\mathfrak{S}_k \times \mathfrak{S}_{n-k}}^{\mathfrak{S}_n} V_\lambda \boxtimes 1_{n-k},$$

where 1_n is the trivial representation of \mathfrak{S}_n and the dimension is $\binom{n}{k} \langle s_\lambda, p_{1^k} \rangle$. The representation D_0 is defined by $D_0([0]) = \mathbb{Q}$ and $D_0([n]) = 0$ for all $n > 0$. Finally, D_k is defined by $D_k([n]) = 0$ for $n < k$ and by

$$D_k([n]) = V_{(n-k+1, 1^{k-1})}.$$

Moreover $D_k([n])$ has dimension $\binom{n-1}{k-1}$ for $n, k > 0$.

Definition 1.5. A finitely generated representation V is of degree k if there exists a surjection $\oplus_i \mathbb{P}_{\lambda_i} \twoheadrightarrow V$ with $\lambda_i \vdash n_i$ and $n_i \leq k$.

Since there exist surjections $\mathbb{P}_\lambda \twoheadrightarrow C_\lambda$ and $\mathbb{P}_{1^k} \twoheadrightarrow D_k$, D_k has degree k and C_λ has degree $|\lambda|$.

Proposition 1.6 ([Wil14, Theorem 6.39]). Every finitely generated representation V of degree k has a finite resolution

$$0 \rightarrow \mathbb{P}_0 \oplus \mathbb{D}_0 \rightarrow \cdots \rightarrow \mathbb{P}_{k-1} \oplus \mathbb{D}_{k-1} \rightarrow \mathbb{P}_k \rightarrow V \rightarrow 0,$$

where \mathbb{P}_i is a direct sum of Schur projective of pure degree $\leq i$ and \mathbb{D}_i is direct sum of some copies of D_{i+1} and D_{i+2} . Moreover, D_k has an infinite projective resolution given by

$$\cdots \rightarrow \mathbb{P}_{1^{k+2}} \rightarrow \mathbb{P}_{1^{k+1}} \rightarrow \mathbb{P}_{1^k} \rightarrow D_k \rightarrow 0.$$

Lemma 1.7. We have the following:

- (1) if $h \neq k + i$ then $\text{Ext}^i(D_k, D_h) = 0$, and $\text{Ext}^i(D_k, D_{k+i}) = \mathbb{Q}$,
- (2) let V a representation of degree k , then $\text{Ext}^i(V, C_\lambda) = 0$ if $k < |\lambda| + i$.

Proof.

- (1) Consider the resolution \mathbb{P}_\bullet of D_k from Proposition 1.6, we compute the Ext functor using the projective resolution $\text{Ext}^i(D_k, D_h) = H^i(\text{Hom}(\mathbb{P}_\bullet, D_h))$. The complex $\text{Hom}(\mathbb{P}_\bullet, D_h)$ is almost trivial because

$$\text{Hom}_F(\mathbb{P}_{1^{k+i}}, D_h) \cong \text{Hom}_{\mathfrak{S}_{k+i}}(V_{1^{k+i}}, V_{(k+i-h+1, 1^{h-1})})$$

is zero for $k + i \neq h$ and for $h = k + i$ is \mathbb{Q} by the Schur Lemma.

- (2) Consider the resolution $\mathbb{P}_\bullet \oplus \mathbb{D}_\bullet$ of V from Proposition 1.6. It is enough to verify that $\text{Hom}_F(\mathbb{P}_{k-i} \oplus \mathbb{D}_{k-i}, C_\lambda) = 0$, indeed $\text{Hom}(D_a, C_\lambda) = 0$ because they are two different simple representations and

$$\text{Hom}_F(\mathbb{P}_\mu, C_\lambda) = \text{Hom}_{\mathfrak{S}_{|\mu|}}(V_\mu, 0) = 0,$$

since $|\mu| \leq k - i < |\lambda|$. \square

Let V be a finitely generated representation of F of degree k . Define a canonical filtration F_\bullet such that $F_i \subseteq V$ is the submodule generated by $V([j])$ for all $j \leq i$.

Example 1.8. Consider the short exact sequence

$$0 \rightarrow D_2 \rightarrow \mathbb{P}_1 \rightarrow D_1 \rightarrow 0,$$

on the object $[n]$, $\mathbb{P}_1([n]) = \mathbb{Q}^n$, $D_2([n]) \cong \mathbb{Q}^{n-1}$ is the subobject of vectors whose coordinates have zero sum, and $D_1([n]) \cong \mathbb{Q}$ is their quotient. The sequence does not split and the degree of the subject D_2 (equals to 2) is bigger then the degree of \mathbb{P}_1 (equals to 1).

The inclusion $j: D_2([n]) \hookrightarrow \mathbb{P}_1([n])$ has rank $n - 1$, but the graded map $\text{gr}_F j$ is zero.

Example 1.9. In [Ryb18], minimal projective resolutions of C_λ are described. The minimal resolution of $C_{(4)}$ is

$$0 \rightarrow \mathbb{P}_{(1,1)} \rightarrow \mathbb{P}_{(2,1)} \oplus \mathbb{P}_{(3)} \rightarrow \mathbb{P}_{(4)} \rightarrow C_{(4)} \rightarrow 0.$$

It follows from eq. (2) that $\dim \text{Ext}_F^1(C_{(4)}, C_{(2)}) = 1$.

Lemma 1.10. Let V a representation of degree k , then the composition factors of the Jordan Hölder filtration are C_λ for $|\lambda| \leq k$ or D_i for $i \leq k+1$.

Proof. It is enough to prove the claim for $V = \mathbb{P}_k$ a Schur projective representation of degree k . The homology functor H_0 is defined by $H_0(V)([n]) = V/F_{n-1}V([n])$, this functor is right exact.

Consider a Jordan Hölder filtration G_\bullet of \mathbb{P}_k . Proposition 6.32 of [Wil14] shows that $H_0(G_i)[n] = 0$ for $n > k+1$ and that $H_0(G_i)[k+1]$ is sum of sign representations of \mathfrak{S}_{k+1} . The short exact sequence

$$H_0(G_{i-1})[n] \rightarrow H_0(G_i)[n] \rightarrow H_0(G_i/G_{i-1})[n] \rightarrow 0$$

implies $H_0(G_i/G_{i-1})[n] = 0$ for $n > k+1$ and $H_0(G_i/G_{i-1})[k+1]$ contains only the sign representation. Now G_i/G_{i-1} is simple and the homology of simple modules of degree j is concentrate in degree j . The equalities $H_0(C_\lambda)[|\lambda|] = V_\lambda$ and $H_0(D_k)[k] = V_{1^k}$ imply the result. \square

Theorem 1.11. Let $f: V \rightarrow W$ be a morphism between two finitely generated F -representations and $\text{gr}_F f: \text{gr}_F V \rightarrow \text{gr}_F W$ be the corresponding graded map. Suppose that the composition factors of the Jordan Hölder filtration of V are different from D_i , for $i \in \mathbb{N}$. Then:

- (1) $\text{gr}_F V$ is semisimple and the addenda of F_n/F_{n-1} are of the type C_λ for $\lambda \vdash n$,
- (2) $\text{rk}(f) = \text{rk}(\text{gr } f)$, i.e the map f is strict with respect the filtration F_\bullet .

Proof. Consider the composition factors of $F_n V/F_{n-1} V$ have degree at least n because $F_n V/F_{n-1} V([n-1]) = 0$ and at most n by Lemma 1.10 using the hypothesis that D_{n+1} does not appear. Therefore the composition factors are C_λ for $\lambda \vdash n$. Since $\text{Ext}^1(C_\lambda, C_\mu) = 0$ for $\lambda, \mu \vdash n$, an inductive reasoning proves that F_n/F_{n-1} is semisimple.

Suppose that $f(y) = x$ with $y \in F_n V \setminus F_{n-1} V$ and $x \in F_{n-1} W$, we show that $x \in f(F_{n-1} V)$. Let (x) and (y) be the subrepresentation generated by x and y , f induces a surjective morphism

$$\bar{f}: (y)/(y) \cap F_{n-1} V \rightarrow (x)/(x) \cap f(F_{n-1} V).$$

As noticed above $H_0((y)/(y) \cap F_{n-1} V)$ is concentrate in degree n and does not contain the sign representation, instead $H_0((x)/(x) \cap f(F_{n-1} V))[n]$ contains only some copies of the sign representation. Thus, $H_0(\bar{f})$ is zero and surjective because \bar{f} does. This implies $H_0((x)/(x) \cap f(F_{n-1} V)) = 0$, so $(x)/(x) \cap f(F_{n-1} V) = 0$ and hence $x \in f(F_{n-1} V)$. \square

Representation theory of $E(X)$. From now on, we assume that $\chi(X) = 0$. Let $F(k, n)$ be the set of all maps from $[k]$ to $[n]$. The increasing filtration $F_\bullet E(X)$ defined in the previous section can be described as

$$F_k E(X) = \sum_{f \in F(k, n)} \text{Im } f_*.$$

The module $F_k E(X)[n]$ is, by definition, the submodule of $E(X, n)$ generated by all monomials in $G_{i,j}$ and x_i (for $i, j \leq n$ and $x \in H^\bullet(X)$) with at most k different indices.

Example 1.12. Let $x \in H^\bullet(X)$ be any element. We have $G_{1,2} \in F_2 E(X) \setminus F_1 E(X)$, $x_1 - x_2 \in F_1 E(X)$ and $x_1 x_2 \in F_2 E(X) \setminus F_1 E(X)$.

We fix a graded basis $\{b_i\}_i$ of $H^\bullet(X)$ and assume $b_1 = 1 \in H^0(X)$. A *labelled partition* λ of n is $\lambda \vdash n$ whose blocks are decorated with elements of the fixed basis. We define $q(\lambda) = n - l(\lambda)$ the difference between n and the number of blocks $l(\lambda)$, and $p(\lambda) = \sum_{i=1}^{l(\lambda)} \deg(\lambda_i)$ the sum of the cohomological degrees of all the labels. Let $L(\lambda) < \mathfrak{S}_n$ be a subgroup generated by disjoint cycles of length λ_i for $i = 1, \dots, l(\lambda)$, $L(\lambda)$ is isomorphic to $\times_{i=1}^{l(\lambda)} C_{\lambda_i}$. Let $N(\lambda) < \mathfrak{S}_n$ be the subgroup of the stabilizer of $L(\lambda)$ that permutes the cycles with the same labels, it is isomorphic to $\times_j \mathfrak{S}_{n_j}$ for some n_j . Finally, define $Z(\lambda) < \mathfrak{S}_n$ as the subgroup generated by $L(\lambda)$ and $N(\lambda)$, i.e. the semidirect product $L(\lambda) \rtimes N(\lambda)$.

We denote the sign representation of \mathfrak{S}_i by ϵ_i and a faithful representation of the cyclic group C_i by φ_i . Let $\varphi_\lambda, \alpha_\lambda, \xi_\lambda$ be the one dimensional representations of $L(\lambda), N(\lambda), Z(\lambda)$ defined by:

$$\begin{aligned}\varphi_\lambda &= \epsilon_{n|L(\lambda)} \otimes (\boxtimes_{i=1}^{l(\lambda)} \varphi_{\lambda_i}), \\ \alpha_\lambda &= \boxtimes_j \epsilon_j^{\otimes m_j}, \\ \xi_\lambda &= \varphi_\lambda \boxtimes \alpha_\lambda,\end{aligned}$$

where $m_j = \lambda_i + \deg(\lambda_i) + 1$ where λ_i is any block permuted by \mathfrak{S}_{n_j} .

Example 1.13. Consider an elliptic curve $C = (S^1)^2$ and the basis of $H^\bullet(C)$ given by $1, x, y, xy$. Let $\lambda = (4, 4, 4, 4, 1, 1, 1) \vdash 19$ with labels $(xy, xy, xy, 1, x, x, x)$. We have $l(\lambda) = 7$, $q(\lambda) = 12$, $p(\lambda) = 9$. The associated groups are $L(\lambda) \cong (C_5)^{\times 4}$ generated by $(1, 2, 3, 4), (5, 6, 7, 8), (9, 10, 11, 12), (13, 14, 15, 16)$, $N(\lambda) \cong \mathfrak{S}_3 \times \mathfrak{S}_3$ generated by $(1, 5)(2, 6)(3, 7)(4, 8), (1, 9)(2, 10)(3, 11)(4, 12), (17, 18), (17, 19)$, and $Z(\lambda) = (C_4 \wr \mathfrak{S}_3) \times C_4 \times \mathfrak{S}_3 < \mathfrak{S}_{19}$. The representations are $\varphi_\lambda = \varphi_4 \boxtimes \varphi_4 \boxtimes \varphi_4 \boxtimes \varphi_4$ (because $\epsilon_{19|C(\lambda)} = 1$), $\alpha = \epsilon_3 \boxtimes \epsilon_3$ (because $4 + 2 + 1$ and $1 + 1 + 1$ are odd) and $\xi_\lambda = (\varphi_4 \wr \epsilon_3) \boxtimes \varphi_4 \boxtimes \epsilon_3$.

A decomposition of $E(X, n)$ into \mathfrak{S}_n representations is provided in [AAB14].

Theorem 1.14. Let X be a smooth projective variety. The Križ model decomposes as

$$E^{p,q}(X, n) \cong \bigoplus_{\substack{q(\lambda)=q \\ p(\lambda)=p}} \text{Ind}_{Z(\lambda)}^{\mathfrak{S}_n} \xi_\lambda.$$

For each representation $V = \bigoplus_\lambda V_\lambda^{n_\lambda}$ of \mathfrak{S}_n , we define the F -representation $C_V = \bigoplus_\lambda C_\lambda^{n_\lambda}$. For any labelled partition λ let $f(\lambda)$ the number of blocks of size 1 labelled with $1 \in H^\bullet(X)$.

Theorem 1.15. Let X be a smooth projective algebraic variety with $\chi(X) = 0$. Then

$$\text{gr}_F^\bullet H^{\bullet,\bullet}(E(X), d) = H^{\bullet,\bullet}(\text{gr}_F^\bullet E(X), \text{gr}_F^\bullet d)$$

and for $q > 0$ the F -representation $E^{p,q}(X)$ has associated graded:

$$\text{gr}_F^r E^{p,q}(X) = \bigoplus_\lambda C_{\text{Ind}_{Z(\lambda)}^{\mathfrak{S}_n} \xi_\lambda},$$

where the sum is taken over all labelled partitions $\lambda \vdash r$ with $p(\lambda) = p$, $q(\lambda) = q$, and $f(\lambda) = 0$.

Proof. We first prove that the sign representation ϵ_n does not appear in $E^{p,q}(X, n)$ for $q > 0$. We have

$$\begin{aligned} \langle \epsilon_n, E^{p,q}(X, n) \rangle_{\mathfrak{S}_n} &= \sum_{\lambda} \langle \epsilon_n, \text{Ind}_{Z(\lambda)}^{\mathfrak{S}_n} \xi_{\lambda} \rangle_{\mathfrak{S}_n} \\ &= \sum_{\lambda} \langle \epsilon_n|_{Z(\lambda)}, \xi_{\lambda} \rangle_{Z(\lambda)} \\ &= \sum_{\lambda} \langle \epsilon_n|_{L(\lambda)}, \varphi_{\lambda} \rangle_{L(\lambda)} \langle \epsilon_n|_{N(\lambda)}, \alpha_{\lambda} \rangle_{N(\lambda)} \\ &= \sum_{\lambda} \langle 1_{L(\lambda)}, \boxtimes_{i=1}^{\|\lambda\|} \varphi_{\lambda_i} \rangle_{L(\lambda)} \langle \epsilon_n|_{N(\lambda)}, \alpha_{\lambda} \rangle_{N(\lambda)} \\ &= \sum_{\lambda} \langle \epsilon_n|_{N(\lambda)}, \alpha_{\lambda} \rangle_{N(\lambda)} \prod_{i=1}^{\|\lambda\|} \langle 1_{\lambda_i}, \varphi_{\lambda_i} \rangle_{C_{\lambda_i}} = 0, \end{aligned}$$

because $\langle 1_{\lambda_i}, \varphi_{\lambda_i} \rangle_{C_{\lambda_i}} \neq 0$ if and only if $1_{\lambda_i} = \varphi_{\lambda_i}$ if and only if $\lambda_i = 1$. The sign representation appears only in $E^{p,0}(X, n)$.

It follows that D_i cannot be a composition factor of $E^{p,q}(X)$ for $q > 0$, so by Theorem 1.11 we obtain that d is strict.

The second part of the statement follows from the fact that the submodule $\text{Ind}_{Z(\lambda)}^{\mathfrak{S}_n} \xi_{\lambda}$ is contained in $F_r E^{p,q}(X)([n])$ if and only if $f(\lambda) \geq n - r$. Therefore,

$$F_r E^{p,q}(X) / F_{r-1} E^{p,q}(X)([r]) = \bigoplus_{\lambda \text{ s.t. } f(\lambda)=0} \text{Ind}_{Z(\lambda)}^{\mathfrak{S}_n} \xi_{\lambda},$$

and since $\text{gr}_F^r E^{p,q}(X)$ is semisimple, we obtain the claimed equality. \square

Example 1.16. In the case $X = C$, we have $d(G_{i,j}) = (x_i - x_j)(y_i - y_j)$ and $\text{gr}_F d(G_{i,j}) = -x_i y_j - x_j y_i$. Moreover $\text{gr}_F d(x_i G_{i,j}) = x_i x_j y_j - x_i x_j y_i$ and notice that the product of x_i and $G_{i,j}$ in $\text{gr}_F^{\bullet} E(C)$ is zero.

Behaviour of mixed Hodge numbers. Recall that every polynomial $P(t)$ in $\mathbb{Q}[t]$ such that $P(n) \in \mathbb{N}$ for all $n \in \mathbb{N}$ can be written uniquely as

$$P(t) = \sum_{i=0}^{\deg P} a_i \binom{t}{i},$$

for some $a_i \in \mathbb{Z}$. The value a_i not need to be positive.

Example 1.17. Consider $\dim H^2(\text{Conf}(S^3, n); \mathbb{Q})$ for $n > 0$, it is known that is polynomial in n and its value is $\frac{(n-1)(n-2)}{2}$. However $a_1 = -1$ because

$$\dim H^2(\text{Conf}(S^3, n); \mathbb{Q}) = \frac{(n-1)(n-2)}{2} = \binom{n}{2} - \binom{n}{1} + \binom{n}{0}.$$

Indeed $H^2(\text{Conf}(S^3)) \cong D_3$ as F-module (included in position $(0, 2)$ of the Leray spectral sequence).

We collect the information about $\dim H^{p,q}(E(X, n), d)$ in a polynomial

$$P^{p,q}(n) := \dim H^{p,q}(E(X, n), d) = \sum_{i=0}^{p+2q} a_i^{p,q} \binom{n}{i},$$

for some unique integers $a_i^{p,q}$.

Proposition 1.18. For $q > 0$, the coefficients $a_i^{p,q}$ are positive and coincide with

$$a_i^{p,q} = \dim H^{p,q} \left(E(X, i) /_{F_{i-1}} E(X, i), d \right).$$

Proof. We use Theorem 1.15 and the fact that $\mathrm{gr}_F^i E^{p,q}(X)$ has not composition factors of type D_k to obtain:

$$\begin{aligned} \dim H^{p,q}(E(X, n), d) &= \sum_{i \leq p+2q} \dim \mathrm{gr}_F^i H^{p,q}(E(X, n)) \\ &= \sum_{i \leq p+2q} \dim H^{p,q}(\mathrm{gr}_F^i E(X, n)) \\ &= \sum_{i \leq p+2q} \dim H^{p,q} \left(\mathrm{Ind}_{\mathfrak{S}_i \times \mathfrak{S}_{n-i}}^{\mathfrak{S}_n} E(X, i) /_{F_{i-1}} E(X, i) \right) \\ &= \sum_{i \leq p+2q} \dim \mathrm{Ind}_{\mathfrak{S}_i \times \mathfrak{S}_{n-i}}^{\mathfrak{S}_n} H^{p,q} \left(E(X, i) /_{F_{i-1}} E(X, i) \right) \\ &= \sum_{i \leq p+2q} \binom{n}{i} \dim H^{p,q} \left(E(X, i) /_{F_{i-1}} E(X, i) \right) \end{aligned}$$

Since \mathbb{N} is infinite we obtain the corresponding equality between polynomials. \square

2. THE ELLIPTIC CASE

Let C be an elliptic curve, topologically $C = (S^1)^2$, the cohomology $H^\bullet(C; \mathbb{Q})$ is the exterior algebra on two generator x and y such that $x \smile y = [C]$. The construction of the previous section is compatible with the action of $SL_2(\mathbb{Q})$.

We recall two results from [Pag18, Lemma 1.6, Theorem 3.9].

Proposition 2.1. The cohomology $H^k(\mathrm{Conf}(C, n))$ vanishes for $k > n + 1$ and so $a_i^{p,q} = 0$ for $p + q > i + 1$ or $p + 2q < i$. Moreover the mixed Hodge numbers for $q = 0$ are given by:

$$P^{p,0}(t) = (p+1) \binom{t}{p} + (p-1) \binom{t}{p-1}.$$

Indeed, an easy computation shows that

$$H^{p,0}(\mathrm{Conf}(C)) = (\mathbb{P}_{1^p} \boxtimes \mathbb{V}_p) \oplus (\mathbb{P}_{1^{p-1}} \boxtimes \mathbb{V}_{p-2}),$$

as representation of $F \times SL_2(\mathbb{Q})$.

Remark 2.2. The multiplicity of the representation $C_{(k)}$ for $k > 1$ in the graded cohomology $\mathrm{gr}_F^\bullet H^\bullet(\mathrm{Conf}(C))$ can be deduced from [Pag19].

For the sake of notation, we denote $\mathrm{Ind}_{\mathfrak{S}_n \times \mathfrak{S}_m}^{\mathfrak{S}_{n+m}} V \boxtimes W$ by $V \cdot W$.

Lemma 2.3. The following decomposition of $\mathfrak{S}_{p+2q} \times SL_2(\mathbb{Q})$ -representations holds:

$$\mathrm{gr}_F^{p+2q} E^{p,q}(C, p+2q) \cong \bigoplus_{a=0}^{\lfloor \frac{p}{2} \rfloor} \left(\mathrm{Ind}_{\mathfrak{S}_{2^a} \wr \mathfrak{S}_q}^{\mathfrak{S}_{2q}} V_{(2)}^{\boxtimes q} \otimes V_{(1^q)} \right) \cdot V_{(2^a, 1^k)} \boxtimes \mathbb{V}_k,$$

where $p = 2a + k$.

Proof. From Theorem 1.15, we have

$$\mathrm{gr}_F^{p+2q} E^{p,q}(C, p+2q) \cong \bigoplus_{b=0}^p \left(\mathrm{Ind}_{C_{2^b} \wr \mathfrak{S}_q}^{\mathfrak{S}_{2q}} V_{(2)}^{\boxtimes q} \otimes V_{(1^q)} \right) \cdot V_{(1^{p-b})} \cdot V_{(1^b)}$$

because the labelled partitions $\lambda \vdash p+2q$ with $p(\lambda) = p$, $q(\lambda) = q$ and $f(\lambda) = 0$ are $(2^q, 1^{p-b}, 1^b)$ for some b with blocks labelled respectively by 1, x , and y . Moreover for such λ we have $\varphi_\lambda = V_{(2)}^{\boxtimes q}$ and $\alpha_\lambda = V_{(1^q)} \boxtimes V_{(1^{p-b})} \boxtimes V_{(1^b)}$. By definition the maximal torus of $SL_2(\mathbb{Q})$ acts with weight $p - 2b$ on the b -th addendum, so

$$\mathrm{gr}_F^{p+2q} E^{p,q}(C, p+2q) \cong \bigoplus_{a=0}^{\lfloor \frac{p}{2} \rfloor} W \cdot \left(V_{(1^{a+k})} \cdot V_{(1^a)} \ominus V_{(1^{a+k+1})} \cdot V_{(1^{a-1})} \right) \boxtimes \mathbb{V}_k,$$

where $k = p - 2a$ and $W = \mathrm{Ind}_{C_{2^a} \wr \mathfrak{S}_q}^{\mathfrak{S}_{2q}} V_{(2)}^{\boxtimes q} \otimes V_{(1^q)}$. The representation $V_{(1^{a+k})} \cdot V_{(1^a)} \ominus V_{(1^{a+k+1})} \cdot V_{(1^{a-1})}$ has dimension $\binom{p}{a} - \binom{p}{a-1}$. Using the Littlewood-Richardson rule (see [Ful97]), we observe that the representation $V_{(2^a, 1^k)}$ appears in $V_{(1^{a+k})} \cdot V_{(1^a)}$ but not in $V_{(1^{a+k+1})} \cdot V_{(1^{a-1})}$. The hook formula shows that $\dim V_{(2^a, 1^k)} = \frac{p!(k+1)}{a!a+k+1!} = \frac{k+1}{a+k+1} \binom{p}{a}$ and an easy computation show that

$$\dim V_{(2^a, 1^k)} = \dim (V_{(1^{a+k})} \cdot V_{(1^a)} \ominus V_{(1^{a+k+1})} \cdot V_{(1^{a-1})}).$$

Since the first representation is contained in the second one, we complete the proof. \square

Oyster partitions. We need the Frobenius notation for partitions: for any sequences $a_1 > a_2 > \dots > a_k > 0$ and $b_1 > b_2 > \dots > b_k > 0$ let $(a_1, \dots, a_k \mid b_1, \dots, b_k)$ be the partition of $\sum_{i=1}^k (a_i + b_i - 1)$ such that the i -th row has length $a_i + i - 1$ and the i -th column has length $b_i + i - 1$ for $i \leq k$. Let $Q(n)$ be the set of all the partitions of n of the form $(a_1, \dots, a_k \mid a_1 - 1, \dots, a_k - 1)$. From [Wey03, Proposition 2.3.9 (a)], [Sta99, Theorem A2.8], or [Mac15, Appendix A eq. (6.2)], we have

$$\mathrm{Ind}_{\mathfrak{S}_{2^a} \wr \mathfrak{S}_q}^{\mathfrak{S}_{2q}} V_{(2)}^{\boxtimes q} \otimes V_{(1^q)} \cong \bigoplus_{\lambda \in Q(2q)} V_\lambda.$$

Definition 2.4. A k -core partition of $2q+k$ is any partition of the form $(\lambda_1, \dots, \lambda_k)$ such that $(\lambda_1 - 1, \dots, \lambda_k - 1)$ is in $Q(2q)$. A (k, a) -shell partition is any partition of the form $(b_1 + 3, \dots, b_a + 3 \mid b_1, \dots, b_a)$ with $b_a > k$. A (k, a) -oyster partition is a partition $(c_1, \dots, c_{a+k} \mid d_1, \dots, d_{a+k})$ such that $(c_1, \dots, c_a \mid d_1, \dots, d_a)$ is a (k, a) -shell partition and $(c_{a+1}, \dots, c_{a+k} \mid d_{a+1}, \dots, d_{a+k})$ is a k -core partition.

Example 2.5. Let $k = 2$ and consider the 2-core partition $(4, 4)$ obtained from $(3, 2 \mid 2, 1) \in Q(6)$ and the $(2, 1)$ -shell partition $(6, 1, 1)$. The union of the shell and

the core give the following oyster partition $\lambda = (6, 5, 5)$:

				1	1
				2	
				3	

The representation $V_\lambda \boxtimes \mathbb{V}_2$ is an addendum of

$$V_{(4,4,4)} \cdot V_{(2,1,1)} \boxtimes \mathbb{V}_2 \subset \left(\text{Ind}_{\mathfrak{S}_2 \wr \mathfrak{S}_6}^{\mathfrak{S}_{12}} V_{(2)}^{\boxtimes 6} \otimes V_{(1^6)} \right) \cdot V_{(2,1,1)} \boxtimes \mathbb{V}_2 \subset \text{gr}_F^{16} E^{4,6}(C, 16).$$

Indeed, $6 = 3 + 2 + 1$ and the partition $(4, 4, 4)$ is in Frobenius notation $(4, 3, 2 \mid 3, 2, 1)$. The Littlewood-Richardson rule shows that the multiplicity of V_λ in $V_{(4,4,4)} \cdot V_{(2,1,1)}$ is one and correspond to the skew semistandard Young tableaux of shape $(6, 5, 5)/(4, 4, 4)$ of content $(2, 1, 1)$ shown above.

Lemma 2.6. The module $\text{gr}_F^{p+2q} H^{p,q}(\text{Conf}(C))$ contains

$$\bigoplus_{a=0}^{\lfloor \frac{p}{2} \rfloor} \bigoplus_{\lambda(k,a)\text{-oyster}} C_{V_\lambda} \boxtimes \mathbb{V}_k,$$

where $k = p - 2a$ and the sum is taken over all (k, a) -oyster partitions of $p + 2q$.

Proof. It is enough to prove that for every (k, a) -oyster partition $\lambda \vdash p + 2q$ the representation $V_\lambda \boxtimes \mathbb{V}_k$ appears in $\text{gr}_F^{p+2q} E^{p,q}(C, p + 2q)$ but not in $\text{gr}_F^{p+2q} E^{p+2,q-1}(C, p + 2q)$ neither in $\text{gr}_F^{p+2q} E^{p-2,q+1}(C, p + 2q)$.

For $\lambda = (\lambda_1, \dots, \lambda_r)$ consider the partition $\mu = (\mu_1, \dots, \mu_r)$ such that $\mu_i = \lambda_i - 2$ for $i \leq a$, $\mu_i = \lambda_i - 1$ for $a < i \leq a + k$, and $\mu_i = \lambda_i$ for $a + k < i$. By construction μ belongs to $Q(2q)$ and so V_μ appears in $V_{(1^q)}[V_2]$. Applying the Littlewood-Richardson rule to $V_\mu \cdot V_{(2^a, 1^k)}$, we obtain that V_λ has multiplicity one in $V_\mu \cdot V_{(2^a, 1^k)}$. We have proven $V_\lambda \boxtimes \mathbb{V}_k \subset \text{gr}_F^{p+2q} E^{p,q}(C)$.

In order to show that $\text{gr}_F^{p+2q} E^{p-2,q+1}(C, p + 2q)$ does not contain $V_\lambda \boxtimes \mathbb{V}_k$ is equivalent to show that V_λ is not contained in $V_\eta \cdot V_{(2^{a-1}, 1^k)}$ for all $\eta \in Q(2q + 2)$. Suppose by contradiction that there exist $\eta \in Q(2q + 2)$ and a Littlewood-Richardson skew tableau of shape λ/η and content $(2^{a-1}, 1^k)$. The i -th column of λ has length $\lambda_i - 3$ for $i \leq a$ and the one of μ has length $\mu_i - 1$ for all i . Since μ is contained in λ then $\mu_i \leq \lambda_i - 2$ for all $i \leq a$. In particular λ/η has at least $2a$ boxes in the first a rows. From the reverse lattice word property of λ/η the number j cannot appear in the first $j - 1$ rows, hence λ/η has at most $2a - 1$ boxes in the first a rows. We have obtained a contradiction.

Suppose that V_λ is contained in $V_\eta \cdot V_{(2^{a+1}, 1^k)}$ for some $\eta \in Q(2q - 2)$. The restrictions of $\eta \in Q(2q - 2)$ and the existence of a Littlewood-Richardson skew tableau of shape λ/η and content $(2^{a+1}, 1^k)$ imply that $\eta_i = \lambda_i - 2$ for all $i \leq a$. The inequality $\eta_i \leq \lambda_i - 2$ holds because columns of η are shorter than the corresponding ones of λ , the other inequality $\eta_i \geq \lambda_i - 2$ holds because the skew tableaux λ/η has at most $2i$ elements in the first i -th rows for all i . Therefore λ/η has at most $a + k$ nonempty rows, but each Littlewood-Richardson skew tableau of content $(2^{a+1}, 1^k)$ must have at least $a + k + 1$ nonempty rows. This is contrary to the hypothesis that V_λ is contained in $V_\eta \cdot V_{(2^{a+1}, 1^k)}$ for some $\eta \in Q(2q - 2)$. \square

Corollary 2.7. We have the following lower bounds for the bi-graded Betti numbers:

$$\begin{aligned} \mathrm{gr}_F^{2q+2} H^{2,q}(E(C), d) &\supseteq C_{(q+3|q)} \boxtimes \mathbb{V}_0, \\ \mathrm{gr}_F^8 H^{2,3}(E(C), d) &= C_{(6|3)} \boxtimes \mathbb{V}_0 \oplus C_{(4,3|2,1)} \boxtimes \mathbb{V}_2, \\ \mathrm{gr}_F^6 H^{2,2}(E(C), d) &= C_{(5|2)} \boxtimes \mathbb{V}_0 \oplus C_{(4,1|2,1)} \boxtimes \mathbb{V}_2, \\ \mathrm{gr}_F^4 H^{2,1}(E(C), d) &= C_{(4|1)} \boxtimes \mathbb{V}_0 \oplus C_{(3|2)} \boxtimes \mathbb{V}_2, \\ \mathrm{gr}_F^3 H^{1,1}(E(C), d) &= C_{(3|1)} \boxtimes \mathbb{V}_1. \end{aligned}$$

In particular $\dim \mathrm{gr}_F^{2q+2} H^{2,q}(E(C, n), d) \geq \binom{2q+1}{q-1} \binom{n}{2q+2}$.

Proof. The first inclusion follows from Lemma 2.6 and the fact that $(q+3|q)$ is an $(0, 2)$ -oyster partition of $2q+2$ for $q > 0$. The following oyster partitions (with empty shells):

			1
			2

			1
	2		

		1
2		

		1

imply that the right hand sides are contained in the left hand sides. The other containment follows from a dimensional argument: the dimensions of the left hand sides are computed in Tables 7, 8 and 10 and eq. (11) and coincide with the dimension of the representations on the right. Finally, since $\dim V_{(q+3, 1^{q-1})} = \binom{2q+1}{q-1}$, it follows that $\dim C_{V_{(q+3, 1^{q-1})}}([n]) \boxtimes \mathbb{V}_0 = \binom{2q+1}{q-1} \binom{n}{2q+2}$. \square

Upper bounds for Betti numbers. We denote by $E^{p,q}(C)_k$ the subspace in $E^{p,q}(C)$ of highest vectors for $SL_2(\mathbb{Q})$ of weight k and similar for $\mathrm{gr}_F E^{p,q}(C)_k$ and for $H^{p,q}(E(C), d)_k$. Therefore

$$E^{p,q}(C) \cong \bigoplus_{a=0}^{\lfloor \frac{p}{2} \rfloor} E^{p,q}(C)_{p-2a} \boxtimes \mathbb{V}_{p-2a}.$$

In order to give upper bounds for Betti numbers, we need the following result.

Lemma 2.8. The following cohomology groups are zero:

$$H^{0,q}(E(C), d) = 0 \quad q > 0, \quad (3)$$

$$\mathrm{gr}_F^{2q+1} H^{1,q}(E(C), d) = 0 \quad q > 1, \quad (4)$$

$$\mathrm{gr}_F^{2q} H^{1,q}(E(C), d) = 0 \quad q > 2, \quad (5)$$

$$\mathrm{gr}_F^{2q+2} H^{2,q}(E(C), d)_2 = 0 \quad q > 3. \quad (6)$$

Proof. Equation (3) is proven in [AAB14, Proposition 1.2].

For eq. (4) we proceed by induction, the base case follows from the entry $(1, 2)$ of Table 9. Let G_\bullet the filtration of the complex $D_r = \bigoplus_q \mathrm{gr}_F^r E^{r-2q,q}(C)$ for fixed r defined by $G_0 = 0$, $G_2 = D_r$ and

$$G_1^q = \langle x_1 \alpha, y_1 \alpha \mid \alpha \in E^{r-1-2q,q}(C, \{2, \dots, r\}) \rangle.$$

The complex G_1 is isomorphic to two copies of $\mathrm{gr}_F^{r-1} E^{r-1-2q,q}(C)$ and the quotient complex G_2/G_1 is identified with $2q$ copies of $\mathrm{gr}_F^{r-2} E^{r-2q,q-1}(C)$ (one for each G_{1i}). For $r = 2q+1$ we have:

$$\dim \mathrm{gr}_F^{2q+1} H^{1,q}(E(C), d) \leq \dim H^q(G_1) + \dim H^q(G_2/G_1)$$

$$= 2 \dim \operatorname{gr}_F^{2q} H^{0,q}(E(C), d) + 2q \dim \operatorname{gr}_F^{2q-1} H^{1,q-1}(E(C), d).$$

The first addendum is zero by eq. (3) and the second one by inductive step.

For eq. (5) we proceed by induction, the base case follows from the entry (1, 3) of Table 10. Consider the filtration G'_\bullet of $D'_r = \oplus_q \operatorname{gr}_F^r E^{r-2q+1,q}(C)$ defined by $G'_0 = 0$, $G'_3 = D'_r$,

$$G'_1 = \langle x_1 \alpha, y_1 \alpha, x_1 y_1 \beta \mid \alpha, \beta \text{ w/o index } 1 \rangle,$$

$$G'_2 = G'_1 + \langle G_{1,i} \alpha, G_{1,i} x_1 \beta, G_{1,i} y_1 \beta, G_{1,i} x_1 y_1 \gamma \mid \alpha, \beta, \gamma \text{ w/o indices } 1, i \rangle_{i=2,\dots,r}.$$

For $r = 2q$, we have

$$H^q(G'_1) \cong \operatorname{gr}_F^{2q-1} H^{0,q}(E(C)) \oplus y_1 \operatorname{gr}_F^{2q-1} H^{0,q}(E(C)) = 0 \quad (7)$$

by eq. (3). Similarly, $H^q(G'_2/G'_1)$ is equal to

$$\operatorname{gr}_F^{2q-2} H^{1,q-1}(E(C))^{\oplus 2q-1} \oplus \operatorname{gr}_F^{2q-2} H^{0,q-1}(E(C))^{\oplus 4q-2} = 0 \quad (8)$$

by inductive step and by eq. (3). The top cohomology vanishes:

$$H^q(G'_3/G'_2) \cong \bigoplus_{1 < i < j} G_{1,i} G_{1,j} \operatorname{gr}_F^{2q-3} H^{1,q-2}(E(C)) = 0 \quad (9)$$

because of eq. (4). Putting together eq. (7), (8) and (9), we obtain the claimed equality $\operatorname{gr}_F^{2q} H^{1,q}(E(C), d) = 0$.

For eq. (6) we proceed by induction, the base case $\operatorname{gr}_F^{10} H^{2,4}(E(C), d)_2 = 0$ is computed with the Python code available at https://www.dm.unibo.it/~roberto.pagaria/Top_graded_cohom_order_config_elliptic_curve.py. The computation involves only the 2-weight space for $T \subset SL_2(\mathbb{Q})$, i.e. the subspace of the graded module $\operatorname{gr}_F^r E^{r-2q,q}(C, r)$ generated by the monomials with 2 more occurrences of x than y . Consider the filtration G''_\bullet defined as the filtration G_\bullet restricted to the subspace of weight 2 for the torus action. We need to prove for $r = 2q + 2$ that $H^q(G''_1) = 0$ and $H^q(G''_2/G''_1) = 0$. The first equality follows from $H^q(G''_1) = \operatorname{gr}_F^{2q+1} H^{1,q}(E(C), d)_1 = 0$ by eq. (4). The complex G''_2/G''_1 is a direct sum of $2q + 1$ copies of the complex $\operatorname{gr}_F^{2q} E(C)_2$ and so $H^q(G''_2/G''_1) \cong \operatorname{gr}_F^{2q+1} H^{2,q}(E(C), d)_2^{\oplus 2q+1} = 0$ by inductive step. \square

The below Corollary follows from the previous Theorem and the numerical computation exposes in the next section.

Corollary 2.9. The Betti numbers of $\operatorname{Conf}(C, n)$ are:

$$\begin{aligned} b_0 &= 1, \\ b_1 &= 2n, \\ b_2 &= 2 \binom{n}{3} + 3 \binom{n}{2} + n, \\ b_3 &= 14 \binom{n}{4} + 8 \binom{n}{3} + 2 \binom{n}{2}, \\ b_4 &= 32 \binom{n}{6} + 74 \binom{n}{5} + 32 \binom{n}{4} + 5 \binom{n}{3}, \\ b_5 &= 63 \binom{n}{8} + 427 \binom{n}{7} + 490 \binom{n}{6} + 154 \binom{n}{5} + 18 \binom{n}{4}, \end{aligned}$$

$$b_k = c_k \binom{n}{2k-2} + o(n^{2k-2}),$$

where $c_k \geq \binom{2k-3}{k-3}$.

Proof. The case b_i for $i \leq 5$ are computed from Tables 7 to 11. Recall that the Poincaré polynomial of $\text{Conf}(C, n)$ and of $\text{Conf}(C, n)/C$ differ by a factor $(1+t)^2$ and the case $q = 0$ follows from Proposition 2.1. The case b_5 need the vanishing results of Lemma 2.8 and eq. (11).

For general k , we have

$$b_k(n) = \sum_{p+q=k} \sum_{i=p+q-1}^{p+2q} \binom{n}{i} \dim H^{p,q} \left(E(C, i) /_{F_{i-1}} E(C, i) \right).$$

Eq. (3) and (4) ensure that the polynomial has degree at most $2k-2$. Eq. (5) and (6) implies that

$$b_k(n) = \binom{n}{2k-2} \dim H^{2,k-1} \left(E(C, 2k-2) /_{F_{2k-3}} E(C, 2k-2) \right)_0 + o(n^{2k-2}).$$

Finally, Corollary 2.7 implies the desired result. \square

Conjecture 2.10. We claim that

$$b_k = \binom{2k-3}{k-3} \binom{n}{2k-2} + o(n^{2k-2}).$$

APPENDIX A. SMALL CASES

The elliptic curve C acts on $\text{Conf}(C, n)$ by $p \cdot (p_1, \dots, p_n) = (p_1 + p, \dots, p_n + p)$ where $+$ is the group operation on the elliptic curve C . This action is compatible with the structure of F-module and the fibration

$$C \rightarrow \text{Conf}(C, n) \rightarrow \text{Conf}(C, n)/C$$

has a non-canonical section $s: \text{Conf}(C, n)/C \rightarrow \text{Conf}(C, n)$. This induces an isomorphism

$$H^\bullet(\text{Conf}(C, n)) \cong H^\bullet(C) \otimes H^\bullet(\text{Conf}(C, n)/C) \quad (10)$$

as rings, but not as F-modules because the section cannot be chosen in an equivariant way. We used a Python3 to compute the cohomology of $\text{Conf}(C, n)/C$ for $n \leq 7$. The code is available at https://www.dm.unibo.it/~roberto.pagarria/Cohom_order_config_elliptic_curve.py. and the bigraded Betti numbers are presented in Tables 1 to 6.

Since $\text{gr}_F^r H^{p,q}(\text{Conf}(C))$ is semisimple for $q > 0$, we have a decomposition analogous to eq. (10):

$$\text{gr}_F^\bullet H^{\bullet,q}(\text{Conf}(C, n)) \cong \text{gr}_F^\bullet H^\bullet(C) \otimes \text{gr}_F^\bullet H^{\bullet,q}(\text{Conf}(C, n)/C)$$

Tables 7 to 11 report the numbers $a_i^{p,q}$ for $q > 0$ associated to the F-module $\text{gr}_F^\bullet H^{p,q}(\text{Conf}(C, n)/C)$. These entries are computed from the corresponding ones of Tables 1 to 6 as convolution with binomial coefficients.

As an example we consider $\dim H^{2,2}(\text{Conf}(C, n)/C)$, the values in tables Tables 1 to 6 are 0, 0, 0, 38, 260, 1022 and coincides with the evaluation of the polynomial

$$38 \binom{n}{5} + 32 \binom{n}{6}$$

0
1 2

TABLE 1. The dimension of the cohomology $H^{p,q}(\text{Conf}(C, 2)/C)$.

0
0 2
1 4 3

TABLE 2. The dimension of the cohomology $H^{p,q}(\text{Conf}(C, 3)/C)$.

0
0 4
0 8 10
1 6 9 4

TABLE 3. The dimension of the cohomology $H^{p,q}(\text{Conf}(C, 4)/C)$.

0
0 12
0 20 38
0 20 50 24
1 8 18 16 5

TABLE 4. The dimension of the cohomology $H^{p,q}(\text{Conf}(C, 5)/C)$.

at $n = 2, 3, 4, 5, 6, 7$. The coefficient of this polynomial are reported in the corresponding entries (2, 2) of Tables 7 to 11. (Notice that in those tables the 0-th row is omitted.)

The values of Tables 1 and 2 corresponding to the cases $n = 2, 3$ were previously computed in [Aza15].

Moreover, using the same code we have:

$$H^{2,3}(\text{Conf}(C, 8)/C; \mathbb{Q}) = 7063, \quad (11)$$

and so $a_8^{2,3} = 63$.

Acknowledgement. I would thank Gian Marco Pezzoli for the useful discussions and John Wiltshire-Gordon for notifying me the reference [Ryb18].

REFERENCES

- [AAB14] Samia Ashraf, Haniya Azam, and Barbu Berceanu, *Representation theory for the Križ model*, Algebr. Geom. Topol. **14** (2014), no. 1, 57–90. MR 3158753
- [Aza15] Haniya Azam, *Cohomology groups of configuration spaces of Riemann surfaces*, Bull. Math. Soc. Sci. Math. Roumanie (N.S.) **58(106)** (2015), no. 1, 33–47.
- [CF13] Thomas Church and Benson Farb, *Representation theory and homological stability*, Adv. Math. **245** (2013), 250–314. MR 3084430
- [Chu12] Thomas Church, *Homological stability for configuration spaces of manifolds*, Invent. Math. **188** (2012), no. 2, 465–504. MR 2909770
- [EW15] Jordan S. Ellenberg and John D. Wiltshire-Gordon, *Algebraic structures on cohomology of configuration spaces of manifolds with flows*, arXiv e-prints (2015), arXiv:1508.02430.

0
0 48
0 72 176
0 60 260 152
0 40 150 144 50
1 10 30 40 25 6

 TABLE 5. The dimension of the cohomology $H^{p,q}(\text{Conf}(C, 6)/C)$.

0
0 240
0 336 976
0 252 1491 1040
0 140 1022 1232 425
0 70 350 504 350 90
1 12 45 80 75 36 7

 TABLE 6. The dimension of the cohomology $H^{p,q}(\text{Conf}(C, 7)/C)$.

0
0 2

 TABLE 7. The coefficients $a_3^{p,q}$ of the F-module $H^{p,q}(\text{Conf}(C)/C)$ ($q > 0$).

0
0 4
0 0 10

 TABLE 8. The coefficients $a_4^{p,q}$ of the F-module $H^{p,q}(\text{Conf}(C)/C)$ ($q > 0$).

0
0 12
0 0 38
0 0 0 24

 TABLE 9. The coefficients $a_5^{p,q}$ of the F-module $H^{p,q}(\text{Conf}(C)/C)$ ($q > 0$).

0
0 48
0 0 176
0 0 32 152
0 0 0 0 50

 TABLE 10. The coefficients $a_6^{p,q}$ of the F-module $H^{p,q}(\text{Conf}(C)/C)$ ($q > 0$).

0					
0	240				
0	0	976			
0	0	259	1040		
0	0	0	168	425	
0	0	0	0	0	90

TABLE 11. The coefficients $a_7^{p,q}$ of the F-module $H^{p,q}(\text{Conf}(C)/C)$ ($q > 0$).

- [Ful97] William Fulton, *Young tableaux*, London Mathematical Society Student Texts, vol. 35, Cambridge University Press, Cambridge, 1997, With applications to representation theory and geometry. MR 1464693
- [Kri94] Igor Križ, *On the rational homotopy type of configuration spaces*, Ann. of Math. (2) **139** (1994), no. 2, 227–237. MR 1274092
- [Mac15] I. G. Macdonald, *Symmetric functions and Hall polynomials*, second ed., Oxford Classic Texts in the Physical Sciences, The Clarendon Press, Oxford University Press, New York, 2015, With contribution by A. V. Zelevinsky and a foreword by Richard Stanley, Reprint of the 2008 paperback edition [MR1354144]. MR 3443860
- [Pag18] Roberto Pagaria, *The connection between combinatorics and cohomology of elliptic arrangements*, arXiv e-prints (2018), arXiv:1805.04906.
- [Pag19] ———, *The cohomology rings of the unordered configuration spaces of the torus*, arXiv e-prints (2019), arXiv:1901.01171.
- [Ryb18] Christopher Ryba, *Resolving Irreducible $\mathbb{C}S_n$ -Modules by Modules Restricted from $GL_n(\mathbb{C})$* , arXiv e-prints (2018), arXiv:1812.07212.
- [SS17] Steven V. Sam and Andrew Snowden, *Gröbner methods for representations of combinatorial categories*, J. Amer. Math. Soc. **30** (2017), no. 1, 159–203. MR 3556290
- [Sta99] Richard P. Stanley, *Enumerative combinatorics. Vol. 2*, Cambridge Studies in Advanced Mathematics, vol. 62, Cambridge University Press, Cambridge, 1999, With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin. MR 1676282
- [Tot96] Burt Totaro, *Configuration spaces of algebraic varieties*, Topology **35** (1996), no. 4, 1057–1067. MR 1404924
- [Wey03] Jerzy Weyman, *Cohomology of vector bundles and syzygies*, Cambridge Tracts in Mathematics, vol. 149, Cambridge University Press, Cambridge, 2003. MR 1988690
- [Wil14] John D. Wiltshire-Gordon, *Uniformly Presented Vector Spaces*, arXiv e-prints (2014), arXiv:1406.0786.

ROBERTO PAGARIA

Università di Bologna, Dipartimento di Matematica, PIAZZA DI PORTA SAN DONATO 5 - 40126 BOLOGNA, ITALY.

Email address: roberto.pagaria@unibo.it