

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Safety and technological issues of dry fermented sausages produced without nitrate and nitrite

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Safety and technological issues of dry fermented sausages produced without nitrate and nitrite / Tabanelli G.; Barbieri F.; Soglia F.; Magnani R.; Gardini G.; Petracci M.; Gardini F.; Montanari C.. - In: FOOD RESEARCH INTERNATIONAL. - ISSN 0963-9969. - STAMPA. - 160:October 2022(2022), pp. 111685.1-111685.10. [10.1016/j.foodres.2022.111685]

*Availability:*

This version is available at: <https://hdl.handle.net/11585/893044> since: 2022-08-25

*Published:*

DOI: <http://doi.org/10.1016/j.foodres.2022.111685>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

# Food Research International

## Safety and technological issues of dry fermented sausages produced without nitrate and nitrite

<b>Manuscript Number:</b>	FOODRES-D-22-01442R4
<b>Article Type:</b>	Research Paper
<b>Keywords:</b>	fermented sausages, nitrate, nitrite, microbial safety, starter cultures, clean label
<b>Corresponding Author:</b>	Chiara Montanari Universita di Bologna Cesena, ITALY
<b>First Author:</b>	Giulia Tabanelli
<b>Order of Authors:</b>	Giulia Tabanelli
	Federica Barbieri
	Francesca Soglia
	Rudy Magnani
	Gabriele Gardini
	Massimiliano Petracci
	Fausto Gardini
	Chiara Montanari
<b>Abstract:</b>	<p>The aim of this work was to investigate the possibility to industrially produce fermented sausages without the addition of nitrate and nitrite. Indeed, despite their antimicrobial effect and multiple technological roles, an increasing pressure for their removal has recently raised. To achieve this goal while maintaining an acceptable final product quality, we deeply modified the whole process, that was carried out at 10-15°C (i.e., temperatures lower than traditional Mediterranean products) and by using bioprotective starter cultures at high concentrations (7 log CFU/g) to lead the fermentation. Different glucose amounts (0.2 or 0.4 % w/w) were also tested to optimize the process. The results showed no significant differences between the control (with nitrate/nitrite) and the sausages without preservatives in terms of <math>a_w</math> (value range 0.908-0.914), weight loss (about 38% in all samples), lactic acid bacteria (value range 8.1-8.3 log CFU/g) and coagulase negative cocci (value range 6.8-7.1 log CFU/g). The amount of sugar affected the final characteristics of sausages. Indeed, in the absence of curing salts, lower sugar concentration resulted in better textural features (reduced hardness and gumminess) and lower oxidation (TBARS values 0.80 vs. 1.10 mg MDA/kg of meat product in samples with 0.2% or 0.4% of glucose, respectively). Finally, challenge tests evidenced the inability of selected strains of <i>Listeria innocua</i>, <i>Salmonella enterica</i> sub. <i>enterica</i> and <i>Clostridium botulinum</i> to grow, under the adopted conditions, in fermented sausages. This research highlighted that nitrate/nitrite removal from these meat products requires accurate technological changes to guarantee the final quality.</p>

This is the postprint of: *Safety and technological issues of dry fermented sausages produced without nitrate and nitrite* by Giulia Tabanelli, Federica Barbieri, Massimiliano Petracci, Francesca Soglia, Fausto Gardini, Rudy Magnani, Chiara Montanari, which has been published in final form in *Food Research International* 160 (2022), n. 111685. The final published version is available online at: <https://doi.org/10.1016/j.foodres.2022.111685>.

© 2022 Elsevier. This manuscript version is made available under the Creative Commons Attribution–NonCommercial–NoDerivs (CC BY–NC–ND) 4.0 International License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

## HIGHLIGHTS

- Nitrate/nitrite elimination requires technological changes to produce high quality salamis
- Ripening temperature was lowered for guaranteeing microbiological safety
- Higher concentrations of bioprotective starter cultures were adopted
- Low sugar concentration allowed to obtain better textural features and lower oxidation level
- Salamis were challenged with *Listeria innocua*, *Salmonella enterica* and *Clostridium botulinum*

**Safety and technological issues of dry fermented sausages produced without nitrate and nitrite**

2

3

4

**Giulia Tabanelli<sup>a</sup>, Federica Barbieri<sup>a</sup>, Francesca Soglia<sup>a</sup>, Rudy Magnani<sup>b</sup>, Gabriele Gardini<sup>b</sup>, Massimiliano  
Petracci<sup>a</sup>, Fausto Gardini<sup>a</sup>, Chiara Montanari<sup>a\*</sup>**

7

8

9

<sup>a</sup> Department of Agricultural and Food Sciences, University of Bologna, Bologna, Italy

<sup>b</sup> C.I.a.i. Soc. Coop., Via Gambellara 62/A, 40026 Imola (BO), Italy

12

13

14

15

16

\* Corresponding author:

Chiara Montanari

Centro Interdipartimentale di Ricerca Industriale Agroalimentare,

Università degli Studi di Bologna, Sede di Cesena,

Via Quinto Bucci 336, 47521 Cesena, Italy

Tel. +39 0547338145, Fax +39 0547338103

E-mail: [chiara.montanari8@unibo.it](mailto:chiara.montanari8@unibo.it)

24

## ABSTRACT

The aim of this work was to investigate the possibility to industrially produce fermented sausages without the addition of nitrate and nitrite. Indeed, despite their antimicrobial effect and multiple technological roles, an increasing pressure for their removal has recently raised. To achieve this goal while maintaining an acceptable final product quality, we deeply modified the whole process, that was carried out at 10-15°C (*i.e.*, temperatures lower than traditional Mediterranean products) and by using bioprotective starter cultures at high concentrations (7 log CFU/g) to lead the fermentation. Different glucose amounts (0.2 or 0.4 % w/w) were also tested to optimize the process. The results showed no significant differences between the control (with nitrate/nitrite) and the sausages without preservatives in terms of  $a_w$  (value range 0.908-0.914), weight loss (about 38% in all samples), lactic acid bacteria (value range 8.1-8.3 log CFU/g) and coagulase negative cocci (value range 6.8-7.1 log CFU/g). The amount of sugar affected the final characteristics of sausages. Indeed, in the absence of curing salts, lower sugar concentration resulted in better textural features (reduced hardness and gumminess) and lower oxidation (TBARS values 0.80 vs. 1.10 mg MDA/kg of meat product in samples with 0.2% or 0.4% of glucose, respectively). Finally, challenge tests evidenced the inability of selected strains of *Listeria innocua*, *Salmonella enterica* sub. *enterica* and *Clostridium botulinum* to grow, under the adopted conditions, in fermented sausages. This research highlighted that nitrate/nitrite removal from these meat products requires accurate technological changes to guarantee the final quality.

**Key words:** fermented sausages, nitrate, nitrite, microbial safety, starter cultures, clean label

## 1. INTRODUCTION

Meat curing consists in the addition of sodium chloride (NaCl) and nitrite/nitrate to meat to assure safety and prolong the shelf-life of foods. It is one of the most ancient preservation strategies and it has been adopted by humanity for long time (Toldrá & Hui, 2014). In the past centuries, nitrates and nitrites were added as impurities of the salt, which was used in considerably higher amounts than today. It is only starting from end of XIX century that the role of these salts in preserving foods has been understood. While NaCl can reduce water availability (*i.e.*  $a_w$ ), nitrate and nitrite are responsible for several effects that today are well clarified and studied (Pegg & Honikel, 2014). In addition, the progressive purification of produced NaCl, makes the addition of nitrates and nitrites a specific step of cured meat manufactory process.

The addition of nitrate and nitrite in meats is explained by at least four fundamental reasons: i) antimicrobial effect exerted through the inhibition of metabolic pathways, limitation of oxygen uptake or iron sequestering; traditionally the effects of these nitrogen compounds are mainly addressed to suppress the outgrowth of clostridia spores, but its effectiveness against enterobacteria, *Staphylococcus aureus*, *Listeria monocytogenes* and *Bacillus cereus* is well documented (Alahakoon, Jayasena, Ramachandra, & Jo, 2015); ii) colour formation, which depends on the formation of nitric oxide able to react with many substrates, including  $Fe^{++}$  of myoglobin and  $Fe^{+++}$  of metmyoglobin, leading to the development of the typical cured meat colour (Honikel, 2004); iii) antioxidant effect, able to retard oxidative reactions and the development of rancidity due to their rapid reaction with oxygen (Ford & Lorkovic, 2002); iv) formation of cured flavour, depending on the formation of nitrite-related flavours and aromas enhanced by the suppression of rancidity (Sindelar & Milkowski, 2011).

Despite the functions exerted and their strictly controlled use in meat (Directive 2006/52/EC), nitrate and nitrite addition in food raised many concerns related to their potential of forming carcinogenic N-nitroso compounds (nitrosamine), which can be produced both in food matrices and human body (Hammes, 2012; Bernardo, Patarata, Lorenzo, & Fraqueza, 2021). This possibility was firstly suggested by Lijinsky and Epstein (1970) and confirmed by Tricker and Preusmann (1987). The reaction is potentiated by severe heat treatments and the effective health risks associated with the consumption of not thermally treated fermented sausages

has not been well elucidated (Hammes, 2012). The debate which followed these alarms showed that cured meats, and particularly fermented sausages, are nitrate sources of minor importance if compared with other potential sources, such as vegetables or water (Sindelar & Milkowski, 2011).

Nevertheless, there are increasing amounts of consumers which require fermented meats with artisanal feel (Leroy, Geyzen, Janssens, De Vuyst, & Scholliers, 2013), in which nitrate and nitrite, independently of their historical presence in sausages and their low concentration, are perceived as extraneous and, consequently, dangerous. Fermented sausage industry has the need to comply with these consumer demands requiring the elimination of nitrate and nitrite. Given the crucial role exerted by these molecules in fermented sausages, their elimination must be carefully evaluated to avoid quality decreases and maintain microbiological safety. A first strategy consists in the change of curing salts with other ingredients able to replace their activities. The use of plant extracts (such as celery, spinach, or Swiss chard) naturally rich in nitrate are a pseudo-solution (Leroy et al., 2013; Bernardo, Patarata, Lorenzo, & Fraqueza, 2021), adopted to obtain a clean label. However, European Commission recently specified that these extracts, when used with technological function, have to be considered and declared as food additives and not flavouring agents (European Commission, 2018). Other strategies are based on the use of other plant derivatives, such as essential oils or other fruit and vegetable derivatives (Ozaki et al., 2021) which, in many cases, can only partially replace nitrate and nitrite in their antimicrobial and antioxidant functions (Alahakoon et al., 2015). Also, the use of organic acids (lactate, citrate, sorbate, etc.) has been considered as an alternative to these compounds. Nevertheless, independently of their effectiveness, the use of acids, mainly due to their effect on pH, is often not compatible with the characteristics required for traditional fermented sausage, especially those produced in Mediterranean area (Van Reckem et al., 2019).

The aim of this work was to exploit the possibility to produce fermented sausages without the addition of nitrate and nitrite. The main risks of their elimination consist in the possible growth of undesired microorganisms such as *Clostridium botulinum*, *Salmonella* and *Listeria monocytogenes* (Bernardo et al., 2021; Patarata, Novais, Fraqueza, & Silva, 2020) and in the formation of undesirable colour of the meat, together

96 with a diverse sensory profile (Fraqueza, Laranjo, Elias, & Patarata, 2021). To obtains this goal, efforts was  
97 posed on the modification of process parameters adopted for the production of typical Italian salamis. In  
98 particular, the factors considered were the amounts of starter cultures and glucose added and, especially, the  
99 condition of fermentation and ripening, *i.e.* temperature and relative humidity (RH). The characteristics of the  
100 product obtained were compared with fermented sausages produced with nitrate and nitrite by analysing  
101 microbial counts, physico-chemical values, oxidation level, aroma profile, colour and texture. Finally, a  
102 challenge test was carried out to test the safety of the productive process.

103

## 104 **2. MATERIALS AND METHODS**

### 105 **2.1 Sausage manufacture**

106 The dry fermented sausages were produced in C.I.a.i. Soc. Coop. (Imola, Italy) with fresh pork meat used within  
107 48 h from slaughtering and refrigerated at 0°C. Lean meat and fat (shoulder and neck, respectively) were  
108 minced (3.5 mm) and mixed (ratio 3:1) at approx. 0°C, added with NaCl (2.5% w/w), spices (garlic powder  
109 0.01% and black pepper powder 0.15% w/w) and a commercial starter culture containing *Latilactobacillus*  
110 *sakei*, *Pediococcus acidilactici*, *Staphylococcus carnosus* and *Staphylococcus xylosus* (Chr. Hansen, Parma, Italy),  
111 at an initial concentration of approx. 7 log CFU/g for each species. Then the meat batter (300 kg) was divided in  
112 nine batches (approx. 30 kg each one): three (representing the control group, C) were added with KNO<sub>3</sub> (150  
113 mg/kg) and NaNO<sub>2</sub> (50 mg/kg) and glucose 0.2% (w/w), three batches were prepared without preservatives  
114 and adding 0.2% glucose (F02) whereas the last three batches were prepared without preservatives and adding  
115 0.4% glucose (F04). The meat mixtures were stuffed in a natural reconstituted hog casing (Varani, Castelpiano,  
116 Italy) with a diameter of 50 mm to obtain an initial weight of about 500 g. A spore suspension of *Penicillium*  
117 *nalgiovense* (Kerry Ingredient, Ireland) was sprayed on the casings. The fermentation and ripening process was  
118 carried out at temperature ranging from 15 to 10°C, at relative humidity (UR) from 65 to 90% for 50 days.

119

### 120 **2.2 pH, a<sub>w</sub> and weight loss**



121 The analyses of pH and  $a_w$  of the fermented sausages during fermentation and ripening (0, 2, 6, 13, 24, 31 and  
122 50 days from the production) were performed in triplicate by using a pH-meter Basic 20 (Crison Instruments,  
123 Barcelona, Spain) and an Aqualab CX3-TE (Labo-Scientifica, Parma, Italy), respectively. At each sampling time,  
124 sausages were also weighed to calculate the mean weight loss (%) with respect to the initial one. The analyses  
125 were performed in triplicate (three different sausages) and the results were expressed as mean value.

126

### 127 **2.3 Microbial counts**

128 The minced meat used to prepare sausages, the meat batter at time 0 and fermented sausages at the end of  
129 ripening (50 days) were analysed to determine microbial counts. After aseptic removal of the casing, a slice of  
130 approx. 10 g of sausage was transferred into a stomacher bag, mixed with 90 ml of 0.9% (w/v) NaCl sterile  
131 solution and homogenized in a Lab Blender Stomacher (Seward Medical, London, UK) for 2 min. Decimal  
132 dilutions were prepared in physiological solution and plated onto selective media to detect specific microbial  
133 groups. Lactic acid bacteria (LAB) were enumerated on de Man-Rogosa-Sharpe (MRS) agar incubated at 30°C  
134 for 48 h in anaerobic conditions. **Coagulase negative cocci (CNC)** were counted on Mannitol Salt Agar (MSA)  
135 incubated at 30°C for 72 h. Enterococci and *Enterobacteriaceae* were enumerated on Slanetz and Bartley  
136 medium and Violet Red Bile Glucose agar incubated for 24 h at 42°C and 37°C, respectively. **Pseudomonads**  
137 **were counted on Pseudomonas Agar Base, supplemented with CFC Supplement and incubated for 48 h at 30°C.**  
138 Sabouraud Dextrose Agar added with 0.2 g/l of chloramphenicol was used to determine yeasts by incubating  
139 plates at 28°C for 72 h. All media were provided by Oxoid (Basingstoke, UK). The analyses were performed in  
140 triplicate (three different sausages) and the results were expressed as mean value.

141

### 142 **2.4 Colour and Texture Profile Analysis**

143 Colour was analysed at the end of ripening (50 days). Colour [lightness ( $L^*$ ), redness ( $a^*$ ), and yellowness ( $b^*$ )]  
144 was measured in five replicates on slices having a height of approximately 1 cm by a reflectance colorimeter

145 (Minolta Chroma Meter CR-400, Minolta Italia S.p.A., Milan, Italy), previously calibrated with a standard white  
146 ceramic tile, using illuminant source C.

147 Texture Profile Analysis (TPA) was assessed at 22°C using a TA-Hdi® texture analyser (StableMicro Systems, UK)  
148 equipped with a 25 kg loading cell. The test was performed on a 1 cm-high and 1.5 cm-wide cylindrical-shaped  
149 sample compressed up to 40% of its initial height by using a 5 cm-diameter aluminium probe. A time of 20 sec  
150 was set to elapse between two compression cycles. Force-time deformation curves were obtained and  
151 Hardness (kg), Springiness, Cohesiveness, Chewiness (kg), and Gumminess (kg) were calculated according to  
152 Bourne (1978). The analyses were performed in triplicate (three different sausages) and the results were  
153 expressed as mean value.

154

## 155 **2.5 Thiobarbituric Acid Reactive Substances (TBARS)**

156 TBARS were measured at the end of ripening according to the procedure described by Bao and Ertbjerg (2015)  
157 with slight modifications. Briefly, a 5 g sample was homogenized by Ultra-Turrax (IKA, Labortechnik, Staufen,  
158 Germany) in 15 ml trichloroacetic acid (5%, w/v) and 0.5 ml butylated hydroxytoluene (4.2% in ethanol, w/v) in  
159 ice. Then, the homogenate was filtered (Whatman 1, GE Healthcare), and the reaction prepared by boiling in  
160 water bath (100°C) for 40 min, 2 ml of filtrate mixed with 2 ml thiobarbituric acid (0.02 M). After cooling the  
161 samples, absorbance was read at 532 nm and TBARS content, expressed as mg malondialdehyde/kg of meat  
162 product, calculated from a standard curve prepared with 1,1,3,3-tetraethoxypropane. The analyses were  
163 performed in triplicate (three different sausages) and the results were expressed as mean value.

164

## 165 **2.6 Aroma profile analysis**

166 The volatile profile of sausages was analysed after 50 days (end of ripening). Gas-chromatography-mass  
167 spectrometry coupled with solid phase microextraction (GC-MS-SPME) technique was used to assess the aroma  
168 profile of the sausages at the end of ripening. 3 g of samples were added with known amount of 4 methyl-2-  
169 pentanol (Sigma-Aldrich, Steinheim, Germany) as internal standard and analysed according to the protocol

170 reported by Montanari et al. (2016). Volatile peak identification was carried out using Agilent Hewlett–Packard  
171 NIST 2011 mass spectral library (Gaithersburg, MD, United States) (NIST, 2011). The mass spectrum  
172 identification was confirmed in the same conditions by injection of the pure standards (Sigma- Aldrich, St.  
173 Louis, MO). Data are expressed as ratio between each molecule peak area and the peak area of internal  
174 standard. The analyses were performed in triplicate (three different sausages) and the results were expressed  
175 as mean value.

176

## 177 **2.7 Challenge test**

178 To assess the safety of fermented sausages produced without preservatives, a challenge test was performed in  
179 the sausages added with 0.2% of glucose (F02) and the control with nitrate/nitrite (C). The sausages containing  
180 0.4% glucose were not considered in the challenge test due to the results of the previous trials, in particular  
181 higher oxidation level and hardness of the sausages. The target microorganisms were *Listeria innocua* as  
182 surrogate of *L. monocytogenes* (da Silva, de Oliveira Pena, Pflanzner, & da Silva do Nascimento, 2019),  
183 *Salmonella enterica* subspecies *enterica* and *Clostridium botulinum*. The three microbial groups considered  
184 were separately inoculated in the different batches.

185 For *L. innocua*, a cocktail of five strains of different origin was used (four isolated from sausage production  
186 environment and a collection strain – ATCC33090). Strains were precultured twice in Brain Heart Infusion (BHI,  
187 Oxoid, Basingstoke, UK) at 30°C for 24 h (first step) and at 12°C for 72 h (second step), then mixed and this  
188 suspension was used to inoculate the meat batter (10 kg, 20 sausages) at a concentration of approx. 7 log  
189 CFU/g. As far as *Salmonella*, a mixture of 3 strains (*Salmonella enterica* subspecies *enterica* serovar  
190 Typhimurium ATCC14028, *Salmonella enterica* subspecies *enterica* serovar Typhimurium monophasic variant  
191 isolated from pork sausage and *S. enterica* subspecies *enterica* serovar Derby isolated from pork meat) was  
192 used, after two 24 h precultures in BHI at 30°C for 24 h. The precultures were added to reach a final  
193 concentration in the meat batter (10 kg, 20 sausages) of approx. 7 log CFU/g. Finally, for *Cl. botulinum* a  
194 cocktail of four strains was used and namely two non-proteolytic (type B and type E) and two proteolytic (type

195 A and type B), including a collection strain – ATCC 19397. In this case, strains were cultivated in TPGYT medium  
196 (Trypticase, Peptone, Glucose, Yeast-extract, Trypsin) at 30°C for 15 days under anaerobiosis to allow  
197 sporification. Then cell suspensions were thermally treated (80°C for 20 min), and spores were enumerated to  
198 be inoculated in the meat batter (10 kg, 20 sausages) at a concentration of approx. 3 log CFU/g.  
199 The different meat batters were then stuffed and processed in the same conditions reported in section 1.1 and  
200 analysed during fermentation and ripening (0, 2, 6, 13, 31 and 50 days) to monitor the behaviour of these  
201 microbial species in the products. The enumeration or the detection of *Listeria* were performed according to  
202 the methods EN ISO 11290-2 and EN ISO 11290-1 (ISO, 2017a, b). *S. enterica* subsp. *enterica* was counted on  
203 Hektoen Enteric Agar (Oxoid, Basingstoke, UK) incubated at 37°C for 24 h, while the detection was performed  
204 according to ISO 6579-1 (ISO, 2017c). Finally, *Cl. botulinum* was counted following the procedure described in  
205 ISO 15213 (ISO, 2013).

206

## 207 **2.8 Statistical analysis**

208 Data were analysed through One-way ANOVA considering the absence of preservatives together with the  
209 eventual addition of a higher glucose content in the formulation as the main effect. Means were subsequently  
210 analysed through the parametric Tukey-HSD test.

211 To further investigate the effect of the absence of nitrite and nitrate on the oxidation, colour and texture  
212 profile, planned orthogonal contrasts were performed to compare the findings obtained within the control  
213 group (C) with those found in F02 and F04 samples. All statistical differences were considered significant at a  
214 level of  $p \leq 0.05$ . Analyses were carried out by using Statistica software (StatSoft Italy srl, Vigonza, Italy).

215

## 216 **3. RESULTS AND DISCUSSION**

### 217 **3.1 Process and ripening conditions and microbial counts**

218 Safety and quality of fermented sausages is the result of the application of several hurdles addressed to the  
219 inhibition of undesirable microorganisms. The removal of one of these hurdles (such as nitrate/nitrite) must be

220 counterbalanced by more drastic conditions for the other ones or the addition of new factors. In this trial, we  
221 firstly decided to lower fermentation and ripening temperatures 10°C vs. 15-20°C of the traditional program  
222 (Montanari et al., 2018; Tabanelli, Montanari, Grazia, Lanciotti, & Gardini, 2013) with the aim to reduce, in the  
223 absence of nitrate and nitrite, the growth potential of *Enterobacteriaceae* during the first days, when  $a_w$  is still  
224 high (Hospital et al., 2015). The second modification concerned the use of commercial starter cultures  
225 (containing selected strains of *Lat. sakei*, *P. acidilactici*, *Staph. carnosus* and *Staph. xylosus*) added at higher  
226 concentration (approx. 7 log CFU/g) with respect to the standard procedures adopted for Mediterranean  
227 sausages, in which the cultures are added at about 6 log CFU/g (Montanari et al., 2018; Tabanelli et al., 2013).  
228 Besides, the LAB strains used were characterized by antagonistic and bioprotective activity against *L.*  
229 *monocytogenes* (Raimondi, Popovic, Amaretti, Di Gioia, & Rossi, 2014; Stahnke, 2008), which could take  
230 advantage by the low temperature adopted during fermentation and ripening, especially in low contaminated  
231 meats (Patarata, Novais, Fraqueza, & Silva, 2020). In fact, in order to limit the growth potential of wild  
232 undesirable microbial population the sausages were manufactured using with fresh (not frozen) pork meat,  
233 maintained under strictly controlled refrigeration temperature (0°C) and used within 48 h after slaughtering.  
234 Three different typologies were considered in the trials: sausages without nitrate/nitrite produced with glucose  
235 at 0.2% (F02), sausages without nitrate/nitrite produced with glucose at 0.4% (F04) and the control, *i.e.*  
236 sausages with nitrate/nitrite produced with glucose at 0.2% (C).  
237 The results of the microbial counts (Table 1) showed that LAB and CNC amounts in the meat batter  
238 immediately after casing were highly dependent on the addition of the starter cultures, **as expected**. Indeed,  
239 the counts of both these microbial groups in the minced meat before the addition of the cultures were lower  
240 than 4 log CFU/g. At the end of ripening, the counts of LAB were higher than 8 log CFU/g in all the samples,  
241 while CNC were approx. 7 log CFU/g and no significant difference ( $p > 0.05$ ) in relation to the type of sausage  
242 was observed. These levels of LAB and CNC are comparable with those found in similar Italian and Portuguese  
243 fermented sausages at the end of ripening (Belleggia et al., 2022a, b; Tabanelli et al., 2015; Van Reckem et al.,  
244 2019). In other studies, CNC were more susceptible to the presence of variable amounts of nitrate/nitrite.

Hospital et al. (2015) reported a double contrasting effect of nitrate/nitrite on these bacteria. On one side, their presence can favour CNC growth, especially in the central part of sausages in which O<sub>2</sub> is scarce and nitrate can act as final electron acceptor, accelerating their multiplication. On the other side, excessive accumulation of nitrite can result in growth inhibition.

Enterococci were present at low concentration (2.5 log CFU/g) after casing and their final number never exceeded 2.7 log CFU/g. Interestingly, enterococci counts in the sausages added with nitrate/nitrite were significantly lower than in the sausages obtained without these preservatives.

A similar behaviour was observed for *Enterobacteriaceae*. Their initial count (2.6 log CFU/g) decreased in all the sausages, but at a higher significant extent in the control, confirming the important role exerted by nitrate/nitrite in the inhibition of these bacteria. This fact emphasizes the importance of enterobacteria control in the first steps of fermentation when a<sub>w</sub> and pH can still allow their multiplication (Christieans, Picgirard, Parafita, Lebert, & Gregori, 2018; Hospital, Hierro, & Fernández, 2014; Hospital et al., 2015). Pseudomonads were always below the detection limit.

Yeasts, initially present at 1.9 log CFU/g, grew up to more than 4 log CFU/g in all the samples without significant differences. These counts are in accordance with data previously reported by other studies (Belleggia et al., 2022b; Greppi et al., 2015; Montanari et al., 2018; Selgas & García, 2014).

261

### 262 3.2 Chemico-physical features

The results concerning pH, a<sub>w</sub> and weight loss of the sausages during ripening are reported in Figure 1. The evolution of pH in the different samples (Figure 1A) was clearly influenced by the amounts of glucose added. In fact, in the sausages containing 0.4% glucose the pH decrease observed during fermentation is more relevant (about 0.25 pH units) if compared with the control (C) and the sausages containing 0.2% glucose and without nitrate/nitrite (F02). This difference remained constant during the ripening, and the pH increased up to 5.50 in the sausages F02 and to 5.25 in the samples F04.

269 No significant differences were observed for  $a_w$  and weight loss during ripening (Figures 1B and 1C). The final  $a_w$   
270 ranged between 0.908 and 0.914 and the weight losses between 37.7 and 38.7%.

271

### 272 **3.3 Colour and oxidative stability of the lipid fraction**

273 Overall, the addition of different glucose levels in the formulation exerted some effects on both the colour  
274 parameters and the textural traits of the dried fermented sausages (Table 2). In detail, although no significant  
275 differences were found in lightness ( $L^*$ ) and yellowness ( $b^*$ ), redness ( $a^*$ ) was remarkably affected by  
276 preservatives as well as by the glucose content added in the formulation. Indeed, if compared to C, a  
277 significantly lower  $a^*$  value was found in F02 (13.51 vs. 11.58;  $p < 0.01$ ), whereas F04 exhibited an intermediate  
278 value (12.57). This finding may be primarily ascribed to the absence of nitrate and nitrite in the formulation  
279 regardless of the glucose content added. Indeed, by means of planned orthogonal contrasts performed to  
280 assess the possible effect of the absence of preservatives (regardless of the glucose content added), a  
281 significant difference ( $p < 0.05$ ) in  $a^*$  was found between C and those experimental groups in which no  
282 preservatives were added (*i.e.*, F02 and F04) (LSM: 13.51 vs. 11.58 and 12.57). Within this context, it is worth  
283 mentioning that curing colour development is strongly affected by the acidification process taking place during  
284 product's fermentation (Campbell-Platt & Cook, 1995). Also, the colour parameter Chroma ( $C^*$ ) exhibited a  
285 similar trend: if compared to C, a significantly lower value was found in F02 (14.60 vs. 12.42;  $p < 0.01$ ), whereas  
286 F04 exhibited an intermediate value (13.69). As previously observed for  $a^*$ , planned orthogonal contrasts  
287 suggest that this outcome may be due to the absence of nitrate and nitrite in dry fermented sausages  
288 formulation. If compared to C, a significantly lower ( $p < 0.05$ ) saturation index was observed, regardless of the  
289 glucose content added, in those experimental groups in which no preservatives were added (LSM:14.60 vs.  
290 12.42 and 13.69). This finding may be explained by considering that the rate at which total heme pigments are  
291 converted in their nitric oxide form sharply increases as the pH decreases, especially within the pH range from  
292 5.5 to 4.5 (Fox & Thompson, 1963).

293 As for the textural parameters, if compared to C, significantly higher Hardness, Gumminess and Chewness  
294 values were found in F02 and F04 by means of both analysis of variance and planned orthogonal contrasts.  
295 Regarding hardness, the values increased with the glucose content in the formulation in which no preservatives  
296 were added, with F04 exhibiting the highest value (3.74 kg). This outcome may be explained by considering  
297 that during fermentation, pH values close to the isoelectric point of the proteins are reached (Figure 1A), thus  
298 leading to a consequent increased protein aggregation that determines the development of firmer sausages  
299 (Fretheim, Egeland, Harbiz, & Samejima, 1985; Gonzalez-Fernandez, Santos, Rovira, & Jaime, 2006). Indeed,  
300 a strong acidification is essential to reduce the water holding capacity of the meat proteins and thus promote  
301 water loss and evaporation (Lorenzo, Gómez, & Fonseca, 2014). In addition, under a fast acidification process,  
302 acid solubilization of collagen is also likely induced (Aktas & Kaya, 2001) and may account for the trend  
303 observed for Cohesiveness, which exhibited remarkably higher values in C samples rather than in F02 and F04  
304 (1.83 vs. 1.72 and 1.73;  $p < 0.001$ ).

305 Lipid oxidation level was assessed by measuring TBARS developed at the end of the ripening process. Overall,  
306 the amount of secondary products of lipid oxidation significantly differ among the experimental groups (as  
307 shown in Table 2), due to the antioxidant effect of nitrate/nitrite. In detail, as expected, the lower TBARS were  
308 observed in the sausages added with nitrate/nitrite. These findings are in agreement with the widely known  
309 antioxidant properties of nitrate/nitrite in sausages (Honikel, 2008). Indeed, nitrite can limit the development  
310 of oxidative reactions affecting the lipid fraction in several ways: i) sequestering oxygen molecules (Honikel,  
311 2008), ii) stabilizing heme iron and sequestering free iron (Bergamaschi and Pizza, 2011), and iii) reacting with  
312 lipid radicals thus breaking the oxidative chain reaction upon solubilization of nitric oxide in fats (Skibsted,  
313 2011). On the other hand, fermented sausages formulated with the addition of 0.2% glucose exhibited a lower  
314 oxidation level of the lipid fraction if compared to those containing the higher glucose concentration (*i.e.*,  
315 0.4%). This outcome may be explained by considering the differences in pH directly associated to the amount  
316 of sugar added into products' formulation. Indeed, in light of the increased solubility of iron under acidic  
317 conditions (Chaijan & Panpipat, 2017; Domínguez et al., 2019) the lower pH resulting from the addition of a



318 higher glucose content may have promoted the development of oxidative reactions affecting the lipid fraction.  
319 In addition, it has been recently reported that reducing sugars may be involved in the oxidative deamination of  
320 the amino group residues in protein (Akagawa, Sasaki, & Suyama, 2002; Luna & Estèvez, 2018). Thus, the  
321 significantly higher lipid oxidation observed in F04 may be explained by considering the pro-oxidant potential  
322 of these compounds along with the strong interdependence existing between the oxidative modifications  
323 affecting the protein and the lipid fractions.

324

### 325 **3.4 Aroma profile**

326 In Table 3, the main volatile molecules detected in the sausages at the end of ripening through SPME-GC-MS  
327 analysis are reported. The molecules deriving from spices, added in the same amounts in the meat batter, are  
328 not included in the table. Nevertheless, the compounds detected were limonene,  $\beta$ -phellandrene,  
329 caryophyllene,  $\alpha$ -phellandrene, *o*-cymene, copaene, terpinene-4-ol, linalool (in decreasing order, deriving from  
330 black pepper) and allyl-methyl-sulfide (from garlic). The remaining molecules were grouped according to their  
331 chemical characteristics in ketones, aldehydes, alcohols, acids, and esters.

332 Relevant differences related to the amount of sugar added and the presence of nitrate/nitrite were observed  
333 among ketones. These differences mainly concerned acetone, 2,3-butanedione (diacetyl) and 3-hydroxy-2-  
334 butanone (acetoin). According to Flores (2018), the origin of these compounds can be related to fermentation  
335 processes. In particular, the metabolism of pyruvate is the starting point for the production these molecules.  
336 LAB, including *Lat. sakei*, are particularly active in using pyruvate through alternative metabolic routes,  
337 especially when fermentable sugars are scarce or depleted (von Wright & Axelsson, 2011). Besides, they can  
338 also obtain pyruvate from amino acids (Barbieri, Laghi, Gardini, Montanari, & Tabanelli, 2020; Barbieri et al.,  
339 2022). This may explain their lower accumulation in the sausages produced without nitrate/nitrite containing  
340 0.2% glucose with respect to the sausages with 0.4% of the same sugar. However, the presence of  
341 nitrate/nitrite determined the higher accumulation of diacetyl and acetoin in the control, even if at the lower  
342 sugar concentration. In effect, the reduction of nitrate/nitrite can favour the growth of coagulase positive cocci

343 (CNC) (Perea-Sanz, Montero, Belloch, & Flores, 2019) with a concomitant limitation of the metabolic potential  
344 of *Lat. sakei*.

345 Hospital et al. (2015) described an opposite trend in *chorizo*, in which both diacetyl and acetoin increased with  
346 the diminution of nitrate/nitrite and attributed this outcome to the higher concentration of staphylococci in  
347 the sausages without nitrate/nitrite added, but in this case they used a strain of *Pediococcus pentosaceus* as  
348 starter LAB. In this trial at the end of ripening, no significant difference was observed in CNC in the different  
349 sausages produced (Table 1). Nevertheless, the presence of nitrate can be an alternative source for NADH  
350 regeneration making pyruvate available for other metabolic pathways which can bring to the accumulation of  
351 these molecules (Hammes, 2012; Sánchez-Mainar & Leroy, 2015). Perea-Sanz et al. (2019) found significantly  
352 lower concentrations of diacetyl (but not of acetoin) in sausages produced with lowered nitrate addition. These  
353 discrepancies may be explained also in relation to the process adopted and, in particular, to the final pH of the  
354 sausages which, in the case object of this study, was rather high if compared with pH reached in the cited  
355 studies. The presence of methyl ketones in sausages can be the result of microbial  $\beta$ -oxidation of lipids,  
356 deriving from  $\beta$ -ketoacids produced during  $\beta$ -oxidation carried out in first instance by moulds and staphylococci  
357 (Lorenzo, Gómez, Purriños, & Fonseca, 2016; Ordóñez, Hierro, Bruna, & de la Hoz, 1999). In particular, 2-  
358 heptanone, 3-hexen-2-one and 2-pentanone concentration was higher in the samples without nitrate/nitrite,  
359 indicating a possible higher activity of staphylococci.

360 Aldehydes reached their highest concentration in the sausages without nitrate/nitrite added with 0.4% glucose,  
361 whilst in the control they were detected at the minimum level. A positive correlation between TBARS and linear  
362 aldehydes C3-C7 detected using SPME-GC-MS has been described in literature (Olivares et al., 2011). The data  
363 reported here confirmed this observation (see Table 2). In general, in fact, the quantity of aliphatic aldehydes,  
364 accumulated through the action of hydroperoxydases (Ordóñez et al., 1999) was significantly lower in the  
365 control. The amount of 3-methylbutanal, deriving from leucine and with a positive and important effect on  
366 sausages aroma profile (Flores, 2018; Carballo, 2012) did not show differences in relation to the condition  
367 considered.

368 The sausages produced with glucose at 0.4% were characterized by the highest alcohol accumulation. Ethanol  
369 was the major molecule of this chemical group. Ethanol can be produced during mixed acid LAB fermentation  
370 starting from pyruvate when fermentable sugars are scarce or completely depleted through mixed acid  
371 fermentation (von Wright & Axelsson, 2011) The higher content of ethanol in the sausages produced with 0.4  
372 glucose confirm this hypothesis. Besides, other alcohols presented a similar trend, and in particular 1-octen-3-  
373 ol, 2-octen-1-ol, heptanol and hexanol.

374 Total acids were similar in the control (containing glucose at 0.2%) and in the samples added with 0.4% of  
375 glucose without nitrate/nitrite. These values were significantly higher if compared with the sausages produced  
376 in the absence of preservatives produced adding glucose at 0.2%. Among acids, acetic acid was the most  
377 representative. It can be produced by several pathways from LAB and staphylococci starting from pyruvate  
378 provided by sugar fermentation and other metabolisms, which can involve amino acids (Barbieri et al., 2020;  
379 Gänzle, 2015; Sánchez-Mainar & Leroy, 2015). As observed for ethanol, acetic acid is one of the final products  
380 of the metabolism of mixed acid fermentation in LAB. In particular, under defined conditions, the activity of the  
381 enzyme acetate kinase allows the accumulation of a supplementary ATP from acetyl phosphate obtained from  
382 pyruvate (von Wright & Axelsson, 2011; Zotta, Parente, & Ricciardi, 2017). This could confirm a LAB reduced  
383 activity observed determined by a higher metabolic competitiveness of staphylococci. In contrast with this  
384 data, other authors (Hospital et al., 2015; Perea-Sanz, Montero, Belloch, & Flores, 2018) found an increase of  
385 this acid in the sausages without or with lower nitrate/nitrite content. The other organic acids followed a  
386 similar behaviour, showing lower amounts in the sausages produced without nitrate/nitrite and added with  
387 glucose at 0.2%. This trend characterized also 3-methyl-butanoic (isovaleric) acid, derived from leucine  
388 metabolism. This molecule is often detected in sausages, but when its concentration is too high can be  
389 responsible for the formation of severe off-odours, due to its strong sensory impact (Montanari, Barbieri,  
390 Gardini, & Tabanelli, 2021).

391

### 392 **3.5 Challenge test**

Challenge tests were performed with the aim to evaluate the response of *Salmonella enterica* subsp. *enterica*, *Listeria innocua* (used as surrogate of *Listeria monocytogenes*) and *Clostridium botulinum* under the process conditions adopted. In this part of the work, only the sausages containing 0.2% of glucose, added or not with nitrate/nitrite, have been considered due to its textural and oxidative characteristics. In fact, the oxidation level increased in F02 and F04, as expected, but at a lesser extent. The lower pH of F04 caused also a relevant increase in hardness, if compared with the control, which was not observed in F02. A cocktail of strains belonging to the three microbial groups were separately inoculated to obtain initial concentration of approx. 7 log CFU/g for *S. enterica* subsp. *enterica* and *L. innocua* and 3 log CFU/g for *Cl. botulinum*.

Figure 2A represents the counts of *L. innocua* during fermentation and ripening. After 50 days of ripening, they were 2 log units below the initial inoculum of the strains of *L. innocua*, regardless the presence of nitrate/nitrite. However, the data of the sampling after 2 and 6 days showed significant higher counts in the sausages without nitrate/nitrite. It is noteworthy the increase of the counts observed after 2 days in F02. Even after 13 days, the counts are lower in the control. Only after this period *L. innocua* concentration is similar (without significant differences) in both sausage type. Similar trends for *L. monocytogenes* were observed by Christeians et al. (2018) using reduced levels of nitrate/nitrite. Differently from the observation of these authors, the concentration of *L. innocua* at the end of ripening in the present work did not depend on the level of curing salts added, probably due to the antilisterial activity of starter cultures. Nevertheless, the initial increase of counts observed, although the strains of *L. innocua* were inoculated at a challenging concentration, underlined the need to strictly control the fermentation step.

Regarding the challenge test carried out with *S. enterica* subsp. *enterica* (serovars Typhimurium and Derby), the results reported in Figure 2B showed no significant difference in relation to the presence of nitrate/nitrite and to the sampling time. In both the trials, the ripening conditions applied allowed the decrease of the challenging concentration of *S. enterica* subsp. *enterica* of about 2.5 log units after 50 days. A previous study (Hospital et al., 2014) indicated that *Salmonella* Typhimurium can grow and improve its survival in the absence of nitrate/nitrite. This discrepancy can be explained by the low temperature (<10°C) applied in this industrial

production during fermentation and ripening. In fact, even if the minimum growth temperature for *Salmonella* is approx. 5°C (ANSES, 2011), the growth rate and survival of this pathogen are greatly reduced below 15°C (Bell & Kyriakides, 2002). On the other hand, the effects of nitrate/nitrite on *Salmonella* spp. are rather controversial, differently from other pathogens such as *Listeria* spp. and *Clostridium botulinum*, in which a significant inhibition of the growth is reported (Hospital et al., 2014). Christieans et al. (2018) found that nitrite is a relevant hurdle for *Salmonella* Thyphimurium in sausages. This bacterium has recently been associated with several outbreaks due to fermented sausages consumption (Omer et al., 2018) and its control is becoming crucial for the fermented meats. Independently on the addition of nitrate/nitrite, the results obtained in these trials at the end of ripening were more relevant than those of Mataragas et al. (2015), in which both *L. monocytogenes* and *Salmonella enterica* were reduced less than 2 log units at the end of ripening of Cacciatore and Felino type sausages. In these studies, traditional temperature profiles were adopted.

Figure 2C represents the results of the challenge test with *Cl. botulinum*. In this case, the amount inoculated was lower (about 2.7 log CFU/g). No relevant count increase was observed until the end of ripening, and, at the different sampling times, no difference was observed in relation to the addition of nitrate/nitrite. Undoubtedly, this toxin producer microorganism has been historically viewed as the main target of nitrate and nitrite in fermented sausages. However, under industrial conditions, the number of spores present in raw materials is rather low and the processes adopted are sufficient to limit its growth potential. In this case, the adoption of low temperatures likely inhibits the growth of proteolytic *Cl. botulinum* while the non-proteolytic psychrotrophic members of this species cannot develop when the  $a_w$  is lower than 0.97, which approximately corresponded to the initial mixture  $a_w$  of the sausages. This can explain lack of registered outbreaks from fermented sausages or the low risk of food poisoning from these products (Holck, Axelsson, McLeod, Rode, & Heir, 2017).

440

#### 441 4. CONCLUSIONS

442 In this research a process for producing high quality fermented sausages without the use of nitrate and nitrite  
443 or any other substitute was studied. The removal of this important hurdle to the proliferation of undesirable  
444 microorganisms was balanced by the modification of the process, that was longer and at lower temperature  
445 (10-15°C) with respect to the conditions traditionally adopted for these Mediterranean products. Moreover,  
446 high concentrations of bioprotective starter cultures were used to control the first crucial fermentation  
447 process. The absence of nitrate/nitrite did not significantly affect some chemico-physical and microbiological  
448 parameters, such as  $a_w$ , weight loss and LAB and CNC counts. On the other hand, the amount of sugar added  
449 had a relevant effect on some characteristics of the final products. Indeed, in the absence of curing salts, lower  
450 sugar concentration resulted in better textural features and lower oxidation level. These factors also affected  
451 the aroma profile, since the final product obtained without nitrate/nitrite and with 0.2% of glucose was  
452 characterized by lower amounts of ketones and acids. Among the latter, it is noteworthy the reduced  
453 accumulation of 3-methyl-butanoic acid, whose high presence in fermented sausages can be responsible for off  
454 odours. The challenge tests evidenced the inability of selected strains of *Listeria innocua*, *Salmonella enterica*  
455 sub. *enterica* and *Clostridium botulinum* to grow, under the adopted conditions, in fermented sausages.

456

457

## 458 **FUNDING**

459 The work was supported by Regione Emilia-Romagna in the framework of the project “Qualità e sicurezza di  
460 salami prodotti senza l’impiego di nitrati/nitriti”, PSR 2014-2020 – Progetti di filiera – misura 16.2.01-progetti  
461 pilota e sviluppo dell’innovazione, CUP E23E18000080007.

462

## 463 **REFERENCES**

464 Akagawa, M., Sasaki, T., & Suyama, K. (2002). Oxidative deamination of lysine residue in plasma protein of  
465 diabetic rats. Novel mechanism via the Maillard reaction. *European Journal of Biochemistry*, 269(22),  
466 5451-5458. <https://doi.org/10.1046/j.1432-1033.2002.03243.x>

467 Aktas, N., & Kaya, X. (2001). Influence of weak organic acids and salts on the denaturation characteristics of  
 468 intramuscular connective tissue. A differential scanning calorimetry study. *Meat Science*, 58, 413-419.  
 469 [https://doi.org/10.1016/s0309-1740\(01\)00044-4](https://doi.org/10.1016/s0309-1740(01)00044-4)  
 470 Alahakoon, A. U., Jayasena, D. D., Ramachandra, S., & Jo, C. (2015). Alternatives to nitrite in processed meat: up  
 471 to date. *Trends in Food Science & Technology*, 45, 37-49. <https://doi.org/10.1016/j.tifs.2015.05.008>  
 472 ANSES (2011). Data sheet on foodborne biological hazards / *Salmonella* spp.  
 473 <https://www.anses.fr/en/system/files/MIC2011sa0057FiEN.pdf>. Accessed 11 March 2022.  
 474 Bao, Y., & Ertbjerg, P. (2015). Relationship between oxygen concentration, shear force and protein oxidation in  
 475 modified atmosphere packaged pork. *Meat Science*, 110, 174-179.  
 476 <https://doi.org/10.1016/j.meatsci.2015.07.022>  
 477 Barbieri, F., Laghi, L., Gardini, F., Montanari, C., & Tabanelli, G. (2020). Metabolism of *Lactobacillus sakei* Chr82  
 478 in the presence of different amounts of fermentable sugars. *Foods*, 9, 720.  
 479 <https://doi.org/10.3390/foods9060720>  
 480 Barbieri, F., Laghi, L., Montanari, C., Lan, Q., Levante, A., Gardini, F., & Tabanelli, G. (2022). Insights into the  
 481 metabolomic diversity of *Latilactobacillus sakei*. *Foods*, 11, 477.  
 482 <https://doi.org/10.3390/foods11030477>  
 483 Bell, C., & Kyriakides, A. (2002). *Salmonella: a practical approach to the organism and its control in foods*.  
 484 Oxford: Blackwell Publishing.  
 485 Belleggia, L., Ferrocino, I., Reale, A., Naceur Haouet, M., Corvaglia, M.R., Milanović, V., Boscaino, F., Di Renzo,  
 486 T., Di Bella, S., Borghi, M., Farneti, S., Cesaro, C., Garofalo, C., Cardinali, F., Aquilanti, L., Musari, E.,  
 487 Cocolin, L., & Osimani, A. (2022a). Unravelling microbial populations and volatile organic compounds of  
 488 artisan fermented liver sausages manufactured in Central Italy. *Food Research International*, 154,  
 489 111019. <https://doi.org/10.1016/j.foodres.2022.111019>  
 490 Belleggia, L., Ferrocino, I., Reale, A., Corvaglia, M.R., Milanović, V., Cesaro, C., Boscaino, F., Di Renzo, T.,  
 491 Garofalo, C., Cardinali, F., Aquilanti, L., Cocolin, L., & Osimani, A. (2022b). Unfolding microbiota and

492 volatile organic compounds of Portuguese Painho de Porco Preto fermented sausages. *Food Research*  
 493 *International*, 155, 111063. <https://doi.org/10.1016/j.foodres.2022.111063>

494 Bergamaschi, M., & Pizza, A. (2011). Effect of pork meat pH on iron release from heme molecule during  
 495 cooking. *Journal of Life Sciences*, 5, 376-380.

496 Bernardo, P., Patarata, L., Lorenzo, J.M., & Fraqueza, M.J. (2021) Nitrate is nitrate: the status quo of using  
 497 nitrate through vegetable extracts in meat products. *Foods*, 10, 3019.  
 498 <https://doi.org/10.3390/foods10123019>

499 Bourne, M. C. (1978). Texture Profile Analysis. *Food Technology*, 32, 62-66, 72. Campbell-Platt, G., & Cook, P. E.  
 500 (1995). *Fermented meats* (1st ed.). London: Chapman & Hall.

501 Carballo, J. (2012). The role of fermentation reactions in the generation of flavor and aroma of foods. In B.M.  
 502 Mentha, A. Kama-Eldin, & R.Z. Iwanski (Eds.). *Fermentation, effects on food properties*, 1st Edition (pp  
 503 51-83). Boca Raton: CRC Press.

504 Chaijan, M., & Panpipat, W. (2017). Mechanism of oxidation in foods of animal origin. In R. Banerjee, A. K.  
 505 Verma, & M. W. Siddiqui (Eds.), *Natural Antioxidants. Applications in Foods of Animal Origin* (pp. 1-38).  
 506 Boca Raton: Apple Academic Press, Inc.

507 Christeians, S., Picgirard, L., Parafita, E., Lebert, A., & Gregori, T. (2018). Impact of reducing nitrate/nitrite levels  
 508 on the behavior of *Salmonella typhimurium* and *Listeria monocytogenes* in French dry fermented  
 509 sausages. *Meat Science*, 137, 160-167. <https://doi.org/10.1016/j.meatsci.2017.11.028>

510 da Silva, A.C.M., de Oliveira Pena, P., Pflanzner, S.B. Júnior, da Silva do Nascimento, M. (2019). Effect of different  
 511 dry aging temperatures on *Listeria innocua* as surrogate for *Listeria monocytogenes*. *Meat Science*, 157,  
 512 107884. <https://doi.org/10.1016/j.meatsci.2019.107884>

513 Directive 2006/52/EC of The European Parliament and of the Council of 5 July 2006 amending Directive  
 514 95/2/EC on food additives other than colours and sweeteners and Directive 94/35/EC on sweeteners  
 515 for use in foodstuffs. (2006). *Official Journal of the European Union* L 204/10, 26.7.2006.



516 Domínguez, R., Pateiro, M., Gagaoua, M., Barba, F. J., Zhang, W., & Lorenzo, J. M. (2019). A comprehensive  
 517 review on lipid oxidation in meat and meat products. *Antioxidants*, 8, 429.  
 518 <https://doi.org/10.3390/antiox8100429>

519 European Commission. (2018). Summary report of the standing committee on plants, animals, food and feed  
 520 held in Brussels on 17 September 2018. *Health and Food Safety Directorate General*.

521 Flores, M. (2018). Understanding the implications of current health trends on the aroma of wet and dry cured  
 522 meat products. *Meat Science*, 144, 53-61. <https://doi.org/10.1016/j.meatsci.2018.04.016>

523 Ford, P. C., & Lorkovic, I. M. (2002). Mechanistic aspects of the reactions of nitric oxide with transition-metal  
 524 complexes. *Chemical Reviews*, 102, 993-1018. <https://doi.org/10.1021/cr0000271>

525 Fox, J. B., Jr., & J. S. Thomson, J. S. (1963). Formation of bovine nitrosylmyoglobin. I. pH 4.5-6.5. *Biochemistry*, 2,  
 526 465-470. <https://doi.org/10.1021/bi00903a012>

527 Fraqueza, M.J., Laranjo, M., Elias, M., & Patarata, L. (2021). Microbiological hazards associated with salt and  
 528 nitrite reduction in cured meat products: control strategies based on antimicrobial effect of natural  
 529 ingredients and protective microbiota. *Current Opinion in Food Science*, 38, 32-39.  
 530 <https://doi.org/10.1016/j.cofs.2020.10.027>

531 Fretheim, K., Egelanddal, B., Harbiz, O., & Samejima, K. (1985). Slow lowering of pH induces gel formation of  
 532 myosin. *Food Chemistry*, 18, 169-178. [https://doi.org/10.1016/0308-8146\(85\)90044-5](https://doi.org/10.1016/0308-8146(85)90044-5)

533 Gänzle, M.G. (2015). Lactic metabolism revisited: metabolism of lactic acid bacteria in food fermentations and  
 534 food spoilage. *Current Opinion in Food Science*, 2, 106-117. <https://doi.org/10.1016/j.cofs.2015.03.001>

535 Gonzalez-Fernandez, C., Santos, E. M., Rovira, J., & Jaime, I. (2006). The effect of sugar concentration and  
 536 starter culture on instrumental and sensory textural properties of *chorizo*-Spanish dry-cured sausage.  
 537 *Meat Science*, 74, 467-475. <https://doi.org/10.1016/j.meatsci.2006.04.019>

538 Greppi, A., Ferrocino, I., La Stora, A., Rantsiou, K., Ercolini, D., & Cocolin, L. (2015). Monitoring of the  
 539 microbiota of fermented sausages by culture independent rRNA-based approaches. *International*  
 540 *Journal of Food Microbiology*, 212, 67-75. <https://doi.org/10.1016/j.ijfoodmicro.2015.01.016>

541 Hammes, W. P. (2012). Metabolism of nitrate in fermented meats: the characteristic feature of a specific group  
 542 of fermented foods. *Food Microbiology*, 29, 151-156. <https://doi.org/10.1016/j.fm.2011.06.016>

543 Holck, A., Axelsson, L., McLeod, A., Rode, T. M., & Heir, E. (2017). Health and safety considerations of  
 544 fermented sausages. *Journal of Food Quality*, 2017, 1-25. <https://doi.org/10.1155/2017/9753894>

545 Honikel, K. O. (2004). Curing agents. In W. K. Jensen, C. Devine, & M. Dikeman (Eds.), *Encyclopedia of meat*  
 546 *sciences* (pp. 195e201). Oxford: Elsevier Ltd.

547 Honikel, K. O. (2008). The use and control of nitrate and nitrite for the processing of meat products. *Meat*  
 548 *Science*, 78, 68-76. <https://doi.org/10.1016/j.meatsci.2007.05.030>

549 Hospital, X. F., Hierro, E., & Fernández, M. (2014). Effect of reducing nitrate and nitrite added to dry fermented  
 550 sausages on the survival of *Salmonella typhimurium*. *Food Research International*, 62, 410-415.  
 551 <https://doi.org/10.1016/j.foodres.2014.03.055>

552 Hospital, X. F., Carballo, J., Fernández, M., Arnau, J., Gratacós, M., & Hierro, E. (2015). Technological  
 553 implications of reducing nitrate and nitrite levels in dry-fermented sausages: typical microbiota,  
 554 residual nitrate and nitrite and volatile profile. *Food Control*, 57, 275-581.  
 555 <https://doi.org/10.1016/j.foodcont.2015.04.024>

556 ISO (2013). Microbiology of food and animal feeding stuffs — horizontal method for the enumeration of sulfite-  
 557 reducing. bacteria growing under anaerobic conditions. Geneva: International Organization for  
 558 Standardization.

559 ISO (2017a). ISO 11290-1:2017. Microbiology of the food chain — horizontal method for the detection and  
 560 enumeration of *Listeria monocytogenes* and of *Listeria* spp. Part 1: Detection method. Geneva:  
 561 International Organization for Standardization.

562 ISO (2017b). ISO 11290-2:2017 Microbiology of the food chain — horizontal method for the detection and  
 563 enumeration of *Listeria monocytogenes* and of *Listeria* spp. — Part 2: Enumeration method. Geneva:  
 564 International Organization for Standardization.

ISO (2017c). ISO 6579-1:2017. Microbiology of the food chain — horizontal method for the detection, enumeration and serotyping of *Salmonella*. Part 1: Detection of *Salmonella* spp. method. Geneva: International Organization for Standardization.

Leroy, F., Geyzen, A., Janssens, M., De Vuyst, L., & Scholliers, P. (2013). Meat fermentation at the crossroads of innovation and tradition. A historical outlook. *Trends in Food Science and Technology*, 31, 130-137. <https://doi.org/10.1016/j.tifs.2013.03.008>

Lijinsky, W., & Epstein, S. (1970). Nitrosamines as environmental carcinogens. *Nature*, 225, 21-23. <https://doi.org/10.1038/225021a0>

Lorenzo, J. M., Gomez, M., & Fonseca, S. (2014). Effect of commercial starter cultures on physicochemical characteristics, microbial counts and free fatty acid composition of dry-cured foal sausage. *Food Control*, 46, 382-389. <https://doi.org/10.1016/j.foodcont.2014.05.025>

Lorenzo, J. M., Gómez, M., Purriños, L., & Fonseca, S. (2016). Effect of commercial starter cultures on volatile compound profile and sensory characteristics of dry-cured foal sausage. *Journal of the Science of Food and Agriculture*, 96(4), 1194-1201. <https://doi.org/10.1002/jsfa.7203>

Luna, C., & Estévez, M. (2018). Oxidative damage to food and human serum proteins: Radical-mediated oxidation vs. glyco-oxidation. *Food Chemistry*, 267, 111-118. <https://doi.org/10.1016/j.foodchem.2017.06.154>

Mataragas, M., Bellio, A., Rovetto, F., Astegiano, S., Decastelli, L., & Cocolin, L. (2015). Risk-based control of food-borne pathogens *Listeria monocytogenes* and *Salmonella enterica* in the Italian fermented sausages Cacciatore and Felino. *Meat Science*, 103, 39-45. <https://doi.org/10.1016/j.meatsci.2015.01.002>

Montanari, C., Bargossi, E., Gardini, A., Lanciotti, R., Magnani, R., Gardini, F., et al. (2016). Correlation between volatile profiles of Italian fermented sausages and their size and starter culture. *Food Chemistry*, 192, 736-744. <https://doi.org/10.1016/j.foodchem.2015.07.062>

589 Montanari, C., Gatto, V., Torriani, S., Barbieri, F., Bargossi, E., Lanciotti, R., et al. (2018). Effects of the diameter  
590 on physicochemical, microbiological and volatile profile in dry fermented sausages produced with two  
591 different starter cultures. *Food Bioscience*, 22, 9-18. <https://doi.org/10.1016/j.fbio.2017.12.013>

592 Montanari, C., Barbieri, F., Gardini, F., & Tabanelli, G. (2021). Competition between starter cultures and wild  
593 microbial population in sausage fermentation: a case study regarding a typical Italian salami  
594 (Ventricina). *Foods*, 10, 2138. <https://doi.org/10.3390/foods10092138>

595 NIST. (2011). *NIST/NIH/EPA Mass Spectral Library, Standard Reference Database 1, NIST 11. Standard Reference*  
596 *Data Program*. Gaithersburg, MD, USA: National Institute of Standards and Technology.

597 Olivares, A., Dryahina, K., Navarro, J. L., Smith, D., Španěl, P., & Flores, M. (2011). SPME-GC-MS versus Selected  
598 Ion Flow Tube Mass Spectrometry (SIFT-MS) analyses for the study of volatile compound generation  
599 and oxidation status during dry fermented sausage processing. *Journal of Agricultural and Food*  
600 *Chemistry*, 59, 1931-1938. <https://doi.org/10.1021/jf104281a>

601 Omer, M. K., Álvarez-Ordoñez, A., Prieto, M., Skjerve, E., Asehun, T., & Alvseike, O. A. (2018). A systematic  
602 review of bacterial foodborne outbreaks related to red meat and meat products. *Foodborne Pathogens*  
603 *and Disease*, 15, 598-611. <https://doi.org/10.1089/fpd.2017.2393>

604 Ordóñez, J.A., Hierro, E.M., Bruna, L., & de la Hoz, L. (1999). Changes in the composition of dry-fermented  
605 sausages during ripening. *Critical Reviews in Food Science and Nutrition*, 39(4), 329-367.  
606 <https://doi.org/10.1080/10408699991279204>

607 Ozaki, M.M., Santos, M.D., Ribeiro, W.O, Azambuja Ferreira, N.C.D, Picone C.F.S., Domínguez, R., Lorenzo, J.M.,  
608 & Pollonio, M.A.R (2021). Radish powder and oregano essential oil as nitrite substitutes in fermented  
609 cooked sausages. *Food Research International*, 140, 109855.  
610 <https://doi.org/10.1016/j.foodres.2020.109855>

611 Patarata, L., Novais, M., Fraqueza, M. J., & Silva, J. A. (2020). Influence of meat spoilage microbiota initial load  
612 on the growth and survival of three pathogens on a naturally fermented sausage. *Foods (Basel,*  
613 *Switzerland)*, 9(5), 676. <https://doi.org/10.3390/foods9050676>

614 Pegg, R. B., & Honikel, K. O. (2014). Principles of curing. In F. Toldrá, Y. H. Hui, I. Astiasarán, J. G. Sebranek, & R.  
 615 Talon (Eds.), *Handbook of Fermented Meat and Poultry*, 2nd Edition (pp. 19-30). Hoboken: John Wiley  
 616 & Sons, Inc.

617 Perea-Sanz, L., Montero, R., Belloch, C., Flores, M. (2018). Nitrate reduction in the fermentation process of salt  
 618 reduced dry sausages: Impact on microbial and physicochemical parameters and aroma profile.  
 619 *International Journal of Food Microbiology*, 282, 84-91.  
 620 <https://doi.org/10.1016/j.ijfoodmicro.2018.06.004>

621 Perea-Sanz, L., Montero, R., Belloch, C., & Flores, M. (2019). Microbial changes and aroma profile of nitrate  
 622 reduced dry sausages during vacuum storage. *Meat Science*, 147, 100-107.  
 623 <https://doi.org/10.1016/j.meatsci.2018.08.026>

624 Raimondi, S., Popovic, M., Amaretti, A., Di Gioia, D., & Rossi, M. (2014). Anti-Listeria starters: in vitro selection  
 625 and production plant evaluation. *Journal of Food Protection*, 77(5), 837–842.  
 626 <https://doi.org/10.4315/0362-028X.JFP-13-297>

627 Sánchez Mainar, M., & F. Leroy (2015). Process-driven bacterial community dynamics are key to cured meat  
 628 colour formation by coagulase-negative staphylococci via nitrate reductase or nitric oxide synthase  
 629 activities. *International Journal of Food Microbiology*, 212, 60-66.  
 630 <https://doi.org/10.1016/j.ijfoodmicro.2015.03.009>

631 Selgas M. D., & Garcia M. L. (2014). Yeasts. In F. Toldrá, Y. H. Hui, I. Astiasarán, J. G. Sebranek, & R. Talon (Eds.),  
 632 *Handbook of Fermented Meat and Poultry*, 2nd Edition (pp. 139-146). Hoboken: John Wiley & Sons, Inc.

633 Skibsted, L. H. (2011). Nitric oxide and quality and safety of muscle-based foods. *Nitric Oxide*, 24(4), 176-183.  
 634 <https://doi.org/10.1016/j.niox.2011.03.307>

635 Sindelar, J. J., & Milkowski, A. L. (2011). Sodium nitrite in processed meat and poultry meats: a review of curing  
 636 and examining the risk/benefit of its use. *American Meat Science Association White Paper Series*, 3, 1-  
 637 14.

638 Stahnke, M.L.H. (2008). U.S. Patent No. 2008/0102159 A1. Method for reducing the content of pathogenic  
639 organisms present in food materials. Washington, DC: U.S. Patent and Trademark Office.

640 Tabanelli, G., Montanari, C., Grazia, L., Lanciotti, R., & Gardini, F. (2013). Effects of  $a_w$  at packaging time and  
641 atmosphere composition on aroma profile, biogenic amine content and microbiological features of dry  
642 fermented sausages. *Meat Science*, 94, 177-186. <https://doi.org/10.1016/j.meatsci.2013.01.018>

643 Toldrá, F., & Hui, Y. H. (2014). Dry-fermented sausages and ripened meats: an overview. In F. Toldrá, Y. H. Hui, I.  
644 Astiasarán, J. G. Sebranek, & R. Talon (Eds.), *Handbook of Fermented Meat and Poultry*, 2nd Edition (pp.  
645 1-6). Hoboken: John Wiley & Sons, Inc.

646 Tricker, A. R., & Preusmann, R. (1987). Influence of cysteine and nitrate on the endogenous formation of N-  
647 nitrosoamino acids. *Cancer Letters*, 34, 39-47. [https://doi.org/10.1016/0304-3835\(87\)90071-1](https://doi.org/10.1016/0304-3835(87)90071-1)

648 Van Reckem, E., Geeraerts, W., Champi, C., Van der Veken, D., De Vuyst, L., & Leroy, F. (2019). Exploring the  
649 link between the geographical origin of European fermented foods and the diversity of their bacterial  
650 communities: the case of fermented meats. *Frontiers in Microbiology*, 10, 2302.  
651 <https://doi.org/10.3389/fmicb.2019.02302>

652 von Wright, A., & Axelsson, L. (2011). Lactic acid bacteria: an introduction. In S. Lahtinen, A. C. Ouwehand, S.  
653 Salminen, & A. von Wright (Eds.), *Lactic Acid Bacteria, Microbial and Functional Aspects*, 4th Edition  
654 (pp. 1-16). Boca Raton, FL: CRC Press.

655 Zotta, T., Parente, E., & Ricciardi, A. (2017). Aerobic metabolism in the genus *Lactobacillus*: impact on stress  
656 response and potential applications in the food industry. *Journal of Applied Microbiology*, 122, 857-  
657 869. <https://doi.org/10.1111/jam.13399>

658 **Table 1.** Microbiological analyses (log CFU/g) of meat, meat batter and ripened fermented sausages produced  
 659 with preservatives (C) or without nitrate and nitrite using two different glucose concentrations, (0.2% (F02) or  
 660 0.4% (F04)). Results are the mean of three independent repetitions (standard deviation is reported). For each  
 661 microbial group, significant differences among samples at the end of ripening according to ANOVA are  
 662 indicated by the presence of different letters.

663

Microbial group	Meat	Meat batter	End of ripening (50 days)		
			C	F02	F04
Lactic acid bacteria	3.2 ± 0.2	7.2 ± 0.2	8.2 ± 0.1	8.1 ± 0.2	8.3 ± 0.1
Coagulase negative cocci	3.8 ± 0.2	6.5 ± 0.2	6.8 ± 0.2	7.1 ± 0.1	7.0 ± 0.1
Enterococci	2.3 ± 0.1	2.5 ± 0.2	2.1 <sup>a</sup> ± 0.2	2.7 <sup>b</sup> ± 0.2	2.6 <sup>b</sup> ± 0.1
Enterobacteriaceae	2.5 ± 0.2	2.6 ± 0.2	1.5 <sup>a</sup> ± 0.1	2.3 <sup>b</sup> ± 0.1	2.3 <sup>b</sup> ± 0.1
Yeasts	1.8 ± 0.1	1.9 ± 0.1	4.6 ± 0.1	4.3 ± 0.1	4.4 ± 0.2

664

**Table 2.** Colour parameters, textural features and lipid oxidation level (TBARS) of ripened fermented sausages produced with (C) or without the addition of preservatives (*i.e.*, nitrate and nitrite) using two different glucose concentrations, namely 0.2% (F02) or 0.4% (F04). Results belong to three independent repetitions and are expressed as mean values  $\pm$  standard deviation.

Parameter	End of ripening (50 days)			p-value	
	C	F02	F04	Analysis of variance	Planned orthogonal contrasts <sup>1</sup>
<b>Color</b>					
Lightness - L*	42.27 $\pm$ 4.46	40.26 $\pm$ 3.00	40.81 $\pm$ 2.05	ns	ns
Redness - a*	13.51 <sup>a</sup> $\pm$ 1.22	11.58 <sup>b</sup> $\pm$ 1.68	12.57 <sup>ab</sup> $\pm$ 1.13	**	*
Yellowness - b*	5.50 $\pm$ 1.01	4.44 $\pm$ 0.50	5.38 $\pm$ 0.93	ns	ns
Chroma - C*	14.6 <sup>a</sup> $\pm$ 1.4	12.4 <sup>b</sup> $\pm$ 1.6	13.7 <sup>ab</sup> $\pm$ 1.2	**	*
Hue angle - h	0.39 $\pm$ 0.05	0.37 $\pm$ 0.05	0.40 $\pm$ 0.06	ns	ns
<b>Texture Profile Analyses (TPA)</b>					
Hardness (kg)	2.07 <sup>c</sup> $\pm$ 0.04	3.20 <sup>b</sup> $\pm$ 0.12	3.74 <sup>a</sup> $\pm$ 0.23	***	***
Cohesiveness	1.83 <sup>a</sup> $\pm$ 0.02	1.72 <sup>b</sup> $\pm$ 0.02	1.73 <sup>b</sup> $\pm$ 0.01	***	***
Gumminess (kg)	3.79 <sup>c</sup> $\pm$ 0.07	5.51 <sup>b</sup> $\pm$ 0.18	6.46 <sup>a</sup> $\pm$ 0.34	***	***
Springiness	1.39 $\pm$ 0.65	1.80 $\pm$ 0.03	1.75 $\pm$ 0.05	ns	ns
Chewiness (kg)	5.27 <sup>b</sup> $\pm$ 2.47	9.94 <sup>a</sup> $\pm$ 0.21	11.32 <sup>a</sup> $\pm$ 0.64	**	**
<b>Oxidation of the lipid fraction</b>					
TBARS (mg MDA/kg of meat product)	0.45 <sup>c</sup> $\pm$ 0.03	0.80 <sup>b</sup> $\pm$ 0.02	1.10 <sup>a</sup> $\pm$ 0.07	**	**

<sup>1</sup>Planned orthogonal contrasts performed to assess the eventual effect of the absence of nitrite and nitrate in dry fermented sausages formulation regardless of the glucose content added (C vs. F02 and F04).  
 \*= $p < 0.05$ , \*\*= $p < 0.01$ , \*\*\*= $p < 0.001$ , ns = not significant; a,c = mean values followed by different letters, significantly different among the groups ( $p < 0.05$ ).



675 **Table 3.** Aroma profile detected by SPME-GC-MS in the ripened sausages produced with preservatives (C) or  
676 without nitrate and nitrite using two different glucose concentrations, namely 0.2% (F02) or 0.4% (F04). Data  
677 are expressed as ratio between peak area of each molecule and peak area of the internal standard (4-methyl-2-  
678 pentanol). Results are the mean of three independent repetitions and standard deviations are reported. For  
679 each molecule significant differences between samples according to ANOVA are indicated by the presence of  
680 different letters.

Volatile compound	C	F02	F04
Acetone	32.2 ± 1.0 <sup>a</sup>	18.6 ± 1.3 <sup>b</sup>	33.4 ± 1.8 <sup>a</sup>
2-butanone	1.06 ± 0.18 <sup>ab</sup>	0.821 ± 0.358 <sup>a</sup>	1.62 ± 0.25 <sup>b</sup>
2-pentanone	0.643 ± 0.211 <sup>a</sup>	0.878 ± 0.164 <sup>a</sup>	2.10 ± 0.45 <sup>b</sup>
2,3-butanedione	4.92 ± 0.95 <sup>a</sup>	0.349 ± 0.098 <sup>b</sup>	0.714 ± 0.228 <sup>b</sup>
Methyl isobutyl ketone	0.449 ± 0.182	0.179 ± 0.209	0.491 ± 0.089
3-hexen-2-one	1.78 ± 0.45 <sup>a</sup>	5.10 ± 1.30 <sup>b</sup>	2.86 ± 0.57 <sup>a</sup>
3-hydroxy-2-butanone	11.7 ± 1.2 <sup>a</sup>	0.601 ± 0.411 <sup>b</sup>	4.26 ± 0.99 <sup>c</sup>
2-heptanone	1.90 ± 0.66 <sup>a</sup>	3.51 ± 0.91 <sup>b</sup>	4.07 ± 0.19 <sup>b</sup>
2-nonanone	0.914 ± 0.342	1.07 ± 0.14	1.41 ± 0.33
<b>Ketones</b>	<b>55.6 ± 2.8<sup>a</sup></b>	<b>31.1 ± 2.2<sup>b</sup></b>	<b>50.9 ± 3.3<sup>a</sup></b>
3-methylbutanal	0.175 ± 0.093	0.264 ± 0.133	0.192 ± 0.098
Hexanal	1.43 ± 0.30	1.10 ± 0.23	1.18 ± 0.15
2-heptenal	0.541 ± 0.334 <sup>a</sup>	2.41 ± 0.33 <sup>b</sup>	3.01 ± 0.53 <sup>b</sup>
Nonanal	1.05 ± 0.25 <sup>a</sup>	3.23 ± 0.51 <sup>b</sup>	4.94 ± 0.33 <sup>c</sup>
2-nonenal	1.75 ± 0.53	1.11 ± 0.65	1.83 ± 0.54
Benzaldehyde	0.591 ± 0.402	0.631 ± 0.258	1.29 ± 0.48
Decanal	0.343 ± 0.201	0.539 ± 0.249	0.202 ± 0.212
<b>Aldehydes</b>	<b>5.89 ± 0.41<sup>a</sup></b>	<b>9.28 ± 0.23<sup>b</sup></b>	<b>12.6 ± 0.4<sup>c</sup></b>
Ethanol	4.79 ± 0.89 <sup>a</sup>	5.98 ± 1.01 <sup>a</sup>	9.86 ± 0.75 <sup>b</sup>
Hexanol	0.551 ± 0.319 <sup>a</sup>	0.542 ± 0.291 <sup>a</sup>	1.15 ± 0.34 <sup>b</sup>
1-octen-3-ol	0.404 ± 0.264 <sup>a</sup>	0.810 ± 0.410 <sup>ab</sup>	1.318 ± 0.121 <sup>b</sup>
1-heptanol	0.414 ± 0.244 <sup>a</sup>	0.338 ± 0.173 <sup>a</sup>	1.04 ± 0.34 <sup>b</sup>
1-octanol	0.713 ± 0.331 <sup>a</sup>	2.94 ± 0.79 <sup>b</sup>	0.968 ± 0.413 <sup>a</sup>
2-octen-1-ol	4.15 ± 0.65 <sup>ab</sup>	3.35 ± 0.82 <sup>a</sup>	5.43 ± 0.54 <sup>b</sup>
<b>Alcohols</b>	<b>11.0 ± 1.0<sup>a</sup></b>	<b>14.0 ± 1.2<sup>b</sup></b>	<b>19.8 ± 1.0<sup>c</sup></b>
Acetic acid	24.0 ± 2.0 <sup>a</sup>	7.44 ± 1.11 <sup>b</sup>	19.8 ± 1.9 <sup>c</sup>
3-methyl butanoic acid	1.86 ± 0.15 <sup>a</sup>	0.724 ± 0.363 <sup>b</sup>	1.33 ± 0.23 <sup>b</sup>
Butanoic acid	1.84 ± 0.31 <sup>a</sup>	0.456 ± 0.329 <sup>b</sup>	2.09 ± 0.53 <sup>a</sup>
Pentanoic acid	1.12 ± 0.22 <sup>a</sup>	1.04 ± 0.57 <sup>a</sup>	2.28 ± 0.33 <sup>b</sup>
Hexanoic acid	1.80 ± 0.18	1.46 ± 0.31	2.05 ± 0.45
Heptanoic acid	0.692 ± 0.331	0.714 ± 0.509	1.54 ± 0.43

681


Octanoic acid	$2.03 \pm 0.28^a$	$2.47 \pm 0.65^{ab}$	$3.52 \pm 0.45^b$
Nonanoic acid	$2.35 \pm 0.64$	$1.68 \pm 0.83$	$2.66 \pm 0.22$
<b>Acids</b>	<b><math>35.7 \pm 2.9^a</math></b>	<b><math>16.0 \pm 1.4^b</math></b>	<b><math>35.3 \pm 2.4^a</math></b>
Ethyl acetate	$0.212 \pm 0.153^{ab}$	$0.149 \pm 0.081^a$	$0.429 \pm 0.222^b$
<b>Esters</b>	<b><math>0.212 \pm 0.153^{ab}</math></b>	<b><math>0.149 \pm 0.081^a</math></b>	<b><math>0.429 \pm 0.222^b</math></b>


682 **FIGURE CAPTIONS**

683 **Figure 1** pH (1A),  $a_w$  (1B) and weight loss (1C) during fermentation and ripening of fermented sausages  
684 produced with preservatives (C) or without nitrate and nitrite using two different glucose concentrations,  
685 namely 0.2% (F02) or 0.4% (F04). The data are the mean of three independent samples and standard error bars  
686 are reported.


687

688 **Figure 2:** Microbial counts (expressed as log CFU/g) of *Listeria innocua* (2A), *Salmonella enterica* subsp. *enterica*  
689 (2B) and *Clostridium botulinum* (2C) during fermentation and ripening of fermented sausages produced with  
690 preservatives (C) or without nitrate and nitrite (F02). The glucose addition was 0.2 % for both samples. The  
691 data are the mean of three independent samples and standard error bars are reported.





# Safety and technological issues of dry fermented sausages produced without nitrate and nitrite



## Credit author statement

**Giulia Tabanelli:** conceptualization, writing-original draft preparation; **Federica Barbieri:** investigation, microbiological and chemico-physical analyses; **Francesca Soglia:** investigation, chemico-physical analyses; **Rudy Magnani:** Funding acquisition and resources; **Gabriele Gardini:** chemico-physical analyses; **Massimiliano Petracci:** writing - review & editing; **Fausto Gardini:** conceptualization writing-original draft preparation, supervision; **Chiara Montanari:** conceptualization, investigation, writing-review and editing.

## **Safety and technological issues of dry fermented sausages produced without nitrate and nitrite**

Giulia Tabanelli<sup>a</sup>, Federica Barbieri<sup>a</sup>, Francesca Soglia<sup>a</sup>, Rudy Magnani<sup>b</sup>, Gabriele Gardini<sup>b</sup>,  
Massimiliano Petracci<sup>a</sup>, Fausto Gardini<sup>a</sup>, Chiara Montanari<sup>a\*</sup>

<sup>a</sup> Department of Agricultural and Food Sciences, University of Bologna, Bologna, Italy

<sup>b</sup> C.I.a.i. Soc. Coop., Via Gambellara 62/A, 40026 Imola (BO), Italy

\* Corresponding author:

Chiara Montanari

Centro Interdipartimentale di Ricerca Industriale Agroalimentare,

Università degli Studi di Bologna, Sede di Cesena,

Via Quinto Bucci 336, 47521 Cesena, Italy

Tel. +39 0547338145, Fax +39 0547338103

### **CONFLICT OF INTEREST**

The authors confirm that they have no conflicts of interest with respect to the work described in this manuscript.