

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Bacterial colonization dynamics of different microplastic types in an anoxic salt marsh sediment and impact of adsorbed polychlorinated biphenyls on the plastisphere

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Availability:

This version is available at: <https://hdl.handle.net/11585/897566> since: 2022-10-26

Published:

DOI: <http://doi.org/10.1016/j.envpol.2022.120411>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the Accepted version of the article:

Rosato A, Barone M, Negroni A, Brigidi P, Fava F, Xu P, Candela M, Zanaroli G. Microbial colonization of different microplastic types and biotransformation of sorbed PCBs by a marine anaerobic bacterial community. *Sci Total Environ.* 2020 Feb 25;705:135790. doi: 10.1016/j.scitotenv.2019.135790. Epub 2019 Nov 26. PMID: 31972939.

The final published version is available online at:

<https://www.sciencedirect.com/science/article/abs/pii/S0048969719357857?via%3Dihub>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

MICROBIAL COLONIZATION OF DIFFERENT MICROPLASTIC TYPES AND BIOTRANSFORMATION OF SORBED PCBs BY A MARINE ANAEROBIC BACTERIAL COMMUNITY

Antonella Rosato^a, Monica Barone^b, Andrea Negroni^a, Patrizia Brigidi^b, Fabio Fava^a, Ping Xu^c, Marco Candela^b, Giulio Zanaroli^{a*}

^aDept. of Civil, Chemical, Environmental and Materials Engineering (DICAM), *Alma Mater Studiorum* University of Bologna, Via Terracini 28, 40131 Bologna, Italy

^bDept. of Pharmacy and Biotechnology (FaBit), *Alma Mater Studiorum* University of Bologna, Via Belmeloro 6, 40126 Bologna, Italy

^cSchool of Life Science & Biotechnology, Shanghai Jiao Tong University, 800 Dong Chuan Road, Shanghai, 200240, People's Republic of China

*corresponding author (giulio.zanaroli@unibo.it; +39 0512090924)

Keywords

Microplastics; biofilm; microbial community; marine sediment; polychlorinated biphenyls; reductive dehalogenation

Abstract

We investigated the colonization dynamics of different microplastic (MP) pellets, namely, polyethylene (PE), polyethylene terephthalate (PET), polystyrene (PS), polypropylene (PP) and polyvinyl chloride (PVC), either pristine or contaminated with polychlorinated biphenyls (PCBs), by an organohalide respiring marine microbial community and its biotransformation

activity towards PCBs sorbed on MPs, in anaerobic laboratory microcosms of a marine sediment.

All MPs were rapidly colonized by the microbial community within 2 weeks of incubation, when approximately 10^{10} 16S rRNA gene copies cm^{-2} were detected on PVC, 10^9 copies cm^{-2} on PE, and 10^8 copies cm^{-2} on PET, PP and PS. A greater biofilm growth on PVC pellets than other MPs was confirmed by quantification of the reducing sugars of the EPS and biofilm staining with crystal violet. Illumina sequencing of the 16S rRNA genes and Principal Coordinate Analysis (PCoA) revealed that the biofilm community on MPs significantly differed from the sediment community, being enriched of chemoorganotrophic fermenting species, and was significantly affected by the type of polymer. The presence of sorbed PCBs did not significantly affect the overall community composition, and mainly resulted in the enrichment of Dehalococcoidia, i.e., of the organohalide respiring members of the community.

Reductive dechlorination of PCBs sorbed to MPs was observed after two weeks of incubation, when the average number of chlorines per biphenyl molecule was reduced from 5.2 to 4.8 - 4.3, and was faster (35.2 ± 1.9 to 61.2 ± 5.8 $\mu\text{moles of Cl removed kg}_{\text{MP}}^{-1} \text{ week}^{-1}$) than that of sediment-sorbed ones (33.9 ± 9.1 $\mu\text{moles of Cl removed kg}_{\text{sediment}}^{-1} \text{ week}^{-1}$), which started only after 10 weeks of incubation. These data suggest that microbial colonization of contaminated MPs might change the composition of sorbed PCB mixtures and therefore the toxicity associated to PCB-polluted MPs.

1 Introduction

Plastics use and production has enormously increased since 1950, reaching around 335 million tonnes in 2016, with 60 million tonnes generated in Europe alone (PlasticsEurope, 2018). The most common polymers are high-density and low-density polyethylene (PE), polyethylene terephthalate (PET), polystyrene (PS), polypropylene (PP) and polyvinyl chloride (PVC), which together represent approximately 90% of the global plastic production (Andrady and Neal, 2009). A huge amount of plastic materials ends up in the marine environment, becoming an ever-increasing problem due to the toxicity, persistence and universal presence of such debris. The estimation of plastic released in the oceans varies from 4.8 to 12.7 million tons annually (Haward, 2018), with size ranging from meters to micrometers (Ryan et al., 2009). The term microplastics (MPs) was introduced within the last decade to describe small particles of plastic, commonly defined as < 5mm in diameter (Frias and Nash, 2019). Once discharged into the marine environment, plastic litter undergoes different processes, such as weathering, fragmentation and fouling (Cole et al., 2011). In particular, biomass accumulation on MPs due to bio(fouling) can lead to an increase of their density and thus their sinking (Morét-Ferguson et al., 2010; Chubarenko et al, 2016; Miao et al., 2019a). The anaerobic sediment has been indicated as a possible long-term sink for MPs (Andrady, 2011; Van Cauwenberghe et al., 2015). Microplastic contamination in marine sediments has been reported by several authors (Thompson et al., 2004; McDermid and McMullen, 2004; Claessens et al., 2011), with some of the highest concentrations of MPs found in sediments located in Nova Scotia and Arctic Ocean (up to 8000 and 6595 MPs/kg of sediment, respectively) (Mathalon and Hill, 2014; Bergmann et al., 2017).

The first evidence of microbial colonization of plastic fragments dates back to the early 70s (Carpenter et al., 1972). Recent studies reported that MPs are readily colonized by environmental microbial communities in few hours or days of incubation in seawater

(Ogonowski et al., 2018) and coastal marine sediments (Harrison et al., 2014). The bacterial assemblages colonizing MPs in freshwater and marine environments have been reported to significantly differ in taxonomic composition and structure from those present in the surrounding water and/or sediment (Miao et al., 2019b; Frère et al., 2018; Ogonowski et al., 2018; Rummel et al., 2017; De Tender et al., 2015). Microplastic-associated microbial communities were also reported to have a lower alpha diversity (richness, evenness, and diversity) than those associated to natural substrates, indicating a remarkable differentiation between microbial communities and a substrate-type-coupled species sorting (Miao et al., 2019b; Ogonowski et al., 2018). MPs can thus be considered a distinct ecological habitat for diverse microbial communities, the “plastisphere” (Zettler et al., 2013), potentially characterized by distinct microbial and ecological functions (Miao et al., 2019b; Arias-Andres et al., 2018). Less conclusive information has been reported on the selective colonization of MPs of different materials, possibly as the consequence of the potential confounding temporal and environmental variability associated to the dynamic exposure conditions typically occurring in situ (Ogonowski et al., 2018). Recently, Li et al. (2019) reported that environmental factors, such as the salinity and the nutrients (total nitrogen and total phosphorus), affected the growth rate of biofilms on five types of plastics debris (polyvinyl chloride, polypropylene, polyethylene, polystyrene, and polyurethane) in the Haihe Estuary. The same authors also showed that some genera in the bacterial communities exhibit selectivity for the different polymer types (Li et al., 2019).

Given their small size, MPs may be ingested and accumulated by a wide range of marine organisms, causing direct effects such as physical damage in their intestinal tract, or in other tissues or organs (Van Cauwenberghe et al., 2015). However, the uptake of MPs might also have indirect impacts, since they can absorb and concentrate persistent organic pollutant (POPs) by partition and then transfer such contaminants to the marine food web through ingestion

(Andrady, 2011; Wang et al., 2018). Polychlorinated biphenyls (PCBs), polycyclic aromatic hydrocarbons (PAHs), dichloro-diphenyl-trichloroethane (DDTs), polybrominated diphenyl ethers (PBDEs), alkylphenols and bisphenol A (BPA), are the main contaminants that have been found on plastics debris in the marine environment, at concentrations ranging from 0.001 to 10 mg/kg (Hirai et al., 2011).

Despite their production ban in the 1980s, PCBs are still common contaminants in marine sediments and, as reported recently, on marine plastic pellets, where they have been detected at concentrations up to 7.5 mg/kg (Taniguchi et al., 2016). Due to their high lipophilicity, PCBs are accumulated through the food chain up to humans, where they act as endocrine disrupters and possible carcinogens. Besseling et al. (2013) showed that an exposure of lugworm (*Arenicola marina*) to low concentrations of polystyrene MPs mixed with PCB-contaminated sediment enhanced the bioaccumulation of PCBs. Similar results were obtained by Koelmans et al. (2013) in their model analysis with *A. marina*, although their experiments showed that the role of plastic in the bioaccumulation of POPs may not be considered a significant hazard. Therefore, the release of MPs in the marine environment can have contrasting effects on the PCBs bioaccumulation in marine organisms, which can depend on different causes, such as the experimental conditions or the type of PCB congeners, and further investigations in this regard are necessary (Ziccardi et al., 2016). The toxicity of PCBs varies with the position and number of chlorine atoms on the biphenyl ring, being the coplanar congeners more toxic (Tanabe et al., 1987; Hashmi et al., 2017). Under anoxic conditions, PCBs can undergo reductive dechlorination by organohalide respiring microorganisms occurring in marine sediments, typically belonging to the Dehalococcoidia class of the Phylum Chloroflexi (Zanaroli et al., 2015; Nuzzo et al., 2017). This process replaces chlorine atoms mainly in the *meta* and *para* positions with hydrogen, thus converting highly chlorinated congeners into less chlorinated products that are often less toxic, less prone to bioaccumulation (less hydrophobic) and more amenable

to degradation by indigenous aerobic bacteria (Fava and Agathos, 2006). If taking place on MP-sorbed PCBs, this microbial process might change the composition of PCBs, and thus the toxicity and bioavailability of the sorbed PCB mixture for consumers ingesting MPs. The effect of microbial colonization of plastics debris on the environmental fate and the biodegradation/biotransformation of the pollutants absorbed on the MPs surface has not been adequately evaluated yet. Microbial colonization of low-density PE pellets incubated in marine and river sediments under aerobic conditions has been shown to increase the biotransformation of MP-sorbed DDTs and PAHs, but not of PCBs (Wu et al., 2017), suggesting that it might affect the toxicity associated with polluted microplastics. However, to the best of our knowledge, no studies investigated the influence of microbial colonization of MPs on MP-sorbed pollutants under anoxic conditions typically present in sediments.

The aim of this study was to investigate the colonization dynamics on different types of MPs (low-density PE, PET, PS, PP and PVC) by an anaerobic marine bacterial community containing organohalide respiring bacteria, and the effect of the microbial biofilm on the biotransformation of sorbed PCBs. The process was studied in anaerobic slurry microcosms consisting of marine sediment suspended in seawater, i.e. under biogeochemical conditions mimicking those occurring in situ and controlled exposure conditions.

2 Materials and Methods

2.1 Microplastics and their contamination

Five types of MPs were selected for this study: low density polyethylene pristine pellets (PE), crystalline poly(ethylene terephthalate) pristine pellets (PET), general purpose polystyrene pristine pellets (PS), homo-polypropylene pristine pellets (PP), and poly(vinyl chloride) soft pristine pellets (PVC). The size of the plastic particles ranged from approximately 2.5 to 3 mm. All MPs were sterilized in 70% ethanol for 15 min under shaking, followed by rinsing 3 times

with sterilized deionized water, before their use. MPs were contaminated in the laboratory with a commercial mixture of PCBs (Aroclor 1254, UltraScientific, Bologna, Italy) at the final concentration of 30 mg_{PCBs}/kg_{MPs}. This concentration, which is approximately the same order of magnitude of the highest PCBs concentration reported on marine plastic debris (7.5 mg/kg; Taniguchi et al., 2016), was selected in order to obtain final concentrations of single PCB congeners (or co-eluting congeners) of the mixture in the range 20 µg/kg_{MPs} - 4.8 mg/kg_{MPs}, and thus to better assess the microbial dechlorination processes of the spiked PCB mixture. The contamination protocol was adapted from Beckingham and Ghosh (2017); synthetic marine water (1.5 L) was spiked with 0.6 mL of Aroclor 1254 stock solution (20000 mg/L in acetone) under shaking (acetone:water 0.04%), 400 g of MPs were added immediately after and incubated under shaking (180 rpm) for 10 days. MPs were recovered by sieving and dried at room temperature under sterile conditions. The actual amount of PCBs sorbed on MPs was indirectly estimated by measuring the residual mass of PCBs in the water phase via solid phase extraction with polydimethylsiloxane (PDMS) fibers. Five-cm PDMS fibers (outer diameter 558.8 µm, inner diameter 486 µm, thickness annulus 35.4 µm, fiber volume 0.597 µL/cm) were incubated in the water phase under mixing (150 rpm) at room temperature for 40 days. PCBs were then eluted for 16 h in 0.1 mL hexane and analysed as described in section 2.3. The water concentration of PCBs was calculated from their concentration in the PDMS fiber using the fiber-water coefficient (K_{PDMS-W}). The latter was calculated, for each PCB congener, from the octanol–water partition coefficient (K_{ow}) using the equation $\log K_{PDMS-W} = 0.725 \log K_{ow} + 0.479$ (Thomas et al., 2014). Octanol–water partition coefficients for all PCB congeners were obtained from Hawker and Connell (1988). An almost negligible mass of PCBs was detected in the water phase, corresponding to 0.71%, 0.15%, 0.16%, 0.01% and 0.01% (in case of PET, PP, PS, PE and PVC, respectively) of the total mass of PCBs initially added for MPs contamination. The residual mass of PCBs was assumed to be sorbed on MPs.

2.2 *Microcosms set up, incubation and sampling*

Sacrificial anaerobic slurry microcosms (200 mL total volume) consisting of sediment (20% w/v) and seawater collected from Piailassa Baiona (Ravenna, Italy) were set up according the procedure described in Nuzzo et al. (2017). Microcosms were autoclave-sterilized for 1 h on three consecutive days, inoculated at 5% (v/v) with a PCB-dechlorinating microbial culture enriched previously from marine sediments (Nuzzo et al., 2017) and supplemented with 5 g of MPs, corresponding to a number of MPs ranging from approximately 175 to approximately 300, depending on the density and size of the MP type. This resulted in a final concentration ranging from approximately 4375 to approximately 7500 MPs/kg of sediment, which is comparable with the concentrations of microplastics reported in the most MP-impacted marine sediments (Bergmann et al., 2017; Mathalon and Hill, 2014). For each type of MPs, the following microcosms sets were prepared: i) supplemented with MPs contaminated by PCBs; ii) supplemented with pristine MPs (not contaminated). A sterile control set was set up for each MP type by supplementing sterile, not inoculated slurry microcosms with MPs contaminated by PCBs. In addition, a MP-free biologically active control was set up by spiking Aroclor 1254 PCBs in the sediment and inoculating the same marine culture.

The microcosms were incubated at 20°C in the dark under static conditions for 28 weeks. After 2, 5, 10, 14, 19 and 28 weeks of incubation, 3 microcosms for each MP type were sacrificed and MPs were recovered through sieving on sterilized 0.5 mm sieves for the analysis of biofilm microbial communities (except from sterile controls) and of the sorbed PCBs (where spiked).

2.3 *PCB extraction and analysis*

PCBs were batch extracted from 1 g of MPs (i.e., approximately 35 to 60 MPs, depending on the MP type) with 4 mL of hexane overnight (30°C, mixing at 150 rpm) and sonication (Hong et al.,

2017). PCBs in the organic extracts were analyzed with a 6890N gas-chromatograph equipped with a ^{63}Ni electron capture detector (μECD) and a 6890 series-automatic sampler (Agilent Technologies) using a $30\text{m} \times 0.25\text{mm}$ HP-5 capillary column (Agilent Technologies) under the conditions described elsewhere (Fava et al., 2003). Calibration curves were obtained and verified monthly using standard mixtures of Aroclor 1254 and Aroclor 1242 (0.5 to 30 mg/L concentration range).

2.4 *Quantification of the bacterial biofilm*

Prior to each analysis, the MPs were separated from the sediment by sieving (mesh 0.5 mm), rinsed three times with sterile water, in order to remove not-attached cells, and air dried under sterile conditions. Three different approaches were used for the quantification of the biofilm growth on MPs: i) metagenomic DNA extraction followed by qPCR of bacterial 16S rRNA genes; ii) cell/biofilm staining with crystal violet; iii) quantification of the reducing sugars after alkaline hydrolysis of the EPS polysaccharides (Costerton et al., 1995).

Metagenomic DNA was extracted from a mixture of replicate samples (0.5 g of MPs, i.e., approximately 18-30 MPs, depending on the MP type) with the UltraClean Soil DNA kit (MoBio Laboratories, Carlsbad, CA, USA) following the procedure described by the provider, with an additional enzymatic cell lysis step before mechanical cell lysis. In particular, MPs were incubated in the bead solution (provided with the kit) in the presence of Lysozyme (11.9 μL of a 100 mg mL^{-1} solution) at $37\text{ }^\circ\text{C}$ on a rotary shaker for 30 min, and then of Proteinase K (3 μL of a 20 mg mL^{-1} solution) at $37\text{ }^\circ\text{C}$ on a rotary shaker for 45 min, prior to addition of SDS and mechanical cell lysis (bead beating on vortex at maximum speed for 10 min). Total DNA was quantified using Qubit® dsDNA HS Assay Kit with a Qubit 3.0 fluorimeter, following the manufacturer's specifications. The 16S rRNA genes were quantified via qPCR with primers 905f (5'-AAACTCAAAGGAATTGACGG-3') and 1044r (5'-GACARCCATGCASCACCTG-3') using the

reactions conditions described in Nuzzo et al. (2017). 7-point standard curves were included in each plate using *E. coli* 16S rRNA gene. Samples and standards were analyzed in triplicate reactions and 16S rRNA gene copy numbers per cm² of MPs were finally calculated.

A colorimetric biofilm quantification protocol using crystal violet was adapted from Burton et al. (2007) and Zanaroli et al. (2011). MPs (0.5 g, i.e., approximately 18-30 MPs, depending on the MP type) were stained 20 minutes with 2 mL of crystal violet solution (0.05% w/v). MPs were then rinsed with sterile distilled water and air-dried for 15 min. The crystal violet on MP-associated biofilm was solubilized by adding 2 mL of 96% (v/v) ethanol and the absorbance of the destaining solution was spectrophotometrically measured at 570 nm versus a blank solution obtained with the same procedure using ethanol-sterilized MPs (see above).

Reducing sugars occurring in the EPS polysaccharides of the biofilm were measured via the colorimetric method described by Bailey (1988), using 3,5 dinitrosalicylic acid, potassium tartrate and NaOH. MPs (0.5 g, i.e., approximately 18-30 MPs, depending on the MP type) were incubated in a boiling water bath for 15 minutes. After cooling in ice, the absorbance of the supernatant was measured at 540 nm versus a blank solution obtained with the same procedure using ethanol-sterilized MPs (see above).

To identify statistically significant differences between MP types, multiple pair-wise comparisons were performed with the Tukey test ($P < 0.05$).

2.5 Analysis of the biofilm community via PCR-DGGE of the 16S rRNA genes

PCR-DGGE analysis of 16S bacterial rRNA genes was used to investigate changes in richness (Rr) and community structure (Co) of the bacterial biofilm on different MPs over time. The V3-V5 variable regions of the bacterial 16S rRNA genes were PCR amplified with primers GC-357f, containing a 40 bp GC-clamp (5'-CGC CCG CCG CGC CCC GCG CCC GGC CCG CCG CCC CCG CCC C CC TAC GGG AGG CAG CAG-3') and 907r (5'-CCG TCA ATT CCT TTG AGT TT-3') (Sass et al.,

2001). The PCR program consisted of an initial denaturation at 95 °C for 5 min, 30 cycles of repeated denaturation at 95 °C for 30 s, annealing at 55 °C for 30 s, and elongation at 72 °C for 1 min, followed by a final elongation at 72 °C for 10 min. The PCR reaction (50 µL) consisted of mixtures containing 1× colorless GoTaq® Flexi buffer (Promega Corporation, Italy), 1.5 mM MgCl₂, 0.2 mM each dNTP, 0.4 µM each primer, 1 U of GoTaq® G2 Flexi DNA Polymerase (Promega Corporation, Italy), and of 4 µL template DNA.

The amplicons were resolved with a D-Code Universal Mutation Detection System (Bio-Rad, Milan, Italy) on a 7% (w/v) polyacrylamide gel (acrylamide-N,N'-methylenebisacrylamide, 37:1) containing a denaturing gradient from 40% (top) to 60% (bottom) denaturant (100% denaturant: 7M urea, 40% v/v formamide) as described in Nuzzo et al. (2017). DGGE image analysis was used to calculate the community richness (Rr) and community organization (Co) indexes (Nuzzo et al., 2017).

2.6 Analysis of the biofilm community via Illumina sequencing of 16S rRNA genes and statistical analysis

DNA extracted from biofilms of selected samples was subjected to sequencing in order to investigate the bacterial community composition. The selection of samples was based on the observed colonization dynamics in terms of i) amount of bacterial biofilm formed and, ii) changes of the community richness and organization over time, detected via PCR-DGGE analysis (section 3.1), as well as on the observed reductive dechlorination of MP-sorbed PCBs (section 3.3). V1-V3 hypervariable regions of 16S rRNA were PCR amplified and sequenced on Illumina Miseq with 300 bp paired-end approach (BMR genomics, Padova, Italy).

The 16S rRNA gene raw sequences were processed using the open source software pipeline Quantitative Insights Into Microbial Ecology 2 (QIIME 2) version 2017.12 (<http://qiime2.org>).

SILVA was used as reference database to classify the representative sequences from our dataset, and Jaccard distance matrices as beta-diversity measures. The Principal Coordinate Analysis (PCoA) plot was created in R (version 3.5.1) through community ecology vegan package. The significance of separation among study groups was determined by permutation test with pseudo- F ratios using the function `adonis` of vegan package. Significant differences in the relative abundance of Dehalococcoidia between MPs groups – pristine and PCB-contaminated – were evaluated with the Wilcoxon test.

3 Results

3.1 Colonization dynamics of different MPs

The colonization dynamics was evaluated during a 28-weeks experiment using different approaches: i) quantification of the Bacterial 16S rRNA genes via qPCR, as an approximation of cell density on the MPs surface (16S rRNA gene copies per cm² of MPs), ii) cell/biofilm staining based on crystal violet, which targets both the bacterial cells and the extracellular polymeric substances (EPS) they produce when growing attached to surfaces, and iii) quantification of reducing sugars occurring in the polysaccharide fraction of EPS.

The qPCR analysis revealed that all MPs were rapidly colonized by the bacterial community within the first 2 weeks of incubation, without any remarkable difference between the PCB-contaminated MPs and the pristine (not-contaminated) MPs (Figure 1A,D). The highest cell density was found on PVC pellets, which reached approximately 10¹⁰ 16S rRNA gene copies per cm² of MPs, followed by PE (approximately 10⁹ 16S rRNA copies cm⁻²), and then PET, PP and PS (approximately 10⁸ 16S rRNA copies cm⁻²). Since the surface area of MPs was estimated using the geometry of each plastic pellet, i.e., not considering the irregularities of the shapes and the porosities, the calculated superficial cell density may be an underestimation of the actual microbial colonization. No remarkable cell density changes were observed along the rest

of the incubation, indicating that cells attachment and growth were not taking place and/or that were balanced by cells death and detachment from the surface (McDougald et al., 2012).

Staining with crystal violet (Figure 1B,E) and quantification of the reducing sugars (Figure 1C,F) revealed a biofilm increase up to weeks 10-15 on PE, PS, PP and PET, and up to the end of incubation (week 28) on PVC, without remarkable differences in the presence and absence of the sorbed pollutants. The biofilm increase over time may indicate the occurrence of a biofilm maturation phase, taking place after a rapid cells adhesion and growth. The biofilm amount then changed less markedly, showing slight decreases followed by small increases that may indicate the succession of detachment and regrowth events involving portions of the biofilm, which are typical of these dynamic structures (McDougald et al., 2012). The greatest biomass concentration, approximately one order of magnitude higher than on the other MP types, was detected on PVC MPs also according to crystal violet and quantification of reducing sugars and was significantly higher than that on other MP types ($P < 0.05$, Tukey test) according to all measured parameters.

A preliminary investigation of the evolution of the biofilm community established on the different MPs was carried out through PCR-DGGE analysis of the 16S rRNA genes. PCR-DGGE profiles (Figure S1) were mainly used to evaluate if major changes were taking place over time in the community developed on each MP type, rather than for an in-depth assessment of the community diversity and structure or to compare the biofilm communities on the different MP types. The following parameters were considered: i) community richness (Rr), i.e., the number of species, and ii) community organization (Co), which indicates if the individuals of the population are homogeneously distributed between the species (even community) or if individuals belonging to few species are predominant over those belonging to all others species (uneven community).

The Rr and Co varied markedly between the different type of MPs, but not between the MPs contaminated by PCBs and those not contaminated (Figure S2). In particular, the biofilm communities on PET, PS and PP did not change remarkably in Rr and Co over time, and were characterized by an average Rr of approximately 25 and 18, for the pristine and PCB-contaminated MPs, respectively. The Co resulted to be approximately 25 or lower for all this plastic samples (PET, PS and PP, contaminated and pristine), which indicates a high degree of community evenness. In contrast, a remarkable change of the biofilm community appeared to occur on PE during the first 10 weeks of incubation, where a noteworthy reduction of Rr and a substantial increase of Co were observed. This indicates that a fraction of the biofilm microbial community on the PE surface was able to become dominant over time, causing a loss of biodiversity. Conversely, an increase of Rr and limited fluctuations of the Co were observed on PVC over time. This indicates that an increase of biodiversity took place without affecting the functional organization of the community.

3.2 Composition of the biofilm community on different MPs

The composition of the biofilm communities was then investigated more in detail through Illumina sequencing of the 16S rRNA genes. Samples were selected based on the observed colonization dynamics in terms of i) amount of bacterial biofilm formed and, ii) changes of the community richness and organization over time (section 3.1), as well as on the observed reductive dechlorination of MP-sorbed PCBs (section 3.3). In particular, the biofilm community composition was investigated for all samples after 2 weeks of incubation, since at this sampling point all MP types were colonized by the maximum cells concentration, which then remained almost constant (Figure 1A,D), and a remarkable reductive dechlorination of sorbed PCBs was observed on all MP types (section 3.3, Figure 5); in addition, the final sampling point (28 weeks of incubation) was analysed only for PE and PVC pellets, since substantial changes in Rr and Co

were observed between 2 and 28 weeks of incubation only on these materials (Figure S2). The PCB-dechlorinating marine culture inoculated in the microcosms was also analysed to investigate if adhesion and biofilm formation on MPs involves specific members of the community or the whole population. In addition, the sediment present in the microcosms supplemented with PCB-contaminated MPs was analysed to evaluate any differences between the microbial communities adhering to the MPs and those associated to the sediment particles. At the phylum level (Figure 2), the microbial inoculum mainly consisted of Proteobacteria (58%), Chloroflexi (30%) and, at lower extent, Firmicutes (6%). After 2 weeks of incubation, Firmicutes became dominant in the biofilm communities on all MP types, having a relative abundance that ranged from 88% (pristine PS) to 49% (PCB-contaminated PVC). Conversely, Proteobacteria represented a minor fraction of the biofilm communities, ranging from 1 to 27%, and the fraction of Chloroflexi decreased on all plastic pellets (0.3-13%), except for PVC contaminated by PCBs (45%). Plastic biofilm communities also included, at quite high relative abundances, the phyla Lentisphaerae (0-26%), Deinococcus-Thermus (0-7%), Actinobacteria (0-13%) and Acidobacteria (0-9%), which were not detected in the inoculum or represented less than 2% of the original microbial community.

Relevant changes in composition were detected after 28 weeks of incubation on PE and PVC, when Chloroflexi became more abundant and Firmicutes decreased.

The bacterial communities associated to the sediment surrounding the different PCB-contaminated MPs were very similar to each other and mainly consisted of bacteria belonging to the phyla Firmicutes (30-43%), Chloroflexi (20-28%), Proteobacteria (12-22%), and to less extent, Acidobacteria (6-9%), Synergistetes (4-8%), Lentisphaerae (3-7%), and Actinobacteria (2-3%). These bacterial communities therefore remarkably differed from the microbial inoculum and those colonizing the MPs.

At lower taxonomic levels (Table S1), the original marine microbial community was dominated by bacteria of the genera *Sulfurovum* (20.4%) and *Sulfurimonas* (15.5%), which are both typically sulphur oxidizing, nitrate-reducing bacteria, *Dethiosulfatibacter* (5.3%), a sulphur and thiosulfate-reducing genus, *Magnetovibrio* (3.6%), which includes members capable of anaerobic thiosulfate oxidation using nitrous oxide, *Celeribacter* and another member (unidentified genus) of the *Rhodobacteraceae* family (12.0%) and *Dehalobium* (17.7%), an organohalide respiring Dehalococcoidia, along with other non-dehalorespiring members of the Chloroflexi phylum (unidentified genus, *Anaerolineacea* family, 12.8%). Most of these genera were not detected in the biofilm communities grown on pristine MPs, or they were at much lower abundances. After 2 weeks of incubation, the biofilm present on PS and PVC was dominated by a Firmicutes bacterium of the *Peptostreptococcaceae* family (unidentified genus), which represented 42.4% and 37.6% of the community, respectively. This bacterium was also present at high relative abundance on PE (19.4%) and, at lower abundances, on PET (9.4%) and PP (8.5%) after 2 weeks of incubation. The microbial community on PE and PP after 2 weeks of incubation was mainly dominated by a different Firmicutes bacterium belonging to the SRB2 family of *Thermoanaerobacterales* (unidentified genus), which represented approximately 29% of the biofilm community on both MPs types. The same bacterium was also detected at high abundance on PET (17.6%) and at much lower abundance on PS (5.6%), while was not detected on PVC. A member of the *Clostridiaceae 1* family (unidentified genus) was also dominant or present at relatively high abundances on PET (20.5%), PP (21.0%) and PS (16.4%), whereas the community on PVC included at high abundances a member of Clostridiales (22.5%), *Celeribacter* (15.0%) and a member of Dehalococcoidia (10.0%). Other community members that enriched on pristine MPs after 2 weeks of incubation were an uncultured *Coriobacteriaceae* (unidentified genus) in PVC (12.5%), PE (6.6%), PET (5.3%) and PP (4.3%), a member of Deinococci in PP (6.8%), PE (5.9%) and PS (5.3%), *Sedimentibacter* in PE (10.2%),

PET (7.3%), PP (6.3%) and PS (3.1%), and *Clostridium sensu strictu 1* only in PET (7.9%) and PS (5.4%).

After 28 weeks of incubation, the SRB2 family of *Thermoanaerobacterales* and *Sedimentibacter* were lost by the biofilm community on PE, where *Peptostreptococcaceae* further enriched (40.3%) along with *Anaeroplasma* (6.5%) and *Anarolinaceae* (17.9%). A remarkable change was also observed in the biofilm composition of PVC after 28 weeks of incubation, when the uncultured *Coriobacteriaceae*, Dehalococcidia, Clostridiales and *Celeribacter* drastically reduced, and uncultured *Anaerolinaceae* remarkably enriched (35.4%).

The presence of sorbed PCBs had very limited effects on the overall composition of the biofilm on MPs, resulting mainly in the enrichment of the class Dehalococcidia, i.e., the organohalide respiring Chloroflexi members of the community (Figure 3). This behavior appears even more marked in the case of PVC, where the increase of Dehalococcidia in the biofilm community in the presence of PCBs reaches the statistics significance (P-value = 0.05, Wilcoxon test).

The PCoA analysis based on Jaccard similarity index was then used to assess the overall OTU-level compositional differences in the microbial communities adhering to the different plastic types (PE, PET, PS, PP, PVC) and sediment. The data show a clear separation of microbial community profiles by MPs type and between MPs and the surrounding sediment (P-value < 0.03, permutation test with pseudo-*F* ratio) (Figure 4A). Permutation test with pseudo-*F* ratio of the significance of pairwise separation between each MP type and MPs and the sediment is reported in Table S2. The PCoA analysis also demonstrated that the communities on PCB-contaminated and pristine MPs did not significantly differ between each other (P-value = 0.4) (Figure 4B). The sorption of PCBs on MPs therefore did not significantly change the taxonomic composition of the overall microbial community adhering to them.

3.3 PCB dechlorination

No dechlorination of MP-sorbed PCBs was observed in sterile controls during the 28 weeks of incubation. On the contrary, a significant reductive dechlorination of PCBs associated to all different types of MPs was observed in the microcosms inoculated with the PCB-dechlorinating culture after only two weeks of incubation, when the average number of chlorines per biphenyl molecule was reduced from 5.2 to 4.8 on PP, 4.7 on PE, 4.6 on PET, 4.4 on PS and 4.3 on PVC (Figure 5), corresponding to dechlorination rates of 35.2 ± 1.9 (PP), 44.6 ± 2.6 (PE), 56.8 ± 6.2 (PET), 57.8 ± 5.4 (PS) and 61.2 ± 5.8 (PVC) $\mu\text{moles of Cl removed kg}_{\text{MP}}^{-1} \text{ week}^{-1}$. The reductive dechlorination of sorbed PCBs then proceeded more slowly, leading to an average number of chlorines per biphenyl molecules after 28 weeks of incubation in the range 4.4-3.9. Remarkably, the biotransformation of PCBs sorbed on MPs was faster than that of PCBs sorbed on sediment, which started only after 10 weeks of incubation and proceeded with a dechlorination rate of 33.9 ± 9.1 $\mu\text{moles of Cl removed kg}_{\text{sediment}}^{-1} \text{ week}^{-1}$ until week 28, when the average number of chlorines per biphenyl molecule was reduced to 4.0 (Figure 5).

In general, highly similar dechlorination patterns were observed for PCBs sorbed on different MP types, as indicated by the depletion of the same highly-chlorinated congeners and the accumulation of the same low-chlorinated ones (Figure 6). The lower reduction of the highly-chlorinated congeners and the lower accumulation of the low-chlorinated ones detected on PE and PP pellets at the end of incubation was in accordance with the less extensive dechlorination process occurred on these MPs. The dechlorination patterns of PCBs sorbed on MPs was similar to that observed on PCBs sorbed on sediment (without MPs), with a main difference: the accumulation in the sediment of the co-eluting hexa-/penta-chlorinated congeners 234-35, 235-34 and 2356-24, which did not accumulate on the MPs, and the concomitant accumulation of the tri-chlorinated congeners 25-3 and 24-3 on MPs and not in the sediment. This indicates

that a *meta* and *para* dechlorination of 235-34 and 234-35 to 25-3 and 24-3, respectively, occurred only on MPs.

4 Discussion

Microbial colonization of MPs in the marine environment may have several implications on their fate and sedimentation behavior, the fate of MP-sorbed pollutants and their toxicity and bioavailability to marine organisms (Cole et al., 2011; Wang et al., 2018; Mohamed Nor and Koelmans, 2019). While the colonization of plastic debris by marine microorganisms have been recently reported in seawater (Dussud et al., 2018; Frère et al., 2018; Xu et al., 2019), very limited information is available on plastic colonization in marine sediments (Harrison et al., 2014), which are the ultimate sink, as well as an entry point in the food chain through benthic organisms, of plastic debris and hydrophobic pollutants in the marine environment (Kaiser et al., 2017). Very limited information is available also on the fate of pollutants sorbed on plastics and, in particular, on the potential role of microbial biofilms in the biodegradation/biotransformation of plastic-sorbed pollutants (Wu et al., 2017). In this study, the colonization dynamics of microplastic (MP) pellets of different materials, namely, PE, PET, PS, PP and PVC, either pristine or contaminated with polychlorinated biphenyls (PCBs), was investigated in laboratory microcosms of an anoxic marine sediment and seawater collected from the same site, i.e., under laboratory biogeochemical conditions mimicking those occurring in situ and controlled exposure conditions. The sediment was inoculated with a marine anaerobic microbial community previously selected for its ability to dehalogenate PCBs (Nuzzo et al., 2017), in order to better assess the potential role of microbial reductive dehalogenation processes on the fate of MP-sorbed PCBs. The use of a well-defined source community was also made to better evaluate the possible effects of the MP material and of MP-sorbed pollutants on

the surface colonization dynamics and on structure and composition of the biofilm (Ogonowski et al. 2018).

All MPs were rapidly colonized during the first 2 weeks of incubation, up to a cell density range of 1.0×10^8 to 1.0×10^{10} cells/cm². Harrison et al. (2014) reported a comparable, rapid colonization of low-density PE by bacteria of coastal marine sediments incubated in Petri dishes, that reached a density of 1.0×10^6 to 1.0×10^9 16S rRNA genes DNA per mm² of PE within 7 days. Our work confirms that MPs are a good anthropogenic substrate for colonization by marine sediment microbial communities also under strictly anoxic conditions, which are typically present in situ few millimeters below the sediment surface.

Remarkable differences were observed in terms of colonization of different MP types. In particular, biofilm formation was remarkably higher on PVC pellets than on other MP types, both in terms of cell density reached and amount of EPS produced. Such a higher biofilm growth on PVC is likely the consequence of a higher availability of energy and carbon sources that might have been released from the MPs. Many organic additives are commonly supplemented to plastic materials in order to improve their properties, such as tensile strength, flexibility and durability (Teuten et al., 2009). In particular, PVC is among the plastics containing the highest amounts of additives, mainly plasticizers and stabilizers, that may reach up to 50% w/w of the plastic material. Phthalates (alkyl/aryl esters of 1,2-benzenedicarboxylic acid), as well as low molecular weight, easily biodegradable citrate, adipate and hexanoate esters, are very commonly used plasticizers in PVC (Markarian, 2007; Babinsky, 2006). The microbial degradation of phthalates to methane and carbon dioxide has been also reported under anaerobic conditions (Chang et al., 2005; Liang et al., 2008). All these additives can thus sustain the growth of anaerobic microbes/microbial communities and could have therefore favored a more extensive biofilm growth on PVC pellets.

All MP types selected specific fractions of the inoculated microbial community. The dominant members of the latter were mainly associated with sulphur cycling, a common process taking place in marine sediments (Wasmund et al., 2017), as well as with organohalide respiration, which was a specific feature of the marine microbial community inoculated in the microcosms (Nuzzo et al., 2017). Most of these members were not detected in any of the biofilm communities grown on MPs, which were dominated by taxa mainly including chemoorganotrophic, fermenting bacteria.

Remarkable differences were also observed in terms of composition between the microbial communities associated to the different MP types and the surrounding sediment. Recent studies have demonstrated that the bacterial assemblages that develop on the surface of plastic material in seawater or freshwater systems can significantly differ from those developing on non-plastic substrates, such as wood, cobblestone, cellulose and glass, as well as from the sediment and water bacterial communities (Miao et al., 2019b, Ogonowski et al., 2018; Frère et al., 2018; De Tender et al., 2015). Some of these studies have also reported that the microbial communities developed on PE and PP were quite similar to each other, and more distinct to those associated with PS pellets (Ogonowski et al., 2018; Frère et al., 2018). The differentiation of biofilm communities on different MP types occurred in this study may be attributed to many factors. For example, Ogonowski et al. (2018) recently reported a significant correlation between the substrate hydrophobicity and bacterial composition on different plastic and non-plastic (glass, cellulose) materials. Since the hydrophobicity of the materials tested in this study typically changes in a relatively narrow range (contact angle 80-100; Ogonowski et al., 2018; Gotoh et al., 2011; McGinty and Brittain, 2008), the difference in biofilm composition could be more probably related to other features, such as the presence of different organic and inorganic additives and the different rates they are released from the plastic material. Indeed, the additives released from different MPs may either promote the growth of some microbial

species able to use them as carbon and energy source, or limit the growth and survival of some other species that are sensitive to these chemicals. However, the very high variety of additives used in different plastics, as well as the use of different additives for the same plastic type by different producers, make impossible to identify a typical additives composition for each plastic and thus to relate the specific compositions of the biofilm communities observed in this study to the release of specific chemicals. Interestingly, Klaeger et al. (2019) recently reported that the release of biodegradable residual monomers and oligomers from polymers can lead to a remarkable overestimation of the plastic materials degradation, in particular during the first 19 days of incubation. Indeed, styrene and vinyl chloride monomers have been widely reported to leach from PS and PVC pellets, respectively (Pilevar et al., 2019; Fayad et al., 1997) and are known to be susceptible, as well as other hydrocarbons, to anaerobic biodegradation (Varjani, 2017). This suggests that the leaching of residual monomers and/or oligomers from the different plastic types may have also contributed to the differentiation of biofilm compositions on the different MPs we observed especially during the first 2 weeks of incubation. This is in agreement also with the selective enrichment of Dehalococcoidia (i.e., of the organohalide respiring Chloroflexi members of the inoculated community) in the biofilm promoted by the sorbed PCBs on all MPs, without any significant change of the overall community.

The PCBs sorbed on the different MP types were rapidly converted into less chlorinated congeners by the anaerobic marine biofilms. Indeed, the rate of PCB dechlorination we observed may be much higher than that potentially taking place in situ, given the extremely low concentration of organohalide respiring bacteria typically present in environmental samples and the specialized enriched community with marked organohalide-respiring activity we used in our study under controlled conditions. However, to our knowledge, this is the first report describing the potential occurrence of reductive dechlorination processes towards PCBs sorbed on MPs. Very few studies investigated the biotransformation/biodegradation of

pollutants sorbed on MPs; Wu et al. (2017) reported for example that colonization of low-density PE pellets increased the biotransformation of dichloro-diphenyl-trichloroethanes (DDTs) and polycyclic aromatic hydrocarbons (PAHs) in marine and river sediments incubated aerobically, although had not significant effect on PCBs. This claims the need of further studies on the biotransformation of pollutants sorbed on MPs under different environmental conditions and by different microbial communities.

Moreover, our study shows that PCBs sorbed on MPs underwent reductive dechlorination much more rapidly than those sorbed on the sediment. While Mato et al. (2001) showed that sorption of organic compounds, such as PCBs and DDE, is approximately two orders of magnitude higher in PP pellets than in marine sediments, probably due to the plastic hydrophobic surfaces, our results suggest that PCBs are more bioavailable to the dechlorinating marine microbes when sorbed on MPs, or that the colonization of MPs might increase the bioavailability of MP-sorbed PCBs, e.g., through biosurfactants production, better than of the sediment-sorbed ones. PCB dechlorination was more extensive on PVC, PS and PET, followed by PE and PP, which might indicate a slight different degree of PCB bioavailability on different MP materials. The most abundant highly-chlorinated PCB congeners were extensively converted into less chlorinated products, that are commonly less toxic and less hydrophobic, thus potentially more bioavailable and less prone to bioaccumulation. The toxicity of contaminated MPs is potentially due to the combination of several factors, e.g., the intrinsic toxicity of the sorbed pollutants, their bioavailability and capability to bioaccumulate; the observed change in congener composition of the PCB mixture associated to MPs might thus have remarkable effects on the overall toxicity associated to polluted MPs, that deserves more investigations.

5 Conclusions

The colonization dynamics of different MP pellet types (PE, PET, PS, PP, and PVC) by an anaerobic marine microbial community was investigated for the first time in anoxic marine sediments, which represent the main sink for MPs in the marine environment. This study showed that microbial colonization took place very rapidly on all MPs and that the biofilm composition differed significantly between the five plastic types and from the surrounding sediment community. The colonization of MPs is thus a selective process that may depend on several factors, such as the different properties of polymer surface and the presence of different additives and/or different residual monomers/oligomers that may be released from the plastics and promote or limit selectively the growth of distinct bacterial species. However, the factors driving the enrichment of different biofilm communities on the MPs may be multiple and diverse and need further investigation. This work also showed that PCBs sorbed on different MP types can be dehalogenated much faster than those sorbed on sediments, suggesting that they are highly bioavailable. In addition, their susceptibility to be bioconverted to even more bioavailable, although less toxic, low-chlorinated PCBs indicate that microbial biofilms growing on contaminated MPs, by changing the congener composition of the sorbed PCB mixture, might remarkably affect the toxicity of contaminated MPs in the marine environment, as well as the uptake and bioaccumulation of MP-associated PCBs by marine organisms. Further investigation should be carried out in order to assess ecotoxicity changes associated to the biotransformation of PCBs sorbed to MPs particles.

Acknowledgements

This research was supported through the Joint Programming Initiative Healthy and Productive Seas and Oceans (JPI- Oceans) PLASTOX project (grant agreement No 696324) by the Italian Ministry of Education, University and Research (MIUR, Project Grant 6962 31/03/2015).

We thank Dr. Stefania Piarulli, Dr. Joanne Wong and Prof. Laura Airoidi (University of Bologna) for assistance in sampling of sediment samples from Piailassa Baiona (Ravenna, Italy).

References

- Andrady, A.L., 2011. Microplastics in the marine environment. *Marine pollution bulletin*, 62(8), 1596-1605.
- Andrady, A.L., Neal, M.A., 2009. Applications and societal benefits of plastics. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1526), 1977-1984.
- Arias-Andres, M., Kettner, M.T., Miki, T., Grossart, H.-P., 2018. Microplastics: New substrates for heterotrophic activity contribute to altering organic matter cycles in aquatic ecosystems. *Science of the Total Environment*, 635, 1152–1159.
- Babinsky, R., 2006. PVC additives: a global review. *Plastics, Additives and Compounding*, 8(1), 38-40.
- Bailey, M.J., 1988. A note on the use of dinitrosalicylic acid for determining the products of enzymatic reactions. *Applied Microbiology and Biotechnology*, 29(5), 494-496.
- Beckingham, B., Ghosh, U., 2017. Differential bioavailability of polychlorinated biphenyls associated with environmental particles: Microplastic in comparison to wood, coal and biochar. *Environmental Pollution*, 220, 150-158.
- Bergmann, M., Wirzberger, V., Krumpfen, T., Lorenz, C., Primpke, S., Tekman, M. B., Gerdtz, G., 2017. High quantities of microplastic in Arctic deep-sea sediments from the HAUSGARTEN observatory. *Environmental science & technology*, 51(19), 11000-11010.
- Besseling, E., Wegner, A., Foekema, E.M., Van Den Heuvel-Greve, M.J., Koelmans, A.A., 2013. Effects of microplastic on fitness and PCB bioaccumulation by the lugworm *Arenicola marina* (L.). *Environmental science and technology*, 47(1), 593-600.

- Burton, E., Yakandawala, N., LoVetri, K., Madhyastha, M.S., 2007. A microplate spectrofluorometric assay for bacterial biofilms. *Journal of industrial microbiology and biotechnology*, 34(1), 1-4.
- Carpenter, E.J., Anderson, S.J., Harvey, G.R., Miklas, H.P., Peck, B.B., 1972. Polystyrene spherules in coastal waters. *Science*, 178(4062), 749-750.
- Chang, B.V., Liao, C.S., Yuan, S.Y., 2005. Anaerobic degradation of diethyl phthalate, di-n-butyl phthalate, and di-(2-ethylhexyl) phthalate from river sediment in Taiwan. *Chemosphere*, 58(11), 1601-1607.
- Chubarenko, I., Bagaev, A., Zobkov, M., Esiukova, E., 2016. On some physical and dynamical properties of microplastic particles in marine environment. *Marine Pollution Bulletin*, 108(1-2), 105-112.
- Claessens, M., De Meester, S., Van Landuyt, L., De Clerck, K., Janssen, C.R., 2011. Occurrence and distribution of microplastics in marine sediments along the Belgian coast. *Marine Pollution Bulletin*, 62(10), 2199-2204.
- Cole, M., Lindeque, P., Halsband, C., Galloway, T.S., 2011. Microplastics as contaminants in the marine environment: a review. *Marine pollution bulletin*, 62(12), 2588-2597.
- Costerton, J.W., Lewandowski, Z., Caldwell, D.E., Korber, D.R., Lappin-Scott, H.M., 1995. Microbial biofilms. *Annual review of microbiology*, 49(1), 711-745.
- De Tender, C.A., Devriese, L.I., Haegeman, A., Maes, S., Ruttink, T., Dawyndt, P., 2015. Bacterial community profiling of plastic litter in the Belgian part of the North Sea. *Environmental science and technology*, 49(16), 9629-9638.
- Dussud, C., Hudec, C., George, M., Fabre, P., Higgs, P., Bruzaud, S., Delort, A.M., Eyheraguibel, B., Meistertzheim, A.L., Jacquin, J., Cheng, J., Callac N., Odobel, C., Rabouille, S., Ghiglione, J.F., 2018. Colonization of non-biodegradable and biodegradable plastics by marine microorganisms. *Frontiers in microbiology*, 9, 1571.

- Fava, F., Agathos, S.N., 2006. Uncertainty and research needs in the area of the biological restoration of contaminated sediments. In *Assessment and remediation of contaminated sediments* (pp. 239-246). Springer, Dordrecht.
- Fava, F., Zanaroli, G., Young, L.Y., 2003. Microbial reductive dechlorination of pre-existing PCBs and spiked 2, 3, 4, 5, 6-pentachlorobiphenyl in anaerobic slurries of a contaminated sediment of Venice Lagoon (Italy). *FEMS microbiology ecology*, 44(3), 309-318.
- Fayad, N.M., Sheikheldin, S.Y., Al - Malack, M.H., El - Mubarak, A.H., Khaja, N., 1997. Migration of vinyl chloride monomer (VCM) and additives into PVC bottled drinking water. *Journal of Environmental Science and Health Part A*, 32(4), 1065-1083.
- Frère, L., Maignien, L., Chalopin, M., Huvet, A., Rinnert, E., Morrison, H., Kerninon, S., Cassone, A.-L, Lambert, C., Reveillaud, J., Paul-Pont, I., 2018. Microplastic bacterial communities in the Bay of Brest: Influence of polymer type and size. *Environmental pollution*, 242, 614-625.
- Frias, J.P.G.L., Nash, R., 2019. Microplastics: Finding a consensus on the definition. *Marine pollution bulletin*, 138, 145-147.
- Gotoh, K., Yasukawa, A., Kobayashi, Y., 2011. Wettability characteristics of poly(ethylene terephthalate) films treated by atmospheric pressure plasma and ultraviolet excimer light. *Polymer Journal*, 43, 545-551.
- Harrison, J.P., Schratzberger, M., Sapp, M., Osborn, A.M., 2014. Rapid bacterial colonization of low-density polyethylene microplastics in coastal sediment microcosms. *BMC microbiology*, 14(1), 232.
- Hashmi, M.Z., Zhang, J., Li, B., Su, X., Tariq, M., Ahmad, N., Malik, R.N., Ullah, K., Chen, C., Shen, C., 2017. Effects of structurally different noncoplanar and coplanar PCBs on HELF cell proliferation, cell cycle, and potential molecular mechanisms. *Environmental toxicology*, 32(4), 1183-1190.

- Haward, M., 2018. Plastic pollution of the world's seas and oceans as a contemporary challenge in ocean governance. *Nature communications*, 9(1), 667.
- Hawker, D.W., Connell, D.W., 1988. Octanol-water partition coefficients of polychlorinated biphenyls congeners. *Environmental Science & Technology*, 22, 382-387.
- Hirai, H., Takada, H., Ogata, Y., Yamashita, R., Mizukawa, K., Saha, M., Kwan, C., Moore, C., Gray, H., Laursen, D., Zettler, E.R., Farrington, J.W., Reddy, C.M., Peacock, E.E., Ward, M.W., 2011. Organic micropollutants in marine plastics debris from the open ocean and remote and urban beaches. *Marine Pollution Bulletin*, 62(8), 1683-1692.
- Hong, S.H., Shim, W.J., Hong, L., 2017. Methods of analysing chemicals associated with microplastics: a review. *Analytical Methods*, 9(9), 1361-1368.
- Kaiser, D., Kowalski, N., Waniek, J.J., 2017. Effects of biofouling on the sinking behavior of microplastics. *Environmental Research Letters*, 12(12), 124003.
- Klaeger, F., Tagg, A.S., Otto, S., Bienmüller, M., Sartorius, I., Labrenz, M., 2019. Residual monomer content affects the interpretation of plastic degradation. *Scientific reports*, 9(1), 2120.
- Koelmans, A.A., Besseling, E., Wegner, A., Foekema, E.M., 2013. Plastic as a carrier of POPs to aquatic organisms: a model analysis. *Environmental science and technology*, 47(14), 7812-7820.
- Li, W., Zhang, Y., Wu, N., Zhao, Z., Xu, W.A., Ma, Y., Niu, Z., 2019. Colonization Characteristics of Bacterial Communities on Plastic Debris Influenced by Environmental Factors and Polymer Types in the Haihe Estuary of Bohai Bay, China. *Environmental science & technology*, 53(18), 10763-10773.
- Liang, D.W., Zhang, T., Fang, H.H., He, J., 2008. Phthalates biodegradation in the environment. *Applied microbiology and Biotechnology*, 80(2), 183.

- Markarian, J., 2007. PVC additives–What lies ahead? *Plastics, Additives and Compounding*, 9(6), 22-25.
- Mathalon, A., Hill, P., 2014. Microplastic fibers in the intertidal ecosystem surrounding Halifax Harbor, Nova Scotia. *Marine pollution bulletin*, 81(1), 69-79.
- Mato, Y., Isobe, T., Takada, H., Kanehiro, H., Ohtake, C., Kaminuma, T., 2001. Plastic resin pellets as a transport medium for toxic chemicals in the marine environment. *Environmental science and technology*, 35(2), 318-324.
- McDermid, K.J., McMullen, T.L., 2004. Quantitative analysis of small-plastic debris on beaches in the Hawaiian archipelago. *Marine pollution bulletin*, 48(7-8), 790-794.
- McDougald, D., Rice, S.A., Barraud, N., Steinberg, P.D., Kjelleberg, S., 2012. Should we stay or should we go: mechanisms and ecological consequences for biofilm dispersal. *Nature Reviews Microbiology*, 10, 39–50.
- McGinty, K.M., Brittain, W.J., 2008. Hydrophilic surface modification of poly (vinyl chloride) film and tubing using physisorbed free radical grafting technique. *Polymer*, 49(20), 4350-4357.
- Miao, L., Hou, J., You, G., Liu, Z., Liu, S., Li, T., Mo, Y., Guo, S., Qu, H., 2019a. Acute effects of nanoplastics and microplastics on periphytic biofilms depending on particle size, concentration and surface modification. *Environmental Pollution*, 255, 113300.
- Miao, L., Wang, P., Hou, J., Yao, Y., Liu, Z., Liu, S., Li, T., 2019b. Distinct community structure and microbial functions of biofilms colonizing microplastics. *Science of the Total Environment*, 650, 2395-2402.
- Mohamed Nor, N.H., Koelmans, A.A., 2019. Transfer of PCBs from microplastics under simulated gut fluid conditions is biphasic and reversible. *Environmental science and technology*, 53(4), 1874-1883.

- Morét-Ferguson, S., Law, K.L., Proskurowski, G., Murphy, E.K., Peacock, E.E., Reddy, C.M., 2010. The size, mass, and composition of plastic debris in the western North Atlantic Ocean. *Marine Pollution Bulletin*, 60(10), 1873-1878.
- Nuzzo, A., Hosseinkhani, B., Boon, N., Zanaroli, G., Fava, F., 2017. Impact of bio-palladium nanoparticles (bio-Pd NPs) on the activity and structure of a marine microbial community. *Environmental Pollution*, 220, 1068-1078.
- Ogonowski, M., Motiei, A., Ininbergs, K., Hell, E., Gerdes, Z., Udekwu, K.I., Bacsik, Z., Gorokhova, E., 2018. Evidence for selective bacterial community structuring on microplastics. *Environmental microbiology*, 20(8), 2796-2808.
- Pilevar, Z., Bahrami, A., Beikzadeh, S., Hosseini, H., Jafari, S.M., 2019. Migration of styrene monomer from polystyrene packaging materials into foods: characterization and safety evaluation. *Trends in Food Science and Technology*.
- PlasticsEurope, 2018. *Plastics – the Facts 2017. An Analysis of European Plastics Production, Demand and Waste Data*. Brussels, Belgium.
- Rummel, C.D., Jahnke, A., Gorokhova, E., Kühnel, D., Schmitt-Jansen, M., 2017. Impacts of biofilm formation on the fate and potential effects of microplastic in the aquatic environment. *Environmental Science and Technology Letters*, 4(7), 258-267.
- Ryan, P.G., Moore, C.J., van Franeker, J.A., Moloney, C.L., 2009. Monitoring the abundance of plastic debris in the marine environment. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1526), 1999-2012.
- Sass, A.M., Sass, H., Coolen, M.J., Cypionka, H., Overmann, J., 2001. Microbial communities in the chemocline of a hypersaline deep-sea basin (Urania basin, Mediterranean Sea). *Appl. Environ. Microbiol.*, 67(12), 5392-5402.

- Tanabe, S., Kannan, N., Subramanian, A., Watanabe, S., Tatsukawa, R., 1987. Highly toxic coplanar PCBs: occurrence, source, persistency and toxic implications to wildlife and humans. *Environmental Pollution*, 47(2), 147-163.
- Taniguchi, S., Colabuono, F.I., Dias, P.S., Oliveira, R., Fisner, M., Turra, A., Izar, G.M., Abessa, D.M.S., Saha, M., Hosoda, J., Yamashita, R., Takada, H., Lourenço, R.A., Magalhães, C.A., Bicego, M.C., Montone, R.C., 2016. Spatial variability in persistent organic pollutants and polycyclic aromatic hydrocarbons found in beach-stranded pellets along the coast of the state of São Paulo, southeastern Brazil. *Marine pollution bulletin*, 106(1-2), 87-94.
- Teuten, E.L., Saquing, J.M., Knappe, D.R., Barlaz, M.A., Jonsson, S., Björn, A., Rowland, S.J., Thompson, R.C., Galloway, T.S., Yamashita, R., Ochi, D., Watanuki, Y., Moore, C., Viet, P.H., Tana, T.S., Prudente, M., Boonyatumanond, R., Zakaria, M.P., Akkhavong, K., Ogata, Y., Hirai, H., Iwasa, S., Mizukawa, K., Hagino, Y., Imamura, A., Saha, M., Takada H., 2009. Transport and release of chemicals from plastics to the environment and to wildlife. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 364(1526), 2027-2045.
- Thomas, C., Lamperta, D., Reible, D., 2014. Remedy performance monitoring at contaminated sediment sites using profiling solid phase microextraction (SPME) polydimethylsiloxane (PDMS) fibers. *Environmental science. Processes & Impacts*, 16, 445-452.
- Thompson, R.C., Olsen, Y., Mitchell, R.P., Davis, A., Rowland, S.J., John, A.W., McGonigle, D., Russell, A.E., 2004. Lost at sea: where is all the plastic? *Science*, 304(5672), 838-838.
- Van Cauwenberghe, L., Devriese, L., Galgani, F., Robbins, J., Janssen, C.R., 2015. Microplastics in sediments: a review of techniques, occurrence and effects. *Marine environmental research*, 111, 5-17.
- Varjani, S.J., 2017. Microbial degradation of petroleum hydrocarbons. *Bioresource Technology*, 223, 277-286.

- Wang, F., Wong, C.S., Chen, D., Lu, X., Wang, F., Zeng, E.Y., 2018. Interaction of toxic chemicals with microplastics: a critical review. *Water research*, 139, 208-219.
- Wasmund, K., Mußmann, M., Loy, A., 2017. The life sulfuric: microbial ecology of sulfur cycling in marine sediments. *Environmental Microbiology Reports*, 9(4), 323-344.
- Wu, C.C., Bao, L.J., Liu, L.Y., Shi, L., Tao, S., Zeng, E.Y., 2017. Impact of polymer colonization on the fate of organic contaminants in sediment. *Environmental science and technology*, 51(18), 10555-10561.
- Xu, X., Wang, S., Gao, F., Li, J., Zheng, L., Sun, C., He, C. Wang, Z. Qu, L., 2019. Marine microplastic-associated bacterial community succession in response to geography, exposure time, and plastic type in China's coastal seawaters. *Marine Pollution Bulletin*, 145, 278-286.
- Zanaroli, G., Negroni, A., Calisti, C., Ruzzi, M., Fava, F., 2011. Selection of commercial hydrolytic enzymes with potential antifouling activity in marine environments. *Enzyme and microbial technology*, 49(6), 574-579.
- Zanaroli, G., Negroni, A., Häggblom, M.M., Fava, F., 2015. Microbial dehalogenation of organohalides in marine and estuarine environments. *Current Opinion in Biotechnology*, 33, 287-295.
- Zettler, E.R., Mincer, T.J., Amaral-Zettler, L.A., 2013. Life in the "Plastisphere": Microbial Communities on Plastic Marine Debris. *Environmental Science and Technology*, 47(13), 7137-7146.
- Ziccardi, L.M., Edgington, A., Hentz, K., Kulacki, K.J., Kane Driscoll, S., 2016. Microplastics as vectors for bioaccumulation of hydrophobic organic chemicals in the marine environment: A state - of - the - science review. *Environmental toxicology and chemistry*, 35(7), 1667-1676.

Figures

Figure 1. Colonization dynamics of PCB-contaminated (A-C) and pristine (D-F) MPs determined via qPCR of 16S rRNA genes (A, D), reducing sugars in the EPS (B, E), and biofilm staining with crystal violet (C, F). Values are the mean of triplicate cultures \pm standard deviation.

Figure 2. Composition at the Phylum level of the inoculated microbial community, the biofilm grown on all PCB-contaminated and pristine MPs after 2 weeks of incubation, the biofilm grown on PCB-contaminated and pristine PE and PVC after 28 weeks of incubation and the microbial community associated to the sediment particles surrounding the PCB-contaminated MPs after 2 weeks of incubation.

Figure 3. Boxplots showing the relative abundance (%) of Dehalococcoidia in pristine and PCB-contaminated MPs. *: P-value = 0.05; Wilcoxon test.

Figure 4. PCoA plots of bacterial communities in the inoculum, in the sediment and associated to MPs (A) and of bacterial communities associated to PCB-contaminated and to pristine MPs (B).

Figure 5. Reductive dechlorination of PCBs sorbed on the sediment (biotic control microcosms) and on the different MP types (biologically active microcosms and sterile controls). Values (average number of chlorines/biphenyl molecule) are the mean of triplicate cultures \pm standard deviation.

Figure 6. PCB congeners and their concentrations on MPs (A) and in the sediment (B) at the beginning and the end of incubation (28 weeks). Value are the mean of triplicate cultures \pm standard deviation.

Supplementary information

Table S1. Composition at the genus level of the inoculated microbial community, of the biofilm grown on all MP types after 2 weeks of incubation and after 28 weeks of incubation on PE and PVC MPs, and of the microbial community associated to the sediment particles.

Table S2. Results of permutation test with pseudo-F ratio statistics applied to ordination analysis based on Jaccard similarity index (related to Figure 4A).

Figure S1. PCR-DGGE gels of the 16S rRNA genes of the inoculated marine microbial culture (A) and the biofilm microbial community grown on pristine MPs (A) and on PCB-contaminated-MPs (B).

Figure S2. Evolution of the biofilm community richness (Rr) and organization (Co) over time on PCB-contaminated (A) and pristine (B) MPs, based on PCR-DGGE analysis of 16S rRNA genes (Figure S1).

Figure 1




Figure 2




Figure 3





Figure 4

A



B




Figure 5




Figure 6

