

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

A rescue transcatheter solution for early sutureless basal ring infolding

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version: A rescue transcatheter solution for early sutureless basal ring infolding / Loforte A.; Comentale G.; Coppola G.; Amodio C.; Botta L.; Saia F.; Taglieri N.; Marrozzini C.; Savini C.; Pacini D.. - In: JOURNAL OF CARDIAC SURGERY. - ISSN 0886-0440. - STAMPA. - 37:3(2022), pp. 697-699. [10.1111/jocs.16196]

Availability: This version is available at: https://hdl.handle.net/11585/902746 since: 2022-11-15

Published:

DOI: http://doi.org/10.1111/jocs.16196

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Loforte A, Comentale G, Coppola G, Amodio C, Botta L, Saia F, Taglieri N, Marrozzini C, Savini C, Pacini D.

A rescue transcatheter solution for early sutureless basal ring infolding.

J Card Surg. 2022 Mar; 37(3): 697-699.

The final published version is available online at: <u>10.1111/jocs.16196</u>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<u>https://cris.unibo.it/</u>)

When citing, please refer to the published version.

A RESCUE TRANSCATHETER SOLUTION FOR EARLY SUTURELESS BASAL RING INFOLDING

Running Head: RESCUE TAVI IN SUTURELESS VALVE

Antonio Loforte¹*, MD, PhD; Giuseppe Comentale^{1,3}, MD; Giuditta Coppola, MD¹; Ciro Amodio, MD¹; Luca Botta, MD¹; Francesco Saia, MD, PhD²; Nevio Taglieri, MD²; Cinzia Marrozzini, MD, PhD²; Carlo Savini, MD¹, PhD¹; Davide Pacini, MD, PhD¹

¹Division of Cardiac Surgery, IRCCS Azienda Ospedaliero-Universitaria di Bologna, S. Orsola Hospital, Bologna, Italy;

²Division of Cardiology, IRCCS Azienda Ospedaliero-Universitaria di Bologna, S. Orsola Hospital, Bologna, Italy;

³ Department of Advanced Biomedical Sciences, University of Naples "Federico II", Naples, Italy.

Word Count: 1499

Ethics: IRB approval and Informed consent was obtained.

Disclosures: None.

*Corresponding author:

Dr. Antonio Loforte, MD, PhD

Division of Cardiac Surgery, IRCCS Azienda Ospedaliero-Universitaria di Bologna, S. Orsola Hospital, ALMA Mater Studiorum University of Bologna,

Via Massarenti 9,40138 Bologna, Italy. Tel: +39 051 214 9043/3361. E-mail: antonioloforte@yahoo.it

ABSTRACT

We report the case of an 83-year-old woman treated with a "rescue" valve-in-valve transcatheter aortic valve implantation because of an early basal ring partial collapse of a sutureless valve, probably due to septal hypertrophy.

Keywords: sutureless valve; TAVI

INTRODUCTION

In elderly patients with severe aortic stenosis (AS), transcatheter aortic valve implantation (TAVI) could not be the treatment of choice in presence of marked left ventricular outflow tract (LVOT) calcification or interventricular septum basal hypertrophy [1-2]. Early prosthesis deterioration due to hemodynamic turbulences, residual LVOT obstruction or LVOT-right atrium fistula have been described as potential complications [2-3]. Therefore, in these patients, surgical aortic valve replacement (AVR) should be preferred because it allows to perform a concomitant septal myectomy [1]. AVR with sutureless prostheses has quickly become a very viable option in older patients, especially in those ones with small aortic annulus, because it merges the usefulness of a biological prosthesis with the chance of reducing the cross-clamp and surgical times as well as the rate of postoperative complications compared to the standard AVR [4]. The presence of an anchorage subvalvular stent, however, requires a careful evaluation of the LVOT anatomy in order to avoid postoperative complications or prosthesis dysfunction. Concomitant septal myectomy, indeed, should be considered in presence of LVOT hypertrophy. However, a cut-off value for the interventricular septum dimensions indicating when that procedure is required during surgical AVR is still missing. For these reasons, we report the case of an 83-year-old woman who had to undergo to a "rescue" valve-in-valve (V-in-V) TAVI because she experimented an early basal ring partial collapse of the sutureless valve after minimally invasive AVR probably related to the untreated septal hypertrophy.

CASE PRESENTATION

An 83-year-old woman with a severe symptomatic AS was admitted to our hospital for surgical AVR as the HeartTeam considered TAVI at high risk due to basal septum hypertrophy (Interventricular septum of 16 mm; Left ventricle end-diastole and end-systole diameters: 38 mm and 22 mm, respectively). Her medical history highlighted very high cardiovascular risk: arterial hypertension, dyslipidemia, 1st grade obesity, peripheral artery disease, hyperthyroidism, and chronic atrial fibrillation. Due to the patient's age and comorbidities, a minimally invasive surgical AVR (upper J mini sternotomy) with sutureless prosthesis was planned. Cardiopulmonary bypass was established through standard Aortic-Right atrium cannulation and Custodiol[®] Solution was antegrade administered to achieve the cardioplegic arrest. A Perceval S, size M (Sorin[®], Salluggia, Italy) sutureless valve was implanted according to the manufactures' indications after an

accurate decalcification of the aortic annulus. In order to prevent atrio-ventricular (AV) conduction system damage and considering the absence of significant narrowing of the LVOT, the septal myectomy was not performed. No complications occurred during the valve deployment and the implanted prosthesis appeared well-functioning at the intraoperative transesophageal echocardiography. Cardiopulmonary bypass weaning as well as the surgery course were uneventful.

Postoperative course was unremarkable until the 3rd day after the surgery when she presented type III AV block and progressive worsening of hemodynamics and respiratory function. The echocardiogram (Figure 1A) revealed the presence of moderate aortic regurgitation and high pressures gradients through the implanted valve (max/mean gradients: 52/31 mmHg; Prosthetic Valve Area: 1,15 cm²) that appeared to be "heart shaped" as a result of a partial collapse of the basal ring at the level of the right coronary sinus; this structural distortion jailed the right coronary cusp and caused moderate to severe aortic regurgitation.

Urgent transcatheter balloon valvuloplasty was successfully performed to reshape the valve and to stabilize the hemodynamics but the patients developed the same imaging and clinical pattern nine days after. The Angio-CT scan (Figure 1 B-C) confirmed the partial infolding of the valve basal ring, therefore, after a careful evaluation by the HeartTeam, it was decided to perform a "rescue" transfemoral V-in-V TAVI (Figure 1 D) with a Sapien 23 mm valve (Edwards Lifesciences LLC, Irvine, CA, USA).

The procedure was successfully performed, and the patient was discharged in good clinical conditions on 16th postoperative day. The transcatheter aortic prosthesis remained still competent during the follow-up and no signs of structural distortion or disfunction are present at 4th year of follow-up.

DISCUSSION

Basal stent infolding is a very rare complication after sutureless valves implantation and it's mainly associated with a wrong sizing or prosthesis deployment [5,6]. No cases of basal ring infolding relapse associated with a mild left ventricle septal hypertrophy have been reported. Such complication was not observed even in the premarket approval trials among over than 700 patients with a freedom from valve reoperation of 97.83% at 1 year [7].

Structural malfunctioning is usually detected during intraoperative echocardiographic check or at the time of implantation during balloon inflating [5-8] therefore it can be easily fixed before leaving the operative room.

On the contrary, spontaneous distortion in the early postoperative period is an extremely rare event and could even require an emergency re-intervention.

In our case, the reason of so much delayed and spontaneous stent recoil remains unclear. We suppose that the preoperative septal hypertrophy, increased by myocardial postoperative edema and by the abnormal motion of the interventricular septum related to the cardiopulmonary bypass and to the cardioplegic solution, might have played a significant role. This condition probably modified the symmetry of the aortic annulus in a such extreme way that even the postoperative balloon valvuloplasty didn't solve the problem and the additional radial force of the transcatheter Sapien valve was necessary to stabilize the surgical prosthesis. When the underlying anatomy is suitable, balloon valvuloplasty, indeed, is almost always effective [6-8].

Left ventricle hypertrophy, therefore, can represent a mined field that can hamper even a theoretically "lowerrisk" procedure. The anatomy of LVOT, indeed, have to be carefully considered not only when TAVI with prosthesis including subvalvular stents is planned but also in surgical procedures with sutureless valves. Surgical AVR, indeed, requires the extracorporeal circulation and cardioplegic arrest of the heart: these two conditions can deeply modify the anatomy of the heart in the early postoperative hours and can make wrong all those decisions taken during the preoperative planning. For this reason, during sutureless valve implantation, partial septal myectomy should be considered even in case of borderline septal hypertrophy in order to avoid structural valve malfunctioning.

REFERENCES

- Vahanian A, Beyersdorf F, Praz F, et al. 2021 ESC/EACTS Guidelines for the management of valvular heart disease. Eur J Cardiothorac Surg 2021; 28:ezab389.
- Christ T, Dohmen PM, Laule M, Stangl K, Konertz W. Sutureless Aortic Valve Replacement in a Patient with Transfemoral Aortic Valve Replacement and Left Ventricular Hypertrophy. Thorac Cardiovasc Surg Rep 2016; 5(1):21-23.
- 3. Al Ayouby A, Gilard M, Hebert T, Didier R. Case report: iatrogenic left ventricular outflow tract to right atrium fistula after trans-femoral transcatheter aortic valve implantation associated with asymmetric septal hypertrophy. Eur Heart J Case Rep 2021; 5(3):ytab020.
- Hanedan MO, Yuruk MA, Parlar AI, et al. Sutureless versus Conventional Aortic Valve Replacement: Outcomes in 70 High-Risk Patients Undergoing Concomitant Cardiac Procedures. Tex Heart Inst J 2018; 45(1):11-16.
- 5. Pfeiffer S, Wilbring M, Kappert U, Santarpino G. The 'entangled' stent: a preventable cause of paravalvular leak of the Perceval bioprosthesis. Interact CardioVasc Thorac Surg. 2017; 25:987–9
- Baert J, Astarci P, Noirhomme P, de Kerchove L. The risk of oversizing with sutureless bioprosthesis in small aortic annulus. J Thorac Cardiovasc Surg 2017; 153:270–2
- Suri RM, Javadikasgari H, Heimansohn DA, et al. Prospective US investigational device exemption trial of a sutureless aortic bioprosthesis: One-year outcomes. J Thorac Cardiovasc Surg 2019;157(5):1773-1782.e3.
- 8. Di Eusanio M, Saia F, Pellicciari G, et al. In the era of the valve-in-valve: is transcatheter aortic valve implantation (TAVI) in sutureless valves feasible?. Ann Cardiothorac Surg 2015;4(2):214-7

FIGURE LEGEND

FIGURE 1. (A) Postoperative echocardiogram that shows the basal ring deformation at the level of hypertrophic septum (on the left) and the presence of the presence of moderate aortic regurgitation and high pressures gradients (on the right) through the implanted valve. (B) Postoperative Angio-CT scan showing the basal ring relapse of the sutureless valve that appeared to be "heart shaped". (C) 3D Volume rendering of the structural distortion of the aortic prosthesis. (D) Angiogram snapshot showing the right positioning of the Sapien TAVI inside the surgically implanted sutureless valve.