Supplementary materials

A Luciferase Mutant with Improved Brightness and Stability for Whole-Cell Bioluminescent Biosensors and *In Vitro* Biosensing

Maria Maddalena Calabretta 1,2, Denise Gregucci 1,2, Héctor Martínez-Pérez-Cejuela 1,3 and Elisa Michelini 1,2,4,*

- Department of Chemistry "Giacomo Ciamician", Alma Mater Studiorum-University of Bologna, Via Selmi 2, 40126 Bologna, Italy
- ² Center for Applied Biomedical Research (CRBA), IRCCS St. Orsola Hospital, Bologna, 40138, Italy
- ³ CLECEM Group, Department of Analytical Chemistry, University of Valencia, C/Dr. Moliner, 50, 46100, Burjassot, Valencia, Spain
- ⁴ Health Sciences and Technologies Interdepartmental Center for Industrial Research (HSTICIR), University of Bologna, 40126 Bologna, Italy
- * Correspondence: elisa.michelini8@unibo.it; +39 051 20 9 9533



Figure S1. Amino acid alignment of the BgLuc and PpyLuc luciferases. The red highlighted residues indicate the mutated amino acids.




Figure S2: Emission spectra of (a) BgLuc and (b) Luc2 luciferase in HEK293T cells obtained with D-LH₂ substrate in citrate buffer (1.0 mM, pH 5.0) (solid line) and BrightGlo substrate (dotted line).




Figure S3: BgLuc mutant emission spectra obtained (a) at different temperatures and (b) at different pH with the commercial BrightGlo substrate.