

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

A descriptive pharmacokinetic/pharmacodynamic analysis of continuous infusion ceftazidime-avibactam in a case series of critically ill renal patients treated for documented carbapenem-resistant Gram-negative bloodstream infections and/or ventilator-associated pneumonia

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Gatti, M., Pascale, R., Cojutti, P.G., Rinaldi, M., Ambretti, S., Conti, M., et al. (2023). A descriptive pharmacokinetic/pharmacodynamic analysis of continuous infusion ceftazidime-avibactam in a case series of critically ill renal patients treated for documented carbapenem-resistant Gram-negative bloodstream infections and/or ventilator-associated pneumonia. *INTERNATIONAL JOURNAL OF ANTIMICROBIAL AGENTS*, 61(106699), 1-7 [10.1016/j.ijantimicag.2022.106699].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/914534> since: 2023-02-10

*Published:*

DOI: <http://doi.org/10.1016/j.ijantimicag.2022.106699>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

1 **A descriptive pharmacokinetic/pharmacodynamic analysis of continuous infusion**  
2 **ceftazidime/avibactam in a case series of renal critically ill patients treated for documented carbapenem-**  
3 **resistant Gram-negative bloodstream infections and/or ventilator-associated pneumonia**

4

5 **Milo Gatti<sup>1,2\*</sup>, Renato Pascale<sup>1,3</sup>, Pier Giorgio Cojutti<sup>2</sup>, Matteo Rinaldi<sup>3</sup>, Simone Ambretti<sup>4</sup>, Matteo**  
6 **Conti<sup>2</sup>, Sara Tedeschi<sup>1,3</sup>, Maddalena Giannella<sup>1,3</sup>, Pierluigi Viale<sup>1,3</sup>, Federico Pea<sup>1,2</sup>**

7 <sup>1</sup> Department of Medical and Surgical Sciences, Alma Mater Studiorum University of Bologna, Bologna, Italy

8 <sup>2</sup> Clinical Pharmacology Unit, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy

9 <sup>3</sup> Infectious Diseases Unit, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy

10 <sup>4</sup> Division of Microbiology, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Italy

11

12

13 **Corresponding author**

14 Dr. Milo Gatti

15 Department of Medical and Surgical Sciences, Alma Mater Studiorum University of Bologna, Via Massarenti  
16 9, 40138 Bologna, Italy.

17 e-mail address: milo.gatti2@unibo.it

18 phone number: +39 051 214 3627

19 **Running title:** PK/PD of CI ceftazidime/avibactam in CR-GN infections

20

21 **Words of abstract:** 250

22 **Words of text:** 2889

23 **Abstract**

24 **Objectives:** To describe the pharmacokinetic/pharmacodynamic (PK/PD) behaviour of continuous infusion  
25 (CI) ceftazidime/avibactam and the microbiological outcome in a case series of renal critically ill patients  
26 treated for documented carbapenem-resistant Gram-negative (CR-GN) bloodstream infections (BSI) and/or  
27 ventilator-associated pneumonia (VAP).

28 **Methods:** Critically ill patients with different degrees of renal function treated with CI ceftazidime-avibactam  
29 for documented CR-GN infections and underwent therapeutic drug monitoring from April 2021 to March 2022  
30 were retrospectively assessed. Ceftazidime and avibactam concentrations were determined at steady-state, and  
31 the free fraction ( $fC_{ss}$ ) was calculated. The joint PK/PD target of ceftazidime-avibactam was considered as  
32 optimal when both  $C_{ss}/MIC$  ratio for ceftazidime  $\geq 4$  (equivalent to  $100\% fT >_{4 \times MIC}$ ) and  $C_{ss}/C_T$  ratio for  
33 avibactam  $> 1$  (equivalent to  $100\% fT > C_T$  of 4.0 mg/L) were simultaneously achieved (quasi-optimal if only  
34 one of the two was achieved, and suboptimal if none of the two was achieved). Relationship  
35 between ceftazidime-avibactam PK/PD targets and microbiological outcome was assessed.

36 **Results:** Ten patients with documented CR-GN infections (5 BSIs, 4 VAP, and one BSI+VAP) were retrieved.  
37 The joint PK/PD targets of ceftazidime-avibactam were optimal and quasi-optimal in 8 and 2 cases,  
38 respectively. Microbiological failure occurred in two patients (one with VAP and the other with BSI+VAP),  
39 and one of these developed ceftazidime-avibactam resistance. Both underwent renal replacement therapy, and  
40 failed despite attaining optimal joint PK/PD target and receiving fosfomycin co-treatment.

41 **Conclusion:** CI administration may allow the attainment of optimal joint PK/PD targets of ceftazidime-  
42 avibactam in most critical renal patients with CR-GN infections, and may be helpful in minimizing the risk of  
43 microbiological failure.

44

45 **Keywords:** ceftazidime-avibactam; continuous infusion; carbapenem-resistant Gram-negative infections;  
46 PK/PD target attainment; critical renal patients; microbiological failure

## 47 **Background**

48           The widespread diffusion of difficult-to-treat resistant (DTR) Gram-negative pathogens is a worrisome  
49 health concern, and represents one of the main causes of hospital morbidity and mortality [1]. Several novel  
50 beta-lactam/beta-lactamase inhibitor (BL/BLI) combinations have been recently licensed for the management  
51 of carbapenem-resistant *Enterobacterales* (CRE) and/or carbapenem-resistant *Pseudomonas aeruginosa* (CR-  
52 PA) infections [2].

53           Ceftazidime-avibactam is a novel BL/BLI combination with *in vitro* and *in vivo* activity against either  
54 *Klebsiella pneumoniae* carbapenemase (KPC)-producing and OXA-48-producing *Enterobacterales* or CR-PA  
55 [2]. Several real-world evidences have confirmed the advantages in terms of clinical outcome that ceftazidime-  
56 avibactam may have over old traditional agents in the management of DTR Gram-negative infections [3].  
57 Preclinical models defined the joint PK/PD target for minimal efficacy of ceftazidime-avibactam (1-log kill)  
58 as the simultaneous achievement of 50% of the dose interval that free ceftazidime concentrations are above  
59 the MIC (50%  $fT > MIC$ ) and free avibactam concentrations are greater than the threshold concentration (CT)  
60 of 1.0 mg/L (50%  $fT > CT$  of 1.0 mg/L) [4,5].

61           However, recent findings suggested that minimum PK/PD targets of beta-lactams may not be adequate  
62 when treating severe Gram-negative deep-seated infections in the critically ill patients. More aggressive  
63 PK/PD targets up to 100%  $T_{>4-8 \times MIC}$  were shown to be helpful for maximizing clinical efficacy and  
64 microbiological eradication and for minimizing resistance development [6,7]. In this scenario, continuous  
65 infusion (CI) administration has been suggested as a valuable strategy for maximizing the achievement of  
66 optimal PK/PD targets with beta-lactams [8].

67           The aim of this study was to describe the pharmacokinetic/pharmacodynamic (PK/PD) behaviour of  
68 continuous infusion (CI) ceftazidime/avibactam and the microbiological outcome in a case series of critically  
69 ill patients with altered renal function who were treated with first-line or rescue therapy for documented  
70 carbapenem-resistant Gram-negative (CR-GN) bloodstream infections (BSI) and/or ventilator-associated  
71 pneumonia (VAP).

## 72 **Methods**

73 This retrospective study included a case series of critically ill patients who were treated with CI  
74 ceftazidime-avibactam for documented CRE or CR-PA infections at the intensive care units (ICUs) of the  
75 IRCCS Azienda Ospedaliero-Universitaria of Bologna between 01<sup>st</sup> April 2021 and 31<sup>st</sup> March 2022. All the  
76 included patients underwent real-time therapeutic drug monitoring (TDM) of ceftazidime-avibactam.  
77 Demographic and clinical/laboratory data were extracted for each single patient. Type/site of infection,  
78 ceftazidime-avibactam dosage, treatment duration, mono or combination therapy, MIC of ceftazidime-  
79 avibactam against CRE or CR-PA, and requirement for dosing adjustments were also collected. Combination  
80 therapy was defined as the concomitant use of other antibiotics active against the DTR Gram-negative clinical  
81 isolates.

82 Documented BSI was defined as the isolation of CRE from blood cultures. Documented VAP was  
83 defined as the isolation of CRE or CR-PA with a bacterial load  $\geq 10^4$  CFU/mL in the bronchoalveolar lavage  
84 (BAL) fluid culture after > 48 hours from endotracheal intubation and start of mechanical ventilation [9].

85 Antimicrobial susceptibility of ceftazidime-avibactam was tested by broth microdilution (panel  
86 provided by Merlin Diagnostika GMBH, Bornheim-Hersel, Germany). The range of MIC values tested for  
87 ceftazidime was from 1 to 64 mg/L with a fixed target avibactam concentration ( $C_T$ ) of 4 mg/L. Quality controls  
88 were performed monthly by using reference isolates provided by ATCC (namely *E. coli* 25922, *P. aeruginosa*  
89 27853, and *K. pneumoniae* 700603). Molecular analysis of CRE isolates was performed. Carbapenemase type  
90 was determined by multiplex immunochromatographic assay NG test CARBA 5 (NG Biotech, Guipry-Messac,  
91 France) for detecting the specific carbapenemase enzyme produced (IMP, VIM, NDM, KPC, OXA-48). MIC  
92 values were interpreted according to the EUCAST guidelines [11], and resistance to ceftazidime-avibactam  
93 was defined whenever values were > 8 mg/L.

94 Ceftazidime-avibactam was prescribed at the discretion of the treating physician or infectious disease  
95 consultant as first-line or rescue therapy according to the current clinical practice guidelines implemented at  
96 the IRCCS Azienda Ospedaliero-Universitaria of Bologna. Treatment was always started with a loading dose  
97 (LD) of 2.5 g over 2-h infusion followed by a maintenance dose (MD) administered by CI. For this purpose,  
98 aqueous solutions were reconstituted every 8h or 12h and infused over 8-12h due to stability restrictions [10].

99 MD regimens were initially selected according to the patient's underlying pathophysiological conditions and  
100 renal function. To maximize the attainment of aggressive PK/PD targets, the maintenance dose of 2.5 g q8h  
101 by CI was administered to all of the patients with the intent of overcoming the major  
102 pathophysiological/iatrogenic factors that might have caused potential underexposure in the early phase of  
103 septic shock, including patients with potentially transient severe acute kidney injury. Lower maintenance  
104 dosing were implemented (i.e., 0.625-1.25 g q12h by CI) only in anuric patients who underwent intermittent  
105 haemodialysis for minimizing the risk of neurotoxicity. Higher than labeled dose (2.5 g q6h by CI) were  
106 recommended in specific clinical scenarios, namely patients affected by augmented renal clearance and/or with  
107 deep-seated infections or in patients with residual renal function undergoing high-intensity continuous renal  
108 replacement therapy.

109 Blood samples for measuring ceftazidime and avibactam steady-state concentrations ( $C_{ss}$ ) were  
110 collected firstly within 72 hours from starting treatment and then reassessed whenever feasible. In patients  
111 requiring CVVHDF, blood samples for TDM were collected at least 24 h after starting CRRT to ensure the  
112 achievement of steady-state. Total ceftazidime and avibactam serum concentrations were determined by means  
113 of a validated liquid chromatography-tandem mass spectrometry method [12].

114 As only total ceftazidime and avibactam concentrations were measured, the free fraction ( $f$ ) was  
115 calculated by considering the plasma protein binding reported in the literature (10% and 7% for ceftazidime  
116 and avibactam, respectively) [13]. The percentage of time with ceftazidime concentrations above the MIC was  
117 selected as PD parameter of ceftazidime efficacy and expressed as  $C_{ss}/MIC$  ratio. The percentage of time with  
118 avibactam concentrations above the  $C_T$  was selected as PD parameter of avibactam efficacy and expressed as  
119  $C_{ss}/C_T$  ratio. The primary goal was the attainment of a joint PK/PD target of ceftazidime-avibactam. The joint  
120 PK/PD target of ceftazidime-avibactam was considered as optimal when both  $C_{ss}/MIC$  ratio for ceftazidime  $\geq$   
121 4 (equivalent to 100%  $fT_{>4 \times MIC}$ ) and  $C_{ss}/C_T$  ratio for avibactam  $> 1$  (equivalent to 100%  $fT > C_T$  of 4.0 mg/L)  
122 were simultaneously achieved; as quasi-optimal if only one of the two thresholds was achieved, and as  
123 suboptimal if none of the two thresholds was achieved. Ceftazidime-avibactam dosing adjustments were  
124 provided on the basis of our current clinical practice, as previously reported [14].

125 Microbiological failure was defined as the persistence of the same bacterial pathogen in blood culture  
126 or in BAL culture after  $\geq 7$  days from starting ceftazidime-avibactam treatment, as previously reported [15].  
127 Resistance development was defined as the MIC increase of ceftazidime-avibactam against the clinical isolate  
128 beyond the EUCAST clinical breakpoint of susceptibility. Primary outcome was microbiological eradication,  
129 defined as the absence of the original pathogens from the blood or BAL culture of the specimens in at least  
130 two subsequent assessments. Follow-up blood cultures (in patients with BSI) and/or BAL cultures (in patients  
131 with VAP) were executed between day 2 and day 7, and between day 5 and day 14, respectively, for assessing  
132 microbiological eradication and defining treatment duration. For each patient, it was investigated the  
133 relationship between ceftazidime-avibactam PK/PD targets and microbiological outcome in relation to the site  
134 of infection (BSI and/or VAP) and to the class of renal function, which was stratified in six scenarios  
135 [intermittent haemodialysis (IHD), acute kidney injury (AKI), continuous renal replacement therapy (CRRT)  
136 in anuric patients, normal renal function, CRRT in patients with residual renal function, and augmented renal  
137 clearance (ARC)]. ARC was defined as a measured creatinine clearance  $\geq 130$  mL/min/1.73m<sup>2</sup> in males and  $\geq$   
138 120 in females coupled with a normal serum creatinine value [16]. AKI was classified according to Acute  
139 Kidney Injury Network criteria, including the need for CRRT or IHD [17]. Secondary outcomes included 30-  
140 day mortality rate and occurrence of adverse events (AEs).

141 Descriptive statistics were used. Continuous data were presented as the mean  $\pm$  standard deviation  
142 (S.D.) or median and interquartile range (IQR), whereas categorial variables were expressed by count and  
143 percentage. The study was approved by the Ethics Committee of IRCCS Azienda Ospedaliero-Universitaria  
144 of Bologna (n. 442/2021/Oss/AOUBo approved on 28th June 2021).

## 145 **Results**

146 Overall, during the study period ten critically ill patients had documented CR Gram-negative BSI (n  
147 = 5) or VAP (n = 4) or BSI plus VAP (n = 1) treated with CI ceftazidime-avibactam and underwent at least  
148 one TDM assessment of ceftazidime-avibactam  $C_{ss}$  (**Table 1**). Mean ( $\pm$ SD) age was  $60.7 \pm 14.5$  years with a  
149 male preponderance (70%). Seven out of 10 patients (70%) were admitted in ICU because of acute respiratory  
150 distress syndrome caused by severe COVID-19 pneumonia. The other three patients were admitted in ICU  
151 because of sepsis, which occurred after solid organ transplant (2/10) or major surgery (1/10). All patients  
152 underwent invasive mechanical ventilation, and all but one had septic shock. In regard to renal classes, six out  
153 of 10 patients (60%) underwent continuous venovenous haemodiafiltration (CVVHDF) (4 anuric and 2 with  
154 residual renal function), two (20%) underwent IHD, one each had normal renal function, AKI or ARC (10%  
155 each). One of the patients who underwent CVVHDF needed also extracorporeal membrane oxygenation  
156 (ECMO).

157 BSI and VAP occurred in five and four cases, respectively, whereas one patient had BSI and VAP  
158 simultaneously. All infections were microbiologically documented. Six (60%) were caused by CRE (KPC- or  
159 OXA-48-producing *Klebsiella pneumoniae* in two cases each, KPC/OXA-48 co-producing *Klebsiella*  
160 *pneumoniae* and AmpC-producing associated with porin loss *Klebsiella aerogenes* in one case each) and four  
161 (40%) by CR-PA. Overall, all isolates were fully susceptible to ceftazidime-avibactam, with MICs ranging  
162 from 2 to 8 mg/L.

163 MD of CI ceftazidime-avibactam was started at the full dosage of 2.5 g q8h over 8h in eight patients,  
164 and at the dose of 2.5 g q6h over 6h and of 0.625 g q12h over 12h in one case each. The median (IQR) duration  
165 of treatment was 13.5 days (8.25-24.75 days). Combination therapy was applied with fosfomycin in five cases  
166 and with aztreonam in another one.

167 Blood samples for first TDM assessment of ceftazidime-avibactam were collected on day 2 in five  
168 cases, and on day 3 in the other five. Turnaround time (TAT) of ceftazidime-avibactam was within 12 hours  
169 in 7 cases and within 48 hours in other 3 cases. The dose was confirmed in 5 out of 10 patients (50%), whereas  
170 in the other 5 cases a dose decrease was recommended. The median (IQR) average  $fC_{ss}$  of ceftazidime and  
171 avibactam were 49.9 mg/L (41.3-58.8 mg/L) and 14.1 mg/L (7.5-17.0 mg/L), respectively.



172 The joint PK/PD targets of ceftazidime-avibactam were optimal in eight patients (80%; 3 BSI, 4 VAP,  
173 1 BSI + VAP), quasi-optimal in two cases (20%; 2 BSI both with avibactam  $fC_{ss}/C_T < 1$ ), and never suboptimal.  
174 The relationships between the level of the joint PK/PD target of CI ceftazidime-avibactam and the  
175 microbiological outcome of BSI and/or VAP due to CRE and CR-PA are summarized in **Figure 1a** and **1b**,  
176 respectively. Microbiological eradication was achieved in 100% of patients with BSIs (5/5) and in 75% of  
177 those with VAPs (3/4). Microbiological failure (20%; 2/10) occurred in one patient with VAP (1/4) and in the  
178 only patient with BSI plus VAP. In this latter case, resistance of *K. pneumoniae* OXA-48-producer to  
179 ceftazidime-avibactam occurred (MIC of 64 mg/L). Both of these patients underwent renal replacement  
180 therapy and microbiologically failed despite attaining optimal joint PK/PD target of ceftazidime-avibactam  
181 and being co-treated with fosfomycin.

182 The overall 30-day mortality rate was 30%. None of the three patients passed away because of the CR  
183 Gram-negative infections. The underlying causes of mortality were severe COVID-19 pneumonia coupled  
184 with pulmonary invasive aspergillosis in two cases, and septic shock due to *Enterococcus faecium* in the other  
185 one. No ceftazidime/avibactam-related AEs emerged during treatment.

186 **Discussion**

187 To the best of our knowledge, this is the first study that described the joint PK/PD target attainment of  
188 ceftazidime-avibactam administered by CI and that assessed its relationships with the microbiological outcome  
189 in a real-world scenario of critical renal patients with documented CRE- or CR-PA- related BSI and/or VAP.

190 For traditional beta-lactams, administration by prolonged infusion, by achieving more aggressive  
191 PK/PD targets, was just shown to grant remarkable advantages over intermittent infusion in terms of better  
192 microbiological outcome and/or clinical outcome in critically ill patients [4,7,8].

193 In regard to ceftazidime-avibactam, a recent retrospective observational study assessed use and  
194 outcomes of mono- and/or combo-therapy among 577 patients with infections caused by KPC-Kp strains [18].  
195 Interestingly, it was shown that 30-day mortality was negatively associated at multivariate regression analysis  
196 with administration by prolonged infusion ( $p = 0.006$ ). These findings may suggest that prolonging infusion  
197 of ceftazidime-avibactam, by maximizing the PK/PD target attainment, may be the way forward for dealing  
198 with severe KPC-Kp infections [4,18].

199 Our results showed firstly that implementing a real-time TDM-guided approach of CI ceftazidime-  
200 avibactam may be helpful in attaining very aggressive joint PK/PD targets of ceftazidime-avibactam over time.  
201 Noteworthy, PK/PD target attainment was optimal in 80% of cases, and never suboptimal. This approach  
202 granted microbiological eradication in most cases both of CRE-related and of CR-PA-related BSI and/or VAP.  
203 It's worth mentioning that some real-world studies hypothesized that the unfavourable clinical outcome  
204 observed with ceftazidime-avibactam in some settings, namely pneumonia and/or during renal replacement  
205 therapy, could have been related to sub-optimal PK/PD target attainment at the infection site [15,18].  
206 Unfortunately, none of these studies directly tested this hypothesis. Our findings, although limited, suggest  
207 that a real-time TDM-guided approach may grant the possibility of coupling optimal PK/PD target attainment  
208 with quite high microbiological eradication rates in BSI and/or VAP, and offer the opportunity to argue that  
209 this approach could be helpful in these scenarios.

210 To the best of our knowledge, there is only one case series that previously assessed the PK/PD target  
211 attainment of CI ceftazidime-avibactam in Outpatient Parenteral Antimicrobial Therapy (OPAT) patients, but

212 it did not allow to draw any reliable conclusion as it lacked measurement of avibactam concentrations [19]. It  
213 should not be overlooked that avibactam, differently from tazobactam, is a reversible beta-lactamase inhibitor  
214 [5]. This means that CI administration may be a very powerful tool in maintaining concentrations steadily over  
215 time above the safeguarded fixed threshold adopted by the EUCAST for avibactam when testing  
216 ceftazidime/avibactam susceptibility, namely 4 mg/L [11].

217 In our case series, the real-time TDM-guided approach was very helpful in optimizing treatment,  
218 especially because our cohort included predominantly critically ill patients with variable and fluctuating  
219 degrees of renal function. The failure rate among patients who underwent renal replacement therapy in our  
220 study was quite low compared to those observed in other previous studies (29% vs. 60-83.3%) [15,20]. This  
221 could be explained by the fact that our TDM-guided approach was focused at achieving early aggressive joint  
222 PK/PD targets of CI ceftazidime-avibactam even in renal patients [21]. Of note, we are used to start  
223 ceftazidime-avibactam therapy with full maintenance dose for ensuring aggressive joint PK/PD targets also in  
224 patients with sepsis-associated transient AKI. This approach is thought for preventing the risk of  
225 underexposure in the eventuality that renal function should recover promptly in the first 24-48h, that is before  
226 performing the first TDM assessment [22]. In this scenario, implementation of real-time TDM should be  
227 pursued for granting prompt dosing adjustments in patients with fluctuating renal function.

228 It should be noticed that the only two episodes of microbiological failure occurred, and concerned  
229 VAP patients. This is in agreement with previous real-world evidence showing that pneumonia may be a major  
230 risk factor of clinical failure and mortality among patients treated with ceftazidime-avibactam [15,18]. We  
231 believe that in these two patients poor underlying conditions more than inappropriate antimicrobial therapy  
232 might have played a major role in determining the negative microbiological outcome, even if we could not rule  
233 out that suboptimal PK/PD target attainment might have occurred at the infection site. One occurred in a lung  
234 transplant recipient after long ICU stay and the other in a patient with severe COVID-19 pneumonia. The  
235 occurrence of suboptimal PK/PD target in the epithelial lining fluid (ELF) seems unlikely since in both cases  
236 the joint PK/PD targets achieved in plasma were very aggressive. Considering that the ELF/plasma ratio of  
237 ceftazidime-avibactam reported in the literature is around 0.30-0.35, this approach should have ensured  
238 optimal exposure even at the infection site [23]. Development of resistance to ceftazidime-avibactam occurred

239 only in one case (10%), and this may support the contention that this approach could also concur in minimizing  
240 the risk of emergence of difficult-to-treat phenotypes. Notably, no death was directly attributable to  
241 antimicrobial treatment failure.

242 The microbiological eradication rate was very similar between patients treated with ceftazidime-  
243 avibactam in monotherapy and those receiving combination therapy, likewise reported in a recent retrospective  
244 observational study [18]. This could suggest that when aggressive joint PK/PD target of CI ceftazidime-  
245 avibactam monotherapy have been pursued, combination therapy could not offer any additional benefit.

246 Our study has some limitations. The retrospective monocentric study design and the limited sample  
247 size should be acknowledged. Only total ceftazidime-avibactam concentrations were measured, and the free  
248 moieties were only estimated. However, this is the first real-life experience that described the joint PK/PD  
249 target attainment of CI ceftazidime-avibactam in the treatment of severe carbapenem-resistant Gram-negative  
250 infections among critical renal patients and that explored the relationship with microbiological outcome.

251 In conclusion, our findings suggest that administering ceftazidime-avibactam by CI and adopting a  
252 strategy of real-time TDM-guided dosing adaptation may be very helpful in attaining very aggressive joint  
253 PK/PD targets. This approach may lead to microbiological eradication in most cases of CRE- and/or CR-PA-  
254 related BSI and/or VAP regardless of mono- or combo- therapy. Large prospective clinical studies are  
255 warranted for confirming our hypothesis.

256

257 **Funding:** None.

258

259 **Competing interests:** M.G. reports grants from Angelini S.p.A., outside the submitted work. F.P. has  
260 participated in speaker's bureau for Angelini, BeiGene, Gilead, Menarini, MSD, Pfizer, Sanofi-Aventis,  
261 Shionogi, and as consultant for Angelini, BeiGene, Gilead, MSD, Pfizer, Shionogi, outside the submitted work.  
262 P.V. has served as a consultant for bioMérieux, Gilead, Merck Sharp & Dohme, Nabriva, Nordic Pharma,  
263 Pfizer, Thermo-Fisher, and Venatorx, and received payment for serving on the speaker's bureaus for Correvio,

264 Gilead, Merck Sharp & Dohme, Nordic Pharma, and Pfizer, outside the submitted work. The other authors  
265 report no potential conflicts of interest for this work.

266

267 **Ethical approval:** The study was conducted according to the guidelines of the Declaration of Helsinki and  
268 approved by the Ethics Committee of IRCCS Azienda Ospedaliero-Universitaria of Bologna (n.  
269 442/2021/Oss/AOUBo approved on 28th June 2021).

270 **References**

- 271 [1] Marston HD, Dixon DM, Knisely JM, Palmore TN, Fauci AS. Antimicrobial Resistance. *JAMA*  
272 2016;316:1193–204. <https://doi.org/10.1001/jama.2016.11764>.
- 273 [2] Yahav D, Giske CG, Grāmatniece A, Abodakpi H, Tam VH, Leibovici L. New  $\beta$ -Lactam- $\beta$ -Lactamase  
274 Inhibitor Combinations. *Clin Microbiol Rev* 2020;34. <https://doi.org/10.1128/CMR.00115-20>.
- 275 [3] Soriano A, Carmeli Y, Omrani AS, Moore LSP, Tawadrous M, Irani P. Ceftazidime-Avibactam for the  
276 Treatment of Serious Gram-Negative Infections with Limited Treatment Options: A Systematic Literature  
277 Review. *Infect Dis Ther* 2021;10:1989–2034. <https://doi.org/10.1007/s40121-021-00507-6>.
- 278 [4] Gatti M, Pea F. Continuous versus intermittent infusion of antibiotics in Gram-negative multidrug-  
279 resistant infections. *Curr Opin Infect Dis* 2021. <https://doi.org/10.1097/QCO.0000000000000755>.
- 280 [5] Crass RL, Pai MP. Pharmacokinetics and Pharmacodynamics of  $\beta$ -Lactamase Inhibitors.  
281 *Pharmacotherapy* 2019;39:182–95. <https://doi.org/10.1002/phar.2210>.
- 282 [6] Guilhaumou R, Benaboud S, Bennis Y, Dahyot-Fizelier C, Dailly E, Gandia P, et al. Optimization of the  
283 treatment with beta-lactam antibiotics in critically ill patients-guidelines from the French Society of  
284 Pharmacology and Therapeutics (Société Française de Pharmacologie et Thérapeutique-SFPT) and the  
285 French Society of Anaesthesia and Intensive Care Medicine (Société Française d’Anesthésie et  
286 Réanimation-SFAR). *Crit Care* 2019;23:104. <https://doi.org/10.1186/s13054-019-2378-9>.
- 287 [7] Gatti M, Cojutti PG, Pascale R, Tonetti T, Laici C, Dell’Olio A, et al. Assessment of a PK/PD Target of  
288 Continuous Infusion Beta-Lactams Useful for Preventing Microbiological Failure and/or Resistance  
289 Development in Critically Ill Patients Affected by Documented Gram-Negative Infections. *Antibiotics*  
290 2021;10:1311. <https://doi.org/10.3390/antibiotics10111311>.
- 291 [8] Vardakas KZ, Voulgaris GL, Maliaros A, Samonis G, Falagas ME. Prolonged versus short-term  
292 intravenous infusion of antipseudomonal  $\beta$ -lactams for patients with sepsis: a systematic review and meta-  
293 analysis of randomised trials. *Lancet Infect Dis* 2018;18:108–20. [https://doi.org/10.1016/S1473-](https://doi.org/10.1016/S1473-3099(17)30615-1)  
294 [3099\(17\)30615-1](https://doi.org/10.1016/S1473-3099(17)30615-1).
- 295 [9] Chastre J, Fagon J-Y. Ventilator-associated pneumonia. *Am J Respir Crit Care Med* 2002;165:867–903.  
296 <https://doi.org/10.1164/ajrccm.165.7.2105078>.
- 297 [10] Ceftazidime-avibactam summary of product characteristics n.d.

- 298 [11]EUCAST - Rationale for EUCAST clinical breakpoints - Ceftazidime-avibactam 2020.
- 299 [12]Sillén H, Mitchell R, Sleight R, Mainwaring G, Catton K, Houghton R, et al. Determination of avibactam  
300 and ceftazidime in human plasma samples by LC-MS. *Bioanalysis* 2015;7:1423–34.  
301 <https://doi.org/10.4155/bio.15.76>.
- 302 [13]Merdjan H, Rangaraju M, Tarral A. Safety and pharmacokinetics of single and multiple ascending doses  
303 of avibactam alone and in combination with ceftazidime in healthy male volunteers: results of two  
304 randomized, placebo-controlled studies. *Clin Drug Investig* 2015;35:307–17.  
305 <https://doi.org/10.1007/s40261-015-0283-9>.
- 306 [14]Gatti M, Cojutti PG, Bartoletti M, Tonetti T, Bianchini A, Ramirez S, et al. Expert clinical  
307 pharmacological advice may make an antimicrobial TDM program for emerging candidates more  
308 clinically useful in tailoring therapy of critically ill patients. *Crit Care* 2022;26:178.  
309 <https://doi.org/10.1186/s13054-022-04050-9>.
- 310 [15]Shields RK, Nguyen MH, Chen L, Press EG, Kreiswirth BN, Clancy CJ. Pneumonia and Renal  
311 Replacement Therapy Are Risk Factors for Ceftazidime-Avibactam Treatment Failures and Resistance  
312 among Patients with Carbapenem-Resistant Enterobacteriaceae Infections. *Antimicrob Agents Chemother*  
313 2018;62. <https://doi.org/10.1128/AAC.02497-17>.
- 314 [16]Cook AM, Hatton-Kolpek J. Augmented Renal Clearance. *Pharmacotherapy* 2019;39:346–54.  
315 <https://doi.org/10.1002/phar.2231>.
- 316 [17]Mehta RL, Kellum JA, Shah SV, Molitoris BA, Ronco C, Warnock DG, et al. Acute Kidney Injury  
317 Network: report of an initiative to improve outcomes in acute kidney injury. *Crit Care* 2007;11:R31.  
318 <https://doi.org/10.1186/cc5713>.
- 319 [18]Tumbarello M, Raffaelli F, Giannella M, Mantengoli E, Mularoni A, Venditti M, et al. Ceftazidime-  
320 avibactam use for KPC-Kp infections: a retrospective observational multicenter study. *Clin Infect Dis*  
321 2021. <https://doi.org/10.1093/cid/ciab176>.
- 322 [19]Goncette V, Layios N, Descy J, Fripiat F. Continuous Infusion, Therapeutic Drug Monitoring and  
323 Outpatient Parenteral Antimicrobial Therapy with Ceftazidime-Avibactam: a Retrospective Cohort Study.  
324 *J Glob Antimicrob Resist* 2021. <https://doi.org/10.1016/j.jgar.2021.04.015>.

- 325 [20] Vena A, Giacobbe DR, Castaldo N, Cattelan A, Mussini C, Luzzati R, et al. Clinical Experience with  
326 Ceftazidime-Avibactam for the Treatment of Infections due to Multidrug-Resistant Gram-Negative  
327 Bacteria Other than Carbapenem-Resistant Enterobacterales. *Antibiotics (Basel)* 2020;9:E71.  
328 <https://doi.org/10.3390/antibiotics9020071>.
- 329 [21] Gatti M, Pea F. Pharmacokinetic/pharmacodynamic target attainment in critically ill renal patients on  
330 antimicrobial usage: focus on novel beta-lactams and beta lactams/beta-lactamase inhibitors. *Expert Rev*  
331 *Clin Pharmacol* 2021. <https://doi.org/10.1080/17512433.2021.1901574>.
- 332 [22] Crass RL, Rodvold KA, Mueller BA, Pai MP. Renal Dosing of Antibiotics: Are We Jumping the Gun?  
333 *Clinical Infectious Diseases* 2019;68:1596–602. <https://doi.org/10.1093/cid/ciy790>.
- 334 [23] Nicolau DP, Siew L, Armstrong J, Li J, Edeki T, Learoyd M, et al. Phase 1 study assessing the steady-  
335 state concentration of ceftazidime and avibactam in plasma and epithelial lining fluid following two dosing  
336 regimens. *J Antimicrob Chemother* 2015;70:2862–9. <https://doi.org/10.1093/jac/dkv170>.
- 337



**Table 1 – Demographic and clinical features of critically ill patients with carbapenem-resistant Gram-negative infections treated with continuous infusion ceftazidime-avibactam**

**Carbapenem-resistant Enterobacteriales**

| Age/sex  | ICU admission                       | Type of infection and pathogen      | CAZ-AVI MIC (mg/L) | CAZ-AVI dosage | CAZ-AVI treatment duration (days) | Average free ceftazidime Css (mg/L) | Average free avibactam Css (mg/L) | Ceftazidime fCss/MIC ratio | Avibactam fCss/C <sub>T</sub> ratio | PK/PD target attainment | Dosing adjustment    | Antibiotic co-treatment | CRRT ECMO CLCr   | ME BSI | ME VAP | Resistance development  | 30-day mortality |
|--|-------------------------------------|-------------------------------------|--------------------|----------------|-----------------------------------|-------------------------------------|-----------------------------------|----------------------------|-------------------------------------|-------------------------|----------------------|-------------------------|--|--------|--------|---|------------------|
| 76/F   | ARDS in COVID-19                    | BSI CR <i>K. aerogenes</i>          | 2                  | 2.5g q8h CI    | 9                                 | 30.8                                | 2.6                               | 15.4                       | 0.6                                 | Quasi-optimal           | No                   | No                      | No CLCr 96 mL/min/1.73m <sup>2</sup>   | Yes    | NA     | No  | No               |
| 31/M   | Sepsis in OLT recipient             | BSI <i>K. pneumoniae</i> KPC        | 2                  | 2.5g q8h CI    | 15                                | 31.5                                | 2.4                               | 15.8                       | 0.6                                 | Quasi-optimal           | No                   | No                      | No ARC CLCr 133.1 mL/min/1.73m <sup>2</sup>  | Yes    | NA     | No  | No               |
| 63/F   | ARDS in COVID-19                    | BSI <i>K. pneumoniae</i> KPC/OXA-48 | 4                  | 2.5g q8h CI    | 8                                 | 58.5                                | 16.7                              | 14.6                       | 4.2                                 | Optimal                 | No                   | Fosfomycin              | CVVHDF Quf 2400 mL/h<br><br>Anuric AKIN 3  | Yes    | NA     | No  | Yes              |
| 47/M   | ARDS in COVID-19                    | VAP <i>K. pneumoniae</i> KPC        | 4                  | 2.5g q6h CI    | 28                                | 52.7                                | 16.0                              | 13.2                       | 4.0                                 | Optimal                 | Yes<br>2.5g q8h CI   | No                      | ECMO + CVVHDF Quf 2500 mL/h<br><br>Residual diuresis 715 mL/day (measured CLCr 14.1 mL/min) AKIN 3 | NA     | Yes    | No  | No               |
| 48/M   | ARDS in COVID-19                    | VAP <i>K. pneumoniae</i> OXA-48     | 2                  | 2.5g q8h CI    | 14                                | 139.9                               | 80.6                              | 70.0                       | 20.1                                | Optimal                 | Yes<br>1.25g q12h CI | Aztreonam               | No CLCr 30 mL/min/1.73m <sup>2</sup> AKIN 2  | NA     | Yes    | No  | No               |
| 64/M   | Sepsis in lung transplant recipient | BSI+VAP <i>K. pneumoniae</i> OXA-48 | 2                  | 2.5g q8h CI    | 13                                | 42.6                                | 17.1                              | 21.3                       | 4.3                                 | Optimal                 | Yes<br>1.25g q12h CI | Fosfomycin              | CVVHDF followed by IHD<br><br>Anuric AKIN 3  | Yes    | No     | Yes (isolation of <i>Kp</i> OXA-48 with MIC = 64 mg/L on BAL 3 days after the end of treatment) | No               |
| <b>Carbapenem-resistant Pseudomonas aeruginosa</b> |                                     |                                     |                    |                |                                   |                                     |                                   |                            |                                     |                         |                      |                         |  |        |        |   |                  |
| 72/M   | ARDS in COVID-19                    | BSI CR <i>P. aeruginosa</i>         | 4                  | 2.5g q8h CI    | 43                                | 59.0                                | 12.2                              | 14.8                       | 3.0                                 | Optimal                 | Yes<br>1.25g q8h CI  | Fosfomycin              | CVVDHF Quf 2806 mL/h<br><br>Anuric   | Yes    | NA     | No  | No               |

|      |  |                                  |   |                      |    |      |      |      |     |         |                            |            |  |     |     |    |     |
|------|--|----------------------------------|---|----------------------|----|------|------|------|-----|---------|----------------------------|------------|--|-----|-----|----|-----|
|      |  |                                  |   |                      |    |      |      |      |     |         |                            |            | AKIN 3   |     |     |    |     |
| 72/M | ARDS in COVID-19                         | BSI<br>CR – <i>P. aeruginosa</i> | 4 | 2.5g<br>q8h CI       | 34 | 78.9 | 18.2 | 19.7 | 4.5 | Optimal | Yes<br><br>2.5g q12h<br>CI | No         | CVVHDF<br>Quf 2505 mL/h<br><br>Residual<br>diuresis > 1000<br>mL/day<br>AKIN 3 | Yes | NA  | No | No  |
| 73/M | Septic shock in multiple bowel resection | VAP<br>CR – <i>P. aeruginosa</i> | 2 | 0.625g<br>q12h<br>CI | 8  | 40.9 | 7.3  | 20.5 | 1.8 | Optimal | No                         | Fosfomycin | IHD<br><br>Anuric<br>AKIN 3  | NA  | Yes | No | Yes |
| 61/F | ARDS in COVID-19                         | VAP<br>CR – <i>P. aeruginosa</i> | 8 | 2.5g<br>q8h CI       | 7  | 47.3 | 8.4  | 5.9  | 2.1 | Optimal | No                         | Fosfomycin | CVVHDF<br>Quf 2700 mL/h<br><br>Anuric<br>AKIN 3                                | NA  | No  | No | Yes |

AKIN: Acute Kidney Injury Network definition; ARC: augmented renal clearance; ARDS: acute respiratory distress syndrome; BAL: bronchoalveolar lavage; BSI: bloodstream infection; CAZ-AVI: ceftazidime-avibactam; CI: continuous infusion; CLCr: creatinine clearance; CR: carbapenem-resistant; C<sub>T</sub> = target concentration of 4 mg/L; CVVHDF: continuous venovenous haemodiafiltration; ECMO: extracorporeal membrane oxygenation; IHD: intermittent haemodialysis; ME: microbiological eradication; MIC: minimum inhibitory concentration; NA: not applicable; OLT: orthotopic liver transplant; Q<sub>uf</sub>: ultrafiltration rate; VAP: ventilator-associated pneumonia

339 **Figure Legends**

340 **Figure 1** – Description of pharmacokinetic/pharmacodynamic target attainment and microbiological outcome  
341 for ceftazidime-avibactam in patients with infections caused by carbapenem-resistant *Enterobacterales* (**Panel**  
342 **a**) and carbapenem-resistant *Pseudomonas aeruginosa* (**Panel b**). Green box, microbiological eradication; red  
343 box, microbiological failure; grey box, absence of the specific type of infection. Each row corresponds to a  
344 single patient. Optimal joint PK/PD targets were considered the simultaneously achievement of 100%  $fT_{>4 \times$   
345  $MIC}$  for ceftazidime and 100%  $fT > 4 \times C_T$  of 1.0 mg/L for avibactam, quasi-optimal if only one of the two  
346 thresholds was achieved, and suboptimal if none of the two thresholds was achieved. BSI, bloodstream  
347 infection; CAZ-AVI, ceftazidime-avibactam; PK/PD, pharmacokinetic/pharmacodynamic; VAP, ventilator-  
348 associated pneumonia.