







*This article has been accepted for publication in Monthly Notices of the Royal Astronomical Society ©: 2023 The Authors. Published by Oxford University Press on behalf of the Royal Astronomical Society. All rights reserved.*

# The contribution of *in situ* and *ex situ* star formation in early-type galaxies: MaNGA versus IllustrisTNG

Carlo Cannarozzo <sup>1,2,3★</sup>, Alexie Leauthaud <sup>4</sup>, Grecco A. Oyarzún,<sup>4</sup> Carlo Nipoti <sup>2</sup>,  
Benedikt Diemer <sup>5</sup>, Song Huang,<sup>6</sup> Vicente Rodriguez-Gomez <sup>7</sup>, Alessandro Sonnenfeld <sup>8</sup>  
and Kevin Bundy<sup>4,9</sup>

<sup>1</sup>*Instituto de Astronomía, Universidad Nacional Autónoma de México, A. P. 70-264, 04510 CDMX, México*

<sup>2</sup>*Dipartimento di Fisica e Astronomia ‘Augusto Righi’, Alma Mater Studiorum Università di Bologna, via Piero Gobetti 93/2, I-40129 Bologna, Italy*

<sup>3</sup>*INAF - Osservatorio di Astrofisica e Scienza dello Spazio di Bologna, Via Piero Gobetti 93/3, I-40129 Bologna, Italy*

<sup>4</sup>*Department of Astronomy and Astrophysics, University of California, UCO/Lick Observatory, 1156 High Street, Santa Cruz, CA 95064, USA*

<sup>5</sup>*Department of Astronomy, University of Maryland, College Park, MD 20742, USA*

<sup>6</sup>*Department of Astrophysical Sciences, Princeton University, 4 Ivy Lane, Princeton, NJ 08544, USA*

<sup>7</sup>*Instituto de Radioastronomía y Astrofísica, Universidad Nacional Autónoma de México, A. P. 72-3, 58089 Morelia, Mexico*

<sup>8</sup>*Leiden Observatory, Leiden University, Niels Bohrweg 2, NL-2333 CA Leiden, the Netherlands*

<sup>9</sup>*UCO/Lick Observatory, University of California, Santa Cruz, 1156 High Street, Santa Cruz, CA 95064, USA*

Accepted 2022 October 14. Received 2022 October 13; in original form 2022 May 2

## ABSTRACT

We compare stellar mass surface density, metallicity, age, and line-of-sight velocity dispersion profiles in massive ( $M_* \geq 10^{10.5} M_\odot$ ) present-day early-type galaxies (ETGs) from the MaNGA survey with simulated galaxies from the TNG100 simulation of the IllustrisTNG suite. We find an excellent agreement between the stellar mass surface density profiles of MaNGA and TNG100 ETGs, both in shape and normalization. Moreover, TNG100 reproduces the shapes of the profiles of stellar metallicity and age, as well as the normalization of velocity dispersion distributions of MaNGA ETGs. We generally also find good agreement when comparing the stellar profiles of central and satellite galaxies between MaNGA and TNG100. An exception is the velocity dispersion profiles of very massive ( $M_* \gtrsim 10^{11.5} M_\odot$ ) central galaxies, which, on average, are significantly higher in TNG100 than in MaNGA ( $\approx 50 \text{ km s}^{-1}$ ). We study the radial profiles of *in situ* and *ex situ* stars in TNG100 and discuss the extent to which each population contributes to the observed MaNGA profiles. Our analysis lends significant support to the idea that high-mass ( $M_* \gtrsim 10^{11} M_\odot$ ) ETGs in the present-day Universe are the result of a merger-driven evolution marked by major mergers that tend to homogenize the stellar populations of the progenitors in the merger remnant.

**Key words:** galaxies: elliptical and lenticular, cD – galaxies: evolution – galaxies: formation – galaxies: interactions – galaxies: stellar content – galaxies: structure.

## 1 INTRODUCTION

In the standard cosmological framework, the formation and evolution of galaxies is thought to be driven by mergers and the accretion of material from the intergalactic medium (e.g. Cimatti, Fraternali & Nipoti 2019). In particular, for the assembly of massive early-type galaxies (ETGs), a *two-phase formation scenario* has been proposed (e.g. Naab, Johansson & Ostriker 2009; Oser et al. 2010; Hilz, Naab & Ostriker 2013). In the first phase of this formation process ( $z \gtrsim 2$ ), ETGs are built from stars formed *in situ*, i.e. within the same galaxy, while later, as a consequence of minor and major mergers, ETGs grow mainly by the accretion of stars formed *ex situ*, i.e. in other galaxies.

A natural outcome of mergers experienced by ETGs is the evolution of scaling relations (e.g. Cimatti et al. 2019), i.e. the observed empirical correlations between global galaxy properties, such as

those relating luminosity (or mass) with stellar velocity dispersion (Faber & Jackson 1976), size (Kormendy 1977), or both (the so-called Fundamental Plane; Djorgovski & Davis 1987; Dressler et al. 1987). The improvement of instrumentation technology together with increasing statistics in recent surveys has enabled studies of those relations at different redshifts. Indeed, massive ETGs at high redshift are found to be compact, with an effective radius  $R_e$  smaller than that of galaxies of similar stellar mass in the present-day Universe (e.g. Ferguson et al. 2004; Van der Wel et al. 2014; Damjanov et al. 2019). Also the stellar mass–central velocity dispersion relation ( $M_* - \sigma_c$ ) evolves: on average, for a given stellar mass, the lower the redshift, the lower the velocity dispersion (e.g. Van de Sande et al. 2013; Belli, Newman & Ellis 2014, 2017; Tanaka et al. 2019; Cannarozzo, Sonnenfeld & Nipoti 2020).

Mergers and accretion not only affect global galaxy properties, but also the internal distributions of stellar properties. The spatial distributions of metallicity, chemical abundances, age, and other properties of stellar populations in a galaxy enclose information on the evolutionary processes that have occurred across cosmic time.

\* E-mail: [ccannarozzo@astro.unam.mx](mailto:ccannarozzo@astro.unam.mx)

One possible way to investigate how progenitor stellar populations come together to form present-day ETGs is to perform galaxy-scale high-resolution simulations, an approach recently adopted, for instance, by Nipoti et al. (2020) to study the effect of mergers on the internal distributions of the stellar initial mass function (IMF) and the velocity dispersion in ETGs. A second approach is the study of the evolution of stellar population properties using hydrodynamical cosmological simulations. For example, Oser et al. (2012), exploiting a set of 40 zoom-in hydrodynamical simulations of individual haloes presented in Oser et al. (2010), and Rodriguez-Gomez et al. (2016), using the Illustris (Genel et al. 2014; Vogelsberger et al. 2014a,b; Sijacki et al. 2015) simulations, studied the radial distributions of *in situ* and *ex situ* stars in ETGs. With a similar approach, Barber, Schaye & Crain (2019) studied IMF radial gradients in ETGs drawn from the Evolution and Assembly of GaLaxies and their Environments (EAGLE; Crain et al. 2015; Schaye et al. 2015; McAlpine et al. 2016) cosmological simulations.

The presence of initial gradients in stellar metallicity is thought to be established during the first episodes of star formation (e.g. Larson 1974; Thomas et al. 2005) with metallicity profiles that decrease towards the external regions of galaxies, but with fairly flat stellar age profiles. As shown in Hirschmann et al. (2015), using a set of 10 high-resolution cosmological zoom simulations presented in Hirschmann et al. (2013), as well as in Cook et al. (2016) taking data from Illustris, the large number of mergers and interactions that occur in galaxies then tend to flatten metallicity profiles and almost flatten (or lead to slightly positive) age gradients both because of the mixing of stars with different metallicities and the accumulation of old stellar populations in the outer regions. On the contrary, galaxies with few mergers may retain their original negative metallicity profiles with metal-poor regions dominating in the outskirts of galaxies (e.g. Kobayashi 2004; Pipino et al. 2010; Taylor & Kobayashi 2017).

In the last few decades, integral field spectroscopy (IFS) has formed the basis of many surveys: SAURON (Spectroscopic Areal Unit for Research on Optical Nebulae; Bacon et al. 2001; de Zeeuw et al. 2002), ATLAS<sup>3D</sup> (Cappellari et al. 2011), CALIFA (Calar Alto Legacy Integral Field Array survey; Sánchez et al. 2016), SAMI (Sydney-Australian-Astronomical-Observatory Multi-object Integral-Field Spectrograph; Croom et al. 2012; Bryant et al. 2015), MASSIVE (Ma et al. 2014), and MaNGA (Mapping Nearby Galaxies at Apache Point Observatory; Bundy et al. 2015). These spatially resolved surveys allow in-depth studies of the properties of stellar populations in individual objects, therefore not limiting analyses only to the study of gradients, but revealing the 2D spatial distribution over the entire galaxy on the plane of the sky. By analysing a set of ETGs with  $\log(M_*/M_\odot) > 10.3$  in SAURON, Kuntschner et al. (2010) found that stellar metallicity gradients become shallower with increasing stellar mass, while stellar age gradients are independent of stellar mass. Li et al. (2018), using MaNGA galaxies with  $9 < \log(M_*/M_\odot) < 12.3$ , found metallicity gradients consistent with those of Kuntschner et al. (2010). Moreover, Li et al. (2018) found that stellar metallicity gradients show a strong dependence on stellar velocity dispersion: they peak (being most negative) at velocity dispersions of around  $100 \text{ km s}^{-1}$ . This radial dependence can be interpreted in terms of different evolutionary scenario for galaxies with different velocity dispersions. In particular, metallicity gradients tend to flatten at high velocity dispersions, perhaps indicating the rising role of mergers that redistribute stellar populations in these galaxies.

However, studies conducted so far that involve IFS surveys are also sometimes in disagreement. For example, Goddard et al. (2017a) selected ETGs from MaNGA with  $9 < \log(M_*/M_\odot) <$

$11.5$ . Although the galaxies were drawn from the same survey used by Li et al. (2018), the authors derived metallicity profiles that become steeper towards higher masses. A similar result was found by Zheng et al. (2017), for ETGs in the MaNGA survey with  $8.5 < \log(M_*/M_\odot) < 11.5$ . In Greene et al. (2015), subsequently extended in Greene et al. (2019) to larger radii, ETGs with  $\log(M_*/M_\odot) > 11.6$  show shallow metallicity gradients and radius-independent age and  $\alpha$ -element abundances relative to iron, i.e.  $[\alpha/\text{Fe}]$ . By analysing a sample of 96 passive brightest cluster galaxies from the SAMI survey, Santucci et al. (2020) found negative metallicity gradients that tend to become shallower as the stellar mass increases, slightly positive age gradients, and almost zero  $[\alpha/\text{Fe}]$  gradients, the latter tending to become slightly more negative with increasing mass. This study also revealed that there are no significant differences in the stellar profiles of the analysed properties between central and satellite galaxies, both at fixed stellar mass and as a function of halo mass, suggesting that the two galaxy populations follow a similar formation scenario, which appears to be independent of the environment. Differences among these various studies (also when using the same galaxy survey) appear to result from a combination of different selection criteria adopted to identify ETGs, the stellar mass ranges considered, and the methods used to retrieve properties and their profiles. In addition, we have found that the radial range adopted to measure the gradients and whether the profiles are stacked in physical units or in units of  $R_e$  can lead to some of these discrepancies.

Despite the relatively large number of IFS surveys, understanding whether a stellar population in a galaxy either formed *in situ* or was accreted from another progenitor is not a trivial task. Oyarzún et al. (2019), analysing more than 1000 ETGs with  $10 < \log(M_*/M_\odot) < 12$  from the MaNGA survey, studied the radial distributions of metallicity adopting three different stellar fitting codes, i.e. FIREFLY (Maraston & Strömbäck 2011; Comparat et al. 2017; Goddard et al. 2017b; Wilkinson et al. 2017; Maraston et al. 2020), PROSPECTOR (Leja et al. 2017; Johnson et al. 2019), and pPXF (Cappellari & Emsellem 2004; Cappellari 2017). As the mass increases, the flattening in the metallicity profiles was found to become more prominent at  $R \gtrsim R_e$ . Oyarzún et al. (2019) interpreted this flattening using a toy model in which they assume that the low-mass tail of galaxies in their sample is representative of galaxies mainly constituted by stars formed *in situ*. For high-mass galaxies, the inner part of the profiles ( $R \lesssim R_e$ ) is associated with an *in situ* stellar population, while the external parts are considered to be dominated by stars accreted from other galaxies. Quantitatively, they infer the contribution of *ex situ* stars within  $R \approx 2R_e$  to be  $\approx 20$  per cent of the total stellar mass in ETGs with  $\log(M_*/M_\odot) < 10.5$ , while in ETGs with  $\log(M_*/M_\odot) > 11.5$  this fraction reaches  $\approx 80$  per cent (consistent results are also presented in the observational works of Edwards et al. 2020 and Davison et al. 2021).

An alternative approach that allows to combine observations and simulations has been recently proposed by Nanni et al. (2022). In that paper, the authors built iMaNGA, a MaNGA-like galaxy sample considering both early- and late-type galaxies extracted from the cosmological simulation TNG50 (Nelson et al. 2019b; Pillepich et al. 2019). Specifically, Nanni et al. (2022), collecting simulated galaxies from the snapshots between  $z = 0.01$  and  $z = 0.15$ , so as to cover the whole MaNGA redshift range, took into account all the instrumental effects and methods employed to acquire data for MaNGA sources. The specific use of TNG50 allowed the authors to take all the advantages of high-spatial-resolution data, and generate corresponding mock galaxy spectra. Along similar lines, Bottrell & Hani (2022) presented RealSim-IFSS, a generalized tool for forward-modelling realistic synthetic IFS observations from hydrodynamical

simulations. RealSim-IFS is able to reproduce cubes similar to those produced by the MaNGA survey Data Reduction Pipeline. Furthermore, extracting around 900 galaxies with  $\log(M_*/M_\odot) > 10$  from TNG50, Bottrell & Hani (2022) applied RealSim-IFS to generate a synthetic MaNGA stellar kinematic survey.

In this work, we propose a physically grounded model to provide an interpretative scenario for the radial distributions of stellar properties in observed ETGs in terms of *in situ* and *ex situ* stellar components. In particular, we compare the radial profiles of stellar mass surface density, metallicity, age, and line-of-sight velocity dispersion of observed galaxies drawn from the data release 15 of the MaNGA survey with those of simulated ETGs extracted from the TNG100 simulation of the *The Next Generation* Illustris project (IllustrisTNG<sup>1</sup>; Marinacci et al. 2018; Naiman et al. 2018; Nelson et al. 2018; Pillepich et al. 2018b; Springel et al. 2018). Simulated galaxies are broken down into *in situ* and *ex situ* stellar populations using the methods presented in Rodriguez-Gomez et al. (2015) which serves as a prediction for the gradients of these two populations in massive ETGs. Indeed, the main scope of this work is to suggest a possible evolutionary scenario of the underlying hierarchical stellar mass assembly history of present-day ETGs. With this goal in mind, we focus the analysis on the study of radial distributions of the above-mentioned stellar properties for both MaNGA and IllustrisTNG galaxies, considering quantities *at face value*, i.e. as those directly derived from the pipelines and stellar fitting codes for MaNGA, and those from the TNG100 simulations.

This paper is organized as follows. In Section 2, we describe the galaxy samples and the criteria adopted to select ETGs. The method used to compute radial profiles from simulations is described in Section 3. Our results are presented in Section 4. In Section 5, we discuss the implications of the analysis and compare with previous studies, while Section 6 presents our conclusions.

Throughout this paper, we assume a  $\Lambda$  cold dark matter ( $\Lambda$ CDM) cosmological framework with cosmological parameters derived from Planck Collaboration XIII (2016), i.e.  $\Omega_{\Lambda,0} = 0.6911$ ,  $\Omega_{m,0} = 0.3089$ ,  $\Omega_b,0 = 0.0486$ , and  $H_0 = 67.74 \text{ km s}^{-1} \text{ Mpc}^{-1}$ .

## 2 OBSERVED AND SIMULATED GALAXY SAMPLES

In this section, we describe the selection criteria and physical properties of the observed (MaNGA) and simulated (TNG100) samples and the methods adopted to compare the two samples.

### 2.1 The MaNGA survey

The MaNGA survey (Bundy et al. 2015; Yan et al. 2016b), one of the three components of the fourth generation of SDSS (York et al. 2000; Gunn et al. 2006; Blanton et al. 2017) mapped with the 2.5 m telescope Apache Point Observatory  $\approx 10\,000$  galaxies with  $\log(M_*/M_\odot) > 9$  in the redshift range  $0.01 \lesssim z \lesssim 0.15$ , providing spatially resolved spectra for each source. The galaxy sample is taken from an extended version of the original NASA-Sloan Atlas (NSA v1.0.1<sup>2</sup>; Blanton et al. 2011) catalogue. By exploiting the IFS technique (Smee et al. 2013; Drory et al. 2015; Law et al. 2015), galaxies in MaNGA are observed with a set of 17 hexagonal bundles, each composed of fibres with a diameter that varies from 12 arcsec (with 19 fibres) to 32 arcsec (with 127 fibres). Each fibre has a

diameter of 2 arcsec. MaNGA achieves a uniform radial coverage of galaxies to  $1.5 R_e$  and  $2.5 R_e$ , for  $\approx 2/3$  (*Primary Sample*) and  $\approx 1/3$  (*Secondary Sample*) of the final sample. The observations provide a wavelength coverage in the range 3600–10 300 Å, with a spectral resolution of  $R \sim 1400$  at  $\lambda \sim 4000$  Å and  $R \sim 2600$  at  $\lambda \sim 9000$  Å (see Smee et al. 2013).

The MaNGA observations used in this work were previously reduced by the Data Reduction Pipeline (Law et al. 2016; Yan et al. 2016a). Both the de-projected distances and stellar kinematic maps are computed using the Data Analysis Pipeline (DAP; Westfall et al. 2019) for MaNGA. The MaNGA galaxies forming our observed sample are taken from the SDSS data release 15 (DR15, hereafter simply MaNGA; Aguado et al. 2019), which corresponds to the first 4675 observed MaNGA galaxies.<sup>3</sup>

To study the behaviour of radial profiles of observed ETGs, we use measurements of stellar mass surface density, metallicity, and age derived from two full spectral fitting codes: FIREFLY<sup>4</sup> (Maraston & Strömbäck 2011; Comparat et al. 2017; Goddard et al. 2017b; Wilkinson et al. 2017; Maraston et al. 2020) and PROSPECTOR<sup>5</sup> (Leja et al. 2017; Johnson et al. 2019). The use of two different methods will help to quantify the presence of systematic biases caused by different assumptions, priors, and fitting methods (Conroy 2013). In addition, we take into account estimates of line-of-sight stellar velocity dispersion obtained by using the pPXF code<sup>6</sup> (Cappellari & Emsellem 2004; Cappellari 2017). In the following text, we briefly summarize the settings adopted for the three stellar population fitting codes.

(i) FIREFLY (Fitting Iteratively For Likelihood analysis) is a  $\chi^2$ -minimization fitting code for deriving the stellar population properties. This code aims at disentangling stars and dust, subtracting the low-order continuum shape before fitting spectra. A set of simple stellar populations (SSPs) with a variety of age and metallicity are considered iteratively, in order to minimize the  $\chi^2$  fitting procedure, allowing FIREFLY to fit non-parametric star formation histories (SFHs). We adopt the stellar population models of Maraston & Strömbäck (2011), the MILES stellar library (Sánchez-Blázquez et al. 2006; Vazdekis et al. 2010), and a Chabrier (2003) IMF. The set of SSPs used are spread over the range 6.5 Myr–15 Gyr in age, while metallicity can assume values in the range  $-2.3 \leq \log(Z_*/Z_\odot) \leq 0.3$ . The wavelength range covered by the library is 4000–7400 Å. We include only spectra with  $S/N > 10$  (see Goddard et al. 2017a; Wilkinson et al. 2017), and we mask emission lines.

(ii) PROSPECTOR is a code able to infer stellar population properties from photometric and/or spectroscopic data with flexible models. It is based on the original stellar population synthesis code FSPS<sup>7</sup> (Conroy, Gunn & White 2009; Conroy & Gunn 2010). PROSPECTOR provides the posterior distribution of a stellar population parameter space (externally defined by users), uncertainties, and degeneracies. We adopt the MILES stellar population library, the MIST isochrones (Choi et al. 2016; Dotter 2016), and a Kroupa (2001) IMF.<sup>8</sup> The fitting procedure explores a 10-dimensional parameter space. In this fit, the dust optical depth in the *V* band, stellar mass, stellar

<sup>3</sup> Available at <https://www.sdss.org/dr15/manga/manga-data/>.

<sup>4</sup> Available at [https://github.com/FireflySpectra/firefly\\_release](https://github.com/FireflySpectra/firefly_release).

<sup>5</sup> Available at <https://github.com/bd-j/Prospector>.

<sup>6</sup> Available at <http://www-astro.physics.ox.ac.uk/~mxc/software/>

<sup>7</sup> Available at <https://github.com/cconroy20/fps>.

<sup>8</sup> For our purpose, the assumption of a Kroupa IMF or a Chabrier IMF to retrieve stellar population properties is almost indistinguishable.

<sup>1</sup> Official website at <https://www.tng-project.org>.

<sup>2</sup> Available at <https://www.sdss.org/dr15/manga/manga-target-selection/nsa/>.

**Table 1.** List of priors used for our PROSPECTOR runs. Column 1: parameter. Column 2: prior.

Parameter	Prior
Star formation history	Continuity
dust2	TopHat (0,1)
Stellar metallicity $\log Z_* [Z_\odot]$	TopHat (-2, 0.3)
Formed stellar mass $M_*/M_\odot$	LogUniform( $10^7$ , $10^{12}$ )
Velocity dispersion $\sigma_* [\text{km s}^{-1}]$	TopHat (0.1, 400)

velocity dispersion, and mass-weighted metallicities are taken into account. Moreover, non-parametric SFHs with a continuity prior are considered. Following the same approach described in Leja et al. (2019), our parameter space considers the star formation rate (SFR) spanning the following time intervals:  $0 < t < 30$  Myr,  $30 \text{ Myr} < t < 100$  Myr,  $100 \text{ Myr} < t < 330$  Myr,  $330 \text{ Myr} < t < 1.1$  Gyr,  $1.1 \text{ Gyr} < t < 3.6$  Gyr,  $3.6 \text{ Gyr} < t < 11.7$  Gyr, and  $11.7 \text{ Gyr} < t < 13.7$  Gyr. The priors used for our PROSPECTOR runs are listed in Table 1. Finally, the posterior distributions are obtained exploiting the Dynamic Nested Sampling package *dynesty* (Speagle 2020).

(iii) The PENALIZED PIXEL-FITTING METHOD (PPXF) code derives the stellar or gas kinematics and stellar population from absorption-line spectra of galaxies, using a maximum penalized likelihood method. The original approach was presented in Cappellari & Emsellem (2004) and then improved in Cappellari (2017). We used PPXF to estimate line-of-sight velocity dispersions for our observed ETGs. The penalization of pixels that are not well fit minimizes the mismatch with the templates employed. We ran PPXF with the MILES library.

## 2.2 TNG100 simulation

In this work, we extract simulated ETGs from IllustrisTNG<sup>9</sup> (Marnacci et al. 2018; Naiman et al. 2018; Nelson et al. 2018; Pillepich et al. 2018b; Springel et al. 2018), the successor to the original Illustris<sup>10</sup> simulation suite (Genel et al. 2014; Vogelsberger et al. 2014a,b; Sijacki et al. 2015). The data are publicly available<sup>11</sup> and presented in Nelson et al. (2019a). IllustrisTNG is a state-of-the-art magnetohydrodynamic cosmological simulation that models the formation and evolution of galaxies within the  $\Lambda$ CDM framework. As its predecessor, IllustrisTNG exploits all the advantages of the *unstructured moving-mesh* hydrodynamic method AREPO (Springel 2010), but improves the numerical methods, the subgrid physical model, and the recipe for galaxy feedback both from stars and active galactic nuclei (AGN). In particular, IllustrisTNG is equipped with a novel *dual mode* (thermal and kinetic) AGN feedback that shapes and regulates the stellar component within massive systems, maintaining a realistic gas fraction (Weinberger et al. 2017). Also the feedback model from galactic winds has been improved to have better representation of low- and intermediate-mass galaxies (Pillepich et al. 2018a).

The IllustrisTNG model was calibrated to significantly reduce tensions between the original Illustris suite and observations. As shown in fig. 4 of Pillepich et al. (2018a), some of these relevant improvements include the SFR density as a function of time, the stellar-to-halo mass relation in the present-day Universe, the stellar

mass function, the black hole mass–stellar mass relation, the black hole mass–halo mass relation, the gas content within virial radii, and galaxy sizes.

The IllustrisTNG simulation suite consists of three simulation volumes: TNG50 (Nelson et al. 2019b; Pillepich et al. 2019), TNG100, and TNG300, corresponding to three different box sizes with sides of about 50, 100, and 300 Mpc, respectively. The project assumes a  $\Lambda$ CDM cosmology with cosmological parameters taken from Planck Collaboration XIII (2016). Each run starts at  $z = 127$  using the Zeldovich approximation and evolves down to  $z = 0$ .

We use the highest resolution version of the medium volume size TNG100, i.e. TNG100-1 (hereafter, simply TNG100). This run includes approximately  $2 \times 1820^3$  resolution elements. The dark matter (DM) and baryonic mass resolutions are  $m_{\text{DM}} = 7.5 \times 10^6 M_\odot$  and  $m_{\text{b}} = 1.4 \times 10^6 M_\odot$ . The softening length employed for this version for both the DM and stellar components is  $\epsilon = 0.74$  kpc, while an adaptive gas gravitational softening is used, with a minimum  $\epsilon_{\text{gas,min}} = 0.185$  kpc. In particular, we take into account the properties of subhaloes from snapshot #91, corresponding to  $z = 0.1$ , close to the mean redshift of galaxies in the MaNGA survey.

### 2.2.1 In situ and ex situ stars in IllustrisTNG galaxies

In the last decade, cosmological simulations have suggested that accretion contributes to the mass and size growth of massive galaxies. The fraction of accreted stars depends on the stellar and DM masses (e.g. Oser et al. 2010; Lackner et al. 2012; Pillepich et al. 2014). From the original Illustris simulation suite, Rodriguez-Gomez et al. (2016) derived the *ex situ* fraction for galaxies with stellar masses between  $10^9 M_\odot$  and  $10^{12} M_\odot$ . They found that this fraction increases from  $\lesssim 10$  per cent in the least massive galaxies to above 80 per cent in the most massive systems. A similar analysis has been conducted on the IllustrisTNG runs: Pillepich et al. (2018b) analysed stellar masses within different apertures and found that, at  $z = 0$ , the low-mass tail of galaxies are mainly formed by *in situ* stellar particles, while central galaxies living in the most massive haloes, i.e.  $\log(M_{200c}/M_\odot) > 14$ , accreted more than 80 per cent of their total stellar mass via mergers. Moreover, by considering stellar masses within an aperture larger than 100 kpc, the *ex situ* fraction is found to be dramatically dominant at these distances, exceeding sometimes 90 per cent of the total mass. The relative contribution in massive systems of the *ex situ* component reaches around 60 per cent in the innermost regions ( $< 10$  kpc).

In this paper, we adopt the same definition of *in situ* and *ex situ* stars as used in Rodriguez-Gomez et al. (2016), Pillepich et al. (2018b), and Tacchella et al. (2019), exploiting the method for reconstructing the baryonic merger trees of Rodriguez-Gomez et al. (2015):

(i) *in situ stars* are those stellar particles that formed in a galaxy belonging to the main progenitor branch of the merger tree.

(ii) *ex situ stars* are those stellar particles that, at the time of their formation, were bound to a galaxy outside the main progenitor branch of the descendant galaxy.

## 2.3 ETG selection

We aim to perform a comparison that is as consistent as possible between TNG100 and MaNGA. The first question at hand is how to select ETGs. We opt for a simple selection based on  $(g - r)$  rest-frame colours, identifying hereafter ETGs as *Red Galaxies*, that is those galaxies with  $(g - r) > 0.6$ , a value that marks the transition between the blue cloud and the red sequence of galaxies. Nelson et al. (2018)

<sup>9</sup>Official website at <https://www.tng-project.org>.

<sup>10</sup>Official website at <https://www.illustris-project.org>.

<sup>11</sup><https://www.illustris-project.org/data/>

consider three models for assigning *ugriz* magnitudes to simulated galaxies, comparing them with those of SDSS galaxies. The authors discuss in detail the results obtained using colours derived through the so-called *resolved dust model* (Model C) which accounts for the presence of dust, following the distribution of neutral gas in galaxies, and adding also the attenuation caused by the presence of metals. The data from Model C are available in the supplementary catalogue SDSS *ugriz* Stellar Photometry/Colors with Dust.<sup>12</sup> In Nelson et al. (2018), the IllustrisTNG colours were compared with the observed colours of SDSS DR12 (Alam et al. 2015) galaxies in the present-day Universe ( $z < 0.1$ ). The distributions of  $(g - r)$  colours recover the colour bimodality of SDSS galaxies. In SDSS, the blue and red galaxy populations show two characteristic peaks at  $(g - r) \approx 0.4$  and  $\approx 0.8$ , respectively. Moreover, above  $M_* \simeq 10^{10.5} M_\odot$  the colour bimodality tends to disappear and red galaxies dominate. For this paper, to select ETGs in our observed MaNGA sample, we retrieved the *ugriz* Petrosian magnitudes from the NSA catalogue. To sum up, in the following text, we will present the results of our analysis for MaNGA and TNG100 ETGs selected as such considering only objects with  $(g - r) > 0.6$ .

## 2.4 Stellar mass estimates

Another question to consider when comparing observations and simulations is the consistency of the stellar mass measurements. Indeed, differences in the measurements of stellar masses can be caused by several factors, such as the fitting method used to derive luminosities and colours, as well as the stellar population synthesis models and libraries assumed. Sonnenfeld, Wang & Bahcall (2019) discuss the differences in deriving luminosity of galaxies in massive ETGs observed with the Hyper-Supreme Cam (HSC; Miyazaki et al. 2018) Subaru Strategic Program (Aihara et al. 2018, DR1), assuming either a simple Sérsic fit or a Sérsic + Exponential fit. The difference between the two methods can cause a variation of around 0.1 dex on the measurements of luminosity for the same object. Moreover, a different assumption of IMF can imply a global shift of stellar masses and the potential presence of IMF radial variations can introduce biases. The radius used to estimate stellar mass, or the quality of the observational data, can also be an important factor (e.g. Huang et al. 2018; Ardila et al. 2021).

In this work, we assume the stellar mass estimates from the UPENN\_PHOTDEC\_MSSTAR<sup>13</sup> catalogue of Meert, Vikram & Bernardi (2015) for MaNGA ETGs. In particular, these stellar masses, obtained by multiplying the stellar mass-to-light ratios ( $M_*/L$ ) from Mendel et al. (2014) by the luminosities from the PyMorph SerExp (Sérsic + Exponential) photometry, assume  $M_*/L$  fitting models that account for the effects of dust extinction (table 3 of Mendel et al. 2014).

For TNG100 galaxies we consider the 2D projected stellar mass defined as the sum of all bound stellar particles within a projected radius  $R = 2R_{\text{hm}}$ , where  $R_{\text{hm}}$  is the radius of a circle containing half of all stellar particles bound to each subhalo. Hereafter, we will refer to *2hm mass* as the projected mass within a radius of  $R = 2R_{\text{hm}}$ .

Table 2 summarizes the properties of the final TNG100 and MaNGA *Red Galaxy samples*, i.e. objects with  $\log(M_*/M_\odot) \geq 10.5$  and  $(g - r) > 0.6$ . Fig. 1 shows the colour–mass diagrams and the stellar mass distributions of both samples. As clearly visible,

**Table 2.** Summary table of the MaNGA and TNG100 samples. Column 1: sample. Column 2: number of ETGs. Column 3: stellar mass range. Column 4: mean stellar mass. Column 5: median stellar mass. Stellar masses are in units of  $M_\odot$ .

Sample	$N_{\text{ETG}}$	$(\log M_{*, \text{min}};$ $\log M_{*, \text{max}})$	$\log M_{*, \text{mean}}$	$\log M_{*, \text{median}}$
MaNGA	1427	(10.50; 12.26)	11.07	11.04
TNG100	1543	(10.50; 12.27)	10.83	10.76

the distribution of the stellar masses for MaNGA galaxies appears almost flat. The reason for such distribution is due to the original MaNGA sample design: as argued in Wake et al. (2017), the MaNGA sample was built in such a way that the most massive galaxies are located at higher redshifts, but, at the same time, both the Primary and Secondary samples are selected to have flat stellar mass distributions. We will account for this effect later in the paper.

## 3 RADIAL PROFILES OF STELLAR PROPERTIES

In Sections 3.1 and 3.2, we describe the method used to compute the radial profiles from the observed and simulated 2D stellar galaxy images. In Sections 3.3 and 3.4, we describe how we compute median profiles in different stellar mass bins.

### 3.1 Radial profiles for MaNGA ETGs

To obtain stellar properties at different galactocentric distances for each MaNGA galaxy we adopt the approach described in Oyarzún et al. (2019), with the difference that we consider radial binning in physical units instead of units of effective radii. Specifically, by considering the axial ratio of each source obtained from *r*-band photometric images, elliptical polar radii are associated with spaxels. We then bin in five concentric elliptical annuli each galaxy map, assuming the following radii as the edges of each bin:  $R/\text{kpc} = \{0; 2; 4; 10; 20; 100\}$ . The choice of using a radial binning in physical units is justified by the fact that effective radius measurements may be affected by the depth of the survey. For example, HSC measures different effective radii than SDSS (Huang et al. 2018).


The next step after radial binning consists in shifting spectra to the rest frame by taking the stellar systemic velocity from DAP as a reference. A Voronoi binning is then applied to the maps, considering a minimum  $S/N = 10$  in each bin. Spectra belonging to the same annulus are co-added and, after running PXP with the MILES library, they are stacked to estimate the line-of-sight stellar mean velocity and velocity dispersion.

### 3.2 Radial profiles for TNG100 ETGs

To obtain the radial profiles of stellar properties for TNG100 galaxies, we apply the same method presented in Ardila et al. (2021). We firstly project the 3D particle distributions of simulated galaxies on a 2D  $X$ – $Y$  plane using the `hydrotools` package (Diemer et al. 2018, 2019). For each subhalo, the 2D map consists of 300 pixels per side, with a resolution of 1 kpc per pixel (for a total physical side length of the map of 300 kpc). To extract the 1D stellar profiles we then use the method presented in Huang et al. (2018) and also used by Ardila et al. (2021), which we now briefly describe. We extract 1D stellar mass surface density profiles using the galaxy surface brightness profile

<sup>12</sup>Available at <https://www.tng-project.org/data/downloads/TNG100-1/>.

<sup>13</sup>Available at [http://alan-meert-website-aws.s3-website-us-east-1.amazonaws.com/fit\\_catalog/download/index.html](http://alan-meert-website-aws.s3-website-us-east-1.amazonaws.com/fit_catalog/download/index.html).



**Figure 1.** Left panel: MaNGA (red dots) and TNG100 (2D grey histogram) ETGs. Right panel: 1D histograms of the mass distribution for the MaNGA (red histogram) and the TNG100 (grey histogram) estimates. The histogram of MaNGA ETGs stellar masses is flat due to the MaNGA selection function (see Wake et al. 2017).

function included in the *kungpao* package.<sup>14</sup> Galaxy centroids are identified by means of *extract*, a function included in the *sep* library, and the *ellipse* algorithm is used to fit concentric elliptical isophotes. The position angle and ellipticity of these isophotes are the mean values from the 2D fitting procedure of the galaxy maps. The isophotes are spread over the range 1–150 kpc, in 20 concentric elliptical annuli of constant width in logarithmic space. For stellar metallicity, age, and line-of-sight velocity dispersion profiles, we use the same centre and ellipticity of the isophotes computed on the stellar mass surface density mass maps. To derive mass-weighted stellar metallicity, age, and velocity dispersion, each pixel is weighted by the corresponding value of stellar mass in that pixel. The entire procedure is applied to both the *in situ* and *ex situ* stellar populations, starting from their 2D stellar property maps.

Another effect we accounted for is about the differences in spatial resolutions. For this, we associate each simulated galaxy an angular diameter distance, assuming a redshift drawn from the  $z$ – $M_*$  distribution of the MaNGA sample (see Fig. A1 in Appendix A). For each TNG100 ETG, we smooth the galaxy map with a 2D Gaussian filter kernel:

$$\sigma_{\text{kernel},i} = \sqrt{\mathcal{R}_{\text{MaNGA},i}^2 - \mathcal{R}_{\text{TNG}}^2}, \quad (1)$$

where  $\mathcal{R}_{\text{TNG}} = 1$  kpc is the resolution of the TNG100 sample,  $\mathcal{R}_{\text{MaNGA},i} = \sin(\text{PSF}_{\text{MaNGA}})d_{A,i}$ , with  $\text{PSF}_{\text{MaNGA}} = 2.5$  arcsec ( $\simeq 1.21 \times 10^{-5}$  in radians), and  $d_{A,i}$  is angular diameter distance (in kpc) for the  $i$ -th galaxy in the TNG100 sample determined as described above. For example, at  $z = 0.05$ ,  $\mathcal{R}_{\text{MaNGA}} \simeq 2.52$  kpc. For a given simulated ETG we compute two types of radial profiles for each stellar physical property (for both the *in situ* and *ex situ* stellar populations): the *uncovolved profile* and the *convolved profile*,

the latter obtained by smoothing the projected maps with the 2D Gaussian filter kernel  $\sigma_{\text{kernel}}$ .

### 3.3 Building stellar mass bins

To compare the profiles of stellar properties between MaNGA and TNG100, we divide galaxies into bins of stellar mass and compute the median profile in each stellar mass bin along with the associated uncertainties. As a fiducial choice, we compare galaxies at *fixed stellar mass*. The three stellar mass bins used are  $10.5 \leq \log(M_*/M_\odot) < 11$ ,  $11 \leq \log(M_*/M_\odot) < 11.5$ , and  $\log(M_*/M_\odot) \geq 11.5$ . In Appendix B, we present the same analysis using number-density-based bins, highlighting the differences with respect to the use of bins at fixed stellar mass.

### 3.4 Building median radial profiles with errors

We use a Bayesian hierarchical approach (e.g. Cannarozzo et al. 2020) to estimate the median values and the associated  $1\sigma$  uncertainties on the observed and simulated radial profiles. We assume that each stellar property  $X$  in any radial bin has a Gaussian distribution, so that its likelihood can be written as

$$P(X|X^{\text{data}}, \sigma_X^{\text{data}}, \mu, \sigma) = \frac{1}{\sqrt{2\pi\sigma_X^2}} \exp\left\{-\frac{(X^{\text{data}} - \mu)^2}{2\sigma_X^2}\right\}, \quad (2)$$

where  $X$  is the quantity that we want to infer in each radial bin (e.g. the logarithm of the stellar mass surface density), while  $X^{\text{data}}$  and  $\sigma_X^{\text{data}}$  are the data values and their related uncertainties, respectively. The variance in equation (2) has the form

$$\sigma_X^2 = \sigma_X^{\text{data}2} + \sigma^2. \quad (3)$$

In equations (2) and (3),  $\mu$  and  $\sigma$  are the two *hyper-parameters* of our Bayesian hierarchical approach and represent the mean value and the intrinsic scatter of the distribution of the quantity  $X$ , respectively. We underline that, in the case of simulated ETGs, equation (3) reduces

<sup>14</sup>The *kungpao* library is available at <https://github.com/dr-guangtou/kungpao/>

**Table 3.** Hyper-parameters used to compute MaNGA and TNG100 profiles of stellar properties. Column 1: stellar property. Column 2: uniform prior on the mean (lower bound; upper bound). Column 3: uniform prior on the intrinsic scatter (lower bound; upper bound).

Stellar property	$(\mu_{\min}; \mu_{\max})$	$(\sigma_{\min}; \sigma_{\max})$
Surface density $\log \Sigma_* [\text{M}_\odot \text{kpc}^{-2}]$	(0; 11)	(0; 2)
Metallicity $\log Z_* [Z_\odot]$	(−1; 1)	(0; 1)
Age [Gyr]	(0; 13)	(0; 5)
Velocity dispersion $\sigma_* [\text{km s}^{-1}]$	(0; 350)	(0; 100)

to  $\sigma_{\bar{x}}^2 = \sigma^2$ , since no uncertainties are associated with simulated properties. These parameters are estimated independently in each bin. In Table 3, we list the priors adopted for each property. The stellar properties of MaNGA and TNG100 galaxies are sampled adopting a Markov chain Monte Carlo approach using 10 random walkers and 300 steps (removing the first 200 steps) for each run to reach the convergence of the hyper-parameter distributions. We use the Python adaptation of the affine-invariant ensemble sampler of Goodman & Weare (2010), EMCEE, by Foreman-Mackey et al. (2013).

### 3.5 MaNGA and TNG100 stellar properties

In our comparison between MaNGA and TNG100 ETGs, we consider the circularized radial distributions of the stellar mass surface density  $\Sigma_*$ , stellar metallicity  $Z_*$ , stellar age, and line-of-sight stellar velocity dispersion  $\sigma_*$ .

As already described in Sections 3.1 and 3.2, in order to construct the radial profiles of the aforementioned properties, we build five concentric radial bins for MaNGA ETGs from the innermost regions out to 100 kpc and 20 log-spaced bins out to 150 kpc for TNG100 objects, for which we are able to split the relative contribution of *in situ* and *ex situ* stellar populations. We use mass-weighted stellar metallicities and ages for both MaNGA and TNG100 galaxies. For these two properties, we also tested the luminosity-weighted measurements for MaNGA galaxies finding no significant difference between the two choices. Velocity dispersions, instead, are mass-weighted for simulated galaxies and luminosity-weighted for observed sources.<sup>15</sup>

The stellar mass surface density, metallicity, and age measurements for MaNGA sources are computed using both FIREFLY and PROSPECTOR, while for line-of-sight velocity dispersions we use pPXF (see Section 2.1). For MaNGA galaxies, the median profiles and related uncertainties in each radial bin are computed as described in Section 3.4, but imposing the condition that the measurements are available for at least 75 per cent of the sample in each stellar mass bin. FIREFLY and PROSPECTOR measurements are derived using stellar population libraries that assume different values of

<sup>15</sup>For TNG100 galaxies, the maps of stellar metallicity, age, and velocity dispersion are all weighted by stellar masses for consistency within the simulated sample. We note that, for the velocity dispersion, the comparison between simulated and observed galaxies is not fully self-consistent, because the velocity dispersion of the MaNGA galaxies is weighted by luminosity. However, as shown in Fig. 2, the simulated galaxies have old stellar populations ( $\gtrsim 7$  Gyr), responsible for their red colours. Moreover, the age distributions of TNG100 galaxies are almost flat at all stellar masses, in particular going towards the high-mass tail, suggesting the presence of similar stellar populations. In light of that, we do not expect for our sample of old and red galaxies (selected as such as described in Section 2.3) a significant discrepancy between mass- and luminosity-weighted velocity dispersion profiles.

solar metallicity ( $Z_{\odot, \text{FIREFLY}} = 0.019$  and  $Z_{\odot, \text{PROSPECTOR}} = 0.0142$ ). For this reason, to homogenize and make comparisons easier, all the observed and simulated stellar metallicity profiles are normalized by their respective values at  $\approx 7$  kpc, which corresponds to the median radius of the third MaNGA bin.

Since our main goal is to provide a possible evolutionary scenario on how present-day Universe ETGs have been formed throughout the cosmic history, all the stellar properties considered in this work (for both observed and simulated objects) are *at face value*, i.e. as directly derived from pipelines and fitting codes for MaNGA, and from output catalogues for TNG100. Thus, no mock observational data of simulated galaxies have been produced, as instead done for instance by Nanni et al. (2022).

## 4 RESULTS

We now compare the stellar mass surface density, metallicity, age, and line-of-sight velocity dispersion profiles obtained for our observed and simulated samples of ETGs. We focus here on the results corresponding to the stellar mass bins. The results for the number-density-based bins are shown in Appendix B.

### 4.1 Stellar mass surface density profiles

Fig. 2 shows the results for the total stellar populations. For TNG100, both the raw profiles and profiles convolved with the MaNGA PSF are displayed. We show profiles both using the median estimates from FIREFLY (yellow dots) and PROSPECTOR (green squares). In each stellar mass bin, we find a satisfying agreement at all radii between the two MaNGA measurements, with a small systematic shift to higher values with PROSPECTOR. This is consistent with the observed systematic shift in the total mass estimates (for the same galaxy, PROSPECTOR infers a stellar mass that, on average, is higher than the estimate obtained by FIREFLY by  $\approx 0.15$ – $0.2$  dex). In each stellar mass bin, we find a remarkable agreement between the stellar mass surface densities of MaNGA ETGs and those from TNG100, both in the shape and normalization of the profiles.


In Fig. 3, we separate the TNG100 profiles into *in situ* (red dashed curves) and *ex situ* (blue dotted curves) components. In the mass range  $10.5 < \log (M_*/M_\odot) < 11$ , on average, the *in situ* stellar component is found to be dominant out to  $\approx 30$  kpc and *ex situ* stars dominate at larger radii. The central mass bin, i.e.  $11 < \log (M_*/M_\odot) < 11.5$ , reveals the increasing contribution from *ex situ* stars – the profiles of the two populations contribute in almost equal proportions over the entire radial range. Above  $\log (M_*/M_\odot) \approx 11.5$ , the *ex situ* stars dominate at all radii. To summarize, we find that below  $\log (M_*/M_\odot) \approx 11$  the most relevant stellar component (out to  $\approx 30$  kpc) is the *in situ* population, whereas at higher stellar masses the *ex situ* stars become dominant across the entire radial range.

### 4.2 Metallicity profiles

The second row of plots in Fig. 2 displays the metallicity profiles in three stellar mass bins. The two MaNGA profiles show similar radial distributions, which differ only beyond 10 kpc for the most massive systems by a factor  $\lesssim 0.2$  dex at  $\approx 20$  kpc.<sup>16</sup>

<sup>16</sup>If the two MaNGA stellar metallicity distributions are not renormalized at their  $\sim 7$  kpc values, the FIREFLY profiles, on average, are shifted up from the PROSPECTOR profiles by a factor of  $\lesssim 0.05$  dex at  $\log (M_*/M_\odot) \lesssim 11.5$ , and can differ even of  $\lesssim 0.16$  dex for galaxies with  $\log (M_*/M_\odot) \gtrsim 11.5$ .






**Figure 2.** Radial profiles of stellar mass surface density, metallicity (normalized by the corresponding values of metallicity at  $\approx 7$  kpc), age, and line-of-sight velocity dispersion (from top to the bottom) in three bins of stellar mass for MaNGA and TNG100 ETGs. The green, yellow, and violet dots represent the median estimates, respectively, for MaNGA galaxies from FIREFLY, PROSPECTOR, and PPXF. Both the stellar metallicity and age measurements in MaNGA and TNG100 ETGs are mass-weighted. Velocity dispersions are luminosity-weighted for MaNGA and mass-weighted for TNG100 sources. The vertical grey dashed lines indicate the five radial bins for MaNGA. The two black curves represent the median values of each stellar property for the total stellar population in TNG100. The intrinsic profiles are shown with the thin curve, whereas the thick curves indicate the results when convolved with the MaNGA PSF. The light grey hatched area ( $R \lesssim 2.1$  kpc) shows three times the gravitational softening length of the stellar particles in TNG100. The grey shaded area is the region that lies in between the profiles obtained from the original and the convolved TNG100 maps. Because we account for the MaNGA PSF but not for the effects of the resolution of the simulation, the grey shaded area gives a sense of the uncertainty in the comparison in the inner regions. The errorbars represent the  $1\sigma$  uncertainties on the median for MaNGA and TNG100 estimates.

Globally, TNG100 profiles reproduce fairly well the shapes of the two MaNGA estimates. In particular, below  $10^{11} M_{\odot}$ , the TNG100 MaNGA-PSF-convolved profile reproduces well the median profiles from FIREFLY and PROSPECTOR. Over the interval between  $10^{11} M_{\odot}$  and  $10^{11.5} M_{\odot}$ , the median profiles of MaNGA ETGs lie in between the two TNG100 profiles derived from the MaNGA-

PSF-unconvolved and convolved maps within around 4 kpc, and almost overlap beyond this distance with the two TNG100 profiles. Above  $10^{11.5} M_{\odot}$ , the FIREFLY profile is quite well represented even in the outermost regions, while the PROSPECTOR distribution tends to assume lower values in metallicity, differing from the TNG100 profiles by a factor of  $\lesssim 0.1$  dex. We stress here the importance of



**Figure 3.** Contribution from *in situ* and *ex situ* stellar populations to the radial profiles of TNG100 ETGs. Top to bottom corresponds to profiles of stellar mass surface density, metallicity (not normalized), age, and line-of-sight velocity dispersion. The stellar metallicity, age, and velocity dispersion measurements are mass-weighted. The black solid, red dashed, and blue dotted curves correspond to the total, *in situ*, and *ex situ* stellar populations, respectively. For clarity reasons, here we only show the median profiles convolved with the MaNGA PSF, and we omit the errorbars for the *in situ* and *ex situ* stellar population profiles. The light grey hatched area ( $R \lesssim 2.1$  kpc) shows three times the gravitational softening length of the stellar particles in TNG100. *Note:* the range shown along the y-axes are different from those of Fig. 2.

applying a smoothing using the MaNGA PSF on the original maps of simulated objects. Indeed, at all bins the stellar metallicity profiles from the original maps are steeper than those obtained from the convolved maps, the latter giving values lower by  $\approx 0.15$ – $0.2$  dex in the innermost regions ( $R \lesssim 4$  kpc).

The necessity of renormalizing metallicity measurements to reconcile observations and simulations has been already highlighted by Nelson et al. (2018). The right panel of fig. 2 of Nelson et al. (2018) shows the stellar mass–metallicity relations for TNG100 (and TNG300) compared with observed estimates in the present-day

Universe from Gallazzi et al. (2005), Woo, Courteau & Dekel (2008), and Kirby et al. (2013). Above  $\log(M_*/M_\odot) \approx 10.5$ , simulations and observations almost agree in shape, showing a weak scaling with stellar mass (with a supersolar metallicity normalization). However, the almost-flat trend of the metallicity as a function of stellar mass in simulations implies a discrepancy of up to  $\approx 0.5$  dex at  $\log(M_*/M_\odot) < 10.5$  from observed estimates. A possible reason for the origin of such a discrepancy can be found in different methods to derive simulated and observed metallicities. Indeed, when opportune corrections and spectral fitting codes similar to those

adopted on observational data are applied to simulated galaxies, the aforementioned discrepancy reduces, making the estimates of TNG100 more consistent with those from observations.

The *in situ* and *ex situ* PSF-convolved stellar metallicity profiles (second row of plots in Fig. 3) are shown without adopting any normalization. TNG100 galaxies are characterized by *ex situ* stars that are more metal rich than the *in situ* population: this metallicity difference increases for increasing stellar mass. This apparently counter-intuitive finding could be in tension with the expected scenario from downsizing. A possible explanation for the presence of such metal-rich *ex situ* stellar populations in these massive systems could be ascribed to the fact that, because of the substantial *ex situ* fraction accreted via major mergers (see Section 5) across their stellar mass assembly histories, many galaxies that were centrals at a given snapshot became satellites of slightly more massive systems soon thereafter.

### 4.3 Age profiles

Fig. 2 also displays a comparison between the radial distributions of stellar ages. The stellar age profiles derived from FIREFLY and PROSPECTOR show a common behaviour in all mass bins, namely, a systematic shift in age is found between the two stellar fitting codes. On average, FIREFLY and PROSPECTOR differ in their age estimates by about 1.5–2.5 Gyr (on average, FIREFLY estimates are  $\approx 20$  per cent younger than those from PROSPECTOR). The systematic difference in age obtained by the two codes might be partially explained in terms of the *age–metallicity degeneracy*: the red colours that characterize old stellar populations can be explained also assuming a higher metallicity, and vice versa (Worthey 1994). Indeed, for the same sample of ETGs, on average, FIREFLY derives more metal-rich and younger stellar populations compared to PROSPECTOR. However, this degeneracy should ideally be reflected in the uncertainty values produced by the stellar population synthesis codes. The fact that the measurements are inconsistent may also suggest that elements of the models used by the two codes, such as the stellar libraries, are themselves inconsistent, and not sufficiently flexible. A crucial point is that deriving stellar ages for such old systems is not trivial (see Conroy 2013). Indeed, stellar age grids for these models are sparse at these ages because they tend to be log-spaced. When building non-parametric SFHs, these codes interpolate over the ages sampled by the stellar libraries and isochrones. The large gap between the two stellar age profiles displayed in Fig. 2 can be taken as a measure of the systematic uncertainty on the age of the observed galaxies in our sample. It is clear that these age profiles have little constraining power on theoretical models: the TNG100 profiles lie in between FIREFLY and PROSPECTOR, reproducing only the quasi-flat distributions of observed data. The analysis of the radial distributions of age for the *in situ* and *ex situ* stellar populations in the simulated ETGs (see Fig. 3) shows that, below  $\log(M_*/M_\odot) \approx 11$ , the *ex situ* component is older (up to +1.5 Gyr) than the *in situ* component over the entire radial range, whereas the inner regions ( $R \lesssim 6$  kpc) of galaxies with  $11 \lesssim \log(M_*/M_\odot) \lesssim 11.5$  are composed of *in situ* and *ex situ* stars with similar ages. Above  $\log(M_*/M_\odot) \approx 11.5$ , *ex situ* stars are found to be older (up to +2.5 Gyr) than the *in situ* population at all radii.

### 4.4 Velocity dispersion profiles

The bottom panels in Fig. 2 compares the radial profiles of line-of-sight stellar velocity dispersions for simulated and observed ETGs. Here, the MaNGA values are derived using the PPF code.

Below  $\log(M_*/M_\odot) \approx 11.5$ , we generally find a good first-order agreement in normalization between MaNGA and TNG100, but the MaNGA profiles are steeper than those of TNG100. More quantitatively, the difference between the two median profiles can be at most of around  $30 \text{ km s}^{-1}$ . Instead, above  $\log(M_*/M_\odot) \approx 11.5$ , the velocity dispersion profiles of both MaNGA and TNG100 galaxies are almost flat out to  $R \approx 40$  kpc. However, we find an essentially radius-independent difference between the two profiles, with velocity dispersions for the simulated ETGs generally higher by  $30\text{--}40 \text{ km s}^{-1}$  (this will be discussed further in Section 4.5). We underline that the effect on the observed velocity dispersion  $\sigma_*$  for MaNGA galaxies is the result of two main contributions, i.e. the instrumental dispersion  $\sigma_{\text{inst}}$  and the intrinsic stellar velocity dispersion  $\sigma_{*,\text{int}}$ :

$$\sigma_*^2 = \sigma_{\text{inst}}^2 + \sigma_{*,\text{int}}^2. \quad (4)$$

In fact, over the rest-frame optical range  $0.36 \lesssim \lambda/\mu\text{m} \lesssim 1.03$ , the spectral resolution is  $R \sim 2000$  and the  $1\sigma$  dispersion of the instrumental spectral line-spread function is about  $70 \text{ km s}^{-1}$  (see Westfall et al. 2019; Law et al. 2021). As reported in Westfall et al. (2019), for  $\sigma_* \gtrsim 100 \text{ km s}^{-1}$ , the uncertainties on velocities can be approximated as  $\delta(\Delta v) \approx \langle \sigma_* \rangle / (S/N)_g$ , where  $\langle \sigma_* \rangle$  is the mean velocity dispersion and  $(S/N)_g$  is the  $g$ -band signal-to-noise ratio. At  $(S/N)_g = 10$ , the typical uncertainties on velocities are around 10 per cent of  $\sigma_*$ . For  $\sigma_* \gtrsim 100 \text{ km s}^{-1}$ , the uncertainties on velocity dispersions are slightly larger than those of velocities, but can be roughly approximated by a single proportionality constant. Given that, our estimates of velocity dispersion profiles are quite robust, because, on average, these measurements are greater than  $100 \text{ km s}^{-1}$ , with the only exception of the last radial bin of the profile for galaxies with  $10.5 \leq \log(M_*/M_\odot) < 11$ , which is around  $90 \text{ km s}^{-1}$ .

The line-of-sight stellar velocity dispersion profiles for the *in situ* and *ex situ* components (bottom panels in Fig. 3) almost coincide in the intermediate- and high-mass bins, while in the low-mass bin the velocity dispersion is higher in the centre for the *in situ* component. As it is well known, in the same gravitational potential, the line-of-sight stellar velocity dispersion profile of a given component depends on both its intrinsic velocity distribution and its density distribution: for instance, for isotropic velocity distribution, the steeper the density profile, the lower the velocity dispersion (see Nipoti et al. 2021). In the low-mass bin, the higher central velocity dispersion of the *in situ* component can be qualitatively explained by its shallower surface density profile (see top-left panel in Fig. 3).

### 4.5 Central versus satellite galaxies

We now consider the radial profiles separately for central and satellite galaxies. Haloes and subhaloes in IllustrisTNG are detected by SUBFIND, the subhalo finder code developed by Springel et al. (2001). Specifically, an IllustrisTNG subhalo is classified as *central* (flag `is_primary==1`) if it is the subhalo with the deepest potential well among those belonging to the same friends-of-friends (FoF) halo. Otherwise, subhaloes are classified as *satellites*. To separate MaNGA ETGs into centrals and satellites we rely on the classification provided by Yang et al. (2007), obtained for a sample of more than 300 000 galaxies from SDSS DR4 (Adelman-McCarthy et al. 2006).

Though they are not shown here, we do find that the MaNGA and TNG100 stellar mass surface density profiles, as well as those for the stellar metallicity and age, are in excellent agreement between central and satellite ETGs. This lack of differences in the radial profiles between these two populations is consistent with previous

outcomes in literature, as for instance shown by Santucci et al. (2020) for stellar age and metallicity properties.

Centrals and satellites display similar trends also for the *in situ* and *ex situ* components over the entire mass range considered. We might expect central galaxies to exhibit larger *ex situ* components compared to satellites. However, a possible explanation for the observed similarity could be that most satellites in the considered mass range were recently accreted onto the main halo and acquired a significant fraction of their *ex situ* component when they were centrals of other haloes.

Fig. 4 compares the line-of-sight stellar velocity dispersion profiles for centrals and satellites separately and displays a key result in this paper. Although for MaNGA we find that massive centrals and satellites show similar velocity dispersion profiles, in contrast, for the most massive bin we find that TNG100 predicts a  $\approx 50 \text{ km s}^{-1}$  offset between centrals and satellites. The amount of DM in central simulated galaxies could provide an explanation for the significant difference in velocity dispersion between TNG100 centrals and satellites. As discussed in Lovell et al. (2018), TNG100 predicts an important enhancement of the DM content in the inner regions of subhaloes. Hence, this high fraction of DM, which dominates galaxies at  $z \approx 0$ , may be the responsible of this high velocity dispersion especially for the most massive central galaxies. Our observations appear to exclude a difference in the velocity dispersion profile between centrals and satellites. However, determining whether a galaxy is a central or a satellite is notoriously difficult, and misclassifications in the Yang et al. (2007) catalogue may erase the observational signal. This possibility warrants further investigation before firm conclusions can be drawn.

#### 4.6 Robustness of results

We now present a discussion on the robustness of the results.

(i) *Sample matching method* – MaNGA galaxies can be matched to simulated galaxies according to stellar mass or number density. We have tried both and found only minor differences. These are discussed further in Appendix B.

(ii) *Different definitions of stellar masses* – Given the wide variety of possible systematic effects on the definition of stellar masses, in chapter 3 of Cannarozzo (2021) the same analysis was performed by testing also the SerExp Dust-free stellar masses (from Mendel et al. 2014), the Sérsic and Petrosian fit estimates from the original NSA catalogue, and the masses defined as the sum of the masses included in the five concentric annuli used to derive the profiles of stellar properties from FIREFLY and PROSPECTOR for MaNGA ETGs, while for TNG100 galaxies we considered also the stellar masses within a projected aperture of 30 kpc. These tests do not reveal significant differences from the analysis presented in this work.

(iii) *Different definitions of ETGs* – Also the ETG selection is a factor that could affect the results of this study. In appendix A of Diemer et al. (2019), the authors compare the ETG fractions derived from diverse selection methods in IllustrisTNG with that from the observed compilation of Calette et al. (2018). In particular, the authors measure the ETG fractions adopting classifications based on the concentration of the 3D stellar mass density profiles,  $C_{82}$ , defined as  $5 \times \log(r_{80}/r_{20})$ , with  $r_{80}$  and  $r_{20}$  as the radii including the 80 per cent and 20 per cent of the total stellar mass, on  $(g-r)$  colours, on spheroid-to-total ratios  $S/T$ , and on the fraction of kinetic energy that is in rotation  $\kappa_{\text{rot}}$ . They found that the best indicator of galaxy morphology able to better reproduce the ETG fraction from Calette et al. (2018) is  $C_{82}$ . Instead, the  $(g-r)$  classification

implies an excess of ETGs, and that the colours correlate weakly with structural parameters (as illustrated also in Rodríguez-Gomez et al. 2019). Tacchella et al. (2019) studied the connection between the star formation activity and morphology of central galaxies in IllustrisTNG, adopting as morphological indicators the parameters  $C_{82}$  and  $S/T$ . They found that the  $S/T$  parameter strongly correlates with  $(g-r)$  colours:  $S/T$  is higher for redder colours and higher stellar masses (while, at fixed mass,  $C_{82}$  is found to be weakly dependent on colour). Our choice of adopting a simple selection on colours is driven by the fact that mock colours in IllustrisTNG are generated consistently with observations (we remind that these mock colours are obtained following the observational prescriptions described in section 3 of Nelson et al. 2018). As done for the different stellar mass definitions, in chapter 3 of Cannarozzo (2021) we adopted another ETG selection for both the MaNGA and TNG100 samples, including only those objects with SFRs below 1 dex from the star-forming main sequence of galaxies (one of the methods to select passive systems presented in Donnari et al. 2019). Based on this extensive exploration, even the results here presented are robust and independent of the specific definitions of ETGs for both the observed and simulated sources.

As mentioned above, the selection of ETGs may involve different criteria, each of which may introduce some selection biases (see also Moresco et al. 2013). Usually, ETGs are characterized by either an elliptical (E) or a lenticular (S0) morphology. One of the historical criteria for morphological selection of galaxies is that based on the T-Type (de Vaucouleurs 1959). According to the T-Type-based classification, E/S0 galaxies have values between  $-6$  and  $-1$ , while the various types of spiral galaxies range between 0 and 9. In light of that, for MaNGA objects we also verified the impact of the adopted selection based on colours (see Section 2.3), checking the morphological type assigned by the MANGA MORPHOLOGY DEEP LEARNING DR15 CATALOG.<sup>17</sup> This catalogue, presented in Fischer, Domínguez Sánchez & Bernardi (2019), is built by exploiting the Deep Learning method for identifying the morphology of galaxies as described in Domínguez Sánchez et al. (2020) for all the objects of MaNGA DR15. In our Red Galaxy sample,  $\approx 66$  per cent of the total effectively shows a clear morphology compatible with an E/S0 type, of which  $\approx 33$  per cent are classified as lenticular galaxies.


## 5 DISCUSSION

In this section, we discuss our results and compare our findings with previous works.


### 5.1 The role of mergers in TNG100 galaxies

Fig. 3 shows that the shapes of the *in situ* and *ex situ* stellar mass surface density radial profiles from TNG100 ETGs are stellar-mass-dependent and that the *ex situ* component dominates the total profile at all radii above  $\log(M_*/M_\odot) \approx 11$ . This agrees with previous results from Pillepich et al. (2018b), Tacchella et al. (2019), and Pulsoni et al. (2021), who find that the stellar mass assembly history of very massive galaxies is driven by major mergers. Indeed, major mergers not only allow *ex situ* stellar populations to settle even in the innermost regions of galaxies, but also homogeneously mix the two stellar components at all radii, causing the formation of stellar mass surface density profiles which are similar in shape and differ only

<sup>17</sup> Available at [https://www.sdss.org/dr15/data\\_access/value-added-catalogs/?vac\\_id=manga-morphology-deep-learning-dr15-catalogue](https://www.sdss.org/dr15/data_access/value-added-catalogs/?vac_id=manga-morphology-deep-learning-dr15-catalogue)



**Figure 4.** Line-of-sight stellar velocity dispersion radial profiles for centrals (upper rows) and satellites (lower rows). The violet-filled black and white-filled violet diamonds represent the median estimates of the mass-weighted velocity dispersion profiles for MaNGA central and satellite ETGs from pPXF code, respectively. The vertical grey dashed lines delimit the five radial intervals within stellar velocity dispersion is computed for MaNGA ETGs. The shaded and hatched areas represent the regions delimited by the median values of the mass-weighted velocity dispersion profiles obtained from the original TNG100 maps and the maps convolved with the MaNGA PSF for centrals and satellites, respectively. The black, red, and blue colours correspond to the total, *in situ*, and *ex situ* stellar populations. For clarity reasons, we show only the errorbars for the MaNGA and TNG100 total stellar population profiles. The light grey hatched area ( $R \lesssim 2.1$  kpc) shows three times the gravitational softening length of the stellar particles in TNG100.



**Figure 5.** Data distributions of eight properties related to the merger histories of TNG100 ETGs. Top-left panel: total fractions of the *in situ* (red distributions) and the *ex situ* (blue distributions) stellar components. Top-right panel: fractions of the *ex situ* stellar component from minor (green) and major (orange) mergers normalized to the total *ex situ* stellar fraction. Bottom-left panel: number of minor (green) and major (orange) mergers across the redshift range  $0.1 \leq z \leq 2$ . Bottom-right panel: redshifts of the last minor (green) and major (orange) mergers. All distributions are shown for the same three stellar mass bins as in Figs 2 and 3. Except for the number of minor and major mergers, which are discrete values, the other properties are displayed as violin plots. For each property, the median value of the corresponding distribution is reported.

in their normalization. However, the results of IllustrisTNG disagree with some previous works finding that the growth of massive ETGs is primarily driven by minor mergers. (e.g. Naab et al. 2009; Oser et al. 2010; Hilz et al. 2013). For example, as argued in Genel et al. (2008) and Khochfar & Silk (2009), massive ( $\gtrsim 10^{11} M_{\odot}$ ) DM haloes undergo typically no more than one major merger in the redshift range  $0 \leq z \leq 2$ . There are also some observational works supporting the idea that massive ETGs may experience few major mergers (e.g. Bell et al. 2006; McIntosh et al. 2008), instead undergoing a high number of minor mergers (e.g. Bundy et al. 2009). On the contrary, other studies lend support to the idea in which the role of major mergers may be more relevant, estimating relatively high mass-weighted merger ratios. For example, Sonnenfeld, Nipoti & Treu (2017) infer for galaxies of  $\log(M_*/M_{\odot}) \approx 11$  a mass-weighted merger ratio greater than 0.4.

Fig. 5 displays how minor and major mergers contribute to the total mass of TNG100 ETGs as a function of redshift and galaxy mass. For each stellar mass bin, the data distributions are presented as *violin plots* (except for the number of mergers), which displays the probability density of the data smoothed by a kernel density estimator. The shape of each violin plot represents the frequency of data, so that the larger the violin’s body, the higher the density of data at a given y-axis value. Specifically, we use 100

data points to evaluate each Gaussian kernel density estimation. The complementary behaviour of the *in situ* and *ex situ* stellar mass fractions in the top-left panel of Fig. 5 confirms the rising importance of accreted stars in higher mass ETGs. In particular, the median of the *ex situ* stellar fraction grows from  $\approx 29$  per cent for galaxies with  $10.5 \leq \log(M_*/M_{\odot}) < 11$ , up to  $\approx 77$  per cent for the most massive systems. Following Rodriguez-Gomez et al. (2015, 2016),  $\mu_*$  is defined as the *stellar mass ratio* between the two progenitors of a given galaxy. A *major merger* is then defined by a  $\mu_* > 1/4$ , while a *minor merger* is defined by  $1/10 < \mu_* < 1/4$ . However, the fraction of accreted stars from other galaxies is not only due to major and minor mergers. It also includes stars from the so-called *very minor mergers*, i.e. with  $\mu_* < 1/10$ , as well as tidally stripped stars from surviving galaxies. As illustrated in the top-right panel of Fig. 5, the analysis of the *ex situ* fraction accreted via minor mergers relative to the total *ex situ* fraction, reveals that, on average, around 6 per cent of stars are accreted via minor mergers in ETGs with  $M_* < 10^{11} M_{\odot}$ , reaching  $\approx 14$  per cent in the most massive galaxies. By isolating the role of major mergers, their contribution presents broad distributions, with median values of around 50 per cent (relative to the whole *ex situ* stellar fraction) for ETGs with  $\log(M_*/M_{\odot}) > 11$ . The analysis of the relative contributions deriving from both minor and major mergers suggests that,

on average, below  $10^{11} M_{\odot}$ , TNG100 galaxies accrete the majority of their *ex situ* stellar population through very minor mergers and by stripping stars from surviving objects. Indeed, for these systems  $(f_{\text{ex-situ, minor merger}} + f_{\text{ex-situ, major merger}}) / f_{\text{ex-situ}} \approx 39$  per cent. Instead, above  $10^{11} M_{\odot}$ , more than 60 per cent of the *ex situ* component comes from minor and major mergers, with a larger contribution from major mergers. Considering the distributions of the number of minor and major mergers in TNG100 ETGs over the redshift range  $0.1 \leq z \leq 2$ , below  $10^{11} M_{\odot}$ , ETGs usually undergo at most one minor and/or major mergers, while above  $10^{11} M_{\odot}$  the distributions are slightly wider, confirming the important role of major mergers in shaping massive galaxies (bottom-left panel of Fig. 5). Finally, TNG100 ETGs experience, on average, their last major mergers slightly more recently than their last minor mergers (bottom-right panel of Fig. 5). However, we stress here that by minor mergers we refer to systems with  $1/10 < \mu_* < 1/4$ , which excludes the very minor mergers (i.e.  $\mu_* < 1/10$ ). In addition, we draw attention to the fact that, at a given stellar mass, both the distributions and their median values of the last minor and major mergers are similar, implying that the differences are fairly small. The difference between the median values for the same subsample are lower than 1 Gyr.

## 5.2 Comparison with recent works

In Section 4, we discussed the comparison of the circularized radial distributions of the stellar mass surface density, metallicity, age, and velocity dispersion between in MaNGA and TNG100 ETGs. In this section, we compare our results with other works from the literature.

Using a sample of 366 ETGs with masses in the range  $9.9 < \log(M_*/M_{\odot}) < 10.8$  selected via the the Galaxy Zoo morphological classification (Lintott et al. 2011; Willett et al. 2013), plus visual inspection, Parikh et al. (2018, 2019) analysed the radial gradients of stellar age and metallicity out to one effective radius. If we consider in our sample only galaxies with stellar mass lower than  $10^{10.8} M_{\odot}$  and we rescale our profiles in units of  $R_e$  (the median  $R_e$  for our MaNGA ETGs with stellar mass lower than  $10^{10.8}$  is  $\lesssim 3$  kpc) to directly compare the results with those from Parikh et al. (2018, 2019), we find a satisfying consistency with their stellar age and metallicity radial distributions.

Bernardi et al. (2019) show stellar population gradients for a sample of MaNGA DR15 ETGs subdivided into slow and fast rotators. These ETGs are identified as in Domínguez Sánchez et al. (2020), i.e. applying a morphological classification based on T-Type  $\leq 0$  (see also Section 2.4), considering both pure ellipticals and lenticulars. The stellar age and metallicity gradients measured by Bernardi et al. (2019) out to  $1 R_e$  are qualitatively compatible with our estimates. One of the most relevant outcomes of Bernardi et al. (2019) is that slow rotators dominate above  $\log(M_*/M_{\odot}) \approx 11.5$ : at this stellar mass, where also the size–mass relation slope changes (see Bernardi et al. 2011), the majority of these ellipticals are central galaxies.<sup>18</sup> As an extension of the Bernardi et al. (2019) work, Domínguez Sánchez et al. (2020) focus on stellar properties of S0 lenticular galaxies, highlighting a bimodality in this galaxy population that depends on stellar mass. Above  $\log(M_*/M_{\odot}) \approx 10.5$ , indeed, these galaxies are characterized by stronger age and velocity dispersion gradients, with, instead, negligible gradients in metallicity.

Recently, Barrera-Ballesteros et al. (2022) analysed the entire set of around 10 000 galaxies from the MaNGA survey, presenting the

radial distributions of several physical properties, selecting in particular a subset of about 1400 sources with optimal spatial coverage, for which the authors studied the impact of a selection based on either stellar mass or morphology. Among the properties derived through the the newest PYPIPE3D pipeline (Sánchez et al. 2022), the authors measured the radial distributions of stellar mass surface density, luminosity-weighted stellar metallicity and age, and velocity dispersion. The negative gradients found from Barrera-Ballesteros et al. (2022) for the stellar mass surface density and metallicity in elliptical and lenticular galaxies are qualitatively in agreement with the measurements obtained in this manuscript, reflecting also the increasing of the normalizations of both profiles as the stellar mass goes up. Even the velocity dispersion profiles agree with those computed by pPXF, finding decreasing distributions towards the outer regions, and central measurements of  $\sim 150 \text{ km s}^{-1}$  for E/S0 galaxies with  $10.5 < \log(M_*/M_{\odot}) < 11$ , and of  $\sim 250 \text{ km s}^{-1}$  for objects with  $\log(M_*/M_{\odot}) > 11$ , similarly to those presented here. Regarding stellar ages, the profiles in Barrera-Ballesteros et al. (2022) show values slightly more consistent with those derived by PROSPECTOR (i.e.  $\gtrsim 9$  Gyr) presented in this paper, though their radial distributions tend to reduce at large distances, implying the presence of younger stellar populations in the outer regions of galaxies. Also Oyarzún et al. (2022) present the radial profiles for stellar mass surface density from PROSPECTOR, as well as for element abundances (i.e. [Fe/H] and [Mg/Fe]) and ages from ALF (Conroy & van Dokkum 2012; Conroy et al. 2018; see also Choi et al. 2014; Conroy, Graves & van Dokkum 2014) for a subset of about 2200 passive centrals from MaNGA to understand the impact of stellar and halo masses in assembling these systems. The radial distributions of stellar mass surface density and [Fe/H] (that we can consider as a measurement of the stellar metallicity) are consistent with those shown in Fig. 2. The ages provided by ALF, on average, lie in between our measurements from FIREFLY and PROSPECTOR, showing also the presence of younger stellar populations in the inner regions.

Using the results from the Illustris simulation, Cook et al. (2016) investigated the stellar population gradients for a sample of more than 500 ETGs with  $10 \leq \log(M_*/M_{\odot}) \leq 12$ . The stellar surface brightness, metallicity, and age gradients are overall in agreement with observables. The gradients are subdivided into three intervals: the inner galaxy ( $0.1 - 1 R_e$ ), the outer galaxy ( $1 - 2 R_e$ ), and the stellar halo ( $2 - 4 R_e$ ). Except for the age gradients which are found to be not so informative about the accretion histories of galaxies, both the surface brightness and metallicity profiles show that, at fixed stellar mass, the *ex situ* stellar component produce flatter profiles. In particular, as the stellar mass increases, the higher the accreted star fraction, the flatter the profiles. Though the flattening at large radii of the stellar mass surface density and metallicity profiles is not apparent in Fig. 2, because of the adopted logarithmic scale, we verified that our profiles are quantitatively consistent with those of Cook et al. (2016) when a linear scale in units of  $R_e$  is adopted. Albeit a qualitative agreement in the behaviour of radial profiles is found using Illustris and IllustrisTNG, the fraction of accreted stars as well as the role of mergers in shaping galaxies are significantly different. As highlighted in fig. 10 of Tacchella et al. (2019), though in both simulations the fraction of *ex situ* stars at  $z = 0$  rapidly increases above  $10^{10.5} M_{\odot}$ , TNG100 predicts a fraction that, on average, is higher by a factor of  $\approx 30$  per cent with respect to the total stellar amount. Moreover, while the main channel for the stellar accretion in Illustris is via minor mergers, as shown in the top-right panel of Fig. 5, TNG100 predicts a more relevant role of major mergers. These two substantial differences between Illustris and IllustrisTNG are primarily due to the different feedback model implemented and

<sup>18</sup>Bernardi et al. (2019) use the Yang et al. (2007) environmental catalogue used also in this work.

the consequent stellar mass functions. In a more recent work, Pulsoni et al. (2020) studied the photometric and kinematic properties out to  $15 R_e$  of ETGs stellar haloes for 1114 objects in TNG100 (together with other 80 sources in TNG50), with stellar masses  $10.3 < \log(M_*/M_\odot) < 12$  and selected in  $(g-r)$  colours (similarly with the selection adopted in this paper) and in the angular momentum–ellipticity plane. Analogously to our findings, the high-mass tail of ETGs are everywhere dominated by the accreted stellar component, mainly acquired through major mergers. In addition, IllustrisTNG ETGs are compared with some observational surveys, including MaNGA galaxies. Looking at the distribution of galaxies in the angular momentum–ellipticity plane within  $1 R_e$ , a percentage of the IllustrisTNG galaxies lie in a region where no observed ETGs are found: these are basically elongated, triaxial systems. However, when simulated galaxies with an intermediate-to-major axial ratio  $< 0.6$  at  $1 R_e$  are removed – the centrally elongated objects –, these ETGs reflect the location in the plane of the observed counterpart, except for a region where a large fraction of MaNGA S0 have a high angular momentum.

Overall, the results presented in this manuscript are generally in agreement with previous works in the literature. In particular, the distributions of the stellar properties for the observed galaxies analysed in this work confirm the common trend of negative gradients for stellar mass surface density, metallicity, and velocity dispersion, whose normalizations increase as the stellar mass increases. The radial distributions of stellar age, however, are not always in agreement with those presented in previous works. The differences, whether in normalization or in shape, may depend on multiple factors, such as the considered samples or the stellar fitting codes and libraries used to estimate the age of stellar populations. The scope of this work is to outline a possible scenario for the merger-driven evolution of observed ETGs in the present-day Universe. The scenario predicted by IllustrisTNG in which major mergers may be crucial in shaping massive galaxies at  $z \approx 0$  is somewhat in contrast with some previous theoretical findings (e.g. Naab et al. 2009; Oser et al. 2010; Hilz et al. 2013) which, conversely, back an evolution mainly driven by a high number of minor mergers.

## 6 SUMMARY AND CONCLUSIONS

In this paper, we studied the radial profiles of stellar mass surface density, metallicity, age, and line-of-sight velocity dispersion in massive ( $M_* \geq 10^{10.5} M_\odot$ ) ETGs, selected in colours with  $(g-r) > 0.6$ , comparing observed galaxies from the MaNGA DR15 survey with simulated galaxies from TNG100 of the IllustrisTNG magnetohydrodynamical cosmological simulation suite. For both galaxy samples, the stellar property profiles have been obtained building concentric elliptical annuli on the bi-dimensional projected maps of each source, disentangling in TNG100 ETGs the *in situ* and *ex situ* stellar populations, and both accounting and not accounting for the effects of convolving the maps by MaNGA PSF. All the presented stellar population properties are *at face value*, i.e. the measurements are directly obtained from pipelines and stellar fitting codes for MaNGA, and from the simulation for TNG100 ETGs.

Our main results are the following.

(i) We find a satisfying agreement between observations and simulations in the stacked radial profiles of the stellar mass surface density of massive ETGs, both in shape and normalization. This agreement is observed at all radii and at all stellar mass bins, and is independent of ETG and stellar mass definitions.

(ii) Overall, TNG100 ETGs have, on average, stellar metallicity and velocity dispersion profiles reasonably similar to those observed in MaNGA ETGs. Concerning metallicity, the shape of MaNGA profiles is well reproduced by TNG100 galaxies, though in some cases the observed and simulated profiles differ in normalization (around 0.15 dex in the outermost parts of ETGs with  $M_* \gtrsim 10^{11.5} M_\odot$ ). For galaxies  $M_* < 10^{11.5} M_\odot$ , we find a reasonable agreement for the radial distributions of velocity dispersion between simulated and observed ETGs, the latter showing steeper profiles that differ at most by  $\approx 30 \text{ km s}^{-1}$  from the simulated ETG distributions. Only the very massive ( $M_* \gtrsim 10^{11.5} M_\odot$ ) systems of TNG100 tend to have, over the entire explored radial range, higher velocity dispersion than the corresponding observed system, by up to  $\approx 50 \text{ km s}^{-1}$ .

(iii) The ages of the stellar populations of observed ETGs are highly uncertain, and significantly different age estimates are obtained using different codes (FIREFLY and PROSPECTOR, which differ of about 2–2.5 Gyr at all radii and at all stellar mass bins). The age profiles of stellar populations in TNG100 are found to lie in between the profiles estimated for the corresponding observed ETGs with FIREFLY and PROSPECTOR.

(iv) By separating central and satellite galaxies for both TNG100 and MaNGA samples, we find that there are not relevant differences in all the profiles between the two galaxy populations, except for the velocity dispersion profiles of massive systems ( $M_* > 10^{11.5} M_\odot$ , see Fig. 4). Indeed, while TNG100 and MaNGA satellites have similar velocity dispersion profiles, central simulated galaxies tend to have velocity dispersion at all radii higher than observed ETGs ( $\approx 50 \text{ km s}^{-1}$ ).

(v) The behaviour of the *in situ* and *ex situ* surface density profiles identifies two different scenarios for the merger-driven history of these objects, corroborating previous outcomes in the literature, such as from Pillepich et al. (2018b) and Tacchella et al. (2019): galaxies with  $M_* \lesssim 10^{11} M_\odot$  are mainly dominated by the *in situ* stellar populations out to  $\approx 30 \text{ kpc}$ ; instead, in ETGs with  $M_* \gtrsim 10^{11} M_\odot$ , the contribution of the *ex situ* stars is at least as important as that of the *in situ* component, and even totally dominating for very massive ETGs ( $M_* \gtrsim 10^{11.5} M_\odot$ ).

(vi) The similar shapes found for the radial distributions of the stellar mass surface densities for both *in situ* and *ex situ* stars (see Fig. 3) as well as the detailed analysis of the merger history (see Fig. 5) in simulated ETGs reveal that especially galaxies with  $M_* \gtrsim 10^{11} M_\odot$  experienced across cosmic time an evolution mainly driven by major mergers. Indeed, major mergers allow both to explain the presence of a significant percentage of *ex situ* stars that are able to penetrate even in the innermost parts of galaxies, and also that the two stellar components are well homogenized at all radii, showing similar surface density profiles. The results from TNG100 illustrated in this paper and in previous works in literature (e.g. Pillepich et al. 2018b; Tacchella et al. 2019; Pulsoni et al. 2021) support the possible scenario in which massive systems assembled across their cosmic histories mainly via major mergers (see also Sonnenfeld et al. 2017), in contrast with some previous theoretical (e.g. Naab et al. 2009; Oser et al. 2010; Hilz et al. 2013) and observational (e.g. Bell et al. 2006; McIntosh et al. 2008) studies which, instead, endorse a minor-merger-driven evolution for ETGs.

For the future, we plan to extend the analysis to other physical properties, like chemical abundances of individual elements. In order to provide a more complete scenario behind the cosmic evolution of the ETGs that we observe in the present-day Universe, in simulations we will study the merger history of individual galaxies, considering the evolution of the spatial distribution of their stellar properties.



Finally, we will use the newest TNG50 simulation of IllustrisTNG which, though characterized by a smaller physical volume and thus a lower statistics, benefits from a higher mass resolution that could allow us to make a more reliable comparison at smaller scales of galaxies with data from current and upcoming surveys.

## ACKNOWLEDGEMENTS

We are grateful to M. Auger and M. Bernardi for helpful pieces of advice and comments, which remarkably enriched the analysis presented in this work. We sincerely thank the referee A. Pillepich for her feedback and suggestions that considerably contributed to improve the quality of this manuscript. CC acknowledges useful discussions with F. Ardila, C. Bacchini, F. Belfiore, C. D'Eugenio, G. Iorio, M. Mingozi, R. Pascale, S. Quai, and S. Tacchella. This material is based upon work supported by the National Science Foundation under Grant No. 1714610.

## DATA AVAILABILITY

The data underlying this article regarding the selection of galaxy samples and the radial profiles will be shared upon request to the corresponding author. IllustrisTNG data are publicly available at <https://www.illustris-project.org/data/> (Nelson et al. 2019a). The MaNGA galaxy sample is drawn from an extended version of the NASA-Sloan Atlas (NSA v1.0.1, <https://www.sdss.org/dr15/manga/manga-target-selection/nsa/>; Blanton et al. 2011) catalogue. MaNGA DR15 data are taken from <https://www.sdss.org/dr15/manga/manga-data/> (Aguado et al. 2019). The stellar masses for MaNGA galaxies are included in the UPENN\_PHOTDEC\_MSSTAR catalogue (<http://alan-meert-website-aws.s3-website-us-east-1.amazonaws.com/fit-catalog/download/index.html>; Meert et al. 2015). To check the impact of the adopted selection for the MaNGA sample, we relied on the morphological type presented in the MANGA MORPHOLOGY DEEP LEARNING DR15 CATALOG ([https://www.sdss.org/dr15/data\\_access/value-added-catalogs/?vac\\_id=manga-morphology-deep-learning-dr15-catalogue](https://www.sdss.org/dr15/data_access/value-added-catalogs/?vac_id=manga-morphology-deep-learning-dr15-catalogue); Fischer et al. 2019).

## REFERENCES

Adelman-McCarthy J. K. et al., 2006, *ApJS*, 162, 38  
 Aguado D. S. et al., 2019, *ApJS*, 240, 23  
 Aihara H. et al., 2018, *PASJ*, 70, 58  
 Alam S. et al., 2015, *ApJS*, 219, 12  
 Ardila F. et al., 2021, *MNRAS*, 500, 432  
 Bacon R. et al., 2001, *MNRAS*, 326, 23  
 Barber C., Schaye J., Crain R. A., 2019, *MNRAS*, 483, 985  
 Barrera-Ballesteros J. K. et al., 2022, preprint (arXiv:2206.07058)  
 Bell E. F. et al., 2006, *ApJ*, 640, 241  
 Belli S., Newman A. B., Ellis R. S., 2014, *ApJ*, 783, 117  
 Belli S., Newman A. B., Ellis R. S., 2017, *ApJ*, 834, 18  
 Bernardi M., Roche N., Shankar F., Sheth R. K., 2011, *MNRAS*, 412, L6  
 Bernardi M., Meert A., Sheth R. K., Fischer J. L., Huertas-Company M., Maraston C., Shankar F., Vikram V., 2017, *MNRAS*, 467, 2217  
 Bernardi M., Domínguez Sánchez H., Brownstein J. R., Drory N., Sheth R. K., 2019, *MNRAS*, 489, 5633  
 Blanton M. R., Kazin E., Muna D., Weaver B. A., Price-Whelan A., 2011, *AJ*, 142, 31  
 Blanton M. R. et al., 2017, *AJ*, 154, 28  
 Bottrell C., Hani M. H., 2022, *MNRAS*, 514, 2821  
 Bryant J. J. et al., 2015, *MNRAS*, 447, 2857  
 Bundy K., Fukugita M., Ellis R. S., Targett T. A., Belli S., Kodama T., 2009, *ApJ*, 697, 1369  
 Bundy K. et al., 2015, *ApJ*, 798, 7

Calette A. R., Avila-Reese V., Rodríguez-Puebla A., Hernández-Toledo H., Papastergis E., 2018, *Rev. Mex. Astron. Astrofis.*, 54, 443  
 Cannarozzo C., 2021, PhD thesis, Alma Mater Studiorum Università di Bologna,  
 Cannarozzo C., Sonnenfeld A., Nipoti C., 2020, *MNRAS*, 498, 1101  
 Cappellari M., 2017, *MNRAS*, 466, 798  
 Cappellari M., Emsellem E., 2004, *PASP*, 116, 138  
 Cappellari M. et al., 2011, *MNRAS*, 413, 813  
 Chabrier G., 2003, *PASP*, 115, 763  
 Choi J., Conroy C., Moustakas J., Graves G. J., Holden B. P., Brodwin M., Brown M. J. I., van Dokkum P. G., 2014, *ApJ*, 792, 95  
 Choi J., Dotter A., Conroy C., Cantiello M., Paxton B., Johnson B. D., 2016, *ApJ*, 823, 102  
 Cimatti A., Fraternali F., Nipoti C., 2019, Introduction to Galaxy Formation and Evolution: From Primordial Gas to Present-Day Galaxies. Cambridge Univ. Press, Cambridge, UK  
 Comparat J. et al., 2017, preprint (arXiv:1711.06575)  
 Conroy C., 2013, *ARA&A*, 51, 393  
 Conroy C., Gunn J. E., 2010, *ApJ*, 712, 833  
 Conroy C., van Dokkum P., 2012, *ApJ*, 747, 69  
 Conroy C., Gunn J. E., White M., 2009, *ApJ*, 699, 486  
 Conroy C., Graves G. J., van Dokkum P. G., 2014, *ApJ*, 780, 33  
 Conroy C., Villaume A., van Dokkum P. G., Lind K., 2018, *ApJ*, 854, 139  
 Cook B. A., Conroy C., Pillepich A., Rodriguez-Gomez V., Hernquist L., 2016, *ApJ*, 833, 158  
 Crain R. A. et al., 2015, *MNRAS*, 450, 1937  
 Croom S. M. et al., 2012, *MNRAS*, 421, 872  
 Damjanov I., Zahid H. J., Geller M. J., Utsuni Y., Sohn J., Souchereau H., 2019, *ApJ*, 872, 91  
 Davison T. A., Norris M. A., Leaman R., Kuntschner H., Boecker A., van de Ven G., 2021, *MNRAS*, 507, 3089  
 de Vaucouleurs G., 1959, *Handbuch der Physik*, 53, 275  
 de Zeeuw P. T. et al., 2002, *MNRAS*, 329, 513  
 Diemer B. et al., 2018, *ApJS*, 238, 33  
 Diemer B. et al., 2019, *MNRAS*, 487, 1529  
 Djorgovski S., Davis M., 1987, *ApJ*, 313, 59  
 Domínguez Sánchez H., Bernardi M., Nikakhtar F., Margalef-Bentabol B., Sheth R. K., 2020, *MNRAS*, 495, 2894  
 Donnari M. et al., 2019, *MNRAS*, 485, 4817  
 Dotter A., 2016, *ApJS*, 222, 8  
 Dressler A., Lynden-Bell D., Burstein D., Davies R. L., Faber S. M., Terlevich R., Wegner G., 1987, *ApJ*, 313, 42  
 Drory N. et al., 2015, *AJ*, 149, 77  
 Edwards L. O. V. et al., 2020, *MNRAS*, 491, 2617  
 Faber S. M., Jackson R. E., 1976, *ApJ*, 204, 668  
 Ferguson H. C. et al., 2004, *ApJ*, 600, L107  
 Fischer J. L., Domínguez Sánchez H., Bernardi M., 2019, *MNRAS*, 483, 2057  
 Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, *PASP*, 125, 306  
 Gallazzi A., Charlot S., Brinchmann J., White S. D. M., Tremonti C. A., 2005, *MNRAS*, 362, 41  
 Genel S. et al., 2008, *ApJ*, 688, 789  
 Genel S. et al., 2014, *MNRAS*, 445, 175  
 Goddard D. et al., 2017a, *MNRAS*, 465, 688  
 Goddard D. et al., 2017b, *MNRAS*, 466, 4731  
 Goodman J., Weare J., 2010, *Commun. Appl. Math. Comput. Sci.*, 5, 65  
 Greene J. E., Janish R., Ma C.-P., McConnell N. J., Blakeslee J. P., Thomas J., Murphy J. D., 2015, *ApJ*, 807, 11  
 Greene J. E. et al., 2019, *ApJ*, 874, 66  
 Gunn J. E. et al., 2006, *AJ*, 131, 2332  
 Hilz M., Naab T., Ostriker J. P., 2013, *MNRAS*, 429, 2924  
 Hirschmann M. et al., 2013, *MNRAS*, 436, 2929  
 Hirschmann M., Naab T., Ostriker J. P., Forbes D. A., Duc P.-A., Davé R., Oser L., Karabal E., 2015, *MNRAS*, 449, 528  
 Huang S., Leauthaud A., Greene J. E., Bundy K., Lin Y.-T., Tanaka M., Miyazaki S., Komiyama Y., 2018, *MNRAS*, 475, 3348

- Johnson B. D., Leja J. L., Conroy C., Speagle J. S., 2019, *Astrophysics Source Code Library*, record ascl:1905.025
- Khochfar S., Silk J., 2009, *MNRAS*, 397, 506
- Kirby E. N., Cohen J. G., Guhathakurta P., Cheng L., Bullock J. S., Gallazzi A., 2013, *ApJ*, 779, 102
- Kobayashi C., 2004, *MNRAS*, 347, 740
- Kormendy J., 1977, *ApJ*, 218, 333
- Kroupa P., 2001, *MNRAS*, 322, 231
- Kuntschner H. et al., 2010, *MNRAS*, 408, 97
- Lackner C. N., Cen R., Ostriker J. P., Joung M. R., 2012, *MNRAS*, 425, 641
- Larson R. B., 1974, *MNRAS*, 166, 585
- Law D. R. et al., 2015, *AJ*, 150, 19
- Law D. R. et al., 2016, *AJ*, 152, 83
- Law D. R. et al., 2021, *AJ*, 161, 52
- Leja J., Johnson B. D., Conroy C., van Dokkum P. G., Byler N., 2017, *ApJ*, 837, 170
- Leja J., Carnall A. C., Johnson B. D., Conroy C., Speagle J. S., 2019, *ApJ*, 876, 3
- Li H. et al., 2018, *MNRAS*, 476, 1765
- Lintott C. et al., 2011, *MNRAS*, 410, 166
- Lovell M. R. et al., 2018, *MNRAS*, 481, 1950
- Ma C.-P., Greene J. E., McConnell N., Janish R., Blakeslee J. P., Thomas J., Murphy J. D., 2014, *ApJ*, 795, 158
- McAlpine S. et al., 2016, *Astron. Comput.*, 15, 72
- McIntosh D. H., Guo Y., Hertzberg J., Katz N., Mo H. J., van den Bosch F. C., Yang X., 2008, *MNRAS*, 388, 1537
- Maraston C., Strömbäck G., 2011, *MNRAS*, 418, 2785
- Maraston C. et al., 2020, *MNRAS*, 496, 2962
- Marinacci F. et al., 2018, *MNRAS*, 480, 5113
- Meert A., Vikram V., Bernardi M., 2015, *MNRAS*, 446, 3943
- Mendel J. T., Simard L., Palmer M., Ellison S. L., Patton D. R., 2014, *ApJS*, 210, 3
- Miyazaki S. et al., 2018, *PASJ*, 70, S1
- Moresco M. et al., 2013, *A&A*, 558, A61
- Naab T., Johansson P. H., Ostriker J. P., 2009, *ApJ*, 699, L178
- Naiman J. P. et al., 2018, *MNRAS*, 477, 1206
- Nanni L. et al., 2022, *MNRAS*, 515, 320
- Nelson D. et al., 2018, *MNRAS*, 475, 624
- Nelson D. et al., 2019a, *Comput. Astrophys. Cosmol.*, 6, 2
- Nelson D. et al., 2019b, *MNRAS*, 490, 3234
- Nipoti C., Cannarozzo C., Calura F., Sonnenfeld A., Treu T., 2020, *MNRAS*, 499, 559
- Nipoti C., Cherchi G., Iorio G., Calura F., 2021, *MNRAS*, 503, 4221
- Oser L., Ostriker J. P., Naab T., Johansson P. H., Burkert A., 2010, *ApJ*, 725, 2312
- Oser L., Naab T., Ostriker J. P., Johansson P. H., 2012, *ApJ*, 744, 63
- Oyarzún G. A. et al., 2019, *ApJ*, 880, 111
- Oyarzún G. A. et al., 2022, *ApJ*, 933, 88
- Parikh T. et al., 2018, *MNRAS*, 477, 3954
- Parikh T. et al., 2019, *MNRAS*, 483, 3420
- Pillepich A. et al., 2014, *MNRAS*, 444, 237
- Pillepich A. et al., 2018a, *MNRAS*, 473, 4077
- Pillepich A. et al., 2018b, *MNRAS*, 475, 648
- Pillepich A. et al., 2019, *MNRAS*, 490, 3196
- Pipino A., D’Ercole A., Chiappini C., Matteucci F., 2010, *MNRAS*, 407, 1347
- Planck Collaboration XIII, 2016, *A&A*, 594, A13
- Pulsoni C., Gerhard O., Arnaboldi M., Pillepich A., Nelson D., Hernquist L., Springel V., 2020, *A&A*, 641, A60
- Pulsoni C., Gerhard O., Arnaboldi M., Pillepich A., Rodriguez-Gomez V., Nelson D., Hernquist L., Springel V., 2021, *A&A*, 647, A95
- Rodriguez-Gomez V. et al., 2015, *MNRAS*, 449, 49
- Rodriguez-Gomez V. et al., 2016, *MNRAS*, 458, 2371
- Rodriguez-Gomez V. et al., 2019, *MNRAS*, 483, 4140
- Sánchez-Blázquez P. et al., 2006, *MNRAS*, 371, 703
- Sánchez S. F. et al., 2016, *A&A*, 594, A36
- Sánchez S. F. et al., 2022, *ApJS*, 262, 36
- Santucci G. et al., 2020, *ApJ*, 896, 75
- Schaye J. et al., 2015, *MNRAS*, 446, 521
- Sijacki D., Vogelsberger M., Genel S., Springel V., Torrey P., Snyder G. F., Nelson D., Hernquist L., 2015, *MNRAS*, 452, 575
- Smee S. A. et al., 2013, *AJ*, 146, 32
- Sonnenfeld A., Nipoti C., Treu T., 2017, *MNRAS*, 465, 2397
- Sonnenfeld A., Wang W., Bahcall N., 2019, *A&A*, 622, A30
- Speagle J. S., 2020, *MNRAS*, 493, 3132
- Springel V., 2010, *MNRAS*, 401, 791
- Springel V., White S. D. M., Tormen G., Kauffmann G., 2001, *MNRAS*, 328, 726
- Springel V. et al., 2018, *MNRAS*, 475, 676
- Tacchella S. et al., 2019, *MNRAS*, 487, 5416
- Tanaka M. et al., 2019, *ApJ*, 885, L34
- Taylor P., Kobayashi C., 2017, *MNRAS*, 471, 3856
- Thomas D., Maraston C., Bender R., Mendes de Oliveira C., 2005, *ApJ*, 621, 673
- Van de Sande J. et al., 2013, *ApJ*, 771, 85
- Van der Wel A. et al., 2014, *ApJ*, 788, 28
- Vazdekis A., Sánchez-Blázquez P., Falcón-Barroso J., Cenarro A. J., Beasley M. A., Cardiel N., Gorgas J., Peletier R. F., 2010, *MNRAS*, 404, 1639
- Vogelsberger M. et al., 2014a, *MNRAS*, 444, 1518
- Vogelsberger M. et al., 2014b, *Nature*, 509, 177
- Wake D. A. et al., 2017, *AJ*, 154, 86
- Weinberger R. et al., 2017, *MNRAS*, 465, 3291
- Westfall K. B. et al., 2019, *AJ*, 158, 231
- Wilkinson D. M., Maraston C., Goddard D., Thomas D., Parikh T., 2017, *MNRAS*, 472, 4297
- Willett K. W. et al., 2013, *MNRAS*, 435, 2835
- Woo J., Courteau S., Dekel A., 2008, *MNRAS*, 390, 1453
- Worthey G., 1994, *ApJS*, 95, 107
- Yan R. et al., 2016a, *AJ*, 151, 8
- Yan R. et al., 2016b, *AJ*, 152, 197
- Yang X., Mo H. J., van den Bosch F. C., Pasquali A., Li C., Barden M., 2007, *ApJ*, 671, 153
- York D. G. et al., 2000, *AJ*, 120, 1579
- Zheng Z. et al., 2017, *MNRAS*, 465, 4572


## APPENDIX A: COMPUTING THE ANGULAR DIAMETER DISTANCES TO CONVOLVE TNG100 MAPS

As described in Section 3.2, in order to account for the effects of the MaNGA resolution on simulated galaxies, for each stellar property we consider two limit-case profiles: for a given simulated ETG, one profile is computed directly from the original 2D stellar property map, i.e. the *unconvolved profile*, while the other profile is derived from a map previously convolved with a 2D Gaussian filter kernel  $\sigma_{\text{kernel}}$ , i.e. the *convolved profile*. To compute the kernel of each simulated ETG, we use equation (1), where  $\mathcal{R}_{\text{TNG}} = 1$  kpc is the TNG100 resolution of the original maps, while  $\mathcal{R}_{\text{MaNGA},i} = \sin(\text{PSF}_{\text{MaNGA}})d_{A,i}$  is the resolution that the  $i$ -th TNG100 ETG map should have if it were observed as a MaNGA galaxy, and depends on  $\text{PSF}_{\text{MaNGA}} = 2.5$  arcsec and the angular diameter distance  $d_{A,i}$  of the  $i$ -th TNG100 galaxy. To measure  $d_{A,i}$  we rely on the `angular_diameter_distance` function of the Python package `ASTROPY`, which takes in input the redshift of the  $i$ -th source.

Since MaNGA was built in such a way that the most massive galaxies are located at higher redshifts, we fit the  $z-M_*$  distribution of the MaNGA sample considered and we assign to each simulated galaxy the corresponding redshift value depending on its stellar mass. The functional form adopted for fitting the  $z-M_*$  distributions in MaNGA is

$$z = a e^{b \log(M_*/M_\odot)}, \quad (\text{A1})$$

where  $a$  and  $b$  are the two parameters used for the fit. In Fig. A1, the  $z-M_*$  scatter distribution with the corresponding fit are shown. Thus,



**Figure A1.** The redshift–stellar mass distribution of the MaNGA ETG sample. The black solid curve traces the fit of the distribution. The corresponding fit functional form is reported. The two stripes of dots trace the Primary sample (the lower cloud) and the Secondary sample (the upper cloud) of the MaNGA sample.


the fit is used to assign redshifts to the simulated ETGs to compute their angular diameter distance, and then the kernel used to convolve their stellar property maps. We remind the reader that simulated galaxies are extracted from the  $z = 0.1$  snapshot of TNG100-1. However, to make the comparison with MaNGA as fair as possible, for our scope, we ignore this information for convolving the maps and we reassign to each simulated ETG a new redshift corresponding to its stellar mass according to the fit of the MaNGA distribution on the  $z-M_*$  plane.

## APPENDIX B: COMPARING PROFILES IN NUMBER-DENSITY-BASED STELLAR MASS BINS

In Section 4, we presented the radial profiles of MaNGA and TNG100 ETG stellar properties in the three stellar mass bins  $10.5 \leq \log(M_*/M_\odot) < 11$ ,  $11 \leq \log(M_*/M_\odot) < 11.5$ , and  $\log(M_*/M_\odot) \geq 11.5$ . We repeated here the same analysis building stellar mass bins at *fixed number density*. Specifically, we compute the stellar mass function (SMF) for our TNG100 galaxy sample, whereas for MaNGA we use table 1 of Bernardi et al. (2017) and adopt the observed (i.e. error-broadened) SMF associated with the Dusty ( $\Phi_{\text{Obs}}^{\text{M14d}}$ ) mass estimates from Mendel et al. (2014) with the SerExp photometry of Meert et al. (2015). These are the same stellar masses as used for our MaNGA ETGs. Thus, we compute the cumulative stellar mass functions (CSMFs) for both MaNGA and TNG100 as the sum of the number counts of galaxies with stellar masses greater than a given value  $M_{*,i}$ :

$$n(> M_{*,i}) = \int_{M_{*,i}}^{+\infty} \Phi(M'_*) dM'_* \quad (\text{B1})$$

Fig. B1 shows the CSMFs for TNG100 and MaNGA samples. We adopt three bins in number density as listed in Table B1. Number




**Figure B1.** CSMFs for the TNG100 (grey solid curve) and the MaNGA (red dashed curve) samples. The horizontal dotted lines indicate number-density-based bins:  $\log(n/\text{Mpc}^{-3}) = -2.75$  (blue),  $\log(n/\text{Mpc}^{-3}) = -3.50$  (green),  $\log(n/\text{Mpc}^{-3}) = -4.25$  (yellow). The vertical lines indicate the corresponding stellar mass values for TNG100 (solid lines) and for MaNGA (dashed lines).

**Table B1.** Stellar mass values corresponding to the bounds of the number density bins. Column 1: number density. Column 2: stellar mass for the MaNGA sample. Column 3: stellar mass for the TNG100 sample. Stellar masses are in units of  $M_\odot$ .

$\log(n/\text{Mpc}^{-3})$	$\log M_*, \text{MaNGA}$	$\log M_*, \text{TNG}$
-2.75	10.72	10.50
-3.50	11.20	11.02
-4.25	11.52	11.50

density bins could be more robust against possible mismatches in the stellar mass measurements between observations and simulations. Table B1 indicates the values of the stellar masses in both TNG100 and MaNGA samples that correspond to our three number density bins:  $-3.50 < \log(n/\text{Mpc}^{-3}) \leq -2.75$ ,  $-4.25 < \log(n/\text{Mpc}^{-3}) \leq -3.50$ , and  $\log(n/\text{Mpc}^{-3}) \leq -4.25$ . The bounds of the number density bins, listed in Table B1, are such that the corresponding stellar mass values for simulated ETGs, i.e.  $\log(M_*/M_\odot) = 10.5$ , 11.02, 11.5, almost coincide with the values of stellar masses used in Section 4.

As illustrated in Fig. B2, the most evident exceptions of adopting the number-density-based stellar mass bins concern the first two bins of the stellar properties analysed. In particular, the discrepancy between the radial profiles of the stellar mass surface density for the observed and simulated ETGs increases, highlighting a tendency of the MaNGA sources to assume slightly higher values, with respect to those shown in Fig. 2. A similar behaviour, but less significant, is also found for the stellar metallicity and age distributions. Instead, concerning velocity dispersion, in the first two bins the MaNGA profiles are systematically shifted up by a factor of around 20–30  $\text{km s}^{-1}$ . All these discrepancies are caused by the fact that, by removing from the




**Figure B2.** Same as Fig. 2, but in number-density-based stellar mass bins.

MaNGA sample ETGs with  $\log(M_*/M_\odot) \leq 10.78$  in the first bin, and considering galaxies with  $\log(M_*/M_\odot) > 11.2$  in the second bin, the median profiles of the observed sources tend to assume higher values than the counterparts presented in Fig. 2.

Since we expect number density bins to more closely approximate a halo-mass-based comparison, the aforementioned differences could be an indication of a mismatch between MaNGA and

TNG100 in the stellar-to-halo mass relation. However, qualitatively speaking, the selection in number density bins does not affect remarkably the overall fashion of the profiles for all the stellar properties, showing similar radial distributions as those illustrated in Fig. 2 (the same conclusion is valid for the *in situ* and *ex situ* radial distributions shown in Fig. B3 when compared with Fig. 3).



**Figure B3.** Same as Fig. 3, but in number-density-based stellar mass bins.

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.