

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

A transfer principle: from periods to isoperiodic foliations

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

A transfer principle: from periods to isoperiodic foliations / Calsamiglia G.; Deroin B.; Francaviglia S.. - In: GEOMETRIC AND FUNCTIONAL ANALYSIS. - ISSN 1016-443X. - STAMPA. - 33:1(2023), pp. 57-169. [10.1007/s00039-023-00627-w]

Availability:

This version is available at: <https://hdl.handle.net/11585/920355> since: 2023-03-13

Published:

DOI: <http://doi.org/10.1007/s00039-023-00627-w>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Calsamiglia, G., Deroin, B. & Francaviglia, S. A transfer principle: from periods to isoperiodic foliations. *Geom. Funct. Anal.* **33**, 57–169 (2023).

The final published version is available online at: <https://doi-org.ezproxy.unibo.it/10.1007/s00039-023-00627-w>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

A TRANSFER PRINCIPLE: FROM PERIODS TO ISOPERIODIC FOLIATIONS

GABRIEL CALSAMIGLIA, BERTRAND DEROIN, AND STEFANO FRANCAVIGLIA

ABSTRACT. We classify the possible closures of leaves of the isoperiodic foliation defined on the Hodge bundle over the moduli space of genus $g \geq 2$ curves and prove that the foliation is ergodic on those sets. The results derive from the connectedness properties of the fibers of the period map defined on the Torelli cover of the moduli space. Some consequences on the topology of Hurwitz spaces of primitive branched coverings over elliptic curves are also obtained. To prove the results we develop the theory of augmented Torelli space, the branched Torelli cover of the Deligne-Mumford compactification of the moduli space of curves.

1. INTRODUCTION

1.1. Overview. Let $\Omega\mathcal{M}_g$ be the moduli space of abelian differentials on compact genus $g \geq 2$ smooth curves. The period of an element $(C, \omega) \in \Omega\mathcal{M}_g$ is the element of $H^1(C, \mathbb{C}) \simeq \text{Hom}(H_1(C, \mathbb{Z}), \mathbb{C})$ that is defined by

$$(1) \quad \text{Per}(C, \omega) : \gamma \in H_1(C, \mathbb{Z}) \mapsto \int_{\gamma} \omega \in \mathbb{C}.$$

The periods of an abelian differential do not allow to recover the abelian differential itself, even infinitesimally. Actually, it is always possible to find non trivial isoperiodic deformations of a given abelian differential, namely an immersed complex submanifold $L \subset \Omega\mathcal{M}_g$ such that the period of a form $(C, \omega) \in L$ is a locally constant function, when we use the local identifications of the $H^*(C, \mathbb{C})$'s given by the Gauss-Manin connection. For instance, if $f_t : C_t \rightarrow C$ is a continuous family of degree d branched coverings over C depending on the parameter t , the period map of the family $\omega_t = f_t^* \omega$ is locally constant.

The case $g = 2$ is instructive: every genus two curve is a double cover of \mathbb{P}^1 ramified over six distinct points, say $0, 1, \infty, x_1, x_2, x_3$. An abelian differential on such a curve can be written as the hyperelliptic integrand

$$\frac{(ax + b)dx}{\sqrt{x(x-1)(x-x_1)(x-x_2)(x-x_3)}}.$$

Picard-Fuchs theory (see [19, p.60] for details) tells us that isoperiodic deformations on $\Omega\mathcal{M}_2$ are integral curves of the following vector field

$$(2) \quad \sum_j \frac{x_j(1-x_j)}{ax_j+b} \frac{\partial}{\partial x_j} - \frac{1}{2} \frac{\partial}{\partial a} - \frac{1}{2} \left(1 + \sum_j \frac{b(x_j-1)}{ax_j+b} \right) \frac{\partial}{\partial b}.$$

2010 *Mathematics Subject Classification.* 57M50, 30F30, 53A30, 14H15, 32G13.

Key words and phrases. Isoperiodic foliation, Hodge bundle, isoperiodic abelian differentials.

Apart from the invariant closed subsets characterized by topological properties of the set of periods, there is an interesting family of closed invariant sets for the isoperiodic foliation in genus two called the Hilbert modular invariant manifolds, introduced by Calta in [18] and McMullen in [52, 53]. The set of curves of genus two whose Jacobian has real multiplication by a quadratic order of discriminant D is a Hilbert modular surface X_D , and a remarkable fact is that the subset of the Hodge bundle ΩX_D over X_D consisting of eigenforms for the complex multiplication is invariant by the isoperiodic foliation.

In any genus, the collection of all maximal isoperiodic deformations defines a holomorphic foliation \mathcal{F}_g of $\Omega^*\mathcal{M}_g$, called the isoperiodic foliation¹ (see [56] for further examples). It has dimension $2g - 3$, and it is also algebraic: its leaves are solutions of a system of algebraic equations analogous to (2) with respect to the Deligne-Mumford algebraic structure on moduli space. It admits a real analytic global first integral defined by the volume of the flat metric induced by the abelian differential. Complex multiplication on the forms induces isomorphisms between the restrictions of the foliation to the volume levels.

It turns out that the restriction of \mathcal{F}_g to the volume levels carries a transverse structure modelled on the homogeneous space

$$(3) \quad \mathrm{Sp}(2g, \mathbb{R})/U, \text{ with } U = I_2 \times \mathrm{Sp}(2g - 2, \mathbb{R})$$

with changes of coordinates acting by right multiplications by the elements of the lattice $\mathrm{Sp}(2g, \mathbb{Z})$. There are some well known examples of closed invariant sets for the given restriction and an integer $d \geq 2$: the moduli spaces $H_{g,d} \subset \Omega\mathcal{M}_g$ of forms that are pull-back of elliptic differentials by primitive branched coverings of degree d . The set $H_{g,d}$ is saturated by the isoperiodic foliation and each leaf there is algebraic.

The main goal of this paper is to investigate the topological properties of the isoperiodic foliation, and derive some dynamical consequences: we prove that any dynamical property satisfied by the action of the lattice $\mathrm{Sp}(2g, \mathbb{Z})$ on the homogeneous space (3) can be transferred to a property satisfied by the isoperiodic foliation. This fact is what we call the transfer principle. Using Ratner's theory, it allows to describe explicitly the closure of *each* leaf and prove that it is a real analytic subset. We also obtain the ergodicity of the isoperiodic foliation on these sets, a fact that has been proven independently by Hamenstädt in [29].

The proof of the transfer principle is achieved by studying the topological properties of the fibers of the period map defined on the moduli space of homologically marked genus g curves, with values in the affine space \mathbb{C}^{2g} : we prove that all the fibers that do not correspond to leaves in $H_{g,2}$ with $g \geq 4$ are connected. This result provides a solution to a problem posed by McMullen in [50, p. 2282] for genus $g \geq 4$. The main idea of the proof of the connectedness of the fibers is to show that any form can be deformed continuously and isoperiodically to a stable form on a nodal curve, and then connect all possible stable forms obtained in this way using an inductive argument on the genus. As for fibers formed by branched double coverings over an elliptic differential, we generalize an invariant described by Arnold [4] for branched double coverings over the Riemann sphere (i.e. hyperelliptic loci) to prove they are disconnected when the genus is at least five.

1.2. Statement of results. A homologically marked genus g curve is a couple (C, m) , where C is a genus g curve and $m : \mathbb{Z}^{2g} \rightarrow H_1(C, \mathbb{Z})$ is a symplectic identification. The moduli space of marked genus g curves is the so called Torelli covering $\mathcal{S}_g \rightarrow \mathcal{M}_g$. Its covering group

¹In the literature, this foliation is also called the kernel foliation, or the absolute period foliation.

$J_g \subset \pi_1(\mathcal{M}_g)$ — called the Torelli group— is identified with the set of classes in the mapping class group of an orientable closed surface of genus g that act trivially on the first homology group of the surface. We denote by $\Omega\mathcal{S}_g$ the pull-back of the Hodge bundle to \mathcal{S}_g . We analyze the topological properties of the period map

$$\text{Per}_g : \Omega\mathcal{S}_g \rightarrow \mathcal{H}_g \cup 0 \text{ defined by } \text{Per}_g(C, m, \omega) = \text{Per}(C, \omega) \circ m.$$

where $\mathcal{H}_g \subset \text{Hom}(\mathbb{Z}^{2g}, \mathbb{C})$ denotes the open subset formed by periods of non zero abelian differentials on a homologically marked genus g curve. The isoperiodic foliation lifts to a foliation of $\Omega\mathcal{S}_g$ whose leaves are the connected components of the level sets of the period map.

Definition 1.1 (Primitive degree). Given $p \in \text{Hom}(\mathbb{Z}^{2g}, \mathbb{C})$ we define

- its volume $\text{vol}(p) = \Re(p) \cdot \Im(p)$, the symplectic product on $\text{Hom}(\mathbb{Z}^{2g}, \mathbb{R})$
- its primitive degree, denoted $\deg(p)$ as ∞ if $\Lambda = p(\mathbb{Z}^{2g}) \subset \mathbb{C}$ is non-discrete and as

$$\deg(p) = \frac{\text{vol}(p)}{\text{vol}(\mathbb{C}/\Lambda)} \text{ if } \Lambda \subset \mathbb{C} \text{ is discrete.}$$

When $p \in \text{Hom}(\mathbb{Z}^{2g}, \mathbb{C})$ is the period of some non-zero abelian differential ω on a homologically marked smooth curve (C, m) , $\text{vol}(p)$ corresponds to its volume $\frac{i}{2} \int \omega \wedge \bar{\omega}$ too, and is therefore positive. When the periods of the form define a lattice $\Lambda \subset \mathbb{C}$, the number $\deg(p)$ corresponds to the topological degree of the primitive branched covering $C \rightarrow \mathbb{C}/\Lambda$ defined by integration of the form (i.e. it induces a surjection at the first homology group level). The degree is a positive integer and it is one only if $g = 1$. The conditions $\text{vol}(p) > 0$ and either $g = 1$ or $\deg(p) > 1$ will be referred to as Haupt conditions.

Our main result is

Theorem 1.2. *For $g \geq 2$, the fibers of the period map Per_g over points of primitive degree at least three are connected.*

Its proof, whose structure is detailed in section [1.3](#), will take up most of the paper.

Regarding the fibers of Per_g over points of primitive degree two, it was already known that they are connected for $g = 2, 3$ (see [\[50\]](#)). We suspect that the equivalent statement is still true in genus four, but we were not able to prove it. Nevertheless, in higher genera we prove

Theorem 1.3. *If $g \geq 5$, the fibers of Per_g over points of primitive degree two are disconnected.*

Theorem [1.3](#) is proved using an adaptation of a method due to Arnold [\[4\]](#) based on an invariant associated to branched double coverings. It enabled him to prove that the symplectic representation $\pi_1(H_{g,2}(\mathbb{P}^1)) \rightarrow \text{Sp}(2g, \mathbb{Z})$, induced by the natural inclusion $H_{g,2}(\mathbb{P}^1) \rightarrow \mathcal{M}_g$ of the Hurwitz space $H_{g,2}(\mathbb{P}^1)$ of genus g branched double coverings over the Riemann sphere, is not onto for $g \geq 3$. In our case the invariant will be extended to branched double coverings over an elliptic curve.

The connectedness of the fibers of the lift of Per_g to the universal cover of $\Omega\mathcal{M}_g$ fails in general. For example, in genus $g = 3$ the fibers of Per_3 over points of primitive degree two are biholomorphic to Siegel space \mathfrak{S}_2 , hence simply connected. Therefore there are infinitely many components of the lift of such a fiber to the universal cover of $\Omega\mathcal{M}_3$. In genus $g = 2$, the generating family of $\pi_1(\mathcal{M}_2)$ given by Mess in [\[54\]](#) allows to prove that the lift of the generic fiber is disconnected (see Corollary [6.20](#)). In both cases the proof relies on the fact that the

projection of the isoperiodic sets to Siegel space via the Torelli map, do not accumulate on some of the components of the boundary of Torelli space.

A first application of these results concerns the topology of Hurwitz spaces of branched coverings over elliptic curves. The connectedness of the moduli space of genus $g > 1$ primitive branched coverings of degree d over an elliptic curve was proven by Berstein and Edmonds (see [10]). Our method allows to retrieve this result, and to get some new information on the fundamental group of these moduli spaces whenever $d > 2$:

Corollary 1.4. *Let $H_{g,d}(E)$ be the Hurwitz space of degree d and genus $g \geq 2$ primitive branched coverings over the elliptic curve E . Let $p : H_1(\Sigma_g) \rightarrow H_1(E)$ be the homology map of any of its elements. For $d \geq 3$ the homomorphism*

$$(4) \quad \pi_1(H_{g,d}(\mathbb{C}/\Lambda)) \rightarrow \text{Stab}(p) \subset \text{Sp}(2g, \mathbb{Z}) \text{ is onto.}$$

This result is in fact more general: the fundamental group of the leaves of the isoperiodic foliation that correspond to a period $p \in \mathcal{H}_g$ when $\deg(p) \neq 2$ surjects onto $\text{Stab}(p) \subset \text{Sp}(2g, \mathbb{Z})$. It is an immediate consequence of Theorem [1.2] (see Remark [3.7]).

Let us mention that, by analogy with the fact that the level sets of the period map might be disconnected in the universal cover of $\Omega\mathcal{M}_g$, the morphism from the fundamental group of a leaf to $\pi_1(\Omega\mathcal{M}_g)$ – which is isomorphic to the mapping class group – has an image that might be strictly contained in the stabilizer of its corresponding period. This is precisely what happens in genus two (see Corollary [6.20]).

A second application concerns the transfer principle: the map Per_g is equivariant with respect to the action of the covering group $\text{Sp}(2g, \mathbb{Z})$ of $\pi : \Omega\mathcal{S}_g \rightarrow \Omega\mathcal{M}_g$ if we consider its action on $\text{Hom}(\mathbb{Z}^{2g}, \mathbb{C})$ by precomposition. This implies that to any \mathcal{F}_g -saturated subset $B \subset \Omega\mathcal{M}_g$ we can associate the $\text{Sp}(2g, \mathbb{Z})$ -invariant subset $A = \text{Per}_g(\pi^{-1}(B)) \subset \mathcal{H}_g$. Reciprocally, for each $\text{Sp}(2g, \mathbb{Z})$ -invariant subset $A \subset \mathcal{H}_g$ we can associate an \mathcal{F}_g saturated subset $B = \pi(\text{Per}_g^{-1}(A))$. These correspondences are inverse one of the other if and only if the fibers of Per_g over A are connected. This correspondence is what allows to transfer properties of the $\text{Sp}(2g, \mathbb{Z})$ action on the homogeneous space [3] to properties of the isoperiodic foliation, and justifies the title of the paper. In view of Theorem [1.2], the transfer principle can be applied on the complement of the primitive degree two points of \mathcal{H}_g . Using Ratner's theory, we deduce the following

Theorem 1.5 (Dynamics of isoperiodic foliations). *Let $g > 2$ and $(C, \omega) \in \Omega^*\mathcal{M}_g$ of volume $V = \frac{i}{2} \int \omega \wedge \bar{\omega}$ and Λ the closure of the image of its periods. Then the closure of the leaf $L(C, \omega)$ passing through (C, ω) is, up to the action of $GL(2, \mathbb{R})$*

- *(Λ is discrete) the component of Hurwitz space consisting of genus g primitive branched coverings over $(\mathbb{C}/\Lambda, dz)$ of volume V .*
- *(Λ is $\mathbb{R} + i\mathbb{Z}$) the set of abelian differentials with periods contained in Λ , with primitive imaginary part, and with volume V ,*
- *($\Lambda = \mathbb{C}$) the subset of $\Omega\mathcal{M}_g$ consisting of abelian differentials of volume V ,*

If $g = 2$ the same statement holds, with an extra possibility occurring when ω is an eigenform for real multiplication by a real quadratic order \mathfrak{o}_D of discriminant $D > 0$. In this case the closure is the Hilbert modular invariant submanifold ΩX_D .

Moreover, the restriction of \mathcal{F}_g to any of these real analytic subsets of $\Omega\mathcal{M}_g$ is ergodic with respect to the Lebesgue class.

Note in particular that this result classifies the algebraic leaves of the isoperiodic foliation: the only closed leaves correspond to Hurwitz spaces and they are known to be algebraic.

1.3. Strategy of the proof of Theorem 1.2. McMullen proved that each fiber of the period map- referred to as isoperiodic moduli space- embeds, via the Torelli map, in Siegel space \mathfrak{S}_g . The image is a Zariski dense subset in a slice of the Schottky locus by a Siegel space of lower genus \mathfrak{S}_{g-1} . In particular, in genus 2 or 3, he deduces that it is connected. In higher genera, the complicated geometry of the Schottky locus makes it difficult to implement this analytical approach.

Our strategy consists in analyzing the isoperiodic degenerations of abelian differentials towards abelian differentials on nodal stable curves. We point out that we are merely interested in degenerating the underlying curve, not the abelian differential (i.e. the form does not degenerate to a zero form on any component). The first step of the proof is to prove that it is always possible to find such degenerations. Adding the obtained nodal forms allows to define a bordification of the isoperiodic moduli space, naturally stratified by the number of nodes of the underlying curves. We prove that this stratification is locally homeomorphic to the pull back by an infinite abelian covering of the stratification associated to a collection of normal crossing divisors on a smooth manifold, ramifying along some components of the divisors. In particular, the connectedness of the fiber is equivalent that of the bordification. For genus at least four, we prove the connectedness of the boundary of a fiber that is not formed by primitive double coverings of an elliptic differential by induction on the genus. This step is essentially achieved by algebraic arguments that encode the combinatorics of the boundary stratification.

As in other instances of this degeneration strategy – see [6, 7] for contemporary examples with strata of differentials– the definition of the bordification requires a comprehensive theory. In the present case the convenient ambient space is the augmented Torelli space, quotient of the augmented Teichmüller space by the Torelli group. For lack of a comprehensive adapted reference we include the theory in section 4 and a guideline for the proof in section 2.

1.4. Notes and references. Our strategy for the proof of the connectedness of the boundary of a fiber of Per works for $g \geq 4$ but leads to difficult algebraic problems due to Haupt's conditions when $g = 2, 3$. The description of the fibers as slices of Schottky space in those cases, is crucial for the inductive argument to work.

The same type of difficulties were already present in the characterization of periods of holomorphic forms on Riemann surfaces by Otto Haupt in [34], namely that those are the periods satisfying Haupt's conditions. Most of his work boiled down to treat the arithmetic involved in the genus two case, caused by forms belonging to the Hilbert modular invariant manifolds X_D described above. From the point of view of slices of Schottky loci, the proof of Haupt consists in finding, by elementary algebraic methods, a point in the intersection of the slice with the boundary strata of Torelli space and then apply a surgery. We actually provide a simpler proof of the result of Otto Haupt, which only differs by the remark that the genus two case can be treated using the Torelli map. Torelli's result (in [64]) was published two years before his death, in 1913, shortly before the outbreak of World War One. We wonder whether Haupt was aware of Torelli's results.

An alternative proof of the result of Otto Haupt with techniques that were completely out of reach at his time was given by Misha Kapovich in [42]. Rather surprisingly, it relies on Ergodic and Ratner's theory.

The main observation of Kapovich is that the set of periods of holomorphic one forms on homologically marked Riemann surfaces is invariant by the action of the integral symplectic group. Its action on the period domain is homogeneous and can be studied through Ratner theory. Our approach pushes the analysis of the period map a bit further to allow to transfer properties of the action to properties of the foliation induced by the map.

At around the time our transfer principle was announced, Hamenstädt proved the ergodicity of the isoperiodic foliation, with a different method. in [29]. She announced analogous results for the intersection of the foliation with connected components of strata of abelian differentials having a simple zero. Weiss and Chaika recently announced, at the Geometry and Dynamics (online) seminar at TAU, a proof of the ergodicity of the isoperiodic foliation on any connected component of any stratum by yet another method. They use recent ongoing research of Eskin, Brown, Filip, Rodriguez-Hertz, etc. on generalizations of the 'Magic Wand Theorem' of Eskin-Mirzakhani (see [24]).

The problem of connectedness of isoperiodic sets has been considered by several authors. Martin Schmoll established in [61], among other things, connectedness of moduli spaces of degree d coverings of a given elliptic curve.

Remark that the same statement fails at the level of the Torelli covering for genus at least five and degree two, as stated in Theorem 3.3. We realized this fact after having pre-published a first version of our work on the arXiv, stating erroneously that those fibers were also connected. Their disconnectedness can be established by generalizing to the context of moduli spaces of double coverings over an elliptic curve a famous work of Arnold, who observed that the inclusion of the hyperelliptic locus in the moduli space of genus $g \geq 3$ curves is not surjective (to the integral symplectic group) at the homological level. See [4].

Fortunately, this exception did not fraud the whole (inductive!) argument, and constitute the single exception of disconnected isoperiodic moduli spaces of abelian differentials.

A problem that arises naturally is the description of the connected components of the intersection of the fibers of Per with the other strata and more generally the affine invariant manifolds that have been brought to light in the work of Eskin and Mirzakhani [24]. Kontsevich and Zorich gave a description of the connected components of strata of abelian differentials without any condition on the periods in [43]; they found cases with up to three components. A direct application of our theorem shows that for generic period, the isoperiodic sets in the stratum will have several components too. Another extreme example of disconnected isoperiodic set: the intersection of a leaf of the isoperiodic foliation corresponding to non-discrete periods with the minimal stratum forms an infinite discrete set. Such phenomena were studied by McMullen in the genus two case [50].

The analogous approach using a transfer principle for studying the dynamical properties of the isoperiodic foliation on strata can *a priori* be considered. There have been recent advances in this direction that point towards understanding the dynamical properties that could eventually be transferred: firstly that of determining the image of each stratum by the period map, and secondly that of determining the monodromy representation, i.e. the image of the symplectic representation of a connected component of a stratum. Concerning the first, a clear obstruction in the case of elements of $H_{g,d}$ is that the order at each branch point

cannot be larger than $d - 1$. In recent preprints [8], [44] the authors show that, together with the positive volume condition, these are the only obstructions to realizing the periods in the stratum. For the second, the monodromy in some connected components of strata has been computed in recent works of Hamenstädt [30, 31] and Calderon and Salter [16, 17]. Recent advances in the description and properties of the closure of strata in spaces of stable forms were achieved by Bainbridge, Chen, Gendron, Grushevsky and Möller in [6, 7]. Nevertheless, we point out that the techniques developed in this article do not seem to be enough to compute the connected components of the fibers of the period map on a stratum. For the time being it is unclear whether the transfer of properties to the isoperiodic foliation on a stratum can be carried.

Some interesting recent works have established similar dynamical properties for the isoperiodic foliations on strata and respectively on affine manifolds. The first contribution is by Hooper-Weiss in [38]; they show that the leaf of the isoperiodic foliation of the Arnoux-Yoccoz surface contained in the stratum $\mathcal{H}(g - 1, g - 1)$ is dense. We have been aware recently that K. Winsor has proven that in case $g = 3$ such a leaf has infinite genus. Also, Ygouf has given an interesting criterion enabling to decide if leaves of isoperiodic foliation are dense in certain affine invariant manifolds of rank one. The rank has been defined in the context of affine manifolds by Wright as half of the codimension of the isoperiodic foliation [70].

We get a nice description of the isoperiodic foliation in the Hilbert invariant submanifolds ΩX_D of \mathcal{F}_2 after taking projectivisation. The foliation corresponds to the horizontal foliation of the uniformization of X_D by the product $\mathbb{H} \times \mathbb{H}$ of two copies of the upper-half plane in \mathbb{C} (see [53]). In that paper McMullen describes precisely the $GL^+(2, \mathbb{R})$ -action after projectivization, and finds some very interesting real analytic foliations by Riemann surfaces on X_D , that are not transversely holomorphic.

It follows from works of Calta (in [18]) and McMullen (in [53]) that the union of leaves of \mathcal{F}_2 intersecting a closed $GL^+(2, R)$ -orbit in the minimal stratum $\mathcal{H}(2)$ has interestingly the structure of a closed analytic subset in the generic stratum $\mathcal{H}(1, 1)$. In the problem paper [40], Problem 12, the authors ask to what extent this phenomenon is general. Our theorem shows that in the generic stratum it holds only for the closed leaves. The question remains completely open in the other strata.

1.5. Organization of the paper. In section 3 we give the proof of the results that do not need the bordification of spaces, and Theorem 1.5 as a consequence of Theorem 1.2. In sections 4 to 8 we prove Theorem 1.2. Along the way, in section 6 we analyze isoperiodic sets with one node and give a proof of Haupt's theorem. In the Appendix I (Section 9) we collect the relevant proofs of the dynamical properties in [42] and extend it to the case of genus two. In Appendix II and Appendix III we give the proofs of technical results needed along the paper that are presumably well known, but for which we have not found specific reference.

1.6. Acknowledgements. We warmly thank the referees of this article for the careful reading and the many improvements that their comments made possible. We also thank U. Hamenstädt, P. Hubert, M. Kapovich, E. Lanneau, F. Loray, D. Margalit, M. Möller, G. Mondello, H. Movasati, A. Putman and A. Wright for useful conversations. This paper was partially supported by the France-Brazil agreement in Mathematics. G. Calsamiglia was partially supported by Faperj/CNPq/CAPES/Mathamsud/ Cofecub, B. Deroin by ANR project LAMBDA ANR-13-BS01-0002, and S. Francaviglia by GNSAGA group of INdAM, and by

PRIN 2017JZ2SW5. It was mainly developed at Universidade Federal Fluminense, IMPA, ENS/Paris, U. Cergy-Pontoise, UPMC, and Università de Bologna, to whom we thank the nice working conditions provided.

2. DETAILED GUIDELINE FOR THE PROOF OF THEOREM 1.2

The Deligne-Mumford-Knudsen orbifold (compact) bordification $\overline{\mathcal{M}}_{g,n}$ of the moduli space $\mathcal{M}_{g,n}$ of smooth genus g curves with n marked points by adding the moduli spaces of stable genus g curves with n marked points. It is a compact orbifold in which the boundary forms a normal crossing divisor. Each stratum of complex codimension $k \geq 0$ of the divisor corresponds to a stratum of curves with k nodes. An intermediate (non-compact) bordification $\mathcal{M}_{g,n} \subset \overline{\mathcal{M}}_{g,n}^c \subset \overline{\mathcal{M}}_{g,n}$ is obtained if we add the moduli spaces of stable curves of compact type, i.e. only with separating nodes. Since the boundary has complex codimension one, it cannot separate the total space.

The Hodge bundle can be naturally extended to the bundle $\Omega\overline{\mathcal{M}}_{g,n}$ of holomorphic stable forms. A stable form can be thought as a meromorphic form on each component of the normalization of a stable curve, called a part, having at worst simple poles at the marked points where they are glued and each pair of points that are glued have opposite residue. The local isoperiodic equivalence relation can be naturally extended to $\Omega\overline{\mathcal{M}}_{g,n}$. As in the smooth case, any stable form admits a non-trivial local isoperiodic deformation. However, the local isoperiodic deformation space is not always a complex manifold, but rather a (possibly singular) analytic set. For generic forms with zero components or residues at the nodes they are strictly contained in the boundary. In each stratum the local isoperiodic deformation is parametrized by a product of local isoperiodic deformations of the parts. In Section 4 we will prove that the local space of isoperiodic deformations at a point of the subset $\Omega_0^*\mathcal{M}_{g,n} \subset \Omega\overline{\mathcal{M}}_{g,n}$ of stable forms with zero residues at the nodes and no zero components is – at the level of the orbifold chart – a smooth complex manifold transverse to each boundary component passing through the point. This implies that the set of points in the local isoperiodic complex manifold that lie on some boundary stratum of the ambient space form a normal crossing divisor. Furthermore the local divisor has precisely one codimension one component for each boundary component of $\Omega\overline{\mathcal{M}}_{g,n}$ passing through the point. In particular the set of points lying on boundary strata do not separate the isoperiodic local space in several components and are stratified by the number of nodes in the underlying curves.

To bordify the fibers of the period map we need to consider the pull back of the bundle of stable forms by the *ramified* cover $\mathcal{S}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$ associated to the Torelli subgroup $\mathcal{I}_{g,n}$. This space can be realized as the quotient of Augmented Teichmüller space, and we call it Augmented Torelli space. The structure and relevant properties of these spaces are detailed in Section 4. We loose the manifold and holomorphic structure around the ramification locus, but we still have a topological stratified space by complex manifolds that lifts the stratification of the boundary divisor of $\mathcal{M}_{g,n}$. The local projection has two properties that we will exploit: over each local connected component of a stratum of the boundary divisor $\overline{\mathcal{M}}_{g,n}$ it is an abelian cover (hence a connected complex manifold of the same dimension). In particular, the open stratum (the complement of the boundary strata) is locally connected around each point on some boundary stratum. Moreover, as happens with the stratification of a normal crossing divisor, every local stratum can be identified by the closures of the codimension one strata where it belongs to or not. This last property is what allows to associate a natural simplicial

complex $\mathcal{C}(\bar{\mathcal{S}}_{g,n})$ to the boundary of $\mathcal{S}_{g,n}$ called the dual boundary complex. A vertex is considered for each connected component of codimension one of the stratification, and among k of them we attach a $k - 1$ simplex for each connected component of the intersection of their closures. These simplices codify all possible connected components of the boundary stratification.

The Hodge bundle $\Omega\bar{\mathcal{S}}_{g,n}$ inherits the same boundary stratification by the number of nodes, with each stratum having the same codimension as in $\bar{\mathcal{S}}_{g,n}$. A word of warning: this stratification is not to be confused with the famous substratification given by the zero and polar sets of the forms. The maximal isoperiodic deformation space on $\Omega\mathcal{S}_{g,n}$ is bordified by considering its closure in $\Omega_0^*\bar{\mathcal{S}}_{g,n}$. It coincides with the maximal isoperiodic deformation there. The transversality of the local isoperiodic deformation space with the boundary components show that we have a stratification of its boundary points by the number of nodes that has similar local properties as $\Omega\bar{\mathcal{S}}_{g,n}$. In particular the set of boundary points does not locally separate the isoperiodic space we can associate a similar simplicial complex to its boundary stratification: a vertex for each connected component of the codimension one stratum and a $k - 1$ -simplex joining k vertices for each connected component of the intersection of the closures of the components.

In Section 5 we will prove that any non-zero form in $\Omega\mathcal{S}_g$ can be isoperiodically deformed to converge to a boundary point of the bordification $\Omega_0^*\bar{\mathcal{S}}_g$. The proof follows by finding families of twin geodesics, that is, parallel geodesic paths of the singular metric defined by a non-zero abelian differential, that start at a saddle point and have the same length. An appropriate surgery along them—namely a Schiffer variation—provide a piece of the desired path in $\Omega\mathcal{S}_g$. For forms with a single zero these pairs of twins are found on the boundary of cylinders of closed geodesics whose existence is guaranteed by Masur’s Annulus Theorem (see [47]).

To prove the connectedness of the bordification of a period fiber we just need to connect all boundary points by isoperiodic deformations. An isoperiodic deformation in a boundary stratum of $\Omega_0^*\bar{\mathcal{S}}_g$ can be thought as a product of isoperiodic deformations of forms on $\Omega^*\mathcal{S}_{h,n}$ with $h < g$. This is how we will use the inductive hypothesis to reduce the proof of the connectedness of the boundary to a combinatorial algebraic problem that we describe next.

Given $p \in \mathcal{H}_g$, we consider the dual boundary complex \mathcal{G}_p of the bordification of $\text{Per}^{-1}(p)$ in $\Omega_0^*\bar{\mathcal{S}}_g$. The definition of the stratifications allow to define a simplicial continuous map

$$(5) \quad \mathcal{G}_p \rightarrow \mathcal{C}(\bar{\mathcal{S}}_g)$$

that associates to each connected component of the boundary stratum of codimension one of the isoperiodic set, the component of the stratum of codimension one of the ambient space where it sits. It extends to simplices in the natural way thanks to the transversality of the local isoperiodic deformation space with boundary components. Its image tells us which components of boundary strata of $\bar{\mathcal{S}}_g$ admit forms of periods p without zero components.

In Section 6 we describe all vertices in the image of (5) by characterizing them algebraically as follows: $\bar{\mathcal{S}}_g$ can be thought as the quotient of augmented Teichmüller space quotiented by the Torelli group \mathcal{J}_g . For this cover the complex is the so called curve complex \mathcal{C}_g of a surface of genus g , introduced by Harvey in [35] (see (see [25] Chapter 4.1] for further details). It has a vertex for each homotopy class of essential simple closed curve on a genus g surface and a $k - 1$ simplex joining k given vertices if the corresponding classes can be realized disjointly by simple closed curves. The subcomplex generated by separating curves is denoted by $\mathcal{C}_g^{\text{sep}}$.

The mapping class group of the surface with marked points at the punctures acts naturally on both complexes and we have

$$\mathcal{C}(\overline{\mathcal{S}}_g) \simeq \mathcal{C}_g/\mathcal{I}_g \text{ and } \mathcal{C}(\overline{\mathcal{S}}_g^c) \simeq \mathcal{C}_g^{\text{sep}}/\mathcal{I}_g$$

The vertex of $\mathcal{C}_g/\mathcal{I}_g$ corresponding to a Torelli class of a simple closed curve c is characterized either by the primitive rank one submodule $\mathbb{Z}[c]$ in the first homology group of the surface \mathbb{Z}^{2g} if this class is non-trivial, or by a splitting of $\mathbb{Z}^{2g} = V_1 \oplus V_2$ into a direct sum of orthogonal symplectic submodules otherwise (i.e. c is separating). Some simplexes connecting those vertices can be identified by simply using this algebraic information. As an instance, a splitting of \mathbb{Z}^{2g} into three factors $V_1 \oplus V_2 \oplus V_3$ that are pairwise orthogonal symplectic submodules, allows to construct a marked stable curve with two separating nodes that induces the given splitting. It determines at least an edge in $\mathcal{C}_g/\mathcal{I}_g$ joining the vertex $V_1 \oplus V_2^\perp$ and the one corresponding to $V_3^\perp \oplus V_3$.

A vertex of non-separating type lies in the image of (5) if and only if $[c] \in \ker p \setminus 0$ and the map induced by p on $[c]^\perp/\mathbb{Z}[c]$ is the period of some non-zero abelian differential on a smooth curve. We say that $[c]$ is pinched by p . A vertex of the second type belongs to the image of (5) if and only if $p|_{V_i}$ is the period of some non-zero abelian differential on a smooth curve for $i = 1, 2$. We say that $V_1 \oplus V_2$ is a p -admissible decomposition. Haupt's conditions is what allows to algebraize the problem, even for other simplexes. We will exploit the following dichotomy to identify vertices: either the image of p has large rank – and we can identify enough p -admissible splittings – or p has large kernel – and we can find enough classes pinched by p .

In Section 7 we find, by using normalization of the node, a homeomorphism of the isoperiodic set of period p in a boundary component of $\Omega\overline{\mathcal{S}}_g$ of codimension one with a product of isoperiodic sets of forms (with marked points!) on lower genera. Using the inductive hypothesis and the cases of genera two and three it allows to determine a family of codimension one boundary components of $\Omega_0^*\overline{\mathcal{S}}_g$ (or vertices of $\mathcal{C}_g/\mathcal{I}_g$), called p -simple, that contain a single isoperiodic connected component of period p (i.e. have a single vertex in the preimage by (5)).

When p is injective every boundary stratum of codimension one of $\Omega\overline{\mathcal{S}}_g$ has at most one isoperiodic component of period p . In fact, all boundary points of $\text{Per}^{-1}(p)$ are of compact type, the bordification is a complex manifold and the boundary a normal crossing divisor in it whose dual complex is \mathcal{G}_p . On the other hand, in subsection 8.1 we prove the connectedness of the image of (5) which actually lies in $\mathcal{C}_g^{\text{sep}}/\mathcal{I}_g$ by analyzing the problem in terms of splittings and the volumes of the parts. Haupt's conditions on the factors of a splitting are reduced to a condition on the volume in this case.

When p is not injective, it is not always true that the boundary divisor of compact type is connected. Examples can be constructed in high genus by taking stable curves with one separating node that have double covers over distinct elliptic curves on each side. When they are double covers of the same elliptic curve and the total degree is two, the curve has necessarily genus two. This is due to the fact that the sum of the degrees is at most two, and the degree of each part is at least one. Therefore both parts have genus one. Hence the need of introducing the boundary components of non-compact type.

Whenever $\deg(p) \geq 3$, we show that the restriction of (5) to a subcomplex \mathcal{G}'_p of \mathcal{G}_p spanned by simple vertices has connected image. Here we need to deal with both of Haupt's conditions.

On the other hand, non-simple vertices of \mathcal{G}_p are always related to degree two coverings of elliptic curves and can be isoperiodically degenerated to pinch enough different non-separating curves to be able to connect them to some p -simple component of the boundary. This implies that \mathcal{G}_p is connected finishing the proof of the inductive step of Theorem 1.2.

The boundary components of $\Omega\bar{\mathcal{S}}_g$ with precisely one non-separating node play a fundamental role in the proof of the general case. One of the main technical difficulties of the paper is to control the number of connected components of the intersection of one of them with the closure of a fiber of $\text{Per}^{-1}(p)$. This is done in Section 7. Here we use once again the bordification strategy, but this time we need to add some strata of the boundary of $\Omega\bar{\mathcal{S}}_g$ where the forms of period $p \in \mathcal{H}_g$ have zero components –of genus one– and where the local isoperiodic deformations have, potentially, wilder singularities. The bordification is best described by normalizing the non-separating node. It is carried in subsections 4.11 and 4.20.




FIGURE 1. The closed cycle b becomes a relative homology class b' after normalization

The normalization map –normalizing the nonseparating node– sends the given set homeomorphically to a subset of $\Omega\mathcal{S}_{g-1,2}$ that lies in a fiber F of the composition

$$(6) \quad \Omega\mathcal{S}_{g-1,2} \xrightarrow{\text{For}} \Omega\mathcal{S}_{g-1} \xrightarrow{\text{Per}_{g-1}^{-1}} \mathcal{H}_{g-1}.$$

In fact this subset is precisely an analytic hypersurface of F , defined by the zeros of a holomorphic function

$$h : \Omega\mathcal{S}_{g-1,2} \rightarrow \mathbb{C}$$

also given by integration– on a relative cycle joining the two marked points as b' in Figure 1. The map h extends holomorphically to the Hodge bundle over a smooth bordification $\mathcal{U}_{g-1,2} \subset \bar{\mathcal{S}}_{g-1,2}$ of $\mathcal{S}_{g-1,2}$ described in subsection 4.11. There is a particular isoperiodic deformation in $\Omega\mathcal{S}_{g-1,2}$ that consists in fixing the underlying marked form (C, m, ω) on a smooth genus $g-1$ curve and moving the marked points in C by preserving the value of the integral along the class b' . If this integral lies in the image of the periods of ω it might happen that both marked points tend to the same point q in C along the isoperiodic deformation. We are then approaching a point in a boundary stratum of $\Omega\bar{\mathcal{S}}_{g-1,2}^c$ whose normalization has a zero component of genus zero with three marked points that is glued to (C, m, ω) at a point $q \in C$ where $\omega(q) = 0$ (see Figure 2 for a representation). Those are the boundary points we add to bordify the isoperiodic set (in subsection 4.11). Each of them has singular local isoperiodic deformation space (see subsection 4.20). Indeed, by varying the point of gluing in the neighbourhood of q we obtain another isoperiodic deformation lying on the boundary. However, as the zeros of ω are isolated, the neighbouring isoperiodic boundary points do not




FIGURE 2. Example of a (singular) point of normal crossing of two regular local isoperiodic components in $\Omega\mathcal{U}_{2,2}$: the marked points lie in a zero component of genus zero that is glued to a simple isolated zero of a genus two component.

have local isoperiodic deformations that leave the boundary. We will need to consider those isoperiodic boundary components to deduce the connectedness of the bordification too.

The problem of relating the topology of a space with that of a hypersurface in it is reminiscent of Lefschetz hyperplane section Theorem. In the present context the lack of an algebraic setting for F and h do not allow to use it directly. However, an application of Simpson's Theorem in [62] –a generalization of Lefschetz theorem for functions defined via integration of a holomorphic one-form– we prove that the hypersurface defined by h on the generic (two dimensional) fiber of the forgetful map $\Omega\bar{\mathcal{S}}_{g-1,2}^c \rightarrow \Omega\bar{\mathcal{S}}_{g-1}$ is connected. Moreover it is a nodal curve whose singular points lie in the boundary and have a local irreducible component completely contained in the boundary and the other transverse to the boundary. Each of those nodal points corresponds the stable forms with a zero component defined above. When the fiber F of the map (6) is over a homomorphism of degree at least three, we can use the inductive hypothesis and the fibration For to deduce that the constructed bordification is contained in a connected analytic set in $\Omega\bar{\mathcal{S}}_{g-1,2}^c$ having possibly some additional smooth irreducible components contained in the boundary. The intersection of each of those boundary components with the other components can be parametrized by a set of isoperiodic forms of genus $g-1$ with a marked simple zero. By analyzing how the zeros of a form can be permuted in the isoperiodic set we deduce that this intersection is connected. This allows to take the boundary irreducible components one by one without changing the total number of connected components, to deduce that the fiber of h in F is connected.

3. THE PERIOD MAP ON THE HODGE BUNDLE OVER TORELLI SPACE

3.1. Torelli map and period fibers. Fix a reference closed connected and oriented surface Σ_g of genus g with $n \geq 0$ ordered marked points $\Sigma_{g,n} = (\Sigma_g, q_1, \dots, q_n)$. A homotopical marking of a smooth genus g compact complex curve C with n pairwise distinct ordered marked points, $P = (p_1, \dots, p_n)$ is a homeomorphism $f : \Sigma_{g,n} \rightarrow (C, P)$ sending q_i to p_i . Two such markings f, f' of (C, P) and (C', P') are equivalent if there exists a biholomorphism

$\varphi : (C, P) \rightarrow (C', P')$ such that $\varphi \circ f$ is isotopic to f' . The Teichmüller space $\mathcal{T}_{g,n} = \mathcal{T}(\Sigma_{g,n})$ for (g, n) satisfying $k = 3g - 3 + n > 0$ is the set of equivalence classes of homotopically marked genus g smooth complex curves with n ordered distinct points endowed with the Teichmüller topology, i.e. the weakest topology for which the length function associated to a homotopy class of closed curve on $\Sigma_{g,n}$ is continuous. A point in $\mathcal{T}_{g,n}$ will be denoted by $[f : \Sigma_{g,n} \rightarrow (C, P)]$.

The mapping class group of $\Sigma_{g,n}$ is the group $\text{Mod}(\Sigma_{g,n})$ of isotopy classes of orientation preserving diffeomorphisms that fix each marked point. It acts on $\mathcal{T}_{g,n}$ by precomposition on the marking.

Bers ([13], Section 16) showed that whenever $k > 0$, the space $\mathcal{T}_{g,n}$ can be embedded in \mathbb{C}^k as a bounded open domain, inheriting a complex structure. Following Ahlfors [2], this complex structure is the only over the given topology for which the coordinate functions of the period matrices of curves depend holomorphically on the curve. In more detail, given a symplectic basis $a_1, b_1, \dots, a_g, b_g$ of $H_1(\Sigma_g, \mathbb{Z})$ (i.e. the only non-zero products of the cycles are $a_i \cdot b_i = 1$ and $b_i \cdot a_i = -1$) we can choose, for each marked curve $f : \Sigma_{g,n} \rightarrow (C, P)$ of genus g the unique basis $\omega_1, \dots, \omega_g$ of $\Omega(C)$ such that

$$\int_{f_* a_i} \omega_j = \delta_{ij}.$$

The maps $\tau_{ij} : \mathcal{T}_{g,n} \rightarrow \mathbb{C}$ defined by

$$(7) \quad \tau_{ij}[f : \Sigma_{g,n} \rightarrow (C, P)] = \int_{f_* b_i} \omega_j$$

are well defined and holomorphic. Let \mathfrak{S}_g denote the Siegel space of genus g , i.e. the set of symmetric $g \times g$ matrices with complex entries whose imaginary part is positive definite. Riemann showed that the squared matrix of functions (τ_{ij}) defines a holomorphic map $\mathcal{T}_{g,n} \rightarrow \mathfrak{S}_g$ that is invariant by the action of the Torelli group $\mathcal{I}_{g,n}$, kernel of the representation

$$\text{Mod}(\Sigma_{g,n}) \rightarrow \text{Aut}(H_1(\Sigma_g, q_1, \dots, q_n, \mathbb{Z})).$$

It induces the Torelli map on the Torelli space $\mathcal{S}_{g,n} := \mathcal{T}_{g,n} / \mathcal{I}_{g,n}$,

$$(8) \quad \mathcal{S}_{g,n} \rightarrow \mathfrak{S}_g.$$

In the case of $n = 0$ it was shown to be injective by Torelli ([64]). Each point in $\mathcal{S}_{g,n}$ is characterized by a triple (C, P, f_*) where f_* is the isomorphism induced by f in homology. Therefore we will denote a point in $\mathcal{S}_{g,n}$ simply as (C, p_1, \dots, p_n, m) where $m : H_1(\Sigma_g, q_1, \dots, q_n, \mathbb{Z}) \rightarrow H_1(C, p_1, \dots, p_n, \mathbb{Z})$ is an isomorphism. The quotient $\mathcal{T}_{g,n} / \text{Mod}(\Sigma_{g,n})$ is the moduli space $\mathcal{M}_{g,n}$ of genus g curves with n marked points and the holomorphic structure induces an orbifold structure on $\mathcal{M}_{g,n}$. The Hodge bundle is a holomorphic vector bundle $\Omega\mathcal{M}_{g,n} \rightarrow \mathcal{M}_{g,n}$ whose fiber over a point (C, P) is the set of abelian differentials $\omega \in \Omega(C)$. It can be pulled back to a holomorphic bundle $\Omega\mathcal{S}_{g,n} \rightarrow \mathcal{S}_{g,n}$.

Definition 3.1. The period map on $\Omega\mathcal{S}_{g,n}$ is the holomorphic map

$$\text{Per}_{g,n} : \Omega\mathcal{S}_{g,n} \rightarrow \text{Hom}(H_1(\Sigma_g, q_1, \dots, q_n, \mathbb{Z}), \mathbb{C})$$

defined by $\text{Per}_{g,n}(C, p_1, \dots, p_n, m, \omega) = \{\gamma \mapsto \int_{m(\gamma)} \omega\}$. When $n = 0$ we write $\text{Per}_g = \text{Per}_{g,0}$. When there is no risk of confusion we omit all subindices and write Per .

For instance given a homomorphism $p : H_1(\Sigma_g, q_1, \dots, q_n, \mathbb{Z}) \rightarrow \mathbb{C}$ we denote $\text{Per}^{-1}(p)$ the fiber of $\text{Per}_{g,n}$ over p .

Multiplying forms by a non-zero constant induces a biholomorphism between fibers of $\text{Per}_{g,n}$.

In the case $n = 0$ the properties of the Torelli map and of the intersection form in $H_1(\Sigma_g, \mathbb{Z})$ have a nice consequence:

Theorem 3.2. [50] *Let $g \geq 2$ and $p : H_1(\Sigma_g, \mathbb{Z}) \rightarrow \mathbb{C}$ be a homomorphism with $\text{vol}(p) > 0$. Then $\text{Per}_g^{-1}(p)$ is biholomorphic to the intersection of a linear Siegel space $\mathfrak{S}_{g-1} \subset \mathfrak{S}_g$ and the image of the Torelli map – the so-called Schottky locus. –*

Proof. We take the choices and notations of the definition of (7). Let $(C, m, \omega) \in \text{Per}^{-1}(p)$. Its expression in the chosen basis reads $\omega = p(a_1)\omega_1 + \dots + p(a_g)\omega_g$ and the following equations are satisfied:

$$(9) \quad m_{ij} = \tau_{ij}(C, m) \text{ for } i, j = 1, \dots, g$$

$$(10) \quad p(b_i) = p(a_1)m_{i1} + \dots + p(a_g)m_{ig} \text{ for } i = 1, \dots, g.$$

A point $Z = (m_{ij}) \in \mathfrak{S}_g$ that satisfies (9) is said to belong to the Schottky locus in \mathfrak{S}_g . If it furthermore satisfies (10), then $\omega := \sum p(a_j)\omega_j$ is the unique abelian differential on (C, m) having periods p . This proves that $\text{Per}^{-1}(p)$ is biholomorphic to the intersection of the Schottky locus with the set of solutions of (10).

It remains to show that the condition $\text{vol}(p) > 0$ implies that the set of all solutions $Z = (m_{ij}) \in \mathfrak{S}_g$ of (10) (that we call the set of matrices that admit p as a period) is biholomorphic to \mathfrak{S}_{g-1} . Up to multiplying all forms in $\text{Per}^{-1}(p)$ by a constant we can suppose that all forms have volume one, i.e. $\text{vol}(p) = 1$.

The symplectic automorphism group of $H_1(\Sigma_g, \mathbb{R})$ acts on Siegel space (by changing the marking). Direct calculation shows that, writing $T \in \text{Sp}(2g, \mathbb{R})$ in the basis $\{a_i, b_i\}$ as block $g \times g$ real matrices with two lines A, B and C, D the action is defined by

$$Z' = T \star Z = (AZ + B)(CZ + D)^{-1}.$$

Moreover denoting $p_{\mathbb{R}} : H_1(\Sigma_g, \mathbb{R}) \rightarrow \mathbb{C}$ the natural extension of $p : H_1(\Sigma_g, \mathbb{Z}) \rightarrow \mathbb{C}$, the given action preserves the condition $p_{\mathbb{R}}$ is a period of the matrix.

By definition, $\Re(p_{\mathbb{R}}) \cdot \Im(p_{\mathbb{R}}) = \Re(p) \cdot \Im(p) = \text{vol}(p) = 1$. On the other hand, by duality there exist elements $a_1, b_1 \in H_1(\Sigma_g, \mathbb{R})$ such that $a_1 \cdot b_1 = 1$, $a_1^* = \Re p_{\mathbb{R}}$ and $b_1^* = \Im p_{\mathbb{R}}$. Now, $\ker p_{\mathbb{R}} = \ker \Re(p_{\mathbb{R}}) \cap \ker \Im(p_{\mathbb{R}})$ and contains the rank $2g - 2$ symplectic orthogonal of $\mathbb{R}a_1 \oplus \mathbb{R}b_1$. We deduce $\ker p$ is symplectic of rank $2g - 2$. Choose a symplectic basis $a_2, b_2, \dots, a_g, b_g$ of $\ker p_{\mathbb{R}}$. The matrix in Siegel space associated to the real basis $\{a_i, b_i\}$ has two diagonal blocks: $1 \in \mathbb{C}$ and a matrix in \mathfrak{S}_{g-1} . Every such matrix admits $p_{\mathbb{R}}$ as period, trivially. Therefore the set of solutions of (10) is also a linear Siegel subspace $\mathfrak{S}_{g-1} \subset \mathfrak{S}_g$. \square

Theorem 3.3. *Let $g = 2, 3$ and $p \in H^1(\Sigma_g, \mathbb{C})$ with $\text{vol}(p) > 0$ and $\deg(p) > 1$. Then $\text{Per}^{-1}(p)$ is non-empty and connected.*

Proof. The Schottky locus is Zariski dense in \mathfrak{S}_g for $g = 2, 3$. Its complement corresponds to period matrices of curves of compact type. By Theorem 3.2 we have that $\text{Per}^{-1}(p)$ is biholomorphic to a Zariski open set of \mathfrak{S}_{g-1} , hence connected.

Suppose $\text{Per}^{-1}(p)$ is empty. Then, the linear subspace of Theorem 3.2 is completely contained in the complement of the Schottky locus. The same construction of forms of the proof of Theorem 3.2 (solutions of (9) and (10)) can be carried on Jacobians of curves of compact type to construct stable forms of periods p .

From a stable form of compact type that has no zero component we can obtain a form on a smooth curve having the same periods. Indeed by taking parallel slits instead of points to glue the different components we smoothen the curve, without changing the period map.

If $g = 2$, the curve has two components of genus one and we know that one of the components of the form is zero. This implies $\deg(p) = 1$. A contradiction.

If $g = 3$ the curve has either a part of genus two and a part of genus one or three parts of genus one and the stable form is zero in one of the parts. If it is zero on a genus two part, it implies $\deg(p) = 1$, contrary to assumption. If it is zero in just one part of genus one we can suppose that we have a form $\omega \neq 0$ on a smooth curve of genus two. It has at least one zero z_1 . Consider the local map $z \mapsto \int_{z_1}^z \omega$ and the pre-image γ of a small segment $[0, \varepsilon]$. It is a path with distinct endpoints satisfying $\int_\gamma \omega = 0$. Gluing the endpoints produces a stable form on a curve that is not of compact type. If we take parallel slits at the endpoints instead, the gluing produces a stable form on a smooth curve of genus three. Marking the curve appropriately we obtain that its period is p . A contradiction. \square

For $g \geq 4$ the image of the Torelli map is an analytic set of positive codimension (as an easy dimension count shows) and determining it is known as the Schottky problem. In fact, as stated in Theorem 1.3, there are some of the intersections given by Theorem 3.2 that will not be connected. As for the non-emptiness of the fiber of Per over points of positive volume and primitive degree at least two, we will show inductively that they can be realized as period of a stable form without zero components as in the proof of Theorem 3.3 and apply the surgeries. These surgeries will be extensively developed in Section 5 to produce continuous isoperiodic deformations of stable forms.

3.2. Strata of holomorphic forms on smooth curves and isoperiodic foliations. It is well known that the spaces $\Omega^* \mathcal{M}_{g,n}$ and $\Omega^* \mathcal{S}_{g,n}$ are stratified by the properties of the zero sets of forms. Two points (C, P, ω) , (C', P', ω') lie in the same stratum if there exists a homeomorphism $(C, P) \rightarrow (C', P')$ sending the zero divisor (ω) to the zero divisor (ω') . In other words, each marked point is sent to a marked point, each zero to a zero, and the orders of the zeros are preserved. The generic stratum $\Omega^{SZ} \mathcal{S}_{g,n}$ is formed by forms with $(2g - 2)$ simple zeros, none of which is a marked point. The minimal stratum $\Omega \mathcal{S}_{g,n}(2g - 2)$ is formed by abelian differentials with a single zero (at some marked point if there are any). Veech ([65]) and Masur ([49]) showed that there are local holomorphic coordinates defined on the stratum of a point (C, P, ω) , with coordinates in

$$\text{Hom}(H_1(C, Z(\omega) \cup P, \mathbb{Z}), \mathbb{C}),$$

defined by integration. As a consequence, the period map restricted to any stratum is a linear projection in the coordinates, and is therefore submersive and open. This implies that the fibers of the map $\text{Per}_{g,n}$ are regular and transverse to all strata different from the generic stratum. In particular, in restriction to $\Omega \mathcal{S}_{g,n}(2g - 2)$, the map $\text{Per}_{g,n}$ is a local biholomorphism. On the other hand it also proves that the restriction of the map $\text{Per}_{g,n}$ to each stratum of stable forms defines a regular holomorphic foliation of codimension $2g$.

The natural action of $\text{Mod}(\Sigma_{g,n})$ on $\Omega\mathcal{S}_{g,n}$ preserves each stratum, and the map $\text{Per}_{g,n}$ is equivariant with respect to the corresponding action on $\text{Hom}(H_1(\Sigma_g, q_1, \dots, q_n; \mathbb{Z}), \mathbb{C})$.

Definition 3.4. The regular foliation induced by $\text{Per}_{g,n}$ on $\Omega^*\mathcal{S}_{g,n}$ (and its restriction to each stratum) descends to $\Omega^*\mathcal{M}_{g,n}$ as a regular holomorphic foliation called the isoperiodic foliation and denoted $\mathcal{F}_{g,n}$ (or $\mathcal{F}_g = \mathcal{F}_{g,0}$ when there are no marked points).

If $L \subset \Omega\mathcal{M}_g$ is the leaf of \mathcal{F}_g corresponding to the periods $p \in \mathcal{H}_g$, the restriction of the Torelli cover to the fiber

$$(11) \quad \text{Per}^{-1}(p) \rightarrow L$$

is a Galois covering. We can give some information about the covering group.

Definition 3.5. Given a symplectic module M over a ring and a homomorphism $p : M \rightarrow \mathbb{C}$, the stabilizer of p is the subgroup of the group of symplectic automorphisms of M defined by

$$\text{Stab}(p) = \{h \in \text{Aut}(M) : p \circ h = p\}$$

When we want to stress the group it sits in we denote it $\text{Stab}_{\text{Aut}(M)}(p)$.

Remark 3.6. Let $p = \text{Per}(C, m, \omega)$ be the period map of a marked abelian differential $(C, m, \omega) \in \Omega\mathcal{S}_{g,0}$. Then the group $\text{Stab}(p)$ also stabilizes the fiber $\text{Per}^{-1}(p)$. The isotopy class of a Dehn twist around any simple closed curve in C defining a primitive element $a \in \ker p \setminus 0$, induces a non-trivial element $\delta_a \in \text{Stab}(p)$ whose action fixes no point.

Remark 3.7. The covering group of the map [\(11\)](#) is $\text{Stab}(p)$. In particular, if $\text{Per}^{-1}(p)$ is connected, the image of $\pi_1(L)$ in $\text{Sp}(2g, \mathbb{Z})$ is precisely $\text{Stab}(p)$, which is non-trivial as soon as $\ker(p) \neq 0$.

3.3. Disconnected fibers: proof of Theorem [1.3](#). Let $p \in \mathcal{H}_g$ with $\deg p = 2$. We will define an invariant that is constant on each component of $\text{Per}^{-1}(p)$. On the other hand we will prove that if $g \geq 5$, the stabilizer of p in the symplectic group $\text{Aut}(H_1(\Sigma_g, \mathbb{Z}))$ is large enough to guarantee that the invariant takes at least two values on any orbit of its action on $\text{Per}^{-1}(p)$.

Consider the elliptic curve $E := \mathbb{C}/\text{Im}(p)$. To any $(C, m, \omega) \in \text{Per}^{-1}(p)$ we can associate a double branched covering $\pi : C \rightarrow E$ defined as the integral of ω based at some point. It is well-defined up to post composition by a translation in E . Denote by $C(\pi)$ (resp. $VC(\pi)$) the set of critical points (resp. critical values) of π .

As observed by Arnold, see [\[4\]](#), the map $\pi_* : H_1(C \setminus C(\pi), \mathbb{Z}/2\mathbb{Z}) \rightarrow H_1(E \setminus VC(\pi), \mathbb{Z}/2\mathbb{Z})$ extends as a homomorphism

$$(12) \quad \Pi : H_1(C, \mathbb{Z}/2\mathbb{Z}) \rightarrow H_1(E \setminus VC(\pi), \mathbb{Z}/2\mathbb{Z}).$$

This is due to the fact that any cycle turning once around a critical point of π is mapped to a cycle turning twice around a critical value of π . The image of Π is the kernel of the monodromy representation $\varepsilon : H_1(E \setminus VC(\pi_0), \mathbb{Z}/2\mathbb{Z}) \rightarrow \mathbb{Z}/2\mathbb{Z}$ of the covering π .

To define an invariant that does not depend on the choices made so far, we need to choose a reference for the homology and the critical values. Fix a reference subset $VC_0 \subset E$ of cardinality $2g - 2$. There exists a homeomorphism φ of E sending $VC(\pi)$ to VC_0 , that is homotopic to the identity. Two such choices differ by post-composition by an element of the braid group of the pair (E, VC_0) . We define Arnold's invariant of (C, m, ω) as the map

$$\text{Ar}(C, m, \omega) := \varphi_* \circ \Pi \circ m \in \text{Hom}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}), H_1(E \setminus VC_0, \mathbb{Z}/2\mathbb{Z}))/G,$$

where $\varphi_* : H_1(E \setminus VC(\pi), \mathbb{Z}/2\mathbb{Z}) \rightarrow H_1(E \setminus VC_0, \mathbb{Z}/2\mathbb{Z})$ is the natural map induced by φ , and where $G \subset \text{Aut}(H_1(E \setminus VC_0, \mathbb{Z}/2\mathbb{Z}))$ is the image of the natural representation of the braid group of the pair (E, VC_0) in the homology group $H_1(E \setminus VC_0, \mathbb{Z}/2\mathbb{Z})$, acting on $\text{Hom}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}), H_1(E \setminus VC_0, \mathbb{Z}/2\mathbb{Z}))$ by post-composition. By construction the Arnold invariant is constant on every connected component of $\text{Per}^{-1}(p)$.

The group $\text{Aut}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}))$ of linear automorphisms preserving the symplectic structure acts on $\text{Hom}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}), H_1(E \setminus VC_0, \mathbb{Z}/2\mathbb{Z}))/G$ by precomposition. Now, if the image of the stabilizer of p in $\text{Aut}(H_1(\Sigma_g, \mathbb{Z}))$ by the mod 2 reduction homomorphism

$$\text{Aut}(H_1(\Sigma_g, \mathbb{Z})) \rightarrow \text{Aut}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}))$$

does not stabilize the Arnold invariant of an element, it means that there are at least two values of the invariant in the orbit. We will compare the sizes of the stabilizer and the image to conclude.

Lemma 3.8. *For any $(C, m, \omega) \in \text{Per}^{-1}(p)$ the stabilizer of $Ar(C, m, \omega)$ in $\text{Aut}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}))$ has order at most $2^{4g-5}(2g-2)!$, i.e.*

$$|\text{Stab}_{\text{Aut}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}))}(Ar(C, m, \omega))| \leq 2^{4g-5}(2g-2)!$$

Proof. Let φ be a homeomorphism of E sending $VC(\pi)$ to VC_0 , that is homotopic to the identity, and define $A := \varphi_* \circ \Pi \circ m$, where $\varphi_* : H_1(E \setminus VC(\pi); \mathbb{Z}/2\mathbb{Z}) \rightarrow H_1(E \setminus VC_0, \mathbb{Z}/2\mathbb{Z})$ is the map induced by φ on homology. The Arnold's invariant is the class of A modulo post composition by an element of G . We need to count the number of elements $M \in \text{Aut}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}))$ such that there exists an element $g \in G$ satisfying

$$A \circ M = g \circ A.$$

The map g preserves the image of A , and $g|_{\text{Im}(A)}$ is determined by M . So we have a well-defined representation

$$\rho : \text{Stab}_{\text{Aut}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}))}(A) \rightarrow \text{End}(\text{Im}(A)) \text{ given by } \rho(M) := g|_{\text{Im}(A)}.$$

To bound the cardinality of $\text{Stab}_{\text{Aut}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}))}(A)$ we proceed to bound the cardinality of the image and kernel of ρ .

To bound the cardinality of the image of ρ , let us analyze the action of G on $H_1(E \setminus VC_0, \mathbb{Z}/2\mathbb{Z})$. Consider the natural map $i_* : H_1(E \setminus VC_0, \mathbb{Z}/2\mathbb{Z}) \rightarrow H_1(E, \mathbb{Z}/2\mathbb{Z})$ given by the inclusion. Its kernel can be identified with the space $(\oplus_{v \in CV_0} (\mathbb{Z}/2\mathbb{Z})v) / (\mathbb{Z}/2\mathbb{Z})\sigma$, with $\sigma = \sum_{v \in CV_0} v$, where $v \in CV_0$ corresponds to the cycle turning once around v . We have the exact sequence

$$(13) \quad 0 \rightarrow (\oplus_{v \in CV_0} (\mathbb{Z}/2\mathbb{Z})v) / (\mathbb{Z}/2\mathbb{Z})\sigma \rightarrow H_1(E \setminus VC_0, \mathbb{Z}/2\mathbb{Z}) \xrightarrow{i_*} H_1(E, \mathbb{Z}/2\mathbb{Z}) \rightarrow 0.$$

The group G preserves $\ker i_*$ and acts on it by permutations. Moreover, it acts trivially on the quotient $H_1(E \setminus VC_0, \mathbb{Z}/2\mathbb{Z}) / \ker i_* \simeq H_1(E, \mathbb{Z}/2\mathbb{Z})$ because any element of the braid group is isotopic to the identity once marked points are forgotten.

Since the image of Π is the kernel of the monodromy representation of the branched covering $\pi : C \rightarrow E$, the subspace $\text{Im}(A) \subset H_1(E \setminus VC_0, \mathbb{Z}/2\mathbb{Z})$ is a hyperplane, whose intersection with $\ker i_*$ is the $(2g-4)$ -dimensional vector space $\ker i_* \cap (\text{Im}(A)) \simeq (\oplus_{v \in CV(\pi)} (\mathbb{Z}/2\mathbb{Z})v)^{\text{even}} / (\mathbb{Z}/2\mathbb{Z})\sigma$

(i.e. the formal sums of an even number of critical values of π). Moreover, $i_*(\text{Im}(A)) = H_1(E, \mathbb{Z}/2\mathbb{Z})$. So the exact sequence (13) induces an exact sequence

$$(14) \quad 0 \rightarrow (\oplus_{v \in CV_0} (\mathbb{Z}/2\mathbb{Z})v)^{\text{even}} / (\mathbb{Z}/2\mathbb{Z})\sigma \rightarrow \text{Im}(A) \xrightarrow{i_*} H_1(E, \mathbb{Z}/2\mathbb{Z}) \rightarrow 0.,$$

which is invariant by ρ , the action of this latter on the left-hand module being made by permutations.

Since the group of automorphisms of the exact sequence (14) acting on its left hand side by permutations has cardinality at most $(2g-2)! \times 2^{2(2g-4)}$, we get the bound

$$(15) \quad |\text{Im}(\rho)| \leq (2g-2)! \times 2^{2(2g-4)}.$$

Let us now bound the number of elements of the kernel of ρ . Let $M \in \ker(\rho)$, which means that $A \circ M = A$. Write $M = I + \psi$ where $\psi : H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}) \rightarrow \ker(A)$.

Claim: $\ker(A)$ is a two-dimensional isotropic subspace of $H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z})$.

Proof of the claim. From [10] we deduce that, up to composition by homeomorphisms in source and target, there is only one double branched covering from a genus g closed oriented connected surface to E . It is therefore sufficient to verify the statement on an example that we depict in Figure 3.




FIGURE 3. Example of double cover with bi-dimensional isotropic $\ker A$

First we construct a $2 : 1$ branched covering $\pi' : \Sigma' \rightarrow \mathbb{P}^1$ on a connected closed oriented surface of genus $g-2$, whose associated Arnold's map $A' : H_1(\Sigma', \mathbb{Z}/2\mathbb{Z}) \rightarrow H_1(\mathbb{P}^1 \setminus VC(\pi'), \mathbb{Z}/2\mathbb{Z})$ is injective. Let $z_1, z_2, \dots, z_{2g-2} \in \mathbb{P}^1$ be distinct points in the Riemann sphere, and let $I_0, \dots, I_{g-2} \subset \mathbb{P}^1$ be pairwise disjoint segments, such that $\partial I_k = \{z_{2k+1}, z_{2k+2}\}$. Slit two copies of $\mathbb{P}^1 \setminus (\cup_k I_k)$ and glue them together by the usual rule: the right side of I_k in one copy is glued to the left side of I_k in the second, and vice versa. We get a branched double cover $\pi' : \Sigma' \rightarrow \mathbb{P}^1$ whose critical values are the z_k 's. We let $\alpha_1, \dots, \alpha_{g-2}$ be simple closed curves that are boundaries of small neighborhoods of the I_k 's. Let $\{J_k\}_{k=1, \dots, g-2}$ be a disjoint

family of embedded segments in the Riemann sphere, the boundary of J_k intersecting the union $\cup I_l$ only in its extremities z_{2k} and z_{2k+1} . We denote by β_k the boundary of a small neighborhood of J_k . The curves $\alpha_1, \beta_1, \dots, \alpha_{g-2}, \beta_{g-2}$ lift to curves $a'_1, b'_1, \dots, a'_{g-2}, b'_{g-2} \subset \Sigma'$ forming a symplectic basis of Σ' . Notice that the homology of $\mathbb{P}^1 \setminus \{z_1, \dots, z_{2g-2}\}$ with coefficients in $\mathbb{Z}/2\mathbb{Z}$ is generated by the cycles associated to small simple closed curves c'_k turning around z_k , the only relation being that the sum of these cycles is zero. By construction we have $A'(a'_k) = c'_{2k+1} + c'_{2k+2}$ and $A'(b'_k) = c'_{2k} + c'_{2k+1}$. From this we easily prove that A' is injective.

Now, take an open disc Δ in the Riemann sphere that contains the union of the intervals I_k and J_k , and the curves α_k, β_k, c_k , for $k = 1, \dots, g$. The surface $\Sigma \setminus (\pi')^{-1}(\Delta)$ is a union of two discs corresponding to lifts of $\mathbb{P}^1 \setminus \Delta$. Consider a torus E , and an embedding $i : \Delta \rightarrow E$. Attach to $(\pi')^{-1}(\Delta)$ two copies of $E \setminus i(\Delta)$ along the identification of their boundaries given by the maps i and π . We get a surface Σ of genus g , and a branched covering $\pi : \Sigma \rightarrow E$ defined by π' in $(\pi')^{-1}(\Delta)$ and by the natural identification of the two components of $\Sigma \setminus (\pi')^{-1}(\Delta)$ with $E \setminus i(\Delta)$. Denoting by $\alpha, \beta \subset E$ oriented simple closed curves that do not intersect $i(\Delta)$ and that form a symplectic basis of $H_1(E, \mathbb{Z})$, we let $a_{g-1}, b_{g-1} \subset \Sigma$ and $a_g, b_g \subset \Sigma$ the corresponding cycles in the two copies of $E \setminus i(\Delta)$. We denote by $a_1, b_1, \dots, a_{g-2}, b_{g-2} \subset \Sigma$ the simple closed curves equal to the curves $a'_1, b'_1, \dots, a'_{g-2}, b'_{g-2}$ in $(\pi')^{-1}(\Delta)$. The cycles $a_1, b_1, \dots, a_g, b_g$ in $H_1(\Sigma, \mathbb{Z}/2\mathbb{Z})$ form a symplectic basis, and we have

$$A(a_k) = c_{2k+1} + c_{2k+2}, \quad A(b_k) = c_{2k} + c_{2k+1} \text{ if } k \leq g-2$$

and

$$A(a_{g-1}) = A(a_g) = a, \quad A(b_{g-1}) = A(b_g) = b,$$

where, in these formula, we denote by $c_k = i(c'_k)$. The kernel of A is then the space generated by $a_{g-1} - a_g$ and $b_{g-1} - b_g$, which is isotropic modulo 2. The claim follows. \square

Since M is symplectic and $\ker A$ isotropic of dimension two, ψ satisfies $\psi + \psi^* = 0$, and induces an anti-symmetric map $H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z})/\ker(A)^\perp \rightarrow \ker(A)$ (here the anti-symmetric character is relative to the natural duality between $\ker(A)$ and $H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z})/\ker(A)^\perp$ given by the intersection form). Since $\ker(A)$ is two dimensional, there is a 3-dimensional space of such maps ψ over $\mathbb{Z}/2\mathbb{Z}$. In particular, we have

$$(16) \quad |\ker(\rho)| \leq 2^3.$$

The lemma follows immediately from bounds (15) and (16). \square

We denote by $p[2]$ the reduction of p modulo two, namely the map

$$(17) \quad p[2] : H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}) \rightarrow \text{Im}(p)/2\text{Im}(p) \simeq H_1(E, \mathbb{Z}/2\mathbb{Z}).$$

More geometrically, for an element $(C, m, \omega) \in \text{Per}^{-1}(p)$, we have $p[2] = i_* \circ \varphi_* \circ \Pi \circ m$, where $i_* : H_1(E \setminus VC_0, \mathbb{Z}/2\mathbb{Z}) \rightarrow H_1(E, \mathbb{Z}/2\mathbb{Z})$ is the map induced by inclusion.

To conclude the proof of Theorem 1.3, we use an algebraic result whose proof can be found in Appendix II (see section 10).

Lemma 3.9. *If $g \geq 3$, the image of the mod 2 reduction map*

$$\text{Stab}_{\text{Aut}(H_1(\Sigma_g, \mathbb{Z}))}(p) \rightarrow \text{Aut}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}))$$

is $\text{Stab}_{\text{Aut}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}))}(p[2])$. Its cardinality is $2^{2g-2} \times 2^{2g-3} \times (2^{2g-4} - 1) \times 2^{2g-5} \times \dots \times (2^2 - 1) \times 2$.

The combination of Lemma 3.8 and Lemma 3.9 show that Arnold's invariant takes at least

$$\frac{2^{2g-2} \times 2^{2g-3} \times (2^{2g-4} - 1) \times 2^{2g-5} \times \dots \times (2^2 - 1) \times 2}{2^{4g-5}(2g-2)!}$$

distinct values on any orbit of $\text{Stab}_{\text{Aut}(H_1(\Sigma_g, \mathbb{Z}))}(p)$ in $\text{Per}^{-1}(p)$. This number is strictly larger than 1 if $g \geq 5$, hence the proof of Theorem 1.3 is complete.

In the case of genus $g = 4$, a more detailed study allows to show that the Arnold's invariant takes only one value on the isoperiodic spaces of degree two (and actually, realizes another instance of the exceptional isomorphism between $\text{Sp}(4, \mathbb{Z}/2\mathbb{Z})$ and S_6). It is plausible that these isoperiodic sets are connected indeed, but we could not prove it by using this invariant.

3.4. Dynamics of the action of $\text{Sp}(2g, \mathbb{Z})$ on \mathcal{H}_g . In this subsection, we state the extension of the analysis of the linear action of $\Gamma = \text{Sp}(2g, \mathbb{Z})$ on the set of periods of positive volume, that were presented by Kapovich in [42] for the cases $g \geq 3$ to the case of genus $g = 2$. For the sake of completeness we include the proof in Appendix I (see section 9). This will enable us to derive Theorem 1.5 from Theorem 1.2.

The closure of the orbit of a period p depends heavily on the rationality properties of the real two dimensional symplectic subspace

$$(18) \quad W = \mathbb{R}\Re p + \mathbb{R}\Im p \subset \mathbb{R}^{2g}$$

defined over smaller fields. Except for the case where W is defined over a quadratic field and $W^\sigma = W^\perp$ for the Galois involution, the different cases can be characterized by the dimension of the maximal subspace of W defined over the rationals. This dimension can also be detected by the topological properties of the closure of the \mathbb{Z} -submodule $\Lambda(p)$ of \mathbb{C} generated by the entries of $p \in \mathbb{C}^{2g}$. It is two if $\Lambda(p)$ is discrete, one if it is isomorphic to $\mathbb{R} + i\mathbb{Z}$ and zero if it is dense. Notice that this submodule is invariant under the action of Γ on X . The different possibilities in Theorem 1.5 arise from this analysis at the level of this submodule. We resume the analysis in the following

Proposition 3.10. *Assume $g > 2$. For any $p \in \mathbb{C}^{2g}$ of positive volume, we have the trichotomy for $W = \mathbb{R}\Re p + \mathbb{R}\Im p$*

- *W is defined over \mathbb{Q} . In this case, $\Lambda(p)$ is discrete and either p is the period of a finite branched covering of the elliptic differential $(\mathbb{C}/\Lambda(p), dz)$ or it is a collapse of $g - 1$ handles. The set $\overline{\Gamma \cdot p}$ is the set of periods $q \in \mathbb{C}^{2g}$ of volume $V(q) = V(p)$ such that $\Lambda(q) = \Lambda(p)$.*
- *W is not rational but contains a rational line. In this case, up to the action of $GL(2, \mathbb{R})$, the set $\overline{\Lambda(p)}$ is $\mathbb{R} + i\mathbb{Z}$, and $\overline{\Gamma \cdot p}$ is the set of periods $q \in \mathbb{C}^{2g}$ of volume $V(q) = V(p)$ whose imaginary part are integer valued and primitive.*
- *W does not contain any rational subspace of positive dimension. In this case, $\overline{\Lambda(p)} = \mathbb{C}$, and $\overline{\Gamma \cdot p}$ is the set of periods $q \in \mathbb{C}^{2g}$ such that $V(q) = V(p)$.*

In genus $g = 2$, there is another possibility:

- *W is defined over a quadratic field K , and $W^\perp = W^\sigma$, where σ is the Galois involution of K . In this case $\overline{\Gamma \cdot p}$ is the set of periods q which differ from p by post-composition by an element of $GL(2, \mathbb{R})$, and by pre-composition by an element of $Sp(4, \mathbb{Z})$.*

In any case, the action of Γ is ergodic in $\overline{\Gamma \cdot p}$.

3.5. Proof of (Theorem 1.2)_g \Rightarrow (Theorem 1.5)_g. We suppose that we know that Theorem 1.2 is true for some genus $g \geq 2$ and hence by the Transfer Principle, invariant sets of the isoperiodic foliation $\mathcal{F}_g := \mathcal{F}_{g,0}$ correspond bijectively to invariant sets of the $\mathrm{Sp}(2g, \mathbb{Z})$ action on \mathcal{H}_g . We want to deduce Theorem 1.5 for genus g . For the sake of simplicity we omit the sub-index of $\mathrm{Per}_{g,0}$ and write Per .

The closure \bar{L} of the leaf $L = L(C, \omega)$ of \mathcal{F}_g through $(C, \omega) \in \Omega\mathcal{M}_g$ is a closed \mathcal{F}_g -invariant set. By the Transfer principle (see Theorem 1.2), after choosing a marking m of C , it corresponds to the projection from $\Omega\mathcal{S}_g$ to $\Omega\mathcal{M}_g$ of all marked abelian differentials having periods in the closure of the $\mathrm{Sp}(2g, \mathbb{Z})$ -orbit of $p = \mathrm{Per}(C, m, \omega) \in \mathcal{H}_g$. The final ergodicity part of the statement of Theorem 1.5 follows from the final statement in Proposition 3.10.

Case 1: If ω is a genus two eigenform for real multiplication by a real quadratic order \mathfrak{o}_D of discriminant $D > 0$. Then the closure of L is one of the Hilbert modular manifolds ΩX_D (see [18] or [52]). As shown in [52, Case 3. of Theorem 5.1] this case occurs if and only if p has the properties described in the last possibility of Proposition 3.10.

In what follows we suppose that p does not satisfy the last condition of Proposition 3.10. The other possibilities can be characterized in terms of the closure of $\Lambda(p)$ in \mathbb{C} :

Case 2: $\Lambda(p)$ is discrete. The image of the periods of any $(C', \omega') \in \bar{L}$ are precisely $\Lambda(p)$.

Each element in the fibre $\mathrm{Per}^{-1}(p) \subset \Omega\mathcal{S}_g$ corresponds, by integration, to a unique primitive branched covering over $\mathbb{C}/\Lambda(p)$ of volume V . If we find a marking m' of C' where $\mathrm{Per}(C', m', \omega') = p$, then by Theorem 1.2 we have $(C', \omega') \in L$. Therefore the leaf L is closed. The first item of Proposition 3.10 (or equivalently Lemma 9.1) shows that such a marking m' on (C', ω') exists.

In particular this provides an alternative proof of the connectedness of the Hurwitz space of primitive branched coverings over $\mathbb{C}/\Lambda(p)$ of volume V (see [10] for the original proof).

Case 3: $\overline{\Lambda(p)}$ is neither discrete nor dense in \mathbb{C} . Then its closure is an infinite union of parallel real lines and we fall in the second case of Proposition 3.10. The conclusion is that in the closure \bar{L} we find all pairs (C', ω') of volume V whose periods belong to $\Lambda(p)$ and contain elements in every real line where $\Lambda(p)$ is dense.

Case 4: $\Lambda(p)$ is dense in \mathbb{C} . We fall in the third case of Proposition 3.10. In \bar{L} we find all pairs (C', ω') of volume V .

3.6. Remark on the transfer principle on strata. The generic stratum is the complement of an analytic subset, and therefore its intersection with each fiber of Per_g has the same number of connected components as the whole fiber. We can thus apply the Transfer principle to the generic stratum as well. On the other strata, the question seems to be much more delicate (see the discussion in section 1.4). At the other extreme, on the minimal stratum, the isoperiodic foliation has dimension zero, so we cannot expect to transfer dynamical properties of the symplectic group on the period domain to dynamical properties of the isoperiodic foliation. We even show here that

Proposition 3.11. *The intersection of a fiber of Per_g with the minimal stratum is either empty or infinite discrete.*

The connectedness of the intersection of the fibers of Per_g with the connected components of all other strata remains open.

Proof. The restriction of the map Per_g to the minimal stratum is a local diffeomorphism onto an open set in \mathcal{H}_g . Hence its fibers are discrete. Let $(C, m, \omega) \in \Omega\mathcal{S}_g$ be a point in the minimal stratum and $p = \text{Per}(C, m, \omega)$. Remark that the image of $\text{Aut}(C, \omega) \rightarrow \text{Sp}(H_1(C, \mathbb{Z}))$ is a finite subgroup G . Therefore the group

$$\text{Stab}(C, m, \omega) = \{M \in \text{Sp}(H_1(\Sigma_g, \mathbb{Z})) : (C, m \circ M, \omega) \sim (C, m, \omega)\}$$

is finite. If p has discrete image, then $\text{Stab}_{\text{Sp}(H_1(\Sigma_g, \mathbb{Z}))}(p)$ is infinite and contains G . Hence there is an infinite number of points in the orbit of (C, m, ω) under this group. All of them lie in $\text{Per}^{-1}(p)$, so the result follows in this case.

Next suppose p has non-discrete image. By Proposition 3.10 there exists an infinite sequence of $M_n \in \text{Sp}(H_1(\Sigma_g, \mathbb{Z}))$ such that $p_n = p \circ M_n$ is a sequence of pairwise distinct points converging to p in \mathcal{H}_g . Up to taking a subsequence, these points correspond to points (C_n, m_n, ω_n) in the minimal stratum via Per_g , that project to an infinite family of pairwise distinct points $(C_n, \omega_n) \in \Omega\mathcal{M}_g$. By construction, all points $(C_n, m_n \circ M_n^{-1}, \omega_n)$ belong to the intersection of $\text{Per}^{-1}(p)$ with the minimal stratum and are pairwise distinct in $\Omega\mathcal{S}_g$. \square

3.7. Monodromy of zeros along a fiber of $\text{Per}_{g,n}$ in the generic stratum.

Definition 3.12. Given a connected component K of the intersection of a fiber of $\text{Per}_{g,n}$ with a stratum of $\Omega\mathcal{S}_{g,n}$ having l distinct zeros, define the monodromy group of zeros associated to K as the conjugacy class of the image group G_K of the representation

$$\pi_1(K, (C_0, P_0, m_0, \omega_0)) \rightarrow S_l$$

that associates to each loop in K the permutation induced on the zero set z_1, \dots, z_l of ω_0 .

Our main interest will be the monodromy group of the zeros in the generic stratum. Recall that it is a Zariski open set in the fiber of the isoperiodic map and is therefore connected if and only if the fiber is. Turning around the divisors associated to other strata in the fiber already produces some monodromy of the zeros:

Lemma 3.13. *Let $g \geq 2$ and L be a connected component of a fiber of $\text{Per}_{g,n}$. Suppose it contains a point whose zero divisor is $d_1 p_1 + \dots + d_l p_l$. Then, the monodromy group of the zeros in the generic stratum L^{SZ} of L , contains all transpositions of pairs of elements that belong to the same part of a partition $Z_1 \sqcup \dots \sqcup Z_l$ of the set of $2g - 2$ elements with $|Z_i| = d_i$.*

Before we proceed to the proof we recall the interpretation of an element $(C, p_1, \dots, p_n, \omega)$ in $\Omega\mathcal{M}_g$ as a branched projective structure (see [15] for more details), or, more precisely, a branched translation structure. Indeed, around any point $q \in C$ we can define a local branched covering of degree $\text{ord}_q(\omega) + 1$ by $\phi_q(z) = \int_q^z \omega$. At the intersection of domains of two such branched coverings differ by a translation, and therefore define a branched atlas on the topological curve C with transitions in the set of translations (which are actually holomorphic). The branch points correspond to the zeros of the form. Reciprocally, if we are given an atlas $\{D_q, \phi_q\}$ on a compact topological surface of genus g of (topological) branched coverings $\phi_q : D_q \rightarrow \mathbb{C}$ with a finite number of branch points whose transition maps are translations, it defines a complex structure on the surface and the form defined locally by $\omega := d\phi_q$ is globally defined. In section 5.1 we will exploit this interpretation further. For the moment we will use it to deform a given point without changing the corresponding periods by changing a given atlas locally around a branched point.

Proof of Lemma 3.13: Let $(C, r_1, \dots, r_n, m, \omega)$ be the point satisfying $(\omega)_0 = d_1 p_1 + \dots + d_l p_l$. If all $d_i = 1$ there is nothing to prove. Otherwise, L^{SZ} is the complement of a normal crossing divisor in L (see section 3.2). Choose a local chart $\phi_i : D_i \rightarrow \mathbb{C}$ of degree $d_i + 1$ of the translation structure of ω around p_i . We can suppose that none of the zeros of ω is a marked point r_i and take discs D_i not containing any marked point.

Recall from the appendix in [15] that we can construct a germ of continuous map

$$(H(\phi_1) \times \dots \times H(\phi_{d_l}), (\phi_1, \dots, \phi_l)) \rightarrow (L, \omega)$$

where each $H(\phi_i)$ is a Hurwitz space of coverings of degree d_i over a disc $\phi_i(D_i)$ up to an equivalence in the boundary that allows to glue each element with the translation surface defined by ω on $C \setminus (\cup D_i)$ to obtain a branched translation structure on a closed marked surface of genus g . Since we can keep the marked points and a marking of the homology by avoiding the discs D_i , the gluing preserves all integrals over cycles of $H_1(\Sigma_g, q_1, \dots, q_n; \mathbb{Z})$.

By [15] [Lemma A7, p. 439], each $H(\phi_i)$ can be parametrized by the space of polynomials $P_a(z) = z^{d_i} + a_{d_i-1}z^{d_i-1} + \dots + a_0$ with $\sum a_j = 0$ having critical values in the unit disc. The point with all coordinates $a_j = 0$ corresponds to the initial point. Any choice on each Hurwitz space of a point that has d_i distinct critical values $\{v_1, \dots, v_{d_i}\}$ (therefore determining simple critical points on some set Z_i) has as image in L a point with simple zeros. On the other hand, the set of critical values determines the values of the a_i 's in the parameter space. Choose one such point $(C_0, \omega_0) \in L^{SZ}$. Since the germ of $\mathbb{D}^{d_i} \setminus \{(z_1, \dots, z_{d_i}) : \text{with } z_j = z_k \text{ for some } j \neq k\}$ at the origin is connected, there exists a path in it joining (v_1, \dots, v_{d_i}) to any point obtained by permuting the order of the coordinates. Such a path will determine a closed path in the parameter space whose image in L^{SZ} will permute the zeros of ω_0 in each part Z_i as desired.

□

Lemma 3.14. *Let $g \geq 3$ and $p \in \mathcal{H}_g$ such that $\text{Per}^{-1}(p)$ is connected. Then, the monodromy group of the zeros in the generic stratum of $\text{Per}^{-1}(p)$ is S_{2g-2} if $\deg(p) \geq 3$ and trivial if $\deg(p) = 2$.*

Proof. Assume first that p has infinite primitive degree, or equivalently non discrete image. In this case we claim there exists a form in $\text{Per}^{-1}(p)$ in the minimal stratum, namely having a unique zero of multiplicity $2g - 2$. From Lemma 3.13 we deduce that the whole group S_{2g-2} is in the image of the representation. To prove the claim, observe that up to the action of the group $\text{GL}^+(2, \mathbb{R})$, we can assume that the closure of the orbit of p under the action of the group $\text{Aut}(H_1(\Sigma_g, \mathbb{Z}))$ contains the set P of all periods $q : H_1(\Sigma_g, \mathbb{Z}) \rightarrow \mathbb{R} + i\mathbb{Z}$ of volume $\text{vol}(q) = \text{vol}(p)$ (see Proposition 3.10). Since the period map restricted to the minimal stratum is a local biholomorphism (see subsection 3.2), its image is an open set. It therefore contains points with discrete values in $\mathbb{Q} + i\mathbb{Q}$ that necessarily have finite degree. Up to the action of $\text{GL}(2, \mathbb{R})$ we can find one whose periods lie in P . Moreover, there is a neighborhood of P in $\text{Hom}(H_1(\Sigma_g, \mathbb{Z}), \mathbb{C})$ which consist of periods of forms in the minimal stratum. In particular, the orbit of p under $\text{Aut}(H_1(\Sigma_g, \mathbb{Z}))$ intersects this neighborhood, and thus p itself is the period of a form in the generic stratum. Hence the result follows in the case $\deg(p) = \infty$.

Assume now that $\deg(p) < \infty$. In this case, consider the map $R : \text{Per}^{-1}(p) \rightarrow H_{g,d}(E)$ taking values in the Hurwitz space $H_{g,d}(E)$ of equivalence classes of primitive branched coverings over the elliptic curve $E = \mathbb{C}/\text{Im}(p)$ of primitive degree $d = \deg(p)$, the equivalence being the pre- and post-composition by automorphisms. The map R sends a form

$(C, \omega, m) \in \text{Per}^{-1}(p)$ to the equivalence class of the covering from C to E given by integrating ω . Denoting by $U \subset \text{Per}^{-1}(p)^{SZ}$ the Zariski open subset consisting of forms with simple zeros whose integral between any pair of distinct zeros does not belong to the lattice $\text{Im}(p)$, and by $H_{g,d}(E)^{DC} \subset H_{g,d}(E)$ the Zariski open subset consisting of coverings with $2g - 2$ distinct critical values, the map R induces a Galois covering $U \rightarrow H_{g,d}(E)^{DC}$ with group $\text{Aut}(H_1(\Sigma_g, \mathbb{Z}))$. Consider the representation of permutations of the critical values

$$(19) \quad \pi_1(H_{g,d}(E)^{DC}) \rightarrow S_{2g-2}.$$

It is well-known that this representation is surjective, but by lack of references we reproduce a proof in the following paragraph. With this at hand, let $H \subset S_{2g-2}$ be the normal subgroup corresponding via (19) to the image of the Galois covering $\pi_1(U) \rightarrow \pi_1(H_{g,d}(E)^{DC})$. We claim that if $\deg(p) \geq 3$, and $g \geq 3$, then H contains a transposition, and hence $H = S_{2g-2}$. We will find a transposition by applying Lemma 3.13 to a particular form with a double zero. By the first item of Proposition 3.10, or equivalently Lemma 9.1, it suffices to prove the existence of primitive genus g and degree d branched covering over E , branched over a set $\{e_1, \dots, e_{2g-3}\}$ of $2g - 3$ elements with transitive (covering!) monodromy representation $\pi_1(E \setminus \{e_1, \dots, e_{2g-3}\}) \rightarrow S_d$ taking one of the peripherals to a cycle of order three (which corresponds to the local monodromy of the branched cover defined by integration around a double zero of a holomorphic form) and every other peripheral to a transposition (which corresponds to the local monodromy generator of the branched cover defined by integration around a simple zero of a holomorphic form). This is certainly possible if $d \geq 3$ and $2g - 3 \geq 2$, i.e. $g \geq 3$.

Let us now prove that the representation (19) is onto. Let $c : E' \rightarrow E$ be a non ramified cyclic $d : 1$ covering of E . We denote by $\tau \in \text{Aut}(E')$ a generator of the Galois group of c . Given $g - 1$ smooth curves s_1, \dots, s_{g-1} in E' , diffeomorphic to closed intervals, such that all the intervals $c(s_1), \dots, c(s_{g-1})$ are disjoint we can construct an associated element in $H_{g,d}(E)^{DC}$. Indeed, all curves in the family $s_1, \dots, s_{g-1}, \tau(s_1), \dots, \tau(s_{g-1})$ are disjoint, and we can slit E' along all the curves of this later family, and glue the right side of s_k to the left side of $\tau(s_k)$, and vice versa, for every $k = 1, \dots, g - 1$. We obtain a Riemann surface C of genus g , on which the map c induces a well-defined degree d branched covering $r : C \rightarrow E$, which is primitive. Observe that the critical values of r are the $2g - 2$ points lying at the extremities ι_k, e_k of the curves $c(s_k)$ for $k = 1, \dots, g - 1$.

The previous construction can be used to describe paths in $H_{g,d}(E)^{DC}$ by deforming the chosen smooth curves continuously. Take a choice of $g - 1$ smooth curves s_1, \dots, s_{g-1} in E' and consider the associated covering $r : C \rightarrow E$. Observe, for instance, that moving the curve s_k in E without touching the other curves, and returning back to the same curve s_k but with opposite orientation, we draw a closed loop in $H_{g,d}(E)^{DC}$ based at r , and the permutation on the critical values is the transposition exchanging ι_k and e_k . It remains to show that transpositions of critical values, involving endpoints of distinct segments are also possible. Now, since the elements of $H_{g,d}(E)^{DC}$ have $2g - 2$ critical values, all critical points of r are simple and there is a one-to-one correspondence between critical values and critical points of r . Each critical point corresponds to a zero of $\omega = dr$, so if for each pair of distinct k, l we manage to find a loop in U with base point ω whose associated monodromy of the zeros exchanges the zero associated to the endpoint ι_k of s_k with the zero associated to an endpoint e_l of s_l and fixes every other zero, we will be done. As before, we will invoke Lemma 3.13 which allows to deduce the property if we manage to find a path in $\text{Per}^{-1}(p)$ starting at ω

and finishing at a point in a stratum of forms with precisely $2g - 4$ simple zeros corresponding to the endpoints distinct from ι_k and e_l and a double zero. This path can be constructed in $H_{g,d}(E)^{DC}$ by deforming s_k in such a way that it does not touch any other s_h , apart at the very end of the movement where the only two points of distinct segments that coincide are ι_k and e_l . The set of permutations constructed in this way generate the whole symmetric group of the set $\{\iota_1, e_1, \dots, \iota_{g-1}, e_{g-1}\}$.

It remains to prove the last statement, namely that if p has primitive degree two, the monodromy of the zeros on $\text{Per}^{-1}(p)$ is trivial. Remark that all forms in that set have simple zeros. We can use done using the invariant of Arnold's introduced in the proof of Theorem 1.3. Consider the (locally flat) fiber bundle $\mathcal{E} \rightarrow \text{Per}^{-1}(p)$ by $\mathbb{Z}/2\mathbb{Z}$ -vector spaces, whose fiber over an element (C, m, ω) is the $\mathbb{Z}/2\mathbb{Z}$ vector space generated by the zeros of ω , the only relation being that the sum of all zeros is trivial. Since the number of zeros of ω is $2g - 2 \geq 4$, we need to prove that the monodromy of \mathcal{E} is trivial when the primitive degree of p is equal to two.

The fiber bundle \mathcal{E} embeds naturally in the locally flat $\mathbb{Z}/2\mathbb{Z}$ -vector bundle \mathcal{G} over $\text{Per}^{-1}(p)$ whose fiber over a point (C, m, ω) is the space $H_1(E \setminus \text{VC}(\pi), \mathbb{Z}/2\mathbb{Z})$; the embedding sends a zero of ω to the peripheral cycle turning around its π -image in E .

Fiberwise, the family of maps $\Pi \circ m$, where (C, m, ω) varies in $\text{Per}^{-1}(p)$ and Π is the map (12), induce a map from the constant bundle $H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z})$ over $\text{Per}^{-1}(p)$ to the bundle \mathcal{G} . The image of the constant subbundle $\text{Ker}(p[2])$ of $H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z})$ over $\text{Per}^{-1}(p)$ (recall that $p[2]$ is the reduction of p modulo two, see (17)) fall inside the subbundle \mathcal{E} of \mathcal{G} . Moreover, it consists fiberwise of the subspaces of formal sums of an even number of peripheral cycles in $H_1(E \setminus \text{VC}(\pi), \mathbb{Z}/2\mathbb{Z})$.

We infer that the monodromy of \mathcal{E} acts trivially on this subspace. Since a permutation of a set of at least three elements is determined by its action on the subsets of an even number of elements, we are done. \square

4. AUGMENTED TORELLI SPACE AND EXTENSION OF THE PERIOD MAP

In this section we review the topological and analytical properties of moduli spaces of curves and some of their coverings and (partial) bordifications. We will adopt the analytic point of view first developed by Abikoff in [1]. It is based on the Augmented Teichmüller space (see [3, 13, 11, 12, 26, 32, 36, 39, 48, 67, 68, 69, 71] for details and further references) and the bundles of stable forms over it (see the two recent papers [6, 7]). Most of the section will be devoted to describe the properties of the quotient of Augmented Teichmüller space by the Torelli group, the so-called Augmented Torelli space. We will also analyze the extension of the period map to the points lying in the closure of fibers of $\text{Per}_{g,n}$.

4.1. Marked stable curves.

Definition 4.1. A connected complex curve C with n marked distinct points $r_1, \dots, r_n \in C$ is said to be stable if its singularities are nodes and do not coincide with any of the marked points, and the closure C_i of each component of $C^* := C \setminus \text{Sing}(C)$, called a part of C , has a group of automorphisms that fix the marked points and the boundary points that is finite. The normalization of C is the smooth curve $\hat{C} = \sqcup C_i$. A stable curve C is said of compact type if every node separates C in two components. Otherwise C is said to be of non-compact type.

The arithmetic genus of a stable curve is $g = h^1(C, \mathcal{O})$. When C has δ nodes and its normalization has ν components of genera g_1, \dots, g_ν , the arithmetic genus satisfies (see [32] [p. 48])

$$g = \sum_{i=1}^{\nu} (g_i - 1) + \delta + 1$$

As in section 3.1, for $g, n \geq 0$ we fix a reference surface with marked points $\Sigma_{g,n} = (\Sigma_g, q_1, \dots, q_n)$: a closed connected oriented surface Σ_g of genus g with a set of n distinct ordered marked points $Q = (q_1, \dots, q_n) \in \Sigma_g$. When $n = 0$ we omit the subindex and write $\Sigma_g = \Sigma_{g,0}$.

Definition 4.2. A homotopical marking (or sometimes a collapse) of a connected genus g stable curve C with n ordered pairwise distinct marked points $R = (r_1, \dots, r_n) \in C^*$ is a continuous surjection $f : \Sigma_{g,n} \rightarrow (C, r_1, \dots, r_n)$ such that $f(q_i) = r_i$, the preimage of each node is a simple closed curve on $\Sigma_g \setminus Q$ and on each component of $\Sigma_g \setminus f^{-1}(N)$ where N is the set of nodes, the map f is a homeomorphism onto a part of C that preserves the orientation.

When C is non-singular a collapse is a homeomorphism and the definition coincides with the one given in subsection 3.1.

Definition 4.3. A homotopically marked stable curve with n marked points is a marked stable curve (C, r_1, \dots, r_n) together with a homotopical marking $f : (\Sigma_g, Q) \rightarrow (C, R)$. Two homotopically marked stable curves $f_i : (\Sigma_g, Q) \rightarrow (C_i, R_i)$ for $i = 1, 2$ are said to be equivalent if there exists a conformal isomorphism $g : C_1 \rightarrow C_2$ such that $g \circ f_1$ is homotopic to f_2 relative to Q . The class of a $\Sigma_{g,n}$ marked stable curve will be denoted by $[f : \Sigma_{g,n} \rightarrow (C, r_1, \dots, r_n)]$.

Remark 4.4. If $\Delta : \Sigma_{g,n} \rightarrow \Sigma_{g,n}$ is a Dehn twist around a curve in Σ_g that is collapsed by the marking $f_1 : (\Sigma_g, Q) \rightarrow (C, R)$ to a point, then (C, R) marked by $f_1 \circ \Delta$ is equivalent to the same curve marked by f_1 .

4.2. Augmented Teichmüller space and its stratification.

Definition 4.5. The augmented Teichmüller space $\overline{\mathcal{T}}_{g,n}$ is the set of all homotopically marked stable genus g curves with n marked points up to equivalence.

The Teichmüller space $\mathcal{T}_{g,n}$ is the subset of $\overline{\mathcal{T}}_{g,n}$ formed by curves without nodes. Its complement, denoted by $\partial\mathcal{T}_{g,n} = \overline{\mathcal{T}}_{g,n} \setminus \mathcal{T}_{g,n}$ is called the boundary.

Definition 4.6. A curve system $c = \sqcup c_i$ in $\Sigma_{g,n} = (\Sigma_g, q_1, \dots, q_n)$ is a disjoint collection of simple closed curves c_i on $\Sigma_g \setminus \{q_1, \dots, q_n\}$ none of which is isotopic to any other, to a point or to a cylinder in $\Sigma_g \setminus \{q_1, \dots, q_n\}$. To a curve system c we can associate the subset $B_c \subset \partial\mathcal{T}_g$ of the boundary consisting of homotopically marked stable curves topologically equivalent to a collapse $\Sigma_g \rightarrow \Sigma_g/c$ obtained by identifying each curve in c to a point.

Given a curve system c , for each component Σ^i of $\Sigma_g \setminus c$ we define (Σ_{g_i}, Q_i) to be the closed surface of genus g_i with a set of marked points Q_i obtained by collapsing each boundary component of Σ^i to a (marked) point and keeping the marked points of Σ_g lying on Σ^i in Q_i . As we will see in subsection 4.10 there is a natural identification

$$(20) \quad B_c \cong \Pi_i \mathcal{T}_{g_i, n_i}.$$

The boundary $\partial\mathcal{T}_{g,n}$ is the disjoint union of all boundary strata $\sqcup_c B_c$ where c varies in the set of nonempty curve systems. $\mathcal{T}_{g,n}$ corresponds to the empty curve system. Each stratum B_c has a topology and complex structure given by the bijection (20). All the curves that appear in a stratum have the same number of separating and non-separating nodes. When all the nodes are non-separating we say that the stratum is of compact type.

Given a simple closed curve $c' \subset \Sigma_{g,n}$ we denote by

$$D_{c'} = \bigsqcup_{c' \subset c} B_c$$

the union of all strata that collapse c' to a node.

Definition 4.7. Given $\mathcal{K} \subset \overline{\mathcal{T}}_{g,n}$, a stratum of \mathcal{K} is, by definition, the intersection $\mathcal{K} \cap B_c$ of \mathcal{K} with a stratum $B_c \subset \overline{\mathcal{T}}_{g,n}$. If the stratum lies in the boundary of $\mathcal{T}_{g,n}$ we call it a boundary stratum. The boundary strata of \mathcal{K} is the set $\partial\mathcal{K} := \mathcal{K} \cap \partial\mathcal{T}_{g,n}$.

4.3. Topology and stratification of $\overline{\mathcal{T}}_{g,n}$. For a detailed description of the topology we refer to [3, pp. 485-493] and references therein.

The restriction of the given topology to $\mathcal{T}_{g,n}$ produces the so-called conformal topology. Abikoff showed (in [1][Theorem 1]) that this topology is equivalent to the Teichmüller topology introduced in subsection 3.1.

The restriction of the topology to the boundary stratum B_c corresponding to a curve system c is equivalent to the product topology obtained from (20).

The topology is not locally compact around any boundary point. Indeed, if U is a neighbourhood of a point in B_c , the action of the Dehn twist $\Delta_a : \Sigma_g \rightarrow \Sigma_g$ around a simple closed curve $a \in c$ fixes all the points in U for which the marking collapses a to a point, but has infinite orbits at any other point in U . Therefore, there is no manifold structure in $\overline{\mathcal{T}}_{g,n}$ compatible with the given topology.

Definition 4.8. Given a curve system c the distinguished neighbourhood of the stratum B_c is the set

$$U_c = \bigsqcup_{c' \subset c} B_{c'}.$$

4.4. Local stratification and Deligne-Mumford compactification. The mapping class group of $\Sigma_{g,n}$, i.e. the group $\text{Mod}(\Sigma_{g,n})$ of isotopy classes of orientation preserving diffeomorphisms that fix each marked point, acts on $\overline{\mathcal{T}}_{g,n}$ by homeomorphisms that preserve the stratification and are holomorphic in restriction to any stratum. The action is defined by pre-composition on the marking. The quotient

$$\overline{\mathcal{M}}_{g,n} = \overline{\mathcal{T}}_{g,n} / \text{Mod}(\Sigma_{g,n})$$

is a compact topological space. It can be endowed with a complex orbifold structure.

Consider a curve system c and define Γ_c the abelian group generated by Dehn twists around the curves in c . Following [13], the quotient U_c / Γ_c is equivalent to a bounded domain in \mathbb{C}^{3g-3} . Under this equivalence, each stratum $B_{c'}$ associated to a simple closed curve $c' \subset c$ has image contained in a regular divisor $D_{c'}$. These divisors intersect normally and their intersections define the other different strata: the stratum associated to $c' \subset c$ is the intersection of all the divisors associated to the simple curves in c' . The complement of this divisor is the stratum $B_\emptyset = \mathcal{T}_{g,n}$ formed by smooth marked curves. The union of all natural maps $U_c / \Gamma_c \rightarrow \overline{\mathcal{M}}_{g,n}$

induce a system of (orbifold) charts with holomorphic transition maps on $\overline{\mathcal{M}}_{g,n}$. A more precise local description of the quotient map $U_c \rightarrow U_c/\Gamma_c$ in the neighbourhood of a point of B_c up to local homeomorphisms on source and target that can be found in [3] pp. 485-493]. We recall the model here because it will be useful to determine the local properties of the isoperiodic deformations of stable forms.

Consider the set $X = \frac{\mathbb{R}_+ \times \mathbb{R}}{0 \times \mathbb{R}}$ endowed with the (non-locally compact, first countable, Hausdorff) topology whose basis of open sets around $[0, 0]$ are the sets of the form $\{[\rho, \theta] : \rho < \varepsilon\}$ with $\varepsilon > 0$ and the usual round discs around any other point. Let $n \geq 2$ and $0 \leq l \leq n$ and the the normal crossing divisor

$$D_l = \{(z_1, \dots, z_n) \in \mathbb{C}^n : z_1 \cdots z_l = 0\}.$$

Consider the map with infinite ramification on D_l

$$(21) \quad \varphi_{l,n} : X^l \times \mathbb{C}^{n-l} \rightarrow \mathbb{C}^n \text{ defined by}$$

$(\rho_1, \theta_1, \dots, \rho_l, \theta_l, z_{l+1}, \dots, z_n) \mapsto (\rho_1 e^{2i\pi\theta_1}, \dots, \rho_l e^{2i\pi\theta_l}, z_{l+1}, \dots, z_n)$. It is a continuous map with respect to the product topologies on source and target. On the complement of the preimage of D_l the map $\varphi_{l,n}$ is a topological cover with free abelian covering group— that corresponds to the action of \mathbb{Z}^l on the θ variables by translations, componentwise. This implies that it is a connected complex manifold. Given a subset $I \subset \{1, \dots, l\}$ with k elements, the restriction of $\varphi_{l,n}$ to the set $\{\rho_j = 0 : j \in I\}$ produces map that is equivalent to some other $\varphi_{l',n'}$, so the set $\{\rho_j = 0 : j \in I\} \cap \{\rho_j \neq 0 : j \notin I\}$ is a connected complex manifold of codimension k accumulating the origin. They fit together to form a stratification of $X^l \times \mathbb{C}^{n-l}$. The action of \mathbb{Z}^l leaves *each connected component* of a stratum invariant.

Remark 4.9. Each connected component of a local stratum of positive codimension close to the origin is characterized by the set of local components of codimension one that accumulate to it. This property persists under quotients of $X^l \times \mathbb{C}^{n-l}$ by subgroups $H \subset \mathbb{Z}^l$.

Coming back to the local descriptions of the map $U_c \rightarrow U_c/\Gamma_c$, around a point of the curve system c with l elements, it is locally equivalent to $\varphi_{l,3g-3}$ at the origin.

Definition 4.10. A stratified space is a local abelian ramified cover of a normal crossing divisor if the stratification is locally homeomorphic to a quotient stratification of $X^l \times \mathbb{C}^{n-l}$ by some subgroup $H \subset \mathbb{Z}^l$ for some n and $l \leq n$ at the origin.

Definition 4.11. To any stratified space X that is a local abelian ramified cover of a normal crossing divisor on some fixed dimension $n \geq 2$ we can associate a boundary complex $\mathcal{C}(X)$ having a vertex for each connected component of the codimension one stratum and a simplex joining k vertices for each connected component of the stratum of codimension k contained in the closure of the corresponding components.

By the description of connected component of strata by classes of curve systems, we get an isomorphism of this dual boundary complex with the curve complex $\mathcal{C}(\overline{\mathcal{T}}_{g,n}) \simeq \mathcal{C}_{g,n}$ (see subsection 1.3) for the definition. The group $\text{Mod}(\Sigma_{g,n})$ acts on $\mathcal{T}_{g,n}$ (resp. $\mathcal{C}_{g,n}$) preserving strata (resp. the simplicial structure).

4.5. Augmented Torelli space: a ramified covering of $\overline{\mathcal{M}}_{g,n}$. The map induced in homology by an element of $\text{Mod}(\Sigma_{g,n})$ provides an exact sequence

$$0 \rightarrow \mathcal{J}_{g,n} \rightarrow \text{Mod}(\Sigma_{g,n}) \rightarrow \text{Aut}(H_1(\Sigma_g, q_1, \dots, q_n; \mathbb{Z}))$$

where $\mathcal{J}_{g,n}$ is the Torelli group of $\Sigma_{g,n}$ formed by isotopy classes of diffeomorphisms that act trivially on relative homology. The Augmented Torelli space is the topological quotient

$$\bar{\mathcal{S}}_{g,n} = \bar{\mathcal{T}}_{g,n} / \mathcal{J}_{g,n}.$$

Again, $\bar{\mathcal{S}}_{g,n}$ contains the Torelli space $\mathcal{S}_{g,n} = \mathcal{T}_{g,n} / \mathcal{J}_{g,n}$ as a proper set. Its boundary is the set $\partial\mathcal{S}_{g,n} = \bar{\mathcal{S}}_{g,n} \setminus \mathcal{S}_{g,n}$. Two strata B_c and $B_{c'}$ lie on the same class if and only if there exists an element in the Torelli group that sends the curve system c to the curve system c' . Therefore, the boundary inherits a stratification

Definition 4.12. The equivalence class under the action of the Torelli group of a curve system c in $\Sigma_{g,n}$ will be denoted by \bar{c} .

The boundary inherits a partition induced by the stratification

$$\partial\mathcal{S}_{g,n} = \bigsqcup_{\bar{c} \neq \emptyset} B_{\bar{c}}$$

where \bar{c} runs over all equivalence classes of non-empty curve systems c under the action of the Torelli group $\mathcal{J}_{g,n}$.

The complex structure in restriction to the stratum $B_{\bar{c}}$ induces a complex structure on $B_{\bar{c}}$. However, the action of the Dehn twist around a simple closed curve of non-trivial class in $H_1(\Sigma_{g,n})$ can act non-trivially on $H_1(\Sigma_{g,n})$. For instance, in the case where $n = 0$ and the simple closed curve is non-separating of class $a \in H_1(\Sigma_g) \setminus 0$ the action is on homology is the non-trivial isomorphism $\delta_a : H_1(\Sigma_g, \mathbb{Z}) \rightarrow H_1(\Sigma_g, \mathbb{Z})$ given by

$$\delta_a(b) = b + (a \cdot b)a.$$

The topology of $\bar{\mathcal{S}}_{g,n}$ is still non-locally compact around such points. Therefore, $\bar{\mathcal{S}}_{g,n}$ does not admit a compatible manifold structure.

Lemma 4.13. (Local structure around a stratum $B_{\bar{c}}$ and boundary complex of $\bar{\mathcal{S}}_{g,n}$) *The stratification of $\bar{\mathcal{T}}_{g,n}$ induces a stratification of $\bar{\mathcal{S}}_{g,n}$ that is a local abelian ramified covering of a normal crossing divisor. The dual boundary complex of $\bar{\mathcal{S}}_{g,n}$ is*

$$\mathcal{C}(\bar{\mathcal{S}}_{g,n}) \simeq \mathcal{C}_{g,n} / \mathcal{J}_{g,n}.$$

A neighbourhood of a stratum $B_{\bar{c}}$ in $\bar{\mathcal{S}}_{g,n}$ admits a (compatible complex) manifold structure if and only if $\Gamma_c \subset \mathcal{J}_{g,n}$, i.e. the Dehn twist around any simple closed curve of \bar{c} acts trivially on $H_1(\Sigma_{g,n})$.

Proof. The stratification of the open set $U_c \subset \bar{\mathcal{T}}_{g,n}$ is invariant under the action of the subgroup $\Gamma_c \cap \mathcal{J}_{g,n} \subset \Gamma_c$ and a local abelian ramified cover of a normal crossing divisor. Therefore, the quotient on $U_c / (\Gamma_c \cap \mathcal{J}_{g,n})$ has the same local property. On the other hand, as a consequence of the fact that any automorphism of a stable curve that acts trivially on homology is equivalent to some Dehn twist around the pinched curves, $U_c / (\Gamma_c \cap \mathcal{J}_{g,n}) \rightarrow U_c / \mathcal{J}_{g,n} \subset \bar{\mathcal{S}}_{g,n}$ is a local homeomorphism at every point of the stratum $B_{\bar{c}}$. This proves all statements about the local structure around a stratum.

A connected component of a stratum of $\bar{\mathcal{S}}_{g,n}$ corresponds to one orbit of connected components of a stratum of $\bar{\mathcal{T}}_{g,n}$ under the action of $\mathcal{J}_{g,n}$. The result follows from the isomorphism $\mathcal{C}(\bar{\mathcal{T}}_{g,n}) \simeq \mathcal{C}_{g,n}$.

□

Given a simple closed curve c_1 , there might be distinct strata in the boundary component D_{c_1} of $\mathcal{T}_{g,n}$ that are identified in the quotient. For instance, suppose c_2 is a non-separating curve determining the same homology class as c_1 but distinct homotopy classes in $\Sigma_{g,0}$. By [9][Section 1.3] there exists an element ϕ in the Torelli group $\mathcal{I}_{g,0}$ sending c_2 to c_1 . The two curve systems $c = c_1 \sqcup c_2$ and $c' = \phi(c) = c_1 \sqcup \phi(c_1)$ are equivalent and define two distinct strata contained in D_{c_1} provided $\phi(c_1)$ and c_2 define distinct isotopy classes in $\Sigma_{g,0}$. In terms of the dual boundary complex this means that there are edges that join the same vertex.

4.6. Extension of the Torelli map $\mathcal{S}_g \rightarrow \mathfrak{S}_g$. The holomorphic orbifold structure of $\overline{\mathcal{M}}_g$ can be pulled back to a complex manifold structure in the open dense subset $\overline{\mathcal{S}}_g^c$ of marked curves of compact type in $\overline{\mathcal{S}}_g$. Indeed, all the Dehn twists around the curves that are collapsed to the nodes lie already in the Torelli group and the ramification disappears.

The definition of the coordinate functions via (7) can be done word by word for stable curves of compact type. In this manner we define an extension of (8) to a map $\overline{\mathcal{S}}_g^c \rightarrow \mathfrak{S}_g$ that is still holomorphic. Its image is the so-called Schottky locus, an analytic set that has positive codimension as soon as $g \geq 4$. For $g \geq 3$ the extended Torelli map has fibers of positive dimension over boundary points.

Remark 4.14. In genus $g = 2$ the added points $\overline{\mathcal{S}}_2^c \setminus \mathcal{S}_2$ correspond to products of two marked elliptic curves. The extension of the Torelli map (8) in this case is a biholomorphism (see [54]).

4.7. Homological markings. Given $[f : \Sigma_{g,n} \rightarrow (C, r_1, \dots, r_n)] \in \overline{\mathcal{S}}_{g,n}$ we can associate a surjective homomorphism

$$f_* : H_1(\Sigma_g, q_1, \dots, q_n; \mathbb{Z}) \rightarrow H_1(C, r_1, \dots, r_n; \mathbb{Z}).$$

When $\ker f_* = 0$, any element $\phi \in \text{Mod}(\Sigma_{g,n})$ such that $(f \circ \phi)_* = f_*$ satisfies $\phi_* = \text{Id}$. Therefore f_* determines the Torelli class of the marking f . In particular this is the case when each node of C is separating. When there is at most one node that is non-separating we have a similar result:

Lemma 4.15. *Let $f : \Sigma_g \rightarrow C$ be a homotopical marking that pinches a simple closed, non-separating curve γ and some separating simple closed curves. Let $\phi \in \text{Mod}(\Sigma_{g,0})$ such that $(f \circ \phi)_* = f_* : H_1(\Sigma_g; \mathbb{Z}) \rightarrow H_1(C, \mathbb{Z})$. Then, up to Dehn twists along γ , $\phi_* = \text{id}$, that is to say, it belongs to the Torelli group of Σ_g .*

Proof. Set $a = [\gamma]$, then $\ker f_* = \mathbb{Z}a$. Fix $a = a_1, b_1, a_2, b_2, \dots, a_g, b_g$ a symplectic basis of $H_1(\Sigma_g)$. Since $f_* \circ \phi_* = f_*$ we have $f_*(\phi_* - \text{id}) = 0$. Therefore $\psi = \phi_* - \text{id}$ is a morphism from $H_1(\Sigma_g, \mathbb{Z})$ to $\ker f_* = \mathbb{Z}a$. So we have $\psi(b_i) = \lambda_i a_1, \psi(a_i) = \mu_i a_1$ for some $\lambda_i, \mu_i \in \mathbb{Z}$. From the fact that ϕ is symplectic, we deduce $\mu_i = 0$ for all i and $\lambda_i = 0$ for all $i \neq 1$. ($\delta_{i1} = a_i \cdot b_1 = \phi(a_i) \cdot \phi(b_1) = \delta_{i1} + \mu_i$ and $0 = b_i \cdot b_1 = \phi(b_i) \cdot \phi(b_1) = \lambda_i$). Up to Dehn twist along γ we may assume $\lambda_1 = 0$, so $\psi = 0$ and $\phi_* = \text{id}$, that is to say $\phi \in \mathcal{I}_{g,0}$ is a Torelli mapping class. \square

An equivalent statement fails in general for curves C with more than two non-separating nodes. Indeed, suppose f is a marking that collapses two elements a_1 and a_2 of a symplectic basis $a_1, b_1, a_2, b_2, a_3, b_3, \dots, a_g, b_g$ of $H_1(\Sigma_g)$. Define a morphism $\psi : H_1(\Sigma_g) \rightarrow \ker f_*$ by $\psi(b_1) = a_2$ and $\psi(b_2) = a_1$ and $\psi(a_i) = \psi(b_j) = 0$ for all $i \geq 1$ and for all $j > 2$. Then $\text{Id} + \psi \in \text{Sp}(2g, \mathbb{Z})$ and it is induced by a mapping class $\phi \in \text{Mod}(\Sigma_g)$. By construction

$f_* = f_* \circ \phi_*$ but ϕ is not in the Torelli group, nor is a product of twists along loops whose class belong to $\ker f_* = \mathbb{Z}a_1 \oplus \mathbb{Z}a_2$ (because $\delta_{na_1+ma_2}(b_1) = b_1 + na_1$ so all such twists leave $\langle a_1, b_1 \rangle$ invariant, and ϕ does not).

Definition 4.16. A homological marking of a stable curve with n marked points (C, r_1, \dots, r_n) is a surjective homomorphism

$$m : H_1(\Sigma_g, q_1, \dots, q_n; \mathbb{Z}) \rightarrow H_1(C, r_1, \dots, r_n; \mathbb{Z}).$$

The data (C, r_1, \dots, r_n, m) will be referred to as a homologically marked curve.

4.8. Parametrization of the vertices of the dual boundary complex $\mathcal{C}_g/\mathcal{I}_g$ of $\bar{\mathcal{S}}_g$. With the use of homological markings we will provide a homological characterization of the Torelli classes of simple closed curves on Σ_g . This will allow us to give a special parametrization of the vertices of the dual boundary complex of $\bar{\mathcal{S}}_g$.

First we recall some concepts of homology theory. The intersection of two elements $a, b \in H_1(\Sigma_g, \mathbb{Z})$ is denoted by $a \cdot b$ and the associated intersection form defines an integral unimodular symplectic structure on $H_1(\Sigma_g) := H_1(\Sigma_g, \mathbb{Z})$. Two submodules of a unimodular symplectic module M are said to be orthogonal if the intersection of any element of one of them with an element of the other is zero. Given a submodule $N \subset M$ we denote by $N^\perp \subset M$ the orthogonal submodule of N in M , that is, the set of elements in M that have zero intersection with all elements of N .

Definition 4.17. A submodule N of a unimodular symplectic module M is said to be symplectic if the symplectic form of M restricted to N is still unimodular. A splitting $M = N_1 \oplus \dots \oplus N_k$ into pairwise orthogonal submodules is said to be symplectic if every N_i is a symplectic submodule.

Definition 4.18. A submodule N of a \mathbb{Z} -module M is primitive if whenever $zm \in M$ for some $m \in M$ and $z \in \mathbb{Z}$ then also $m \in N$.

Let c be a simple closed curve in Σ_g . If it is non-separating, it determines a primitive class $[c] \in H_1(\Sigma_g)$. If it is separating, the induced homology class is trivial and carries no interesting information. However, c splits Σ_g in two parts, and these induce a symplectic splitting of $H_1(\Sigma_g)$ that characterizes the Torelli class of the curve in the following sense:

Proposition 4.19 ([25], Proposition 6.14). *Let c, c' be two isotopy classes of simple closed curves in Σ_g . If c and c' are separating, then they are equivalent by some element of the Torelli group $\mathcal{I}_{g,0}$ if and only if the associated symplectic splittings coincide up to changing the order of the factors. If c and c' are non-separating, then they are equivalent by some element of the Torelli group $\mathcal{I}_{g,0}$ if and only if, up to sign, they determine the same homology class in $H_1(\Sigma_g)$.*

We can use this information to parametrize vertices of the dual boundary complex $\mathcal{C}_g/\mathcal{I}_g$ of the augmented Torelli space $\bar{\mathcal{S}}_g$.

Corollary 4.20. *The set of unordered pairs $\{V, V^\perp\}$ of non-trivial symplectic submodules of $H_1(\Sigma_g)$ such that $V \oplus V^\perp = H_1(\Sigma_g)$ is in 1-1 correspondence with Torelli classes of separating essential simple closed curves in Σ_g . The set of non-trivial primitive submodules $\mathbb{Z}[c] \subset H_1(\Sigma_g)$ is in 1-1 correspondence with the set of Torelli classes of non-separating simple closed curves.*

4.9. Identifying some simplexes of $\mathcal{C}_g/\mathcal{J}_g$. A separating node on a homologically marked stable curve (C, m) induces a non-trivial symplectic splitting $V \oplus V^\perp$ of $H_1(\Sigma_g)$. A non-separating node induces a cyclic non-trivial primitive submodule $\mathbb{Z}a$ of $\ker m \subset H_1(\Sigma_g)$.

Definition 4.21. Two marked nodal curves (C_i, m_i) for $i = 1, 2$ of genus g are said to share a separating node if for each $i = 1, 2$ there exists a separating node $r_i \in C_i$ that induces the same splitting $V \oplus V^\perp$ of $H_1(\Sigma_g)$ up to the order of the factors. Equivalently, we say that they share a non-separating node if there exist non-separating nodes $r_i \in C_i$ whose associated cyclic primitive submodules of $H_1(\Sigma_g)$ coincide.

Two marked stable curves that share a node define points that lie in the closure of the same boundary stratum of codimension one. In terms of the complex $\mathcal{C}_g/\mathcal{J}_g$ they represent simplexes that intersect at one vertex.

This characterization allows to identify certain simplexes of $\mathcal{C}_g/\mathcal{J}_g$ by looking at the homological information of the vertices. For instance, given a splitting $V_1 \oplus \dots \oplus V_{k+1}$ of $H_1(\Sigma_g)$ into symplectic submodules of rank at least two, the vertices associated to $\{V_i, V_i^\perp\}$ $i = 1 \dots k$ belong to a simplex of $\mathcal{C}_g/\mathcal{J}_g$. It suffices to construct a marked stable curve with k separating nodes inducing the given splitting. If in each V_i we are given a primitive class of $H_1(\Sigma_g)$ we can construct a stable curve with $2k$ nodes, that defines a simplex joining the corresponding $2k$ vertices, etc. These simplexes will be very useful to analyze the topology of the boundary stratification of $\bar{\mathcal{S}}_g$.

4.10. Attaching and forgetful maps on augmented Torelli spaces. It is well known that attaching maps, i.e. identification of distinct marked points, define holomorphic maps at the level of moduli spaces. The same is true for forgetful maps, i.e. forgetting part of the marked points followed by stabilization. In this subsection we will see how these maps can be defined at the level of augmented Teichmüller and Torelli spaces.

Let $\phi : \Sigma_{g,n} \rightarrow \Sigma_{g,n}/c$ be the collapse of a simple closed curve c in $\Sigma_{g,n} \setminus \{q_1, \dots, q_n\}$. We will define an attaching map associated to ϕ .

If c is a separating curve, the surface $\Sigma_{g,n} \setminus c$ has two components, each with a boundary component. Collapsing each boundary component to a new marked point produces two surfaces with marked points Σ_{g_1, n_1} and Σ_{g_2, n_2} where $g_1 + g_2 = g$, $n_1 + n_2 - 2 = n$ and the new points are named q_{n_1} and q_{n_2} respectively. The attaching map

$$A_\phi : \bar{\mathcal{T}}_{g_1, n_1} \times \bar{\mathcal{T}}_{g_2, n_2} \rightarrow \bar{\mathcal{T}}_{g, n}$$

is defined by the isotopy class of the composition $(f_1 \vee f_2) \circ \phi$ where $f_i : \Sigma_{g_i, n_i} \rightarrow (C_i, r_1^1, \dots, r_{n_i}^1)$ is a representative of its class in $\bar{\mathcal{T}}_{g_i, n_i}$ and $f_1 \vee f_2$ denotes the map $\Sigma_{g,n}/c \rightarrow (C_1 \vee C_2, q_1, \dots, q_n)$ by collapsing c to the new node $q_{n_1} = q_{n_2}$ of $C_1 \vee C_2$ and applying the corresponding f_i in each part of the complement of the node.

Remark 4.22. The isomorphism [\(20\)](#) is realized by attaching maps

Lemma 4.23. *The map A_ϕ induces a well defined continuous attaching map*

$$\bar{\mathcal{S}}_{g_1, n_1} \times \bar{\mathcal{S}}_{g_2, n_2} \rightarrow \bar{\mathcal{S}}_{g, n}.$$

Proof. Remark that ϕ allows to define a map

$$\rho_\phi : \mathcal{J}_{g_1, n_1} \times \mathcal{J}_{g_2, n_2} \rightarrow \mathcal{J}_{g, n}$$

between Torelli groups. Indeed, every pair $(\psi_1, \psi_2) \in \mathcal{J}_{g_1, n_1} \times \mathcal{J}_{g_2, n_2}$ defines an automorphism ψ_c of $\Sigma_{g, n}/c$ that acts trivially on its relative homology (with n marked points). We need to lift the element to an element $\psi \in \mathcal{J}_{g, n}$. The lift via ϕ is a well defined map $\Sigma_{g, n} \setminus c \rightarrow \Sigma_{g, n} \setminus c$. Up to changing ψ'_i s in the neighbourhood of the points q_{n_1} and q_{n_2} homotopically, we can guarantee that there exists an extension $\psi = \rho(\psi_1, \psi_2)$ of the lift to $\Sigma_{g, n}$ that fixes c pointwise.

By construction the map A_ϕ is equivariant with respect to ρ and therefore induces the desired map $\bar{\mathcal{S}}_{g_1, n_1} \times \bar{\mathcal{S}}_{g_2, n_2} \rightarrow \bar{\mathcal{S}}_{g, n}$. \square

If c is non-separating, then $\Sigma_{g, n} \setminus c$ is connected and has two boundary curves. Collapsing each boundary curve to a new marked point produces a surface of genus $g - 1$ with $n + 2$ marked points that we denote $\Sigma_{g-1, n+2}$. The new points will be denoted q_{n+1}, q_{n+2} . Remark that $\Sigma_{g, n}/c$ can be identified with $\Sigma_{g-1, n+2}/q_{n+1} \sim q_{n+2}$. The attaching map is

$$A_\phi : \bar{\mathcal{T}}_{g-1, n+2} \rightarrow \bar{\mathcal{T}}_{g, n}$$

defined by $(\vee f) \circ \phi$ where $\vee f : \Sigma_{g, n}/c \rightarrow (\vee C, r_1, \dots, r_n)$ is the map induced by a marking $f : \Sigma_{g-1, n+2} \rightarrow (C, r_1, \dots, r_{n+1}, r_{n+2})$ on the stable curve $\vee C$ (that has an extra non-separating node, compared to C) obtained by identifying r_{n+1} and r_{n+2} in C .

Lemma 4.24. *The map A_ϕ induces a continuous attaching map*

$$\bar{\mathcal{S}}_{g-1, n+2} \rightarrow \bar{\mathcal{S}}_{g, n}.$$

Proof. As before ϕ induces a homomorphism between the relevant Torelli groups

$$\rho_\phi : \mathcal{J}_{g-1, n+2} \rightarrow \mathcal{J}_{g, n}$$

and A_ϕ is ρ_ϕ equivariant by construction.

To construct ρ_ϕ , suppose $\psi : \Sigma_{g-1, n+2} \rightarrow \Sigma_{g-1, n+2}$ represents an element in $\mathcal{J}_{g-1, n+2}$. Then ψ induces an automorphism of $\Sigma_{g-1, n+2}/q_{n+1} \sim q_{n+2}$. The equivalence between this space and $\Sigma_{g, n}/c$ provides an automorphism of $\Sigma_{g, n}/c$ that can be lifted to $\Sigma_{g, n} \setminus c$ via ϕ . Up to changing the initial ψ homotopically in a small neighbourhood of the points q_{n+1} and q_{n+2} we can suppose that the lift extends to an automorphism $\rho_\phi(\psi) : \Sigma_{g, n} \rightarrow \Sigma_{g, n}$ that fixes c pointwise. The action of $\rho(\psi)$ in $H_1(\Sigma_g, \{q_1, \dots, q_n\}; \mathbb{Z})$ is trivial by construction and therefore $\rho_\phi(\psi) \in \mathcal{J}_{g, n}$. \square

As for forgetful maps, given an inclusion $\phi : \Sigma_{g, n} \rightarrow \Sigma_{g, n+1}$ of the marked points, we can define its associated forgetful map

$$(22) \quad \bar{\mathcal{T}}_{g, n+1} \rightarrow \bar{\mathcal{T}}_{g, n}$$

sending the class of $f : \Sigma_{g, n+1} \rightarrow (C, r_1, \dots, r_{n+1})$ to the stabilization of $f \circ \phi$, i.e. if the component of C that contains the forgotten point has an infinite group of automorphisms after deleting the point, we collapse the component of source and target to a point. Again we have

Lemma 4.25. *The map ϕ induces a continuous forgetful map $\bar{\mathcal{S}}_{g, n+1} \rightarrow \bar{\mathcal{S}}_{g, n}$.*

Proof. In this case, ϕ induces an inclusion $\rho_\phi : \mathcal{J}_{g, n+1} \rightarrow \mathcal{J}_{g, n}$ since any diffeomorphism that fixes $n + 1$ points fixes any of the n points that correspond to the image by ϕ of the n marked points of $\Sigma_{g, n}$. The map (22) is ρ_ϕ -equivariant by construction. \square

4.11. A useful bordification $\mathcal{U}_{g,2}$ of $\mathcal{S}_{g,2}$ in $\overline{\mathcal{S}}_{g,2}$. In this subsection we introduce a smooth partial bordification of $\mathcal{S}_{g,2}$ that will be useful in Section [7](#). We also describe its boundary complex.

Consider the composition of two forgetful maps

$$(23) \quad \text{For}_{g,2} : \overline{\mathcal{S}}_{g,2} \rightarrow \overline{\mathcal{S}}_{g,0} = \overline{\mathcal{S}}_g$$

and the open set

$$\mathcal{U}_{g,2} = \text{For}_{g,2}^{-1}(\mathcal{S}_g)$$

of marked curves whose stabilization after forgetting both marked points is a smooth curve. The restriction of $\text{For}_{g,2}$ to $\mathcal{U}_{g,2}$ will be denoted by For .

Let $\mathfrak{C}(\mathcal{S}_g) \rightarrow \mathcal{S}_g$ denote the universal curve bundle over \mathcal{S}_g and $\mathfrak{C}^2(\mathcal{S}_g) \rightarrow \mathcal{S}_g$ denote the pull back of the square of the universal curve bundle $\mathfrak{C}(\mathcal{S}_g) \times \mathfrak{C}(\mathcal{S}_g) \rightarrow \mathcal{S}_g \times \mathcal{S}_g$ by the diagonal map $\mathcal{S}_g \rightarrow \mathcal{S}_g \times \mathcal{S}_g$.

Proposition 4.26. *There exists a covering map $R : \mathcal{U}_{g,2} \rightarrow \mathfrak{C}^2(\mathcal{S}_g)$ such that*

$$(24) \quad \begin{array}{ccc} \mathcal{U}_{g,2} & \xrightarrow{R} & \mathfrak{C}^2(\mathcal{S}_g) \\ & \searrow \text{For} & \downarrow \\ & & \mathcal{S}_g \end{array}$$

commutes. For each $(C, m) \in \mathcal{S}_g$, $R|_{\text{For}^{-1}(C, m)} : \text{For}^{-1}(C, m) \rightarrow C \times C$ is an $H_1(\Sigma_g)$ -cover with monodromy given by

$$(25) \quad (\alpha_1, \alpha_2) \cdot \gamma = \gamma + \alpha_2 - \alpha_1, \quad \text{for } (\alpha_1, \alpha_2) \in H_1(\Sigma_g)^2 \simeq H_1(C \times C), \gamma \in H_1(\Sigma_g).$$

Moreover, the boundary strata in the complex manifold $\mathcal{U}_{g,2}$ form a smooth divisor whose components are in one to one correspondence with the set

$$\mathfrak{b} = \{\beta \in H_1(\Sigma_g, q_1, q_2; \mathbb{Z}) : \partial\beta = q_2 - q_1\}.$$

Proof of Proposition [4.26](#): We split the proof in several parts:

Action of $\text{Mod}_{g,2}$ on relative homology.

Let $\text{Aut}(H_1(\Sigma_g, \{q_1, q_2\}, \mathbb{Z}))$ be the subgroup of linear automorphisms of $H_1(\Sigma_g, \{q_1, q_2\}, \mathbb{Z})$ that preserve the exact sequence

$$(26) \quad 0 \rightarrow H_1(\Sigma_g, \mathbb{Z}) \rightarrow H_1(\Sigma_g, \{q_1, q_2\}, \mathbb{Z}) \xrightarrow{\partial} \mathbb{Z}(q_2 - q_1) \rightarrow 0$$

where ∂ is the boundary operator. The group $\text{Aut}(H_1(\Sigma_g, \{q_1, q_2\}, \mathbb{Z}))$ splits into an exact sequence

$$0 \rightarrow G \rightarrow \text{Aut}(H_1(\Sigma_g, \{q_1, q_2\}, \mathbb{Z})) \rightarrow \text{Aut}(H_1(\Sigma_g, \mathbb{Z})) \rightarrow 0$$

where $\text{Aut}(H_1(\Sigma_g, \mathbb{Z}))$ is the group of linear automorphisms of $H_1(\Sigma_g, \mathbb{Z})$ that preserve the intersection form, and where G is the abelian unipotent subgroup of $\text{Aut}(H_1(\Sigma_g, \{q_1, q_2\}, \mathbb{Z}))$ formed by the transformations

$$G_\alpha(\gamma) := \gamma + n(\gamma)\alpha$$

where $\alpha \in H_1(\Sigma_g, \mathbb{Z})$, and where we write the boundary operator as $\partial = n(q_2 - q_1)$. Notice that G naturally identifies with the absolute homology group $H_1(\Sigma_g, \mathbb{Z})$.

Since the action of $\text{Mod}_{g,2}$ on $H_1(\Sigma_g, \{p, q\}, \mathbb{Z})$ preserves [\(26\)](#), it gives rise to morphism

$$(27) \quad \text{Mod}_{g,2} \rightarrow \text{Aut}(H_1(\Sigma_g, \{q_1, q_2\}, \mathbb{Z})).$$

Action of the braid group.

Consider the exact sequence

$$(28) \quad 0 \rightarrow B(\Sigma_{g,2}) \rightarrow \text{Mod}(\Sigma_{g,2}) \rightarrow \text{Mod}(\Sigma_g) \rightarrow 0$$

where $B(\Sigma_{g,2})$ is the braid group with two strings on Σ_g , and where the right arrow is given by the forgetful map that forgets the fact that the mapping class fixes the two marked points. By Birman's theorem, the braid group is isomorphic to the fundamental group of the square of Σ_g deprived of its diagonal Δ ²

Its action on relative homology group (obtained by restriction of the action (27)) takes values in the abelian unipotent group G , hence it descends to an action defined on the abelianization of $B_{g,2}$, which is isomorphic to $H_1(\Sigma_g^2 \setminus \Delta, \mathbb{Z}) \simeq H_1(\Sigma_g^2, \mathbb{Z}) \simeq H_1(\Sigma_g)^2$; one checks that under these identifications it is given by the morphism

$$(29) \quad (\alpha_1, \alpha_2) \in H_1(\Sigma_g)^2 \mapsto G_{\alpha_2 - \alpha_1} \in \text{Aut}(H_1(\Sigma_g, \{q_1, q_2\}, \mathbb{Z})).$$

A first G -covering of $\mathcal{S}_{g,0}$.

The group G acts on $\mathcal{S}_{g,2}$ by precomposition of the marking, and the action is free and proper (recall that the action of the group $\text{Mod}_{g,2}/\mathcal{J}_{g,2} \simeq \text{Aut}(H_1(\Sigma_g, \{q_1, q_2\}, \mathbb{Z}))$ on $\mathcal{S}_{g,2}$ is free and discontinuous, see subsection 3.1). Denote by R' the covering $\mathcal{S}_{g,2} \rightarrow \mathcal{S}_{g,2}/G$. The fundamental group of $\mathcal{S}_{g,2}$ is isomorphic to $\mathcal{J}_{g,2}$, the one of $G \backslash \mathcal{S}_{g,2}$ is the group generated by the Torelli group $\mathcal{J}_{g,2}$ and the braid group $B(\Sigma_{g,2})$, and the monodromy of R' is given by the quotient map

$$< B(\Sigma_{g,2}), \mathcal{J}_{g,2} > \rightarrow < B(\Sigma_{g,2}), \mathcal{J}_{g,2} > / \mathcal{J}_{g,2} \simeq G \subset \text{Aut}(H_1(\Sigma_g, \{q_1, q_2\}, \mathbb{Z})) \simeq \text{Mod}_{g,2}/\mathcal{J}_{g,2}$$

The G -action preserves the restricted forgetful map $\text{For}_{|\mathcal{S}_{g,2}} : \mathcal{S}_{g,2} \rightarrow \mathcal{S}_g$, so there is a map $\widetilde{\text{For}} : \mathcal{S}_{g,2}/G \rightarrow \mathcal{S}_g$, we have $\text{For}_{|\mathcal{S}_{g,2}} = \widetilde{\text{For}} \circ R'$. The map $\widetilde{\text{For}}$ is a holomorphic fibration, whose fibers $\widetilde{\text{For}}^{-1}(C, m)$ are biholomorphic to the square of C deprived of its diagonal, namely $\widetilde{\text{For}}^{-1}(C, m) \simeq C \times C \setminus \Delta$ where $\Delta := \{(x, x) \mid x \in C\}$. The fibration at the level of the fundamental group gives the exact sequence

$$0 \rightarrow B(\Sigma_{g,2}) \rightarrow < B(\Sigma_{g,2}), \mathcal{J}_{g,2} > \rightarrow \mathcal{J}_g \rightarrow 0.$$

Hence the fibers of $\widetilde{\text{For}}$ are homologically marked by $H_1(\Sigma_g, \mathbb{Z})^2$ and the monodromy of the G -covering R' in restriction to a fiber of $\widetilde{\text{For}}$ is given by (29).

Construction of R as a bordification of the covering R' .

We now bordify the previous covering R' over $\mathcal{S}_{g,2}/G$. The idea is to add in each fiber $\widetilde{\text{For}}^{-1}(C, m) \simeq C \times C \setminus \Delta$ the diagonal Δ .

As before, the action of $G \simeq H_1(\Sigma_g, \mathbb{Z})$ on $\overline{\mathcal{S}}_{g,2}$ preserves each fiber of the forgetful map $\text{For}_{g,2} : \overline{\mathcal{S}}_{g,2} \rightarrow \overline{\mathcal{S}}_g$, hence it preserves the open subset $\mathcal{U}_{g,2} := \text{For}_{g,2}^{-1}(\mathcal{S}_g) \subset \overline{\mathcal{S}}_{g,2}$. The action is free on this subset. Indeed, recall that the stabilizer of a point $(C, r_1, r_2, m) \in \overline{\mathcal{S}}_{g,2}$ in

²Usually what is referred to as Birman's exact sequence is for a surface punctured at only one point, but the idea of the proof is the same in the case of two points: given an element of the braid group $[\varphi] \in B(\Sigma_{g,2})$, represented by an isotopy class of diffeomorphism $\varphi \in \text{Diff}(\Sigma_g, q_1, q_2)$, there exists an isotopy $(\varphi_t)_{t \in [0,1]}$ in $\text{Diff}(\Sigma_g)$ such that $\varphi_0 = \varphi$ and $\varphi_1 = \text{Id}$. The loop $(\varphi_t(q_1), \varphi_t(q_2)) \in \Sigma_g^2 \setminus \Delta$ defines an element $\gamma \in \pi_1(\Sigma_g^2 \setminus \Delta, (q_1, q_2))$ that only depends on $[\varphi]$ and the map $[\varphi] \mapsto \gamma$ induces an isomorphism between $B(\Sigma_{g,2})$ and $\pi_1(\Sigma_g^2 \setminus \Delta, (q_1, q_2))$.

$\text{Aut}(H_1(\Sigma_g, \{q_1, q_2\}, \mathbb{Z}))$ is generated by the action on relative homology of the Dehn twists along the (Torelli classes of) pinched curves of m . In the case of a point $(C, r_1, r_2, m) \in \mathcal{U}_{g,2}$, the only non trivial (Torelli class of) pinched curves of m are the ones that separate the surface $\Sigma_{g,2}$ into two domains, the first of genus zero containing the two marked points q_1, q_2 , the second being of genus g with a disc removed (a representation of the surface of genus two with the pinched curve is depicted in Figure 2). The Dehn twist along such a curve acts trivially on the relative homology group. On the one hand we deduce from Lemma 4.13 that $\mathcal{U}_{g,2}$ is a manifold and on the other that the action of $H_1(\Sigma_g, \mathbb{Z})$ on $\mathcal{U}_{g,2}$ is free.

We claim that the action of G on $\mathcal{U}_{g,2}$ is moreover proper and discontinuous. Since $\mathcal{U}_{g,2}$ is a manifold, and hence is locally compact, it suffices to prove that the action is discontinuous, namely that any point has a neighborhood which is disjoint from its images by G . We already know this is so on the open stratum $\mathcal{S}_{g,2}$ of $\mathcal{U}_{g,2}$ since there the action of the whole Torelli group is. So it remains to prove that the action is proper around any point of the boundary of $\mathcal{U}_{g,2}$. For any point $[(C, q_1, q_2, m)] \in \partial\mathcal{U}_{g,2}$ denote by $\gamma \subset \Sigma_{g,2}$ a loop which is mapped to the node of C by some marking of C in the equivalence class of m . There is a neighborhood $\mathcal{W} \subset \mathcal{U}_{g,2}$ defined by the property that for every $[(C', q'_1, q'_2, m')] \in \mathcal{W}$ the image of the loop γ in C' by some marking in the equivalence class of m' is of minimal length and this is the unique loop having this property. In particular, if an element of G maps an element of \mathcal{W} to an element belonging to \mathcal{W} , then some of its lift to the mapping class group of $\Sigma_{g,2}$ needs to fix γ (by uniqueness) and this can happen only if the original element of G is the identity. Hence the claim follows.

We denote the quotient map by $R : \mathcal{U}_{g,2} \rightarrow \mathcal{V} := \mathcal{U}_{g,2}/H_1(\Sigma_g, \mathbb{Z})$, which is a $H_1(\Sigma_g, \mathbb{Z})$ -covering, the forgetful map $\text{For}_{g,2}$ transits via a map

$$\overline{\text{For}} : \mathcal{V} \rightarrow \mathcal{S}_g$$

satisfying $\text{For} = \overline{\text{For}} \circ R$; it is a holomorphic fibration isomorphic to the square universal bundle $\mathfrak{C}^2(\mathcal{S}_g) \rightarrow \mathcal{S}_g$ by using Riemann's extension theorem applied on the added boundary points. Notice that for a given $(C, m) \in \mathcal{S}_g$ the fiber $\overline{\text{For}}^{-1}(C, m)$ is biholomorphic to the complex surface $C \times C$ and inherits a natural identification of the homology $m \times m : H_1(\Sigma_g) \times H_1(\Sigma_g) \rightarrow H_1(C) \times H_1(C)$. The covering

$$R|_{\text{For}^{-1}(C, m)} : \text{For}^{-1}(C, m) \rightarrow \overline{\text{For}}^{-1}(C, m) \simeq C \times C$$

is a $H_1(\Sigma_g)$ -covering of monodromy given by (25).

Boundary complex of $\mathcal{U}_{g,2}$.

Each boundary stratum $B_{\bar{c}}$ of $\mathcal{U}_{g,2}$ is characterized by a Torelli class of a simple closed curve c in $\Sigma_{g,2}$ that bounds a disc containing the marked points. The following Lemma proves that every such stratum can be characterized by a homology class in \mathfrak{b} .

Lemma 4.27. *Let c be an isotopy class of separating simple closed curve in $\Sigma_{g,2}$ that splits Σ_g into a disc containing both marked points and a genus g component without marked points. Then, c defines a unique class $\beta_c \in H_1(\Sigma_g, q_1, q_2; \mathbb{Z})$ represented by a cycle in the disc such that $\partial\beta_c = q_2 - q_1$. Two such curves c, c' are equivalent under the Torelli group $\mathcal{I}_{g,2}$ if and only if $\beta_c = \beta_{c'}$.*

Proof. If c and c' are equivalent by an element of the Torelli group it is obvious that $\beta_{c'}$ is the image of β_c and since the action is trivial in homology we have $\beta_c = \beta_{c'}$.

Suppose $\beta_c = \beta_{c'}$. Let D and D' be the open discs bounded by representatives of c and c' respectively. Find an orientation preserving diffeomorphism $\phi : \Sigma_g \rightarrow \Sigma_g$ such that $\phi(c) = c'$, $\phi(D) = D'$ and $\phi_1(q_i) = q_i$. By definition the map ϕ_* induced by ϕ in $H_1(\Sigma_g, q_1, q_2, \mathbb{C})$ satisfies $\phi_*(\beta_c) = \beta_{c'}$. We will show that up to pre-composition of ϕ with an diffeomorphisms ψ that fixes the closure of D pointwise, we can suppose that $(\phi \circ \psi)_*$ fixes every class in the homology group. Indeed, the restriction of ϕ_* to the image of the natural inclusion $0 \rightarrow H_1(\Sigma_g, \mathbb{Z}) \rightarrow H_1(\Sigma_g, q_1, q_2, \mathbb{Z})$ defines an element in the group $\text{Aut}(H_1(\Sigma_g, \mathbb{Z}))$. On the other hand, the action of the mapping class group of orientation preserving isotopy classes of diffeomorphisms of $\Sigma_g \setminus D$ that fix the boundary pointwise, provides a map $\text{Mod}(\Sigma_g \setminus D) \rightarrow \text{Aut}(H_1(\Sigma_g, \mathbb{Z}))$ that is surjective (see [25]p.169 and Section 6.3.2 for details). Hence, there exists a $\psi : \Sigma_g \rightarrow \Sigma_g$ that is the identity on D , such that $\psi_* = (\phi_*|_{H_1(\Sigma_g)})^{-1}$ and $\psi_*(\beta_c) = \beta_c$. By construction $(\phi \circ \psi)_* = \text{Id}$ and $\phi \circ \psi(c) = c'$. \square

This finishes the proof of Proposition 4.26. \square

4.12. Stable forms on a stable curve.

Definition 4.28. Let C be a stable nodal curve. A stable one-form on C is a section of its dualizing sheaf. In other words, a holomorphic 1-form on C^* that has at worst simple poles at the nodes and satisfies that the sums of the residues of the branches meeting at each node is zero. $\Omega(C)$ denotes the space of stable forms on C . A stable one form will be sometimes referred to as an abelian differential.

By Riemann Roch's Theorem, the space of meromorphic 1-forms on a compact connected genus g Riemann surface X with at most simple poles at $k > 0$ marked points has dimension $k + g - 1$. If we apply this to each part of a connected stable nodal curve C of genus g , we deduce that the dimension of the complex vector space $\Omega(C)$ is g (see [32]p.82) Remark that the restriction of a stable form $\omega \in \Omega(C)$ to a component C_i of the normalization of C can be the zero form. If this is the case we say that ω has a zero component.

Definition 4.29. Given a stable curve C we denote

- $\Omega^*(C) \subset \Omega(C)$ the set of stable forms without zero components.
- $\Omega_0(C) \subset \Omega(C)$ the vector subspace of stable forms with zero residue at any node
- $\Omega_0^*(C) = \Omega^*(C) \cap \Omega_0(C)$ the set of stable forms without zero components, and whose residues at the nodes are zero.

If C is of compact type, then by the residue theorem applied to each part of C , all residues of all branches at the nodes have to be zero, so $\Omega_0(C) = \Omega(C)$. If C_1, \dots, C_k are the distinct parts of C , a stable form in $\Omega(C)$ will be written as $\omega_1 \vee \dots \vee \omega_k$ where each $\omega_i \in \Omega(C_i)$ and the \vee indicates that we glue them at the points corresponding to the nodes of C .

Definition 4.30. The order of a stable form (C, ω) at a node q is defined to be

$$\text{ord}_q(\omega) = 2 + \text{ord}_q(\omega_1) + \text{ord}_q(\omega_2)$$

where ω_i denotes the restriction of ω to a branch of C through q .

Note that the order of the node cannot be 1 and $\text{ord}_q(\omega) \geq 0$ for any point $q \in C$.

Given a stable form $\omega \in \Omega^*(C)$ of genus g we define its associated divisor

$$(\omega) = \sum_{q \in C} \text{ord}_q(\omega) q$$

whose primitive degree satisfies

$$\deg(\omega) = \sum_{q \in C} \text{ord}_q(\omega) = 2g - 2.$$

The support is a disjoint union of the zeros $Z(\omega)$ and the nodes $N(\omega)$ of ω . As a consequence any stable curve C with more than $g - 1$ nodes has empty $\Omega_0^*(C)$.

4.13. Stable one forms and singular translation structures. On each component C_i where a stable form (C, ω) is not identically zero it defines naturally a singular translation structure. Indeed, around a point $q \in C_i^*$ we can locally define a holomorphic function $\phi_q(z) = \int_q^z \omega$ that is a branched covering of degree $\text{ord}_q(\omega) + 1$, ramified over 0 if the degree is at least two. At the intersection of domains two such maps ϕ_q and ϕ_p satisfy

$$\phi_p = \phi_q + \text{const.}$$

Reciprocally if we are given a cover U_α of a compact (possibly disconnected) topological surface Σ , and finite branched coverings $\phi_\alpha : U_\alpha \rightarrow V_\alpha \subset \mathbb{C}$ satisfying $\phi_\alpha = \phi_\beta + \text{const}$ at the intersections $U_\alpha \cap U_\beta$, we can define a complex structure on Σ by declaring that the ϕ_α 's are holomorphic. The abelian differential ω defined locally by $d\phi_\alpha$ is well defined on the obtained Riemann surface \hat{C} . By identifying pairs of points in \hat{C} , we obtain all nodal curves C that are normalized by \hat{C} .

4.14. Singular flat metric and geodesic foliations. Denote by $Z(\omega) \cup N(\omega)$ the support of zeros and nodes of the divisor $(\omega) = \sum_{q \in C} \text{ord}_q(\omega)q$ of a stable form (C, ω) having components (C_i, ω_{C_i}) . Any object invariant by translations in \mathbb{C} can be pulled back to $C_i^* \setminus Z(\omega_{C_i}) \cup N(\omega_{C_i})$ with singularities at the points of $Z(\omega) \cup N(\omega)$.

In particular the pull back of the flat metric in \mathbb{C} produces a singular flat metric on any non-zero component (C_i, ω_i) defined by $\omega \otimes \bar{\omega}$. At a branch C_i of C around a point $q \in C_i$ of non-negative order for the restriction $\omega|_{C_i}$, the metric is equivalent to a standard conical point of angle $2\pi(\text{ord}_q(\omega_{C_i}) + 1)$. Around a point with non-zero residue $a \in \mathbb{C}^*$, the metric is a semi-infinite cylinder equivalent to one of the ends of $\mathbb{C}^*/a\mathbb{Z}$. The volume of a stable form $\omega \in \Omega(C)$ is defined as

$$(30) \quad \text{vol}(\omega) = \frac{i}{2} \int_C \omega \wedge \bar{\omega}.$$

In particular, $0 \leq \text{vol}(\omega) \leq \infty$ and it is finite if and only if all the residues of ω at the nodes of C are zero. If $\text{vol}(\omega) = 0$ then ω is the zero form.

The oriented geodesic directional foliation of \mathbb{C} given by an angle $\theta \in \mathbb{S}^1$ is also invariant by translations, so we can also lift it to a singular oriented directional foliation \mathcal{G}_θ on C_i . Its leaves are geodesics for the metric induced by the form ω . At a zero q of ω_{C_i} the foliation has a saddle with $2(\text{ord}_q(\omega_{C_i}) + 1)$ separatrices, that alternatively enter and leave the singularity by forming an angle of π (see Figure 4). At any other point the foliation is regular.

Some subsets of stable forms where these geometric objects encode important information will be of special importance for us and we introduce appropriate notation.

Definition 4.31. Given a family \mathcal{K} of (marked or unmarked) stable curves, we define

- $\Omega_0\mathcal{K} \subset \Omega\mathcal{K}$ to be the set of stable forms over curves in \mathcal{K} whose residues at all nodes are zero (or equivalently the volume of the underlying metric is finite) and




FIGURE 4. A saddle point of a directional foliation at a zero of order two

- $\Omega_0^* \mathcal{C} \subset \Omega_0 \mathcal{C}$ the set of stable forms over curves in \mathcal{C} that have no zero components and zero residue at all nodes (or equivalently the underlying metric has finite volume and isolated singularities)

4.15. The Hodge bundle over the Deligne-Mumford compactification of $\mathcal{M}_{g,n}$. Recall (see [32], Chapter 4) that over the Deligne-Mumford compactification $\overline{\mathcal{M}}_{g,n}$ we can define the universal curve bundle

$$\overline{\mathcal{C}}_{\mathcal{M}_{g,n}} \rightarrow \overline{\mathcal{M}}_{g,n}$$

which is a holomorphic map between compact complex orbifolds whose fiber over the point $(C, r_1, \dots, r_n) \in \overline{\mathcal{M}}_{g,n}$ is biholomorphic to the curve with marked points (C, r_1, \dots, r_n) . The cotangent bundle to this fibration is well defined on the complement of the set of nodes. It extends as a line bundle \mathcal{L} to the whole universal curve, called the relative cotangent bundle. A section of the restriction $\mathcal{L}|_C$ corresponds precisely to a stable form on C . The constancy of the dimension of this space of sections implies that the sheaf obtained by pushing it forward to $\overline{\mathcal{M}}_{g,n}$ is locally free of rank g (see [33] [Exercise 5.8]). Thus it defines a holomorphic vector bundle

$$\pi : \Omega \overline{\mathcal{M}}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$$

called the Hodge bundle. The fiber of π over a point $(C, r_1, \dots, r_n) \in \overline{\mathcal{M}}_{g,n}$ corresponds precisely to the set of stable forms $\Omega(C)$. When $n = 0$, the Hodge bundle trivializes over any of the preferred neighbourhoods $U = U_c/\Gamma_c$. A choice of a Lagrangian $\Lambda \subset H_1(\Sigma_g)$ containing all the classes induced by the curves in the curve system c allows to define a trivialization

$$\Omega U \simeq U \times \text{Hom}(\Lambda, \mathbb{C})$$

Definition 4.32. The boundary strata of $\Omega \overline{\mathcal{M}}_{g,n}$ is the set of stable forms over stable curves with marked points lying the boundary strata $\partial \overline{\mathcal{M}}_{g,n}$.

The pull back of the compactified Hodge bundle $\Omega \overline{\mathcal{M}}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$ by the map $\overline{\mathcal{T}}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$ defines the (topological) complex vector bundle $\Omega \overline{\mathcal{T}}_{g,n} \rightarrow \overline{\mathcal{T}}_{g,n}$ of homotopically marked stable forms of genus g with n marked points. An element in $\Omega \overline{\mathcal{T}}_{g,n}$ is a pair $([f : \Sigma_{g,n} \rightarrow (C, r_1, \dots, r_n)], \omega)$ where $[f : \Sigma_{g,n} \rightarrow (C, r_1, \dots, r_n)] \in \overline{\mathcal{T}}_{g,n}$ and $\omega \in \Omega(C)$. The Torelli group $\mathcal{I}_{g,n}$ acts on the bundle $\Omega \overline{\mathcal{T}}_{g,n} \rightarrow \overline{\mathcal{T}}_{g,n}$. The quotient defines a bundle $\Omega \overline{\mathcal{S}}_{g,n} \rightarrow \overline{\mathcal{S}}_{g,n}$. These will be referred to as the Hodge bundle over augmented Teichmüller or augmented Torelli space, depending on the case.

Their boundaries are, by definition, the preimage of the boundary of $\Omega \overline{\mathcal{M}}_g$ by the corresponding projection, i.e. the stable forms over curves with some node.

Given a collection of (marked or unmarked) stable curves \mathcal{K} we define $\Omega \mathcal{K}$ to be the set of stable forms over \mathcal{K} , with the induced topology in the corresponding vector bundle of forms.

Given a subset of some space of forms $L \subset \Omega\mathcal{K}$ we denote by ∂L the intersection of L with the boundary strata of the ambient space.

Endowed with the complex structure in each boundary stratum B , ΩB becomes a holomorphic vector bundle over B . We will refer to it as a boundary stratum of the space of forms where B lies.

Definition 4.33. A homologically marked stable form of genus $g \geq 0$ with n marked points is a tuple $(C, r_1, \dots, r_n, m, \omega) \in \Omega\bar{\mathcal{S}}_{g,n}$ where ω is a stable one-form on a homologically marked stable curve (C, r_1, \dots, r_n, m) of genus g with n marked points. By abuse of language we say that ω pinches $a \in H_1(\Sigma_g, q_1, \dots, q_n, \mathbb{Z}) \setminus 0$ if a is primitive and $m(a) = 0$. We say that $(C, r_1, \dots, r_n, m, \omega)$ belongs to the boundary strata if C has some node.

4.16. Attaching and forgetful maps on Hodge bundles. The continuous attaching maps defined at the level of marked curves in subsection 4.10 can be lifted to attaching maps on the corresponding Hodge bundles.

In particular, using Lemmas 4.23 and 4.24 the attaching maps can be lifted as continuous maps to the corresponding Hodge bundles:

$$(31) \quad \Omega\bar{\mathcal{S}}_{g_1, n_1} \times \Omega\bar{\mathcal{S}}_{g_2, n_2} \rightarrow \Omega\bar{\mathcal{S}}_{g_1+g_2, n_1+n_2-2} \text{ and}$$

$$(32) \quad \Omega\bar{\mathcal{S}}_{g-1, n+2} \rightarrow \Omega\bar{\mathcal{S}}_{g, n}$$

by simply considering the stable form over the image curve that coincides with the given restriction on each part. The image will only contain forms with zero residue at the node produced by the attaching map. More generally, we can use attaching maps and the biholomorphism (20) to define, via the attaching maps a decomposition of forms with zero residues on a stratum B_c of $\bar{\mathcal{T}}_{g,n}$:

$$(33) \quad \Omega_0 B_c \cong \Pi_i \Omega\mathcal{T}_{g_i, n_i}$$

On the other hand we can also lift the forgetful maps defined by Lemma 4.25 to the Hodge bundle. To construct the natural lift

$$\Omega\bar{\mathcal{S}}_{g, n} \rightarrow \Omega\bar{\mathcal{S}}_{g, n-1}$$

it suffices to restrict the form to the image curve. Remark that on the eventually contracted components after forgetting the point the form is of genus zero and uniquely determined by the value of one of the residues.

4.17. Sub-stratification of Hodge bundles and period coordinates on strata. The boundary stratification of the Hodge bundle over $\bar{\mathcal{M}}_{g,n}$ is substratified by the topological properties of the zero divisor and polar set of the forms. Two elements in $\Omega\bar{\mathcal{M}}_{g,n}$ are said to belong to the same substratum if there exists a homeomorphism between the underlying marked curves preserving the following data of the forms:

- (1) zero components
- (2) nodes with non-zero residue
- (3) the associated zero divisor on each non-zero component

In particular, if an element has a zero of order k at a marked point, then any element in its substratum will have a zero at of order k at a marked point. The same happens with poles

with zero residue: if the restriction of the form to one of the branches has a zero of order k , then any form in its stratum will have this property on the corresponding branch of a node.

This stratification can be lifted to $\Omega\bar{\mathcal{S}}_{g,n}$ and $\Omega\bar{\mathcal{T}}_{g,n}$. It is a substratification of the stratification induced by $\sqcup B_c$ on $\bar{\mathcal{T}}_{g,n}$. The generic stratum in ΩB_c is the only substratum that is open in ΩB_c , and it is Zariski open too. Indeed, the union of all other substrata in ΩB_c form an analytic subset of ΩB_c . A substratum in ΩB_c is minimal if it has minimal dimension among substrata of ΩB_c . In particular, the generic stratum in $\Omega B_\emptyset = \Omega\mathcal{T}_{g,n}$ is the stratum having only simple zeros at unmarked points and is also dense in $\Omega\bar{\mathcal{T}}_{g,n}$. The minimal stratum in $\Omega\mathcal{T}_{g,n}$ is formed by forms with a single zero of order $2g-2$ at some marked point (for $n \geq 1$).

The restriction of an isomorphism of type (33) to all possible products of substrata of $\Omega\mathcal{T}_{g_i,n_i}$ describe all the different substrata of forms with zero residues in $\Omega_0 B_c$. Let us define local holomorphic coordinates on them.

The zero substratum of $\Omega\mathcal{M}_{g,n}$ is isomorphic to $\mathcal{M}_{g,n}$. On the other hand, Veech [65] and Masur in [49] proved that local holomorphic coordinates can be given in any substratum R of $\Omega\mathcal{M}_{g,n}$ around a point (C, P, ω) with $\omega \neq 0$. They are defined by integration of the form on cycles in $H_1(C, Z(\omega) \cup P; \mathbb{Z})$ and therefore lie in the vector space $H^1(C, Z(\omega) \cup P, \mathbb{C})$. These endow the stratum with the structure of a complex manifold compatible with the one induced by the complex structure of $\Omega\mathcal{T}_{g,n}$.

We conclude from the equivalence (33) and the description of strata, on smooth curves, that the substratum of stable forms containing a point $(C, P, f, \omega) \in \Omega_0^* \bar{\mathcal{T}}_{g,n}$ without zero components and with zero residues at the nodes is *locally* isomorphic to the product

$$\prod_i H^1(C_i, (Z(\omega) \cup N(C) \cup P) \cap C_i, \mathbb{C})$$

where each C_i corresponds to a connected component of the normalization of C and $N(C)$ is the union of all points of attaching in the components C_i to obtain C . These local coordinates will be referred to as period coordinates on the substratum.

4.18. Periods of marked stable forms with zero residues at the nodes. A stable form on (C, r_1, \dots, r_n) is holomorphic on C^* and can thus be integrated along paths in C^* . For closed paths the value of the integral does only depend on the homology class in $H_1(C^*, r_1, \dots, r_n; \mathbb{Z})$ and it is called the period of the class. If the residues at nodes are all zero, we can also integrate along paths passing *through* the nodes, and the integral along a closed path depends only on its class in $H_1(C, \mathbb{Z})$.

For any marked stable form $([f : \Sigma_{g,n} \rightarrow (C, r_1, \dots, r_n)], \omega) \in \Omega_0 \bar{\mathcal{T}}_{g,n}$ of genus g with n marked points and zero residues at the nodes we have a well defined notion of **period homomorphism**

$$\text{Per}([f : \Sigma_{g,n} \rightarrow (C, r_1, \dots, r_n)], \omega) : H_1(\Sigma_g, q_1, \dots, q_n; \mathbb{Z}) \rightarrow \mathbb{C}$$

defined by

$$(34) \quad \text{Per}([f : \Sigma_{g,n} \rightarrow (C, r_1, \dots, r_n)], \omega)(\gamma) = \int_{f_*(\gamma)} \omega \text{ for } \gamma \in H_1(\Sigma_g, q_1, \dots, q_n; \mathbb{Z}).$$

Since the period homomorphism depends only on the homological marking induced by the homotopical marking, we can also define the period on a homologically marked stable curve $(C, r_1, \dots, r_n, m, \omega)$ with marked points by $\{\gamma \mapsto \int_{m(\gamma)} \omega\}$. Any homologically non-trivial

curve a in Σ_g pinched by the marking, i.e. belonging to $\text{Ker}(f_*)$, also belongs to the kernel of the period homomorphism. Remark that if the period homomorphism is injective then the marking is an isomorphism, and thus the curve C must be of compact type.

Definition 4.34. The period map associated to $(\Sigma_g, q_1, \dots, q_n)$ is the map

$$\Omega_0 \overline{\mathcal{T}}_{g,n} \rightarrow H^1(\Sigma_g, q_1, \dots, q_n, \mathbb{C})$$

that associates to any stable form with zero residues marked by $(\Sigma_g, q_1, \dots, q_n)$, its period homomorphism. It is invariant by the Torelli group $\mathcal{I}_{g,n}$ action and descends to the quotient

$$\Omega_0 \overline{\mathcal{S}}_{g,n} \rightarrow H^1(\Sigma_g, q_1, \dots, q_n, \mathbb{C}).$$

One of the difficulties that we will encounter is related to the fact that the domain of definition of the period map is neither open nor closed. It is a union of substrata of stable forms. It contains the (dense) open subset of stable forms on smooth curves, but in the boundary strata we need to restrict to the closed set of strata that have no residues at the nodes.

Definition 4.35. An isoperiodic deformation of a point in $\Omega_0 \overline{\mathcal{T}}_0$ or $\Omega_0 \overline{\mathcal{S}}_{g,n}$ is a continuous deformation in the ambient space that has constant value for the period map.

The following notation will be useful

Definition 4.36. Given a homomorphism $p : H_1(\Sigma_g, q_1, \dots, q_n; \mathbb{Z}) \rightarrow \mathbb{C}$ and a family $\mathcal{K} \subset \Omega_0 \overline{\mathcal{S}}_{g,n}$ we denote by $\mathcal{K}(p)$ the subset of marked stable forms in \mathcal{K} having period homomorphism p , i.e. the intersection of the the period fiber over p with the set \mathcal{K} .

4.19. Local structure of the fibers of the period map. A local fiber of the period map can be thought as a disjoint union of sets in the different strata of the ambient space. The next theorem shows that this partition is actually a nice stratification

Theorem 4.37. *A local fiber L of the period map in $\Omega_0^* \overline{\mathcal{S}}_{g,n}$ projects to the orbifold chart of $\Omega_0 \overline{\mathcal{M}}_{g,n}$ as a complex manifold transverse to all boundary divisors through the point. Therefore, L is an abelian ramified cover of a normal crossing divisor in $(\mathbb{C}^{2g+n-3}, 0)$ having precisely one component of codimension one in each component of codimension one of the ambient space $\Omega_0^* \overline{\mathcal{S}}_{g,n}$ through the point.*

The statement is not true in general for forms with zero components (see subsection [4.20](#)).

Proof. A local period fiber on $\Omega_0 \overline{\mathcal{S}}_g$ can be lifted to a local period fiber in $\Omega_0 \overline{\mathcal{T}}_g$. If the lift of the point belongs to the stratum ΩB_c , the covering group is the subgroup $\Gamma_c \cap \mathcal{I}_g \subset \Gamma_c$. Hence, we just need to prove that the period fiber at the level of the Hodge bundle over augmented Teichmüller space has a stratification that is locally equivalent to some map $\varphi_{l,2g+n-3}$ as in [\(21\)](#) where l denotes the number of simple closed curves in the curve system c . It suffices in fact to show that at the level of the quotient $\Omega_0(U_c/\Gamma_c)$ the period fiber is a holomorphic manifold transverse to every boundary component passing through the point. Thanks to the normal crossing condition of the ambient space, this is guaranteed if the period map is submersive in restriction to the (regular) stratum of the normal crossing divisor where the point belongs to. In fact we will not use the boundary stratum but a smaller regular submanifold, consisting of the substratum through the point.

Proposition 4.38. *Consider $([f_0 : \Sigma_{g,n} \rightarrow (C_0, P_0)], \omega_0) \in \Omega_0 \overline{\mathcal{T}}_{g,n}$ and ΩU_c its preferred neighbourhood. Denote $m_0 := f_{0*} : H_1(\Sigma_g, q_1, \dots, q_n; \mathbb{Z}) \rightarrow H_1(C_0, P_0; \mathbb{Z})$ and define H as the germ of the zero set of the map $\Omega U_c \rightarrow \text{Hom}(\ker m_0, \mathbb{C})$ defined by integration. Then $H \subset \Omega_0 U_c$, H/Γ_c is a proper holomorphic submanifold of $\Omega_0(U_c/\Gamma_c)$ and the period map induces a holomorphic map $\Omega_0(H/\Gamma_c) \rightarrow H^1(C_0, P_0, \mathbb{C})$. If the form ω_0 has no zero components, the holomorphic map thus defined is submersive, even by restricting it to the substratum containing (C_0, ω_0) . Hence, the isoperiodic foliation extends along H/Γ_c as a regular foliation.*

Proof. We first need a lemma about the holomorphicity of integration over a cycle:

Lemma 4.39. *Let $U_c/\Gamma_c \rightarrow U \subset \overline{\mathcal{M}}_{g,n}$ be a (orbifold) chart around a point $(C_0, P_0) \in \overline{\mathcal{M}}_{g,n}$ and γ be a path in $\Sigma_g \setminus c$. Then, the map $\Omega U_c \rightarrow \mathbb{C}$ defined by*

$$([f : \Sigma_{g,n} \rightarrow (C, P)], \omega) \mapsto \int_{f_*(\gamma)} \omega$$

induces a well defined holomorphic map on $\Omega U \subset \Omega \overline{\mathcal{M}}_{g,n}$.

Proof. The function is well defined on U_c/Γ_c because γ does not intersect any of the simple closed curves in c . It is holomorphic outside the boundary and bounded in the neighbourhood of every boundary point, hence holomorphic by Riemann extension Theorem. \square

Remark first that there is a non-trivial exact sequence

$$(35) \quad 0 \rightarrow \text{Hom}(H_1(C_0, P_0; \mathbb{Z}), \mathbb{C}) \rightarrow \text{Hom}(H_1(\Sigma_g, q_1, \dots, q_n; \mathbb{Z}), \mathbb{C}) \rightarrow \text{Hom}(\ker m_0, \mathbb{C}) \rightarrow 0$$

induced by that defined by the marking

$$0 \rightarrow \ker m_0 \rightarrow H_1(\Sigma_g, q_1, \dots, q_n; \mathbb{Z}) \rightarrow H_1(C_0, P_0; \mathbb{Z}) \rightarrow 0.$$

The map

$$\Omega U_{\overline{c}} \rightarrow \text{Hom}(\ker m_0, \mathbb{C})$$

given by $([f : \Sigma_{g,n} \rightarrow (C, r_1, \dots, r_n)], \omega) \mapsto \{c \mapsto \int_{f_*(c)} \omega\}$ is well defined and invariant by the action of $\Gamma_{\overline{c}}$ on $\Omega U_{\overline{c}}$. Lemma 4.39 implies that it induces a holomorphic map

$$(36) \quad h : \Omega(U_{\overline{c}}/\Gamma_{\overline{c}}) \rightarrow \text{Hom}(\ker m_0, \mathbb{C}).$$

We claim that h is submersive at $[f_0 : \Sigma_{g,n} \rightarrow (C_0, P_0)]$. In fact, its restriction to the Hodge bundle fiber ΩC_0 is already submersive. It associates to each node, the value of its residue. It is therefore linear. The kernel of this restriction is precisely $\Omega_0 C_0$, the space of stable forms with zero residues on C_0 . Let C_1, \dots, C_l be the components of the normalization of C_0 and define $N(C_i) \subset C_i$ the points corresponding to the nodes of C_0 . The restriction of any stable form in ΩC_0 to the component C_i is a meromorphic form with at worst simple poles on $N(C_i)$. It satisfies the residue theorem. On the other hand, by Riemann-Roch theorem applied to C_i with $i \geq 1$ if we denote $m_i = |N(C_i)|$ and $g_i = g(C_i)$ the genus of C_i we get

- (1) the space M_i of meromorphic forms on C_i that have at worst simple poles at the marked points has dimension $g_i + m_i - 1$ and
- (2) the space ΩC_i of holomorphic forms on C_i has dimension g_i .

Denote $L_i \subset \mathbb{C}^{m_i}$ the codimension one space defined by $x_1 + \dots + x_{m_i} = 0$. By the previous dimension calculations, the sequence $0 \rightarrow \Omega C_i \rightarrow M_i \rightarrow L_i \rightarrow 0$ is exact. Recall that $\sum m_i = 2m$ where m denotes the number of nodes of C_0 . Consider $L \subset \mathbb{C}^{2m}$ the codimension l subspace $L_1 \oplus \dots \oplus L_l$.

The sequence

$$(37) \quad 0 \rightarrow \Omega_0 C_0 \rightarrow M_1 \oplus \cdots \oplus M_l \rightarrow L \rightarrow 0$$

is still exact. A necessary and sufficient condition for an element of $M_1 \oplus \cdots \oplus M_l$ to define a form in ΩC_0 is that the residues at the points that are attached to produce a node in C_0 have opposite sign. This corresponds to cutting L with m linear equations of type $x_i - x_j = 0$ in \mathbb{C}^{2m} where m is the number of nodes of C_0 . When we intersect $m - 1$ of those with L , we easily obtain the last of such equations. In fact, the dimension of the intersection is $2m - l - (m - 1) = m - l + 1$. The points in this intersection are in one to one correspondence with elements in $\text{Hom}(\ker m_0, \mathbb{C})$. Indeed, the non-separating curves in \bar{c} generate $\ker m_0$, and the only relations they have to satisfy are the relations given by the fact that some sums of the generators are boundaries of components. These relations coincide with those defining the relevant subspace of L . On the other hand the exactness of (37) tells us that we can find a form in ΩC_0 with any given residue homomorphism $\ker m_0 \rightarrow \mathbb{C}$. This proves that $h|_{\Omega C_0}$ is surjective. Since it is linear it is also submersive, and so are h and the restriction of h to each boundary component containing (C_0, m_0, ω_0) . The levels of h are regular subvarieties transverse to each boundary component and invariant by Per . The zero level set

$$H = \{([f : \Sigma_{g,n} \rightarrow (C, P)], \omega) \in \Omega U_{\bar{c}} : \int_{f_*(c)} \omega = 0 \quad \forall c \in \ker m_0\}$$

contains the marked stable form $([f_0 : \Sigma_{g,n} \rightarrow (C_0, P_0)], \omega_0)$ and is invariant under the action of $\Gamma_{\bar{c}}$. By the exact sequence (35), the restriction of the period map to H can be thought as a map $H \rightarrow H^1(C_0, \mathbb{C})$ that is invariant by $\Gamma_{\bar{c}}$. The induced map

$$h_2 : H/\Gamma_{\bar{c}} \rightarrow \text{Hom}(H_1(C_0, P_0; \mathbb{Z}), \mathbb{C}) \text{ is holomorphic.}$$

This proves that the fibers of the period map in $\Omega_0 \bar{\mathcal{S}}_{g,n}$ are preimages under the branched covering $\Omega U_{\bar{c}} \rightarrow \Omega(U_{\bar{c}}/\Gamma_{\bar{c}})$ of analytic sets.

It remains to prove that h_2 is submersive in restriction to the substratum R containing $([f_0 : \Sigma_{g,n} \rightarrow (C_0, P_0)], \omega_0)$ whenever ω_0 has no zero components.

Recall that C_1, \dots, C_l denote the components of the normalization of C_0 and define

- ω_i the restriction of ω_0 to C_i
- $g_i = g(C_i)$ the genus of C_i
- $P_i = (P \cap C_i) \cup N(C_i) \cup Z(\omega_{0i})$

The local coordinates in the substratum R_i of the stable form (C_i, P_i, ω_i) . The latter are given by vectors in $W_i = H^1(C_i, P_i, \mathbb{C})$.

By using the attaching maps used to obtain C_0 from C_1, \dots, C_l we define a surjective holomorphic map

$$W_1 \oplus \cdots \oplus W_l \rightarrow R.$$

If we prove that the composition of this map with the period map h_2 is submersive onto $H^1(C_0, \mathbb{C})$ we will be done. The said composition is a map

$$W_1 \oplus \cdots \oplus W_l \rightarrow \text{Hom}(H_1(C_0, P_0, \mathbb{Z}), \mathbb{C})$$

that factors through

$$\widehat{W}_1 \oplus \cdots \oplus \widehat{W}_l \text{ where } \widehat{W}_i = H^1(C_i, N(C_i), \mathbb{C}).$$

The map $W_1 \oplus \cdots \oplus W_l \rightarrow \widehat{W}_1 \oplus \cdots \oplus \widehat{W}_l$ and the map $\widehat{W}_1 \oplus \cdots \oplus \widehat{W}_l \rightarrow \text{Hom}(H_1(C_0, P_0; \mathbb{Z}), \mathbb{C})$ are obviously submersive. \square

\square

Definition 4.40. The map

$$\mathcal{P}er_{g,n} : \Omega_0^* \bar{\mathcal{S}}_{g,n} \rightarrow \text{Hom}(H_1(\Sigma_{g,n}); \mathbb{C})$$

denotes the extension to the bordification $\Omega_0^* \bar{\mathcal{S}}_{g,n}$ of the period map $\text{Per}_{g,n}$ defined on $\Omega \mathcal{S}_{g,n}$. When there is no risk of confusion we omit the subindex and write $\mathcal{P}er$ and Per respectively.

Definition 4.41. Let $p \in \mathcal{H}_g$ and denote \mathcal{G}_p the dual boundary complex of the bordification $\text{Per}^{-1}(p)$ of $\text{Per}^{-1}(p)$ defined by its closure in $\Omega_0^* \bar{\mathcal{S}}_g$. The final statement of Theorem 4.37 and Remark 4.9 allow to define a continuous map of dual boundary complexes

$$(38) \quad \mathcal{G}_p \rightarrow \mathcal{C}(\bar{\mathcal{S}}_g) \simeq \mathcal{C}_g / \mathcal{I}_g$$

Corollary 4.42. *The period map $\text{Per}_{g,n}$ induces a regular holomorphic foliation $\bar{\mathcal{F}}_{g,n}$ on the moduli space $\Omega^* \bar{\mathcal{M}}_{g,n}^c$ of stable forms without zero components on curves of compact type called the isoperiodic foliation. It is transverse to all boundary components and to the substratum passing through the point. Its restriction to $\Omega^* \mathcal{M}_{g,n}$ will be denoted by $\mathcal{F}_{g,n}$.*

Proof. All the residues of all forms in a neighbourhood of a point in $\Omega_0^* \bar{\mathcal{S}}_{g,n}^c$ are zero. Hence, the map $\mathcal{P}er$ is holomorphic in some neighbourhood. Proposition 4.38 guarantees that it is also submersive there. On the other hand $\mathcal{P}er_{g,n}$ is equivariant with respect to the natural action of $\text{Mod}(\Sigma_{g,n})$ on source and target. Hence the underlying foliation is well defined in the quotient. \square

Remark 4.43. The isoperiodic foliation $\bar{\mathcal{F}}_{g,n}$ extends to the whole space $\Omega \bar{\mathcal{M}}_{g,n}$ as a singular holomorphic foliation. In fact, the zero section of the Hodge bundle is part of the singular set of this foliation, as are the points of non-compact type in $\Omega_0^* \bar{\mathcal{M}}_{g,n}$, and most strata with zero components (see section 4.44 for an example). To construct the extension isoperiodic foliation to the missing strata (those with non-zero residue at the nodes) we need to define the notion of isoperiodic family of *meromorphic* forms on smooth curves. This subject will be developed in a forthcoming paper.

4.20. Some singularities of the isoperiodic set on regular points of $\Omega \bar{\mathcal{S}}_{g,2}$. In this subsection we analyze the local isoperiodic deformation spaces at some points of the Hodge bundle over the complex manifold $\mathcal{U}_{g,2} \subset \bar{\mathcal{S}}_{g,2}$, bordification of $\mathcal{S}_{g,2}$, introduced in Section 4.11. The contents of this section will be useful in Section 7.

Recall from Section 4.26 that the boundary points in $\mathcal{U}_{g,2}$ consist on stable curves having precisely one separating node that leaves both marked points on a genus zero component (in Figure 2 we depict such a stable curve for $g = 2$). The strata of $\partial \mathcal{U}_{g,2}$ form a smooth divisor having a connected component B_β for each class $\beta \in \mathfrak{b} = \{\beta \in H_1(\Sigma_g, q_1, q_2; \mathbb{Z}) : \partial\beta = q_2 - q_1\}$

The bundle $\Omega \mathcal{U}_{g,2} \rightarrow \mathcal{U}_{g,2}$ is a complex manifold. The stratification of the ambient space induces a stratification on $\Omega \mathcal{U}_{g,2}$ that has the following boundary strata:

$$\bigsqcup_{\beta \in \mathfrak{b}} \Omega B_\beta.$$

Each stable form in ΩB_β has a zero component of genus zero and lies in the domain $\Omega_0 \bar{\mathcal{S}}_{g,2}$ of the period map $\text{Per}_{g,2}$. Moreover the restriction of this map to $\Omega \mathcal{U}_{g,2}$ is holomorphic (see Proposition 4.38). We want to analyze the local analytic set formed by a period fiber around generic points of ΩB_β .

By abuse of language we write $\Omega^{SZ} B_\beta \subset \Omega B_\beta$ the forms that have simple zeros on the smooth genus g component and a zero component of genus zero, and $\Omega^{SZ,1} B_\beta \subset \Omega^{SZ} B_\beta$ those where the zero component is glued at a simple zero of the non-zero component (as in Figure 2).

Proposition 4.44. *Let $(C^0, r_1^0, r_2^0, m^0, \omega^0) \in \Omega^{SZ} B_\beta \subset \Omega \mathcal{U}_{g,2}$. The local isoperiodic deformation space at $(C^0, r_1^0, r_2^0, m^0, \omega^0)$ is*

- (1) *a normal crossing of two smooth manifolds, precisely one of which is contained in the boundary strata, if $(C^0, r_1^0, r_2^0, m^0, \omega^0) \in \Omega^{SZ,1} B_\beta$ or*
- (2) *a smooth manifold contained in the boundary if $(C^0, r_1^0, r_2^0, m^0, \omega^0) \notin \Omega^{SZ,1} B_\beta$*

Proof of Proposition 4.44. Along the proof we write $U = \mathcal{U}_{g,2}$. Let $(C, r_1, r_2, m, \omega) \in \Omega^{SZ} B_\beta$. We consider the contraction of the spherical component of ω , namely the tuple $(\hat{C}, \hat{q}, \hat{m}, \hat{\omega}) \in \Omega^{SZ} \mathcal{S}_{g,1}$ satisfying the following relations

$$(39) \quad \hat{m} = m \circ \iota \text{ where } \iota : H_1(\Sigma_{g,1}) \rightarrow H_1(\Sigma_{g,2}) \text{ is the natural inclusion}$$

$$(40) \quad (C, r_1, r_2, m) = (\hat{C}, \hat{q}, \hat{m}) \underset{\hat{q}=\infty}{\vee} (\mathbb{CP}^1, 0, 1, \infty) \text{ for some } \hat{q} \in \hat{C}$$

and

$$(41) \quad \omega = \hat{\omega} \vee 0 \text{ on } (\hat{C}, \hat{q}, \hat{m}) \underset{\hat{q}=\infty}{\vee} (\mathbb{CP}^1, 0, 1, \infty).$$

The point lies in $\Omega^{SZ,1} B_\beta$ if and only if $\hat{\omega}(\hat{q}) = 0$.

We can define a holomorphic map

$$(42) \quad h_\beta : \Omega^{SZ} U \rightarrow \mathbb{C} \text{ by } h_\beta(C, r_1, r_2, m, \omega) = \int_{m(\beta)} \omega$$

Its zero set $H_\beta = \{h_\beta = 0\}$ contains $\Omega^{SZ} B_\beta$ and is invariant by the isoperiodic foliation on $\Omega^{SZ} U$.

Let us provide adapted coordinates of $\Omega^{SZ} U$ around the point $(C^0, r_1^0, r_2^0, m^0, \omega^0) \in \Omega^{SZ} B_\beta$. Denote

$$(43) \quad \text{For} : \Omega^{SZ} U \rightarrow \Omega^{SZ} \mathcal{S}_g$$

the restriction of the forgetful map $\text{For}_{g,2} : \Omega \bar{\mathcal{S}}_{g,2} \rightarrow \Omega \mathcal{S}_g$ to the open set $\Omega^{SZ} U$.

Given a small neighbourhood V^0 of $\text{For}(C^0, r_1^0, r_2^0, m^0, \omega^0)$ in $\Omega \mathcal{S}_g$ suppose we are given a local holomorphic section $q : V^0 \rightarrow \Omega^{SZ} \mathcal{S}_{g,1}$ of $\Omega^{SZ} \mathcal{S}_{g,1} \rightarrow \Omega^{SZ} \mathcal{S}_g$ such that

- (1) $q(\hat{C}^0, \hat{m}^0, \hat{\omega}^0) = \hat{q}^0 \in \hat{C}^0$ is the point corresponding to the node of C^0 , and
- (2) the order of the form $\hat{\omega}$ at the point $q(C, m, \omega)$ is independent of (C, m, ω) and (therefore) has value $k - 1 \in \{0, 1\}$

The germs of holomorphic functions

$$r_1 \mapsto \int_{q \circ \text{textFor}(C, r_1, r_2, m, \omega)}^{r_1} \widehat{\omega} \quad \text{and} \quad r_2 \mapsto \int_{q \circ \text{For}(C, r_1, r_2, m, \omega)}^{r_2} \widehat{\omega}$$

at $q(C, m, \omega)$ have each degree k and can therefore be written as

$$r_1 \mapsto (\zeta_1(C, r_1, r_2, m, \omega))^k \quad \text{and} \quad r_2 \mapsto (\zeta_2(C, r_1, r_2, m, \omega))^k$$

for some germ of biholomorphisms

$$\zeta_1, \zeta_2 : (C, q(C, m, \omega)) \rightarrow (\mathbb{C}, 0)$$

In a small neighbourhood W^0 of $(C^0, r_1^0, r_2^0, m^0, \omega^0)$ in $\text{For}^{-1}(V^0)$ we consider the holomorphic map $\Psi_q : W^0 \rightarrow V^0 \times \mathbb{C}^2$ defined by

$$(44) \quad \Psi_q(C, r_1, r_2, m, \omega) = (\text{For}(C, r_1, r_2, m, \omega), \zeta_1(C, r_1, r_2, m, \omega), \zeta_2(C, r_1, r_2, m, \omega)).$$

It is locally injective and therefore it is locally a biholomorphism. In these coordinates, the boundary strata of $\Omega^{SZ}U$ are defined by $\{\zeta_1 = \zeta_2\}$ and the map h_β and $\text{Per}_{g,2}$ by

$$h_\beta((C, m, \omega), \zeta_1, \zeta_2) = \zeta_2^k - \zeta_1^k \quad \text{and} \quad \text{Per}_{g,2}((C, m, \omega), \zeta_1, \zeta_2) = (\text{Per}_{g,0}(C, m, \omega), \zeta_2^k - \zeta_1^k).$$

In particular

$$\text{Per}_{g,2}^{-1}(p_0, 0) = \bigcup_{\{\alpha \in \mathbb{C} : \alpha^k = 1\}} \text{Per}_{g,0}^{-1}(p_0) \times \{\zeta_2 = \alpha \zeta_1\}$$

When $k = 1$ the germ of isoperiodic set at $(C^0, r_1^0, r_2^0, m^0, \omega^0)$ is smooth contained in the boundary.

When $k = 2$, the holomorphic section $q : V_0 \rightarrow \Omega^{SZ}\mathcal{S}_{g,1}$ is uniquely defined by the implicit condition $\omega(q(C, m, \omega)) = 0$. There are two smooth isoperiodic components intersecting transversely at $(C^0, r_1^0, r_2^0, m^0, \omega^0)$; precisely one lies in the boundary. \square

5. SURGERIES ON STABLE FORMS AND MODELS OF DEGENERACY

In this section we recall a surgery called Schiffer variation, that allows to construct isoperiodic deformations of forms without zero components on stable curves. We use them mainly to find boundary points that have some node zero residues and no zero components in the closure of any connected component of a fiber of Per .

5.1. Schiffer variations on stable forms. A Schiffer variation is a continuous deformation of stable forms with some branch point or node. It can be best described in terms of the associated translation structures. The surgery changes the translation structure associated to a form on some small neighbourhood in the surface without changing it on the complement. By taking representatives of the cycles in the homology group that avoid that open set, it is easy to see that the period maps before and after the surgery coincide. There are several instances of the surgery that produce differences on the underlying surface and the translation structure. For instance it allows to deform the projective structure without varying the orders and number of zeros and nodes. It also allows to split a multiple branch point into simpler branch points; at last it allows to produce a smooth surface with several branch points from a node. The surgery operation is invertible (the surgery is actually involutive under suitable restrictions) so the inverse operations can be used to join branch points and to produce nodal curves. They were first considered by Schiffer in [60]. A detailed discussion of Schiffer

variations on projective structures the can be found in [15]. We will introduce it only for the case of branched translation structures and will use them to prove that any abelian differential on a smooth curve can be joined to an abelian differential on a stable curve with some node by a sequence of such surgeries.

Let (C_0, m_0, ω_0) be a marked abelian differential on a nodal curve C_0 and q be point where $0 < \text{ord}_q(\omega_0) < \infty$. Remark that the chart $\phi = \phi_q$ defined by ω_0 around q can be analytically continued along any path in C_0 . Let γ_1 and γ_2 be two embedded paths in C_0 starting at q parametrized by $[0, 1]$ whose images do not intersect outside the endpoints. For our purposes it is important to stress that both endpoints *can* coincide with each other, and/or with the starting point. We say that γ_1 and γ_2 are **twin paths** if none passes through two distinct nodes and the continuation ϕ_i of ϕ along γ_i satisfies that $\phi_1 \circ \gamma_1(t) = \phi_2 \circ \gamma_2(t) \in \mathbb{C}$ for all t and $t \mapsto \phi_i \circ \gamma_i(t)$ is a simple path in \mathbb{C} . Since the chart ϕ_q is a branched covering of degree $\text{ord}_q(\omega_0) + 1$ each path starting at q has $\text{ord}_q(\omega_0)$ candidates to be its twin paths. Remark that the restriction of a pair of twin paths parametrized by $[0, 1]$ to some sub-interval $[0, t_0]$ is still a pair of twin paths.

Given a pair of twin paths γ_1, γ_2 in C_0 , we can use the equivalence between abelian differentials and atlases formed by branched coverings over open sets in \mathbb{C} and with transitions in the set of translations $z \mapsto z + \text{const}$ to produce a new abelian differential ω_1 by cutting and pasting the twin paths as follows. Let U_0 be a small neighbourhood of $\gamma_1 \cup \gamma_2$ where the ϕ_i 's are defined. Cut U_0 along the segments γ_1 and γ_2 and glue the boundary on the left (resp. on the right) of γ_1 to the boundary on the right (resp. on the left) of γ_2 by identifying points that have the same image for ϕ_i (see Figure 5). The fact that none of the twins passes through distinct nodes implies that at any point m , the new germ (V, m) of singular surface has the property that $V \setminus m$ has one or two components. This implies that the new singular surface has only regular points and nodes. Let U_1 be the set of points obtained from U_0 after the gluing. The new nodal surface is equipped with a family of local branched coverings. Indeed, on the complement of U_1 we consider the family of branched coverings given by integrating ω . On U_1 we consider the branched covering defined by the ϕ_i 's after the new identification. This family of branched coverings is translation invariant and defines a stable form ω_1 on some stable curve C_1 , that has a non-zero component whenever the component has points belonging to U_1 . The number of points where the local covering has degree at least two is finite. The order of ω_1 at a point that has not been glued is the same as in ω_0 . The zeros or nodes of ω_1 that lie on the glued points, can only appear at points corresponding to endpoints of the twin paths, or to zeros of ω_0 lying on the twins. By construction, the total order of the zero divisor of ω_1 on U_1 is the same as that of ω_0 on U_0 . Therefore, the genus of C_1 is the same as the genus of C_0 .

Remark that the twin paths γ_1, γ_2 that we cut in ω_0 produce a pair of paths in C_1 . After inverting their orientation we get a pair of twin paths $\tilde{\gamma}_1, \tilde{\gamma}_2$ for ω_1 . Cutting and pasting the twin paths $\tilde{\gamma}_1, \tilde{\gamma}_2$ in ω_1 we obtain (C_0, m_0, ω_0) back. Therefore the inverse surgery is naturally defined in the same manner. We will use this inversion of surgery especially when the base point of the twin paths is a node.

Suppose γ_1, γ_2 are twin paths of a stable form ω_0 starting at a point q and ending at points q_1 and q_2 respectively. Suppose that only the starting point and endpoints can be zeros or nodes of ω_0 . We are going to describe the structure of the twin paths $\tilde{\gamma}_1, \tilde{\gamma}_2$ of the form ω_1 in some cases. Denote by \tilde{q} the starting point of $\tilde{\gamma}_i$ and \tilde{q}_i the final endpoint of $\tilde{\gamma}_i$.

Example 1: Suppose q , q_1 and q_2 are pairwise distinct points and q_1 and q_2 lie in the same component of C_0^* . Then the same is true for \tilde{q} , \tilde{q}_1 and \tilde{q}_2 (see Figure 5).




FIGURE 5. Cut and paste along twin paths contained in one part of the stable curve

Comparing the local degrees of the branched coverings at each point we get:

$$\text{ord}_{\tilde{q}}(\omega_1) + 1 = (\text{ord}_{q_1}(\omega_0) + 1) + (\text{ord}_{q_2}(\omega_0) + 1);$$

$$\text{ord}_q(\omega_0) + 1 = (\text{ord}_{\tilde{q}_1}(\omega_1) + 1) + (\text{ord}_{\tilde{q}_2}(\omega_1) + 1).$$

By taking appropriate combinations of twins and zeros we can do several types of changes to the zero divisor. For instance, if q is a simple zero and q_1, q_2 are regular points, the surgery simply changes the position of the simple zero. We can also **split a multiple zero**, meaning that two or three zeros of ω_1 are produced from a single zero of ω_0 . It suffices to take any pair of twins starting at that zero with distinct endpoints.

The inverse operation of the latter will be referred to as **joining** two or three different zeros in one zero. This is the case, for instance, when two twins leaving a simple zero have distinct endpoints and at least one of them is a zero.

Example 2: Suppose that γ_1 and γ_2 leave q following different branches of C_0 at q . Then q is a node and there are three interesting sub-cases that will be referred to as **smoothing of a node**:

- (1) If q_1, q_2 and q are pairwise distinct, then $\tilde{q}_1 = \tilde{q}_2 \neq \tilde{q}$, none of them is a node, and they are zeros of ω_1 . Each $\tilde{\gamma}_i$ joins the same pair of distinct zeros of ω_1 (see Figure 6).




FIGURE 6. Cut and paste along twin paths contained in different branches of a node to smoothen the node and obtain a couple of twins between zeros

- (2) If $q_1 = q \neq q_2$ and γ_1 does not come back to q through a different branch of C_0 , then \tilde{q} is not a node of ω_1 and $\tilde{q} = \tilde{q}_1 = \tilde{q}_2$, i.e. both $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ are closed loops based at a zero of ω_1 (see Figure 7). Moreover, by considering the intersection of the twins with a neighbourhood of \tilde{q} we observe that among the oriented segments there are

two consecutive segments in the cyclic order with the same sign (either both enter or both leave the singularity).




FIGURE 7. γ_1 is a closed path that leaves and arrives to the node through one of the branches. After the surgery, there is no node, and a couple of closed twin paths based at a zero \tilde{q} .

- (3) If $q_1 = q \neq q_2$ and γ_1 returns to q through the other branch of the node, then the conclusion is the same as in case (2). The only difference with that case is that the cyclic order changes (see Figure 8).




FIGURE 8. γ_1 is a closed path based at the node. Its starting segment and end segment lie in different branches. After the surgery, there is no node, and a couple of closed twin paths based at a zero \tilde{q} .

The inverse surgeries of those examples allow to construct stable forms on nodal curves starting from certain stable forms on smooth curves. In Figure 9 we depict an example in genus two joining a form in the minimal stratum with a form with a node from the point of view of the description of a form as a translation surface.

In figure 10 we provide an example of combinatorics of a pair of closed twins whose Schiffer variation does not lead to a nodal curve.

For future reference we state:

Lemma 5.1. *If a pair of twin paths based at a zero share starting point and endpoint then either the cut and paste surgery produces a stable form with an additional node, or both start and endpoint coincide, and in the cyclic order at that point, any pair of consecutive oriented segments of the twins have opposite sign (as in Figure 10).*




FIGURE 9. The dotted lines represent pairs of twins in the smooth genus two surface. After the Schiffer variation along them we obtain a stable form with a node.




FIGURE 10. The Schiffer variation along a pair of closed twins as in Lemma 5.1 exchanges the relative position of β_2 and β_3 , but no node appears.

Proof. The cases with a node correspond to the inverse surgeries of the smoothing of a node (Example 2). The last case described in the lemma does not produce a node, since the base point obtained after the surgery does not separate the germ of surface in two connected components. \square

The surgery can be generalized to families of twin paths. We call a family of paths $\Gamma = \{\gamma_1, \dots, \gamma_k\}$ a family of twin paths for a form ω_0 if every pair forms a pair of twin paths for ω_0 . The cut and paste surgery associated to the family Γ cuts each γ_i producing two sides of each cut, and glues each to its adjacent twin side. When Γ is maximal and all points in the paths are regular except for the initial point, that is a zero of order $k - 1$, the surgery provides a deformation of the form in its substratum. We can use the families of twins to **join** zeros of any order:

Lemma 5.2. *Let $\Gamma = \{\gamma_1, \dots, \gamma_k\}$ be a maximal family of twin paths starting at a zero of order $k - 1$ of a stable form ω_0 . Suppose that all points in the paths are regular except for their common starting point and the endpoint of γ_1 , that is a zero of ω_0 distinct from the starting point. Then the form ω_1 obtained from the cut and paste surgery associated to Γ has less zeros than ω_0 (See Figure 11 for an example)*

Proof. After the surgery all the endpoints of the initial twins are glued together to form a single zero. The angle between any pair of adjacent twins is 2π . Therefore the starting point




FIGURE 11. On the left a maximal family of twin paths at a double zero. Both endpoints of γ_1 are zeros. The maximality implies that, after the Schiffer variation, the points \tilde{q}_j have angle 2π (the circle around them represents this angle). The obtained form has one zero less.

of the twins splits into k regular points after the surgery. All other glued points are regular and the conclusion follows. \square

Since the restriction of a family of twin paths parametrized by $[0, 1]$ to some interval starting at time zero is still a family of twin paths, the previous construction gives a stable form for each $t \in [0, 1]$. We thus obtain a parametrized family $\{(C_t, \omega_t)\}_{t \in [0, 1]}$ of stable forms. Moreover, at any $t \in (0, 1)$ there are two natural families of twin paths on (C_t, ω_t) . On the one hand there is the family that inverts the surgery from (C_0, ω_0) to (C_t, ω_t) ; on the other, there is the family of twin paths that correspond to the segments of the original twin paths parametrized by $[t, 1]$. The cut and paste surgery applied to the latter family of twin paths on (C_t, ω_t) produces the same stable form (C_1, ω_1) . With this description at hand, items (ii) and (iii) of Example 2 are just a concatenation of Example 1 and the inverse of item (i) in Example 2. To compare the periods of the constructed stable forms we need to produce a (homological) marking m_t on each C_t , starting from a marking m_0 on C_0 . It suffices to do that under the hypothesis of Example 1 and of the inverse of item (i) in Example 2.

In the case of Example 1 (see Figure 5), the neighbourhood U_0 of the twins $\gamma_1 \cup \gamma_2$ can be taken as a topological disc. By construction there is a natural identification between $C_0 \setminus U_0$ and $C_1 \setminus U_1$. The map $\partial U_0 \rightarrow \partial U_1$ has a unique extension up to isotopy to a homeomorphism $\overline{U_0} \rightarrow \overline{U_1}$. The gluing of those two identifications produces a homeomorphism $C_0 \rightarrow C_1$ that induces, by postcomposing it at the homology to m_0 , a marking m_1 on C_1 . With this marking it is readily verified that

$$\text{Per}(C_0, m_0, \omega_0) = \text{Per}(C_1, m_1, \omega_1).$$

In the case of the inverse of item (i) in Example 2, we have the couple of twins $\tilde{\gamma}_1, \tilde{\gamma}_2$ that join two distinct zeros of a stable form (C_1, ω_1) on a smooth curve C_1 (the component they belong to). The neighbourhood U_1 is the annulus depicted in Figure 6 surrounding the twins. The neighbourhood U_0 is a neighbourhood of the node that contains both twins $\gamma_0 \cup \gamma_1$. There is a natural identification $C_1 \setminus U_1 \rightarrow C_0 \setminus U_0$. Each boundary component of U_1 is sent to the boundary of a disc contained in one of the branches of the node, giving an orientation preserving homeomorphism $\partial U_1 \rightarrow \partial U_0$. By arguing as before with each side of the pair of twins, up to isotopy there is a unique collapse $\overline{U_1} \rightarrow \overline{U_0}$ that sends the twins to the node and has the given values on the boundary. If there is a marking m_1 on C_1 , it can be postcomposed

to the action of the collapse $C_1 \rightarrow C_0$ on homology, giving a homological marking m_0 of C_0 . Again, the equality $\text{Per}(C_0, m_0, \omega_0) = \text{Per}(C_1, m_1, \omega_1)$ is easily verified.

With the given markings the parametrized family produces a map $[0, 1] \rightarrow \Omega\bar{\mathcal{S}}_g$ whose composition with the projection $\Omega\bar{\mathcal{S}}_g \rightarrow \bar{\mathcal{S}}_g$ to the space of marked curves $t \mapsto (C_t, m_t) \in \bar{\mathcal{S}}_g$ is continuous by construction. On the other hand we claim that there exists a local trivialization of the fiber bundle $\Omega\bar{\mathcal{S}}_g \rightarrow \bar{\mathcal{S}}_g$ around the point (C_t, m_t) where isoperiodic sets are contained in the constant sections of the local trivialization. Indeed, $\ker m_t$ is isotropic and we can find a symplectic basis $\{a_i, b_i : i = 1, \dots, g\}$ of $H_1(\Sigma_g)$ such that $\ker m_t$ is contained in the Lagrangian generated by a_1, \dots, a_g . The map $\Psi : \Omega U \rightarrow U \times \mathbb{C}^g$ defined by

$$\Psi(C, m, \omega) = ((C, m), \int_{m(a_1)} \omega, \dots, \int_{m(a_g)} \omega)$$

is continuous thanks to Lemma 4.39. It preserves the fibers and it is also linear in restriction to each fiber. This restriction is injective since every form satisfying $\Psi(C, m, \omega) = (C, m, 0, \dots, 0)$ has zero residues at all non-separating nodes, hence at all nodes. In section 6.1 we will see that the condition implies that $\text{vol } \omega = 0$ and this volume is also the sum $\sum \text{vol}(\omega_i)$ where $\text{vol}(\omega_i)$ denotes the (positive) volume of the restriction ω_i of ω to the part C_i of C . Hence $\text{vol}(\omega_i) = 0$ for all i . However a holomorphic form ω_i on a smooth curve C_i has zero volume if and only if it is the zero form. Hence, the only possibility is that ω is zero everywhere. The map Ψ is therefore a trivialization of the bundle $\Omega U \rightarrow U$. In this local trivialization the fibers of Per are contained in the constant horizontal sections, which are continuous. Hence, the map $[0, 1] \mapsto \Omega\bar{\mathcal{S}}_g$ is continuous. The same argument works for a family of twin paths.

Definition 5.3. Let $\Gamma = \{\gamma_1, \dots, \gamma_k\}$ be a (not necessarily maximal) family of twin paths for (C_0, m_0, ω_0) parametrized by $t \in [0, 1]$ and set $p = \text{Per}(C_0, m_0, \omega_0)$. The **Schiffer variation** along Γ is the continuous path

$$(45) \quad t \mapsto (C_t, m_t, \omega_t) \in \text{Per}^{-1}(p)$$

where (C_t, m_t, ω_t) is obtained from (C_0, m_0, ω_0) by cutting and pasting the family of twins $\{\gamma_1|_{[0,t]}, \dots, \gamma_k|_{[0,t]}\}$.

In the case of translation structures there is naturally a distinguished family of twin paths, namely those that have image in straight lines in \mathbb{C} . This produces a sub-family of paths in the fibers of Per . However, connectedness by paths in either of the families is equivalent to connectedness in the case of fibers of Per . The advantage of using straight paths is that the candidates to twins are invariant of the associated directional foliation. This allows to have some control on the embedding/intersection properties. In particular, a pair of geodesic segments of leaves of one of the directional foliations leaving the same singularity and having the same length, form a pair of twin paths. Let us analyze with more detail the oriented directional foliations.

5.2. Periodic annuli. The dynamics of each oriented directional foliation \mathcal{G}_θ induced by an abelian differential ω without residues or zero components on a smooth curve C is well known. Indeed, by Maier's Theorem (see [45] or [63]) there exist a finite number of saddle connections, that is, leaves $\gamma_1, \dots, \gamma_n$ such that each γ_i converges to some singular point in the positive direction and to some singular point in the opposite direction. Each component of $C \setminus \cup \gamma_i$ is saturated by \mathcal{F} and is either a **periodic annulus**, i.e. an annulus formed of closed leaves/geodesics, or **minimal**, i.e. each leaf in the component is dense in the component.

The length of all leaves in a periodic annulus is the same and coincides with the length of each of its boundary components. We orient each boundary component with the orientation of the foliation (not that induced by the orientation of the annulus). We call b^+ the boundary component that has the annulus to its right, and b^- the boundary component that has the annulus on its left. Each boundary component of the annulus is identified with an ordered cycle of saddle connections $(\gamma_1^\pm, \dots, \gamma_k^\pm)$ each starting at the singular point where the former ended. The angle that a saddle connection forms with the following is of π to the corresponding side (right for b^+ , left for b^-). Thus the intersection of the annulus with a neighbourhood of a zero of the abelian differential is a (possibly empty) family of sectors of angle π .

We recall an important existence result:

Theorem 5.4 (Masur, [47]). *For any non-zero abelian differential on a smooth curve there exists a periodic annulus.*

5.3. Degeneracy. We will use Schiffer variations to show the following

Proposition 5.5. *Let $g \geq 2$. Given $(C, m, \omega) \in \Omega^* \mathcal{S}_g$ there exists a finite sequence of Schiffer variations joining it to a stable form with one node and no zero components. If ω has a single zero we can suppose that the node separates the stable curve, and has an elliptic component.*

Proof. Let $(C, m, \omega) \in \Omega^* \mathcal{S}_g$. We will apply Schiffer variations to connect it to a point in $\Omega \mathcal{S}_g$ having a configuration of twin paths as in the first alternative of Lemma 5.1: two twin paths joining the same pair of zeros with a pair of non-alternating signs whenever the zeros coincide. In fact the twins that we will find are geodesic saddle connections. The application of the Lemma then allows to connect the latter to a point in the boundary strata via the associated Schiffer variations. Since Schiffer variations are paths in the fibers of $\mathcal{P}er$, the proposition follows.

First we will treat three cases where the boundary of some periodic annulus for ω provides the twins. In those three cases the Schiffer variation leads to a stable form on a curve with one separating node and an elliptic component. Denote by b^+ and b^- the boundary components of some periodic annulus A of ω .

Case 1: If b^+ and b^- are closed saddle connections at a zero q of ω , meaning that each is formed by a single saddle connection starting and ending at q . Then they cannot both coincide. Indeed, if they did, the surface would have genus one, which is not the case. Therefore they are different and the extension of the chart of ω at q along the saddle connections sends each of them to a segments of oriented straight line of the same length and direction. This implies that they form a pair of twin paths for ω . To be able to apply Lemma 5.1 it remains to check that there is a consecutive pair of separatrices at q with the same sign. Since the annulus lies to the right of b^+ and to the left of b^- the pair of twin segments leaving the singularity are consecutive in the cyclic order and we conclude. The situation is similar to the one depicted in Figure 9.

Case 2: Suppose one of the saddle connections of b^+ is closed, and coincides with one of the saddle connections of b^- . Call the saddle connection γ . In this case we can find another periodic annulus satisfying the conditions of Case 1. Think of the universal cover of A as an infinite horizontal band $\tilde{A} \subset \mathbb{C}$. In each boundary component of \tilde{A} we have copies of γ and they all point in the same direction (see Figure 12). Choose a copy on each boundary component of the band, and draw the parallel geodesics (straight lines) in the band joining

corresponding points in the chosen copies of γ . By construction, for the foliation given by the direction of the geodesics we get a periodic annulus in ω whose boundary components are each formed by one closed saddle connection, as in Case 1.




FIGURE 12. Case 2: Universal cover of the initial annulus A

Case 3: If b^+ and b^- only pass through one zero q of ω , and at least one of them is formed by two saddle connections. The only case that is left is the case where none of the saddle connections of b^+ and b^- coincide. We focus on the distribution of π -sectors described by the annulus at the zero q (see Figure 13).




FIGURE 13. Case 3: The π -sectors corresponding to the boundary of A at the saddle q are coloured black if they correspond to sectors of b^- and light grey if they correspond to sectors of b^+ . Consecutive sectors of distinct colour have an even number of white π -sectors between them. If they are coloured the same there is an uneven number of white sectors between them.

Some of the π -sectors correspond to b^+ and some to b^- . Draw them in two different colors, say black and grey. By hypothesis the intersections of the closures of any two such sectors are empty, since otherwise we would be in Case 2. In the cyclic order at q there is a pair of such π -sectors of distinct color that are consecutive (among the coloured sectors). Between two such consecutive sectors of different colours there is necessarily an even number of uncoloured π -sectors. Hence, up to changing the orientation of the directional foliation if needed, we can consider that the consecutive saddle connections γ^+ and γ^- that lie consecutively in boundaries of those two sectors of different colors both leave the singularity. If the lengths of γ^+ and γ^- coincide, they are twin paths satisfying all the conditions of the first alternative of Lemma 5.1. Otherwise, we consider the shortest among γ^+ and γ^- and its twin path starting along the other. The Schiffer variation along this pair of twin paths produces a form that has a periodic annulus as in Case 2.

This said we proceed to prove the Proposition by induction on the number of zeros of the form.

Suppose first that ω has one zero. Then all the saddle connections appearing in the boundary of a periodic annulus start and end at the zero. Therefore we fall in one of Cases 1,2 or 3 described above. Since in all cases we reached a stable form on a curve of compact type this proves the final statement of Proposition 5.5.

Suppose that we have proved Proposition 5.5 for all marked forms with n zeros and let us prove it for a form ω with $n + 1$ zeros on a smooth curve C .

Let z_1, \dots, z_{n+1} be the zeros of ω . Up to moving them via Schiffer variations we can suppose that the number

$$d_{\min} = \min_{j \neq k} \{\text{dist}(z_j, z_k)\}$$

is realized by a single pair of zeros, say z_1, z_2 . We can further move the point z_2 to guarantee that $\int_{z_1}^{z_2} \omega$ is not in the countable set $\{\int_a \omega \in \mathbb{C} : a \in H_1(C)\}$.

Consider one of the geodesics γ_1 between z_1 and z_2 that realizes the distance between them. It is a saddle connection. If the zero z_1 has order k there are $k + 1$ oriented geodesics leaving z_1 in the same direction as γ_1 . We extend each until distance d_{\min} . By construction they do not pass through zeros of ω , except for (maybe) at their endpoint, where they can only reach z_2 . Indeed, if either reached z_j with $j \neq 2$ one of the conditions of the definition of z_1, z_2 would be violated.

If one of the twins ends at z_2 we are done by Lemma 5.1. Otherwise the family of twin paths satisfies the hypothesis of Lemma 5.2. We use the Lemma to connect ω to a form with less zeros. The inductive hypothesis then concludes the proof. \square

For future reference we state

Corollary 5.6. *Let $p : H_1(\Sigma_g) \rightarrow \mathbb{C}$ be a homomorphism. Then there exists an abelian differential (C, m, ω) on a smooth curve C satisfying $p = \text{Per}(C, m, \omega)$ if and only if the same is true for some stable form without zero components defined on a singular curve.*

Proof. Suppose first that there exists a marked abelian differential (C, m, ω) on a smooth curve C satisfying $\text{Per}(C, m, \omega) = p$. Then Proposition 5.5 implies that there also exists an abelian differential with no zero component on a marked stable curve with a node, having periods p .

For the converse, suppose that there exists a point in a boundary stratum contained in $\text{Per}^{-1}(p)$. By smoothening the nodes isoperiodically we obtain a form on a smooth curve. Alternatively, we can argue by Theorem 4.37: the local fiber L of Per around this point satisfies that $L \setminus \partial L$ is nonempty. A point in the latter set provides a marked abelian differential on a smooth curve with period homomorphism given by p . \square

6. PERIOD HOMOMORPHISMS OF STABLE FORMS

6.1. The image of the period map $\text{Per}_{g,0}$. Recall from subsection 4.14 that a stable form (C, ω) induces a flat singular metric $\omega \otimes \bar{\omega}$ on C^* with volume

$$\text{vol}(\omega) = \frac{i}{2} \int_C \omega \wedge \bar{\omega}.$$

In particular, $0 \leq \text{vol}(\omega) \leq \infty$ and it is finite if and only if all the residues of ω at the nodes of C are zero. If $\text{vol}(\omega) = 0$ then ω is the zero form.

A second fact more directly related with periods is that whenever ω has no zero components on C , and the set of periods $\Lambda = \{\int_a \omega \in \mathbb{C} : a \in H_1(C)\}$ is a lattice in \mathbb{C} , integration along paths produces a branched covering

$$(46) \quad \pi : C \rightarrow \mathbb{C}/\Lambda \text{ whose primitive degree satisfies } \deg(\pi) = \frac{\text{vol}(\omega)}{\text{vol}(\mathbb{C}/\Lambda)}.$$

In particular if the genus of C is at least 2, $\deg(\pi) > 1$.

Now, by Riemann's relations, the volume of ω can be calculated by using the periods of ω . Indeed, for any symplectic basis $\{a_1, b_1, \dots, a_g, b_g\}$ of $H_1(C)$ we have

$$\begin{aligned} \text{vol}(\omega) &= \int_C \Re \omega \wedge \Im \omega = \frac{i}{2} \int_C \omega \wedge \bar{\omega} = \frac{i}{2} \sum_{j=1}^g \int_{a_j} \omega \int_{b_j} \bar{\omega} - \int_{a_j} \bar{\omega} \int_{b_j} \omega = \\ &= \sum_j \Im \left(\overline{\int_{a_j} \omega} \int_{b_j} \omega \right) \end{aligned}$$

The positivity of both terms implies that if the periods of a form are discrete, they cannot be contained in a real line and thus define a lattice.

We extend the definitions of volume and degree to homomorphisms on symplectic modules:

Definition 6.1. Given a unimodular symplectic \mathbb{Z} -module M and a homomorphism $p : M \rightarrow \mathbb{C}$ we define

- The volume of p as $\text{vol}(p) := \Re(p) \cdot \Im(p)$ where the intersection is on the dual space $\text{Hom}(M, \mathbb{R})$. Up to a choice of a symplectic basis $\{a_j, b_j\}$ of M , the volume can be calculated as

$$\text{vol}(p) = \sum_j \Im(\overline{p(a_j)} p(b_j))$$

- Define the primitive degree of p , denoted $\deg(p)$ as ∞ if $p(M)$ is non-discrete and as

$$\deg(p) = \frac{\text{vol}(p)}{\text{vol}(\mathbb{C}/p(M))} \text{ if } p(M) \subset \mathbb{C} \text{ is discrete.}$$

For any symplectic submodule $V \subset M$ we denote $\text{vol}_p(V) := \text{vol}(p|_V)$

Remark that if $0 < \deg(p) < \infty$, the kernel of p has necessarily co-rank two.

If (C, m, ω) is a homologically marked stable form with zero residues and no zero components on a marked stable curve (C, m) , then the homomorphism $p = \mathcal{P}er(C, m, \omega) \in H^1(\Sigma_g, \mathbb{C})$ has the following properties:

- (H₁) $\text{vol}(p) > 0$ and
- (H₂) If $g \geq 2$, $\deg(p) > 1$.

Haupt proved in [34] that conditions (H₁) and (H₂) are in fact also sufficient for a homomorphism $p : H_1(\Sigma_g) \rightarrow \mathbb{C}$ to be the period of a non-zero abelian differential on a smooth curve. For genera $g = 2, 3$ we have already given an argument in Theorem 3.3. We will give an alternative proof of this theorem in subsection 6.6.

If we allow the non-zero form to have some zero components, there is a case where (H₂) is not satisfied: when there is a single non-zero component and it has genus one.

6.2. Haupt homomorphisms. Condition (H_2) of a homomorphism has other equivalent statements that will be useful, and show that the said exception is unique:

Proposition 6.2. *Let M be a unimodular symplectic module of rank $2g$ and $p : M \rightarrow \mathbb{C}$ be a homomorphism. Suppose $\text{vol}(p) > 0$ and $\Lambda = p(M) \subset \mathbb{C}$ discrete. Then the following are equivalent:*

- (1) $\text{vol}(p) \leq \text{vol}(\mathbb{C}/\Lambda)$;
- (2) $\deg(p) = 1$;
- (3) $\ker p$ is a symplectic submodule of M (of rank $2g - 2$).

Proof. First remark that by virtue of Riemann's relations we have $\text{vol}(p) \in \mathbb{Z} \text{vol}(\mathbb{C}/\Lambda)$. If we assume (1) then the positivity of $\text{vol}(p)$ implies (2). Obviously (2) implies (1). For the proof of (2) \Rightarrow (3), we can normalize p by post-composing it by a real linear map to suppose that $\Lambda = \mathbb{Z}^2 \subset \mathbb{C}$. The kernels before and after this composition coincide. Up to taking a symplectic basis of M , we have $p : \mathbb{Z}^{2g} \rightarrow \mathbb{Z}^2 \subset \mathbb{C}$ a homomorphism of positive volume. Then $\text{vol}(p) = p_1 \cdot p_2$ where $p = (p_1, p_2)$ and $p_i \in (\mathbb{Z}^{2g})^*$. On the other hand $\ker p = \ker p_1 \cap \ker p_2$. Now, the dual of p_i is an element $u_i \in \mathbb{Z}^{2g}$ and we have the equality $p_1 \cdot p_2 = u_1 \cdot u_2$. If $\text{vol}(p) = \text{vol}(\mathbb{C}/\mathbb{Z}^2) = 1$, then u_1, u_2 generate a symplectic submodule of rank two. Its orthogonal is a symplectic submodule of rank $2g - 2$ that coincides with $\ker p$.

Next suppose (3) and let us prove (2). Recall that $\ker p$ has rank $2g - 2$. Since $\ker p$ is a symplectic unimodular submodule of M we can take a symplectic basis $a_1, b_1, \dots, a_{g-1}, b_{g-1}$ of $\ker p$ and complete it to a symplectic basis of M by adding a_g, b_g . Apply Riemann's relations to deduce that $0 < \text{vol}(p) = \Im(p(a_g)\overline{p(b_g)})$. This implies that $p(a_g)$ and $p(b_g)$ generate a lattice; since they also generate Λ , we have $\text{vol}(p) = \text{vol}(\mathbb{C}/\Lambda)$. □

Definition 6.3. Let M be a unimodular symplectic module of rank $2g$ and $p : M \rightarrow \mathbb{C}$ be a homomorphism. We say that p is a Haupt homomorphism if $\text{vol}(p) > 0$ and either $g = 1$ (and $\deg p = 1$) or $g \geq 2$ and $\deg(p) > 1$. The set of Haupt homomorphisms in $H^1(\Sigma_g, \mathbb{C}) = \text{Hom}(H_1(\Sigma_g), \mathbb{C})$ will be denoted by \mathcal{H}_g .

Corollary 6.4. *A homomorphism $p : M \rightarrow \mathbb{C}$ on a unimodular module of rank $2g$ with $\text{vol}(p) > 0$ is not Haupt if and only if $g \geq 2$ and $\ker p$ is a symplectic module of rank $2g - 2$.*

6.3. Homological invariants of isoperiodic boundary components. The image of the extended period map Per_g is contained in $\mathcal{H}_g \cup 0$. The fundamental remark for the proof of Theorem 1.2 is related to the fact that the normalization of some node of a form in $\Omega_0^* \overline{\mathcal{S}}_g$ is again a collection of forms in some $\Omega_0^* \overline{\mathcal{S}}_{h,n}$ with $h < g$ and $n \geq 1$. The periods of those forms are intimately related and are subject to satisfy Haupt's conditions. In more detail:

Suppose c is a simple closed curve in Σ_g that is collapsed by the marking of a stable form $(C, m, \omega) \in \Omega_0^* \overline{\mathcal{S}}_g$ to a node and set $p = \text{Per}(C, m, \omega) \in \mathcal{H}_g$.

If $[c] = 0$ in homology, then the curve is separating and C is a union $C_1 \vee C_2$ of stable curves and the form $\omega = \omega_1 \vee \omega_2$ where $\omega_i \in \Omega_0^*(C_i)$. On the other hand m and the decomposition $H_1(C) = H_1(C_1) \oplus H_1(C_2)$ define a symplectic splitting $V_1 \oplus V_2$ of $H_1(\Sigma_g)$. The map $m_i = m|_{V_i}$ is a marking of $H_1(C_i)$. The periods of (C_i, m_i, ω_i) are precisely $p_i := p|_{V_i}$, which are Haupt homomorphisms of lower genus. Therefore we have a decomposition

$$p = p_1 \oplus p_2$$

of the Haupt homomorphism p into two Haupt homomorphisms. In this case

$$\text{vol}(p) = \text{vol}_p(V_1) + \text{vol}_p(V_2) = \text{vol}(p_1) + \text{vol}(p_2)$$

and each factor of the sum is positive. On the other hand

$$\deg p = \deg p_1 \cdot \left| \frac{\text{Im}(p)}{\text{Im}(p_1)} \right| + \deg p_2 \cdot \left| \frac{\text{Im}(p)}{\text{Im}(p_2)} \right|.$$

If $[c] \neq 0$ then the curve is non-separating and $p([c]) = 0$. The normalization of the node produces a stable curve C_2 of genus $g-1$ equipped with a form $\omega_2 \in \Omega_0^* \bar{\mathcal{S}}_{g-1}$. Let V_1 be a rank two symplectic submodule of $H_1(\Sigma_g)$ that contains $[c]$ and $V_1 \oplus V_2$ its associated symplectic splitting of $H_1(\Sigma_g)$. Then, $m|_{V_2}$ defines a marking of $H_1(C_2)$. The map $p_2 = p|_{V_2}$ corresponds to the periods of (C_2, m_2, ω_2) and is therefore a Haupt homomorphism. In this case the volume of $p_1 = p|_{V_1}$ is equal to zero and we have

$$\text{vol}(p) = \text{vol}(p_1) + \text{vol}(p_2) = \text{vol}(p_2).$$

The positivity of $\text{vol}(p_2)$ is guaranteed by that of $\text{vol}(p)$. However for p_2 to be a Haupt homomorphism we need to verify the primitive degree condition (H_2) .

Remark 6.5. Given any symplectic submodule V of rank two containing a primitive element $a \in \ker p$, there is a natural symplectic isomorphism between V^\perp and $a^\perp/\mathbb{Z}a$. Under this identification $p|_{V^\perp}$ is equal to the map

$$p_a : a^\perp/\mathbb{Z}a \rightarrow \mathbb{C}$$

induced by p on the quotient. In particular, if one is a Haupt homomorphism then so is the other.

Definition 6.6. Let $p : M \rightarrow \mathbb{C}$ be a Haupt homomorphism defined on a symplectic unimodular module. Given a symplectic splitting $V_1 \oplus \cdots \oplus V_k$ of M , we say that it is p -admissible if every $p|_{V_i}$ is a Haupt homomorphism. We say that a primitive element $a \in \ker(p) \setminus 0$ is pinched by p if the induced homomorphism p_a is a Haupt homomorphism.

Remark 6.7. If $p : M \rightarrow \mathbb{C}$ is a Haupt homomorphism, then every primitive element of a symplectic submodule of $\ker p$ is pinched by p .

In Corollary [6.18](#) we will prove inductively that all those decompositions and elements of $\ker p$ appear as induced by marked stable forms.

Factors of rank two of p -admissible splittings and the primitive elements they contain will be important in the sequel.

Definition 6.8. Given a Haupt homomorphism $p : M \rightarrow \mathbb{C}$ define

$$\mathcal{V}_p := \{V \subset M : \text{rank}(V) = 2, V \oplus V^\perp \text{ is a } p\text{-admissible splitting of } M\}.$$

A primitive element $a \in M$ is said to be p -admissible if it belongs to some $V \in \mathcal{V}_p$.

6.4. Volumes of symplectic submodules and p -admissible elements. A necessary condition for a symplectic splitting $V \oplus V^\perp$ to be p -admissible is that $\text{vol}_p(V) \in (0, \text{vol}(p))$. In this subsection we will analyze the possibilities for the volumes for rank two symplectic submodules containing a given element a from an algebraic point of view. The idea behind the

Lemmas is that we can parametrize such submodules by a^\perp . The volume function is then affine with linear part

$$L_a : a^\perp \rightarrow \mathbb{R} \text{ defined by } L_a(c) = \Im(p(c)\overline{p(a)}).$$

Up to an isomorphism $\mathbb{R}^2 \cong \mathbb{C}$, $L_a(c)$ is the projection of $p(c) \in \mathbb{C}$ along the line $\ell = p(a)\mathbb{R}$ to its orthogonal line. Whenever the image is discrete (which amounts to having many elements having p -image in ℓ) there will be none, or a finite number of values of the affine function in the interval $(0, \text{vol}(p))$. In the other cases, the values are dense in \mathbb{R} .

Lemma 6.9. *Let W be a unimodular symplectic module of rank $2g \geq 4$ and $p : W \rightarrow \mathbb{C}$ a non-trivial homomorphism. Let $a \in W \setminus \ker p$ and suppose that one of the following conditions hold:*

- (1) $\text{rank}(a^\perp \cap p^{-1}(\mathbb{R}p(a))) \leq 2g - 3$ or
- (2) *there exists a real line $\ell \neq \mathbb{R}p(a)$ in \mathbb{C} containing 0 and satisfying*

$$\text{rank}(p(W) \cap \ell) > 2.$$

Then for every $\varepsilon_1 < \varepsilon_2$ there exists a symplectic submodule $V \subset W$ of rank 2 such that $a \in V$ and $\varepsilon_1 < \text{vol}_p(V) < \varepsilon_2$. In particular if a is primitive, it is also p -admissible.

Proof. First suppose a is primitive and choose $b \in W$ be such that $a \cdot b = 1$. For each $e \in a^\perp$ define $b' = b + e$. Denote $\alpha = p(a), \beta = p(b)$. The volume of $V = \mathbb{Z}a + \mathbb{Z}b'$ is given by

$$(47) \quad \text{vol}_p(V) = \Im(\beta\bar{\alpha}) + \Im(p(e)\bar{\alpha}).$$

If 1) holds, the form $e \in a^\perp \mapsto \Im(p(e)\bar{\alpha}) \in \mathbb{R}$ has image a submodule of \mathbb{R} of rank

$$\text{rank}(a^\perp) - \text{rank}(a^\perp \cap p^{-1}(\mathbb{R}\alpha)) \geq 2g - 1 - (2g - 3) \geq 2.$$

Therefore its image is dense in \mathbb{R} and we can find the desired e for any given ε 's. On the other hand, 2) implies 1) so the same conclusion holds.

If a is not primitive, it is an integer multiple of some primitive a_1 . If one of the conditions (1) or (2) is valid for a , then it is also valid for a_1 . Since the lemma is valid for a_1 we obtain V of appropriate volume containing a_1 . By construction $a \in V$.

□

Proposition 6.10. *Let W be a unimodular symplectic module of rank $2g \geq 4$, and $p : W \rightarrow \mathbb{C}$ be a homomorphism whose image is not contained in a real line. Suppose that either p is injective or $\text{rank}(p) \geq 5$. Then at least one of the following possibilities occur*

- (1) *there exists an element $a \in W \setminus \ker p$ such that for any pair of real numbers $\varepsilon_1 < \varepsilon_2$, there exists a rank two symplectic submodule $V \subset W$ containing a such that*

$$\varepsilon_1 < \text{vol}_p(V) < \varepsilon_2.$$

- (2) *$g = 2$, and for every real line $l \subset \mathbb{C}$, the preimage $p^{-1}(l)$ is either $\{0\}$ or a Lagrangian submodule of W .*

Moreover, if $g \geq 3$ there exists a submodule $\ker p \subset I \subset W$ of positive co-rank such that the conclusion is true for every primitive $a \in W \setminus I$. If $I = \ker p$ does not have the property, then there exists a unique real line $\ell \subset \mathbb{C}$ such that $\text{rank}(p(W) \cap \ell) > 2$. In this case, the module $I = p^{-1}(\ell)$ does the job.

Proof. We first treat the case $g \geq 3$. Assume that for every real line $l \subset \mathbb{C}$, $p(W) \cap l$ has rank ≤ 2 . Take $a \in W \setminus \ker p$. Then

$$\begin{aligned} \text{rank}(a^\perp \cap p^{-1}(\mathbb{R}p(a))) &\leq \text{rank } p^{-1}(\mathbb{R}p(a)) \leq \text{rank } \ker p + \text{rank}(p(W) \cap \mathbb{R}p(a)) \leq \\ &\leq 2g - 5 + 2 = 2g - 3 \end{aligned}$$

and we conclude by Lemma 6.9. Therefore in this case the conclusion with $I = \ker p$ is valid. If there exists a real line $l \subset \mathbb{C}$ such that $p(W) \cap l$ has rank > 2 therefore item (1) in Proposition 6.10 follows by item (1) of Lemma 6.9. In this case the submodule $I = p^{-1}(l)$ does the job. For the uniqueness of the module I as defined: suppose that there exists $a \in W \setminus \ker p$ and an interval $(\varepsilon_1, \varepsilon_2)$ in \mathbb{R} such that no symplectic submodule $V \subset W$ containing a satisfies $\varepsilon_1 < \text{vol}_p(V) < \varepsilon_2$. Then there exists a real line $l \subset \mathbb{C}$, $p(W) \cap l$ has rank > 2 . On the other hand, no real line $l \neq \mathbb{R}p(a)$ can have this property, since otherwise a would belong to rank two submodules of W of arbitrary volume. Hence the only possibility is that $l = \mathbb{R}p(a)$. The submodule $I = p^{-1}(l)$ is the only of this type that has the desired property.

Next suppose $g = 2$. Then p is injective by assumption. If there exists a real line $l = \mathbb{R}p(w) \subset \mathbb{C}$ with $\text{rank}(a^\perp \cap p^{-1}(l)) = 1$ or $\text{rank}(p^{-1}(l)) > 2$, we can use Lemma 6.9 to find the desired element $a \in W \setminus 0$. Otherwise we have that for every $a \in W \setminus 0$

$$\text{rank}(a^\perp \cap p^{-1}(\mathbb{R}p(a))) = 2 \text{ and } \text{rank } p^{-1}(\mathbb{R}p(a)) \leq 2.$$

By injectivity of p this means that $p^{-1}(\mathbb{R}p(a)) \subset a^\perp$ for every a , so $p^{-1}(\mathbb{R}p(a))$ is a Lagrangian. \square

Corollary 6.11. *Let W be a unimodular symplectic module of rank $2g \geq 6$, and $p : W \rightarrow \mathbb{C}$ be a homomorphism whose image is not contained in a real line such that $\text{rank}(p) \geq 5$. Then, either all elements in $W \setminus \ker p$ are p -admissible or all elements that are not p -admissible have image in a real line $\ell \subset \mathbb{C}$ containing 0 and lie in a submodule of rank $2g - 2$ or $2g - 1$ of W . Consequently for every real line $l \neq \ell$ passing through 0 we have $\text{rank } p^{-1}(l) \leq 2$.*

Example 6.12. In \mathcal{H}_2 there are examples of injective homomorphisms for which the volume of symplectic submodules can only take discrete values. They necessarily correspond to case (2) in Proposition 6.10. These examples correspond precisely to the periods of forms belonging to Hilbert modular invariant submanifolds (see Theorem 1.5). Let W be a rank 4 unimodular symplectic module and the homomorphism $p : W \rightarrow \mathbb{C}$ given on a symplectic basis a_1, b_1, a_2, b_2 by $\alpha_1 = 1, \beta_1 = i\sqrt{D}, \alpha_2 = \sqrt{D}, \beta_2 = i$, where $D \geq 2$ is an integer. Then, direct calculation shows that for any symplectic submodule V of W we have $\text{vol}_p(V) \in \sqrt{D} + \mathbb{Z}$. Even if the possible volumes of symplectic submodules form a discrete set, there are an infinite number of elements in \mathcal{V}_p , all having volumes in a finite set of values.

6.5. Existence of pinched elements for non-injective $p \in \mathcal{H}_g$.

Lemma 6.13. *Suppose V_1 is a symplectic module of rank ≥ 4 , and V_2 one of rank ≥ 2 . Let $p_i : V_i \rightarrow \mathbb{C}$ be a homomorphism for $i = 1, 2$. Suppose p_1 is Haupt and $\text{vol}(p_2) \geq 0$. Then $p = p_1 \oplus p_2 : V_1 \oplus V_2 \rightarrow \mathbb{C}$ is a Haupt homomorphism.*

Proof. We already have $\text{vol}(p) = \text{vol}(p_1) + \text{vol}(p_2) \geq \text{vol}(p_1) > 0$. If p were not Haupt, then $\text{vol}(p) = \text{vol}(\mathbb{C}/\text{Im}(p))$. On the other hand $\text{Im}(p_1) \subset \text{Im}(p)$ are discrete and therefore

$$\text{vol}(\mathbb{C}/\text{Im}(p)) \leq \text{vol}(\mathbb{C}/\text{Im}(p_1)) < \text{vol}(p_1) \leq \text{vol}(p)$$

where the strict inequality comes from the Haupt condition for p_1 .

□

Lemma 6.14. *Let $p : V \rightarrow \mathbb{C}$ be a homomorphism on a rank $2g \geq 6$ unimodular symplectic module V and suppose $W \subset \ker p$ is a symplectic submodule of rank at most $2g - 4$. Then p is Haupt if and only if $p|_{W^\perp}$ is Haupt. If both are Haupt, then*

- (1) *all primitive elements in W are pinched by p ;*
- (2) *a primitive $a \in \ker(p) \cap W^\perp$ is pinched by p if and only if it is pinched by $p|_{W^\perp}$.*

Whenever p is Haupt and $\deg(p) < \infty$ there exists such a W of (maximal) rank $2g - 4$.

Proof. We have that $V = W \oplus W^\perp$ is a symplectic splitting and $\text{vol}_p(W) = 0$. Hence $\text{vol}_p = \text{vol}_p(W^\perp)$ is positive as soon as one of them is. As for the second of Haupt's conditions, remark that $\ker p$ is a symplectic submodule of rank $2g - 2$ if and only if $\ker p \cap W^\perp$ is a symplectic submodule of rank $\text{rank}(W^\perp) - 2$. By Proposition 6.2 we have that under the positivity of the volume hypothesis, p is not Haupt if and only if $p|_{W^\perp}$ is not Haupt.

To prove item (1), we remark that for any $a \in W$, p_a is a Haupt homomorphism if and only if $p|_{W^\perp}$ is. As for item (2), take $a \in \ker p \cap W^\perp$ and $W_1 \subset W^\perp$ a symplectic submodule of rank two containing a . Take the splitting $W^\perp = W_1 \oplus W_2$. If a is pinched by $p|_{W^\perp}$, then $p|_{W_2}$ is a Haupt homomorphism. By Lemma 6.13 $p|_{W_2 \oplus W}$ is also a Haupt homomorphism which implies that a is pinched by p . If a is not pinched by $p|_{W^\perp}$ then $W_2 \cap \ker p$ is a symplectic submodule of rank $2g - 6$ and thus $\ker p \cap W_1^\perp$ is a symplectic submodule of rank $2g - 4$. This implies that p_a is not Haupt, and thus a is not pinched by p .

Using the normal form for homomorphisms of finite primitive degree, see Lemma 9.1 proved in Appendix 9, we find a rank $2g - 4$ symplectic submodule $W \subset \ker p$. □

Lemma 6.15. *Let $g \geq 2$ and $p \in \mathcal{H}_g$ be a Haupt homomorphism with $\ker p \neq 0$. Then either all primitive elements of $\ker p$ are pinched by p , or, $g \geq 3$, $\ker p$ has rank $\geq 2g - 3$ and contains a symplectic submodule of rank $2g - 4$ (whose primitive elements are pinched by p).*

Proof. Suppose that there exists a primitive element $a \in \ker p$ such that p_a is not Haupt and take a rank two symplectic submodule V containing a . Then $p(V) \neq 0$ and $0 < \text{vol}(p) = \text{vol}_p(V) + \text{vol}_p(V^\perp) = \text{vol}_p(V^\perp)$.

If $g = 2$ then V^\perp has rank two, and p_{V^\perp} is a Haupt homomorphism. A contradiction. Thus in genus $g = 2$ every $a \in \ker p$ satisfies that p_a is Haupt, therefore pinched by p .

If $g \geq 3$, then $p|_{V^\perp}$ is of positive volume and primitive degree one, therefore contains a symplectic submodule $W \subset \ker p$ of rank $2g - 4$. Moreover $W \oplus \mathbb{Z}a \subset \ker p$. □

6.6. Alternative proofs of Haupt's Theorem. In this section we provide an alternative proof of Haupt's Theorem in [34].

Theorem 6.16 (Haupt's Theorem). *A character $p \in H^1(\Sigma_g, \mathbb{C})$ is the period of some marked abelian differential on a smooth curve if and only if it is a Haupt homomorphism.*

Proof. For genus one and $p \in \mathcal{H}_1$ the condition $\text{vol}(p) > 0$ implies that p is injective and the image of p is a lattice $\Lambda \subset \mathbb{C}$. The abelian differential dz on \mathbb{C}/Λ has periods p .

For genus 2 (and even three) we can apply Theorem 3.3.

For $g \geq 3$, we can proceed by induction. Suppose Haupt's Theorem is true for all genera up to $g - 1 \geq 2$ and take $p \in \mathcal{H}_g$.

Case 1: If $\ker p = 0$. By Proposition 6.10 applied for $g \geq 3$ we deduce that there exists a p -admissible splitting $V_1 \oplus V_2$. The restriction $p_i = p|_{V_i}$ is an injective Haupt homomorphism. By inductive hypothesis we can realize p_i as the periods of a marked abelian differential on a smooth curve (C_i, m_i, ω_i) . The marked nodal form $(C_1 \vee C_2, m_1 \oplus m_2, \omega_1 \vee \omega_2)$ has period p , no zero components and induces the splitting $V_1 \oplus V_2$.

Before proceeding to the proof of the non-injective case, we state a well-known lemma from the theory of Riemann surfaces:

Lemma 6.17. *Let (C, ω) be a non-zero abelian differential on a smooth curve of genus $g \geq 1$. For any $z \in \mathbb{C}$ there is an immersed arc δ in C so that $\int_\delta \omega = z$. Moreover, if $g \geq 2$, the arc δ can be chosen to be embedded with distinct endpoints.*

Proof. Let \tilde{C} be the universal cover of C and $\tilde{\omega}$ the lift of ω to \tilde{C} . The map $I : \tilde{C} \rightarrow \mathbb{C}$ defined by $x \mapsto \int_{x_0}^x \tilde{\omega}$ is holomorphic, non-constant and equivariant with respect to the action of $\pi_1(C)$ on \mathbb{C} by translations under elements of $\Lambda = \{\int_\gamma \omega : \gamma \in \pi_1(C)\}$. The image of I is therefore open and invariant by Λ . We claim that the image is \mathbb{C} . Indeed, if Λ is dense we are done. If Λ is discrete, it is a lattice and the map I induces a branched covering $C \rightarrow \mathbb{C}/\Lambda$ onto a torus. The equivariance then tells us that the image of I is the universal cover \mathbb{C} of the torus. The only other possibility is that the closure of $\Lambda \subset \mathbb{C}$ is isomorphic to $\mathbb{R} \times \mathbb{Z}$. Up to post-composing I with a real linear map we can suppose that $\bar{\Lambda} = \mathbb{R} + \mathbb{Z}i$. The imaginary part $\Im I$ induces a surjective map $C \rightarrow \mathbb{R}/\mathbb{Z}$. Therefore the image of I covers a neighbourhood U of the imaginary axis. The Λ invariant set containing U is then \mathbb{C} .

Let $z \in \mathbb{C}$ be the chosen point. By the surjectivity of I , there exists an immersed arc δ in C , starting from x_0 , such that $\int_\delta \omega = z$. If $g \geq 2$ we can require x_0 to be a zero of ω . In this case the map I is a branched covering of degree > 1 near x_0 . Thus, if the endpoints of both δ coincide with x_0 , we can move them a little so that $\int_\delta \omega$ does not change and they become distinct. Now, since δ is an arc, via an homotopy relative to endpoints we can eliminate all self-intersections so that δ becomes embedded. \square

Case 2: If $\ker p \neq 0$. Use Lemma 6.15 to consider a primitive element $a \in \ker p$ such that p_a is Haupt. Choose $b \in H_1(\Sigma_g)$ such that $a \cdot b = 1$, define $V_1 = \mathbb{Z}a \oplus \mathbb{Z}b$, $V_2 = V_1^\perp$, and $p_i = p|_{V_i}$. By construction p_2 is a Haupt homomorphism. By inductive hypothesis, let ω_2 be an abelian differential on a smooth curve of periods p_2 . Since $g - 1 \geq 2$, by Lemma 6.17 applied to ω_2 we can find an embedded arc β with distinct endpoints in ω_2 such that $\int_\beta \omega_2 = p(b)$. Glue the endpoints and mark the obtained nodal curve to guarantee that the period homomorphism of the stable form is p and the class a is pinched to the node.

In either case smoothing the node produces a form of period p on a smooth curve. \square

Kapovich gave a proof of Haupt's Theorem for genus $g \geq 3$ as a corollary of Proposition 3.10 in [42]. Indeed, he remarked that the image of the map Per_g is an open set in $H^1(\Sigma_g, \mathbb{C})$ invariant by the action of $\text{Sp}(2g, \mathbb{Z})$. For genus $g \geq 3$ he deduced that it is the set \mathcal{H}_g by finding forms with periods in each of the closed invariant sets defined by Proposition 3.10, except of course, for the collapse of $g - 1$ handles.

Using the constructions of Cases 1 and 2 of the proof of Haupt's Theorem we can easily construct stable forms with more nodes and given period map:

Corollary 6.18. *Let $p : H_1(\Sigma_g) \rightarrow \mathbb{C}$ be a Haupt homomorphism and $H_1(\Sigma_g) = V_1 \oplus \dots \oplus V_k$ a p -admissible splitting. Then $k \leq g$ and there exists a stable form in $\Omega_0^* \bar{\mathcal{S}}_g(p)$ with*

$k - 1$ separating nodes such that each $p|_{V_j}$ corresponds to the periods of a component of its normalization. Moreover, if we choose up to $g - k \geq 0$ classes $a_j \in \ker p \setminus 0$, each belonging to a distinct V_j and such that a_j is pinched by $p|_{V_j}$, then there exists a stable form as before that shares the separating nodes with the form of the previous case, and has non-separating nodes that pinch precisely the chosen classes a_j .

6.7. Non-admissible splittings for injective $p \in \mathcal{H}_g$ and disconnected covers. Since the primitive degree condition (H_2) of a Haupt homomorphism is automatically satisfied for injective $p \in \mathcal{H}_g$, a symplectic submodule $V \subset H_1(\Sigma_g)$ induces a p -admissible splitting $V \oplus V^\perp$ if and only if the intermediate volume condition $0 < \text{vol}_p(V) < \text{vol}(p)$ is satisfied. Using the bijection between symplectic splittings and boundary components of the ambient space of compact type given in section 4.7 we prove

Proposition 6.19. *Let $g \geq 2$. If $p \in \mathcal{H}_g$ is injective, then $\Omega\bar{\mathcal{S}}_g^c(p)$ cuts an infinite number of boundary strata of $\Omega\bar{\mathcal{S}}_g^c$ and also avoids an infinite number of them.*

Proof. If p falls into case (1) of Proposition 6.10, for each choice of interval $(\varepsilon_1, \varepsilon_2)$ in \mathbb{R} we can construct an example of symplectic splitting $V \oplus V^\perp$ of $H_1(\Sigma_g)$ with V of rank two and $\text{vol}_p(V) \in (\varepsilon_1, \varepsilon_2)$ by using that proposition. If the interval belongs to $(0, \text{vol}(p))$ the splitting is p -admissible. If it belongs to $\mathbb{R} \setminus [0, \text{vol}(p)]$ it is not p -admissible. Since two splittings having factors of different volumes are different and an infinite number of disjoint intervals can be chosen, we conclude.

If p falls into case (2) of Proposition 6.10 we have that the genus $g = 2$. Haupt's Theorem guarantees that $\text{Per}^{-1}(p)$ contains a stable form of period p on a smooth curve. By Corollary 5.6, $\Omega\bar{\mathcal{S}}_g(p)$ contains a point in some boundary stratum of codimension one. It corresponds necessarily to a p -admissible splitting $H_1(\Sigma_g) = V \oplus V^\perp$ with factors of rank two. Furthermore, the detailed description of the isoperiodic set of marked stable forms associated to p can be found in [50, Theorem 1.2, p. 2274]. Among other properties we find an infinite number of elements defined over nodal curves with a (necessarily separating) node, and inducing distinct p -admissible splittings $H_1(\Sigma_g) = V \oplus V^\perp$ with factors of rank two.

To construct splittings that are not p -admissible we take any p -admissible splitting $V \oplus V^\perp$ with $V = \mathbb{Z}a \oplus \mathbb{Z}b$ and $c \in a^\perp$ with $p(c) \notin \mathbb{R}p(a)$. For each $k \in \mathbb{Z}$ consider $V_k = \mathbb{Z}a \oplus \mathbb{Z}(b + kc)$. Then only for a finite number of k 's we have $\text{vol}_p(V_k) \in (0, \text{vol}(p))$. Since the volumes of the V_k 's are all different, we have found an infinite family of distinct splittings $V_k \oplus V_k^\perp$ that are not p -admissible.

Any p -admissible splitting is realized by some singular stable form via Corollary 6.18. □

Corollary 6.20. *The generic fiber of the period map on $\Omega\mathcal{T}_2$ is disconnected.*

Proof. By [54] there is a free generating family of the Torelli group \mathcal{I}_2 that can be thought in Siegel space as the family of cycles around every boundary component of Torelli space in Siegel space. These boundary components are parametrized by symplectic splittings $V \oplus V^\perp$. Let $p \in \mathcal{H}_2$ be injective (which is generic). From the proof of Theorem 3.3, $\text{Per}^{-1}(p)$ is a slice of dimension one of Torelli space that is isomorphic to $\mathbb{D} \setminus B$ where B is the intersection of the slice with the boundary components. Inclusion induces a map $\pi_1(\mathbb{D} \setminus B) \rightarrow \mathcal{I}_2$. By Corollary 6.19 this map is not surjective. Hence the lift of $\text{Per}^{-1}(p)$ to \mathcal{T}_2 is disconnected. □

6.8. Non-admissible splittings for non-injective $p \in \mathcal{H}_g$ and compact type boundary.

Proposition 6.21. *Let $g \geq 2$ and $p \in \mathcal{H}_g$. There exists a p -admissible splitting of $H_1(\Sigma_g)$ if and only if either $g = 2$ or $g \geq 3$ and $\deg(p) > 2$.*

Proof. First remark that if there are no p -admissible splittings, then there do not exist forms of compact type with no zero components and periods p . By Proposition 5.5 there do also not exist forms with a single zero (i.e. in the minimal stratum) of period p .

As was already shown in the beginning of the proof of Lemma 3.14, the image of the minimal stratum contains all points with $\deg(p) = \infty$, so we are left with the cases of finite degree.

For $g = 2$ there are always p -admissible splittings. Indeed, if $\deg(p) = 2$ there cannot exist a form with a single zero of period p , since the local degree of the covering at the zero would be three so we cannot use this argument. However the stable form with a separating node $dz \vee dz$ on $\frac{\mathbb{C}}{\Lambda} \vee \frac{\mathbb{C}}{\Lambda}$ where $\Lambda := p(H_1(\Sigma_g))$ defines a p -admissible splitting. For $\deg(p) \geq 3$ there are forms in the minimal stratum with period p ([44], [8] for constructive proofs).

If $g \geq 3$ and $3 \leq \deg(p) < \infty$ we proceed by induction on the genus to construct p -admissible splittings. There is a symplectic submodule $W \subset \ker p$ of rank two, the map $p|_{W^\perp}$ has the same degree and volume as p . If $\deg(p) \geq 3$ then there is a $p|_{W^\perp}$ -admissible splitting of W^\perp . Adding W to one of the factors produces a p -admissible splitting.

It remains to treat the case $g \geq 3$ and $\deg(p) = 2$. Without loss of generality we can suppose that the image of p is the group of Gaussian integers. Using the normal form of period of finite primitive degree (see Lemma 9.1 proved in Appendix 9) we can find a symplectic basis $\{a_i, b_i\}$ by $p(a_1) = p(a_2) = 1$, $p(b_1) = p(b_2) = i$ and zero elsewhere. In this case $\text{vol}(p) = 2$ and $\deg p = 2$. The p -volume of any symplectic submodule of $H_1(\Sigma_g)$ is an integer. Suppose that there exists a p -admissible splitting $V_1 \oplus V_2$ of $H_1(\Sigma_g)$. The volume of each component is a positive integer. Hence, the only possibility is that each component has volume one and thus $\text{vol}(V_i) = \text{vol}(\mathbb{C}/\text{Im}(p))$. Since $g \geq 3$, one of both factors, say V_1 , has even rank ≥ 4 so $p|_{V_1}$ is not a Haupt homomorphism, contradicting the definition of p -admissible decomposition. \square

Corollary 6.22. *If $p \in \mathcal{H}_3$ has $\deg(p) = 2$ then $\text{Per}_3^{-1}(p) \simeq \mathfrak{S}_2$. Therefore its lift to $\Omega\mathcal{T}_3$ has infinitely many connected components.*

Proof. The proof of Theorem 3.3 together with the fact that there are no p -admissible splittings, imply that $\text{Per}^{-1}(p)$ is isomorphic to Siegel space \mathfrak{S}_2 . By simple connectedness of Siegel space we deduce the final statement. \square

Corollary 6.23. *For $p \in \mathcal{H}_g$ with $\deg(p) \geq 3$ that can be decomposed as a direct sum of primitive degree two homomorphisms on modules of rank at least ten, the set of boundary points of compact type of $\text{Per}^{-1}(p)$ in $\Omega_0^*\mathcal{S}_g$ is disconnected.*

Proof. Let $V_1 \oplus V_2$ be the splitting such that $p_i = p|_{V_i}$ has degree two for $i = 1, 2$. By Proposition 6.21 there are no p_i -admissible decompositions of V_i .

In particular there are no forms of compact type with two separating nodes and periods p such that one of the nodes induces the splitting $V_1 \oplus V_2$ so we cannot approach a point of intersection with another irreducible boundary component of $\text{Per}^{-1}(p)$ of compact type. On the other hand, the intersection of the boundary component with the fiber $\text{Per}^{-1}(p)$ is

homeomorphic to a product of two fibers as in Theorem 1.3 via the normalization map of the node. Hence it is a disconnected set. \square

This corollary indicates that to prove the connectedness of the bordification the boundary of compact type does not solve the problem, since it is not connected.

7. DETECTING CONNECTED COMPONENTS OF ISOPERIODIC FORMS WITH ONE NODE

Attaching and forgetful maps have nice relations with respect to the fibers of period maps. For instance, the restriction of a map of type (31) with $n_1 = n_2 = 1$ to a product of fibers produces a map

$$\Omega\mathcal{S}_{g_1,1}(p_1) \times \Omega\mathcal{S}_{g_2,1}(p_2) \rightarrow \Omega\bar{\mathcal{S}}_{g_1+g_2,0}$$

whose image points have one node and constant periods p that can be decomposed as $p_1 \oplus p_2$. Equivalently, the restriction of a map of type (32) with $n = 0$ to a fiber $\Omega\mathcal{S}_{g-1,2}(p)$ has its image in the set of forms with a unique non-separating node in a fiber of the period map in $\Omega\bar{\mathcal{S}}_{g,0}$ (see Figure 1). In fact, thanks to the characterization given in Corollary 4.20, these maps provide continuous parametrizations of the components of the intersections of fibers of Per with some boundary stratum ΩB_c of stable forms with one node. Remark that the hypothesis of Theorem 1.2 does not apply to isoperiodic spaces of forms *with* some marked points, so to control the number of connected components of the image of the attaching maps, we will first relate isoperiodic sets on spaces of curves with marked points with isoperiodic spaces without marked points.

7.1. Connectedness of the fibers of $\text{Per}_{g,2}$. Recall that the relative homology long exact sequence gives a natural injection

$$0 \rightarrow H_1(\Sigma_g, \mathbb{Z}) \rightarrow H_1(\Sigma_g, q_1, q_2; \mathbb{Z}).$$

Theorem 7.1. *Let $g \geq 1$ and $p : H_1(\Sigma_g, q_1, q_2; \mathbb{Z}) \rightarrow \mathbb{C}$ be a homomorphism such that $0 \neq p_0 = p|_{H_1(\Sigma_g, \mathbb{Z})}$ has degree at least three and $\Omega\mathcal{S}_{g,0}(p_0)$ is connected. Then, $\Omega\mathcal{S}_{g,2}(p)$ is non-empty and connected.*

Since non-generic sub-strata of forms in $\Omega\mathcal{S}_{g,2}(p)$ form analytic subsets, the claim of Theorem 7.1 is equivalent to proving that the open subset

$$\Omega^{SZ}\mathcal{S}_{g,2}(p) = \{(C, m, \omega) \in \Omega\mathcal{S}_{g,2}(p) : \omega \text{ has simple zeros}\}$$

is connected. To prove Theorem 7.1 we will bordify $\Omega^{SZ}\mathcal{S}_{g,2}(p)$ in the complex manifold $\Omega^{SZ}\mathcal{U}_{g,2} \subset \Omega\bar{\mathcal{S}}_{g,2}$ defined in subsection 4.20 by adding the limit points. Thanks to Proposition 4.44 the closure is a smooth complex manifold where the boundary points form a divisor.

Recall that $\mathcal{U}_{g,2} = \text{For}_{g,2}^{-1}(\mathcal{S}_{g,0}) \subset \bar{\mathcal{S}}_{g,2}$ where $\text{For}_{g,2} : \bar{\mathcal{S}}_{g,2} \rightarrow \bar{\mathcal{S}}_{g,0}$ is the map that forgets both marked points and stabilizes the obtained curve. Every point in some boundary stratum in $\mathcal{U}_{g,2}$ has one node and a part of genus zero containing both marked points. To simplify the notations along the proofs we fix g and write $\mathcal{U} := \mathcal{U}_{g,2}$.

The fibration $\Omega^{SZ}\mathcal{U} \rightarrow \mathcal{U}$ of forms having only simple zeros in the smooth genus g part is a complex manifold and the restriction of the period map defined on $\Omega^{SZ}\mathcal{U}$ is holomorphic (see Lemma 4.39).

Given a homomorphism $p : H_1(\Sigma_g, q_1, q_2; \mathbb{Z}) \rightarrow \mathbb{C}$, we define the analytic subset

$$X_p = \{(C, r_1, r_2, m, \omega) \in \Omega^{SZ}\mathcal{U}_{g,2} : \text{Per}_{g,2}(C, r_1, r_2, m, \omega) = p\}.$$

To understand the topology of X_p we will slice it by fibers of the restriction of the forgetful map to $\Omega^{SZ}\mathcal{U}$ introduced in equation (43)

$$(48) \quad \text{For} : \Omega^{SZ}\mathcal{U} \rightarrow \Omega^{SZ}\mathcal{S}_g.$$

The fiber of For over a point $(C, m, \omega) \in \Omega\mathcal{S}_g$ is biholomorphic to the covering of the surface $C \times C$ described in Proposition 4.26

Recall that $p_0 := p|_{H_1(\Sigma_g)}$. Thanks to Lemma 6.17, we have

$$\text{For}(X_p) = \Omega^{SZ}\mathcal{S}_g(p_0).$$

Lemma 7.2. *For any $(C, m_0, \omega_0) \in \Omega^{SZ}\mathcal{S}_g(p_0)$, the analytic subset*

$$X_p \cap \text{For}^{-1}(C, m_0, \omega_0)$$

is a connected singular curve.

Proof. Fix a homology class $\alpha_0 \in H_1(\Sigma_g, \{q_1, q_2\}, \mathbb{Z})$ with boundary $\partial\alpha_0 = q_2 - q_1$. Define the map $f : \text{For}^{-1}(C, m_0, \omega_0) \rightarrow \mathbb{C}$ by

$$f(C', r_1, r_2, m, \omega) = \int_{m_*\alpha_0} \omega \quad \text{where} \quad C' = C \quad \text{or} \quad C' = C \vee \mathbb{P}^1$$

Since $H_1(\Sigma_g, \{q_1, q_2\}, \mathbb{Z})$ is generated by $H_1(\Sigma_g, \mathbb{Z})$ and α_0 , the set $X_p \cap \text{For}^{-1}(C, m_0, \omega_0)$ is equal to the fiber $f^{-1}(z_0)$ with $z_0 = p(\alpha_0) \in \mathbb{C}$. We will actually prove that all fibers of f are connected to conclude the proof.

The function f is a primitive of the holomorphic 1-form $r^*\Omega$ where

- (i) $r = R|_{\text{For}^{-1}(C, m_0, \omega_0)} : \text{For}^{-1}(C, m_0, \omega_0) \rightarrow C \times C$ is the $H_1(\Sigma_g)$ -covering of monodromy

$$(\alpha_1, \alpha_2) \cdot \alpha = \alpha + \alpha_2 - \alpha_1,$$

described in Proposition 4.26 and

- (ii) Ω is the form on $C \times C$ defined by $\Omega = pr_2^*\omega_0 - pr_1^*\omega_0$, where $pr_i : C \times C \rightarrow C$ for $i = 1, 2$ are the projections on the first and second factors respectively.

A result of Simpson [62] shows that either

- (1) the fibers of f are connected, or
- (2) there exists a complex curve E equipped with a holomorphic one form η , and a holomorphic map $h : C \times C \rightarrow E$ with connected fibers such that $\Omega = h^*\eta$.

So it remains to prove the Lemma in the second case.

Since the set of zeros of Ω is finite (it is the square of the set of zeroes of ω_0 in $C \times C$), and that fibers of h are one dimensional subvarieties, the form η does not vanish, hence E is an elliptic curve. So we have $E = \mathbb{C}/\Lambda$ for a certain lattice $\Lambda \subset \mathbb{C}$, and $\eta = dz$ for $z \in \mathbb{C}$ the coordinate.

Notice that the set of absolute periods of ω, Ω and η coincide with Λ ; indeed, it is clear that the set Λ' of absolute periods of ω and Ω coincide and that it is contained in the set of periods of η , which is equal to Λ . Now, assuming for contradiction that Λ' is strictly contained in Λ , the map h could be factored in the form $h = r' \circ h'$ where $h' : C \times C \rightarrow \mathbb{C}/\Lambda'$ and $r' : \mathbb{C}/\Lambda' \rightarrow \mathbb{C}/\Lambda \simeq E$ a covering of degree > 1 . But h' is onto since it is a non constant holomorphic map, so this contradicts the fact that the fibers of h are connected. Hence

$\Lambda' = \Lambda$. This discussion shows that up to changing the coordinate z by a translation if necessary, we have the following commutative diagram

$$(49) \quad \begin{array}{ccc} \text{For}^{-1}(C, m_0, \omega_0) & \xrightarrow{f} & \mathbb{C} \\ \downarrow r & & \downarrow \\ C \times C & \xrightarrow{h} & \mathbb{C}/\Lambda \end{array}$$

Let us fix $z \in \mathbb{C}$ and analyze the restriction

$$(50) \quad r|_{f^{-1}(z)} : f^{-1}(z) \rightarrow h^{-1}(z \bmod \Lambda)$$

Recall from Proposition 4.26 that the covering group $G \simeq H_1(\Sigma_g)$ of r is presented as $G \subset \text{Aut}(H_1(\Sigma, \{q_1, q_2\}, \mathbb{Z}))$ acting on α_0 by $\alpha \cdot \alpha_0 = \alpha_0 + \alpha$. Therefore we have

$$f \circ \alpha = f + p_0(\alpha).$$

Together with the diagram (49), this shows that $r|_{f^{-1}(z)}$ is a $\text{Ker}(p_0)$ -covering. Its monodromy is the composition of the inclusion map $H_1(h^{-1}(z \bmod \Lambda), \mathbb{Z}) \rightarrow H_1(C \times C, \mathbb{Z}) \simeq H_1(C, \mathbb{Z})^2$ with that of the $H_1(\Sigma_g, \mathbb{Z})$ -covering r . The connectedness of $f^{-1}(z)$ is equivalent to the transitivity of the action of the monodromy on the fiber of $r|_{f^{-1}(z)}$.

Picard-Lefschetz theory can be applied to analyze this monodromy. We include a statement of a particular instance of it that is needed here, in Corollary 11.2 of Appendix 11. It shows that the image of $H_1(h^{-1}(z \bmod \Lambda), \mathbb{Z}) \rightarrow H_1(C \times C, \mathbb{Z})$ is the kernel of $h_* =: H_1(C \times C, \mathbb{Z}) \simeq H_1(C, \mathbb{Z})^2 \rightarrow H_1(E, \mathbb{Z}) \simeq \Lambda$. Moreover, the map h at the level of the homology is given by the periods of Ω , namely by $h_* = p_0 \circ (pr_2 - pr_1)$. In conclusion, the monodromy of the $\text{Ker}(p_0)$ -covering $r|_{f^{-1}(z)} : f^{-1}(z) \rightarrow h^{-1}(z \bmod \Lambda)$ acts transitively, and this concludes the proof that $f^{-1}(z)$ is connected, hence Lemma 7.2. \square

Corollary 7.3. *X_p is connected if and only if $\text{For}(X_p)$ is connected.*

Proof of Corollary 7.3. Suppose $\text{For}(X_p) = \Omega^{SZ} \mathcal{S}_g(p_0)$ is connected. By Lemma 7.2, each connected component of X_p is a union of fibers of $\text{For}|_{X_p}$. Suppose $X_p = X^1 \sqcup X^2$ where X^1 and X^2 are disjoint open sets and $X^1 \neq \emptyset$ is connected. Then X^1 is a union of fibers of $\text{For}|_{X_p}$, and so is its complement X^2 . Moreover $\text{For}|_{X_p} : X_p \rightarrow \Omega^{SZ} \mathcal{S}_g(p_0)$ is an open map, since on each fiber there is some point where $\text{For}|_{X_p}$ is a submersion. Then $\text{For}(X_p) = \text{For}(X^1) \sqcup \text{For}(X^2)$. By connectedness of $\text{For}(X_p)$ we deduce $\text{For}(X^2) = \emptyset$, and therefore $X^2 = \emptyset$, which implies that $X_p = X^1$ is connected. The other implication is obvious. \square

When X_p contains a point in some boundary stratum, there are, by Proposition 4.44, irreducible components of X_p strictly contained in the boundary. To deduce the connectedness of the subset of points of X_p that lie outside the boundary (to prove Theorem 7.1) we need to show that taking out those components we still have a connected set. We will apply the following general

Lemma 7.4. *Let X be a connected analytic set on a complex manifold and suppose $X = Y \cup Z$ where both Y and Z are unions of irreducible components of X that satisfy*

- (1) *around each point of $\text{Sing}(X)$ the set X is a normal crossing of two smooth manifolds, one of which lies in Y and the other in Z , and*




FIGURE 14. Real representation for the set X in Lemma 7.4: each plane represents component of Z . Each circle in a component of Z represents a connected component of $\text{Sing}(X)$ that forms a divisor in Y , the closure of $X \setminus Z$.

(2) the intersection of each irreducible component of Z with $\text{Sing}(X)$ is connected.

Then $X \setminus Z \subset Y$ is connected.

Proof. The diagram in Figure 14 helps to understand the ideas behind the proof. The properties of $\text{Sing}(X)$ imply that Y and Z are smooth manifolds. Moreover, each connected component of $\text{Sing}(X)$ is a smooth complex submanifold of Y and of Z that coincides with $Z_i \cap Y$ for some irreducible component Z_i of Z . In particular, the connectedness of $X \setminus Z$ is equivalent to the connectedness of Y .

On the other hand, for X , (and also for the manifolds Y and $\text{Sing}(X)$)) connectedness is equivalent to path connectedness.

Given $y_1, y_2 \in Y \subset X$ we can, by hypothesis, find a path $\gamma : I \rightarrow X$ starting at y_1 and ending at y_2 . If the path lies in Y we are done. Otherwise $I \setminus \gamma^{-1}(Y)$ is a finite family of disjoint segments $I_1, \dots, I_k \subset I$ where $\gamma|_{I_k}$ is a path contained in a unique Z_k with endpoints in $\text{Sing}(X)$. Substitute $\gamma|_{I_k}$ by a path in the connected manifold $Z_k \cap Y \subset \text{Sing}(X)$ joining the same endpoints. The constructed path lies in Y . Therefore, Y is path connected and so is $X \setminus Z$. \square

7.1.1. *Proof of Theorem 7.1.* Recall from subsection 4.20 that the boundary strata in $\Omega^{SZ}\mathcal{U}_{g,2}$ is precisely the disjoint union $\bigsqcup_{\beta \in \mathfrak{b}} \Omega^{SZ}B_\beta$, where $\mathfrak{b} = \{\beta \in H_1(\Sigma_g, q_1, q_2, \mathbb{Z}) : \partial\beta = q_2 - q_1\}$. The boundary strata in X_p are

$$Z_p = \bigsqcup_{\beta \in \mathfrak{b}} \Omega^{SZ}B_\beta(p).$$

Let Y_p be the union of irreducible components of X_p that have some point outside the boundary.

From Proposition 4.44 we deduce that

$$\text{Sing}(X_p) = \bigsqcup_{\beta \in \mathfrak{b}} \Omega^{SZ,1}B_\beta(p).$$

where $\Omega^{SZ,1}B_\beta$ is the substratum of forms where the points corresponding to the node are zeros of the form on both sides of the node. Moreover, the local structure of X_p at a singular point is a normal crossing of two smooth components: one lying in Z_p and the other in Y_p .

Claim 1: X_p is connected.

Proof of Claim 1: Since $\deg(p_0) \geq 3$ and $\text{Per}_{g,0}^{-1}(p_0)$ is connected by hypothesis, so is the Zariski open set $\text{Per}_{g,0}^{-1}(p_0) \cap \Omega^{SZ}\mathcal{S}_{g,0} = \text{For}(X_p)$. Corollary 7.3 then implies that X_p is connected.

Claim 2: Each $\Omega^{SZ}B_\beta(p)$ is a smooth irreducible component of Z_p .

Proof of Claim 2: Proposition 4.44 shows that the set $\Omega^{SZ}B_\beta(p)$ is a smooth manifold. As for connectedness, the attaching map

$$(51) \quad \Omega^{SZ}\mathcal{S}_{g,1} \cong \Omega^{SZ}\mathcal{S}_{g,1} \times \Omega\mathcal{S}_{0,3} \rightarrow \Omega^{SZ}B_\beta$$

that sends a form with simple zeros on a *smooth* genus g curve with a marked point to the form that attaches a zero component of genus zero at the marked point, is a biholomorphism. Define $p_1 = p|_{H_1(\Sigma_{g,1})}$. Then, the set $\Omega^{SZ}B_\beta(p)$ corresponds to $\Omega^{SZ}\mathcal{S}_{g,1}(p_1)$ under this biholomorphism. Remark that the forgetful map $\mathcal{S}_{g,1} \rightarrow \mathcal{S}_g$ is a fiber bundle that is equivalent to the universal curve bundle $\mathcal{CS}_g \rightarrow \mathcal{S}_g$. The restriction of the forgetful map $\mathcal{C}\Omega\mathcal{S}_{g,1} \rightarrow \Omega\mathcal{S}_g$ to $\Omega^{SZ}\mathcal{S}_{g,1}(p_1) \subset \Omega^{SZ}\mathcal{S}_{g,1}$ coincides with the restriction of the universal curve bundle to the open set $\Omega^{SZ}\mathcal{S}_g(p_0) \subset \Omega\mathcal{S}_g$. Since the latter is connected by hypothesis, so is $\Omega^{SZ}\mathcal{S}_{g,1}(p_1) \subset \Omega^{SZ}\mathcal{S}_{g,1}$.

Claim 3: Each $\Omega^{SZ,1}B_\beta(p)$ is a connected set.

Proof of Claim 3: By the biholomorphism (51), the set $\Omega^{SZ,1}B_\beta(p)$ corresponds to the subset $\Omega^{SZ,1}\mathcal{S}_{g,1}(p) \subset \Omega^{SZ}\mathcal{S}_{g,1}$ where the marked point coincides with one of the simple zeros. This set can be interpreted as the multi-section described by the set of $2g - 2$ simple zeros on each fiber of the universal curve bundle

$$\mathcal{C}\Omega^{SZ}\mathcal{S}_{g,0}(p_0) \rightarrow \Omega^{SZ}\mathcal{S}_{g,0}(p_0).$$

Since $\Omega^{SZ}\mathcal{S}_{g,0}(p_0)$ is connected and $\deg p_0 \geq 3$, we deduce from Lemma 3.14 that this multi-section is connected.

From Claims 1, 2 and 3 we deduce that the hypotheses of Lemma 7.4 are satisfied for $X_p = Y_p \cup Z_p$. We deduce from it that $\Omega\mathcal{S}_{g,2}(p) = X_p \setminus Z_p$ is connected.

7.2. Some connected components of isoperiodic sets on curves with one node.

Proposition 7.5. *Let $g \geq 2$, $p \in H^1(\Sigma_g, \mathbb{C})$, and \bar{c} be a Torelli class of a simple separating curve in Σ_g such that $\Omega^*B_{\bar{c}}(p)$ is non empty. Let $H_1(\Sigma_g, \mathbb{Z}) = V_1 \oplus V_2$ the symplectic splitting associated to \bar{c} and denote $p_i = p|_{V_i}$. Suppose that, after a symplectic identification of V_i with $H_1(\Sigma_{g_i}, \mathbb{Z})$ for corresponding g_i 's, the set $\Omega\mathcal{S}_{g_i}(p_i)$ is connected for $i = 1, 2$. Then, $\Omega_0^*B_{\bar{c}}(p)$ is connected as well.*

Proof. First remark that $p_i \neq 0$ and $\Omega\mathcal{S}_{g_i,0}(p_i)$ is non-empty, since the restriction of an element $(C_1 \vee C_2, m, \omega_1 \vee \omega_2) \in \Omega^*B_{\bar{c}}(p)$ to a part $(C_i, m|_{V_i}, \omega_i)$ belongs to that set. By Theorem 7.1 the set $\Omega\mathcal{S}_{g_i,1}(p_i)$ is also non-empty and connected. The natural attaching map

$$\Omega\mathcal{S}_{g_1,1}(p_1) \times \Omega\mathcal{S}_{g_2,1}(p_2) \rightarrow \Omega^*B_{\bar{c}}(p)$$

that attaches the marked curves and the forms at the marked points is a surjective continuous map (see Subsection 4.16). Therefore the image is connected. \square

Proposition 7.6. *Let $g \geq 3$, $p \in H^1(\Sigma_g, \mathbb{C})$, and \bar{c} be a Torelli class of a simple non separating curve in Σ_g such that $\Omega_0 B_{\bar{c}}(p)$ is non empty. Suppose that, after a symplectic identification of $[\bar{c}]^\perp / \mathbb{Z}[\bar{c}]$ with $H_1(\Sigma_{g-1}, \mathbb{Z})$, the primitive degree of $p_{[\bar{c}]}$ is at least three, and $\Omega \mathcal{S}_{g-1}(p_{[\bar{c}]})$ is non-empty and connected. Then, $\Omega_0 B_{\bar{c}}(p)$ is connected as well.*

Proof. Take a representative of \bar{c} and introduce the closed connected oriented topological surface Σ_{g-1} with two marked points $q_1 \neq q_2 \in \Sigma_{g-1}$ defined by cutting Σ_g along \bar{c} and identifying each component of the geometric completion to a point, one defining the point q_1 and the other the point q_2 . Notice that we have a natural isomorphism

$$(52) \quad H_1(\Sigma_{g-1}, q_1, q_2; \mathbb{Z}) \simeq H_1(\Sigma_g, \mathbb{Z}) / \mathbb{Z}[\bar{c}],$$

where $[\bar{c}]$ denotes the homology class of \bar{c} . The inclusion $H_1(\Sigma_{g-1}, \mathbb{Z}) \hookrightarrow H_1(\Sigma_{g-1}, q_1, q_2; \mathbb{Z})$ coming from the long exact sequence of relative homology translates under the isomorphism (52) into the inclusion $[\bar{c}]^\perp / \mathbb{Z}[\bar{c}] \hookrightarrow H_1(\Sigma_g, \mathbb{Z}) / \mathbb{Z}[\bar{c}]$.

Since $[\bar{c}]$ belongs to the kernel of $p : H_1(\Sigma_g, \mathbb{Z}) \rightarrow \mathbb{C}$, this latter defines a period $p_{rel} : H_1(\Sigma_{g-1}, q_1, q_2; \mathbb{Z}) \rightarrow \mathbb{C}$. The attaching map that identifies the two marked points provides an isomorphism

$$\Omega_0 B_{\bar{c}}(p) \simeq \Omega \mathcal{S}_{g-1,2}(p_{rel}).$$

Now, the restriction $(p_{rel})_0$ of p_{rel} to $H_1(\Sigma_{g-1}, \mathbb{Z})$ coincides with $p_{[\bar{c}]}$ up to the given identification. In particular $\deg((p_{rel})_0) \geq 3$ and $\Omega \mathcal{S}_{g-1,0}((p_{rel})_0)$ is (non-empty and) connected. By Theorem 7.1 applied to p_{rel} we deduce that $\Omega \mathcal{S}_{g-1,2}(p_{rel})$ is connected. \square

8. CONNECTEDNESS OF THE BOUNDARY OF A FIBER OF $\mathcal{P}er$

This section will be devoted to show the following

Theorem 8.1. *Let $g \geq 4$ and suppose that Theorem 1.2 is true up to genus $g - 1$. Then for any $p \in \mathcal{H}_g$ with $\deg p \geq 3$ the boundary $\partial \mathcal{P}er^{-1}(p)$ in $\Omega_0^* \bar{\mathcal{S}}_g$ is connected.*

Its proof will be split in two separate parts, depending on whether $p \in \text{Hom}(H_1(\Sigma_g, \mathbb{Z}), \mathbb{C})$ is injective or not.

8.1. Proof of Theorem 8.1 for injective $p \in \mathcal{H}_g$. In the case of injective $p \in \mathcal{H}_g$ all points in $\Omega \bar{\mathcal{S}}_g(p)$ are contained in the smooth manifold $\Omega^* \bar{\mathcal{S}}_g^c$ formed by stable forms without zero components on curves of compact type. Moreover, Corollary 4.42 guarantees that $\Omega \bar{\mathcal{S}}_g(p)$ is a smooth manifold, and the boundary strata $\partial \mathcal{P}er^{-1}(p)$ form a normal crossing divisor. The connectedness of this divisor is equivalent to the connectedness of its dual complex \mathcal{G}_p .

Remark 8.2. If $L \in \text{GL}_2^+(\mathbb{R})$ is an orientation preserving real linear map, the complexes associated to p and to $L \circ p$ are isomorphic. In particular, for any $c \in \mathbb{C}^*$, \mathcal{G}_p and \mathcal{G}_{cp} are isomorphic.

In the rest of this section we will prove the connectedness of the complex \mathcal{G}_p .

Since each component of a boundary stratum of $\text{Per}^{-1}(p)$ lies in a component of a stratum of the boundary of $\Omega\bar{\mathcal{S}}_g^*$ we can define, using the dual boundary complexes, a continuous map of simplicial complexes

$$(53) \quad \mathcal{G}_p \rightarrow \mathcal{C}_g^{\text{sep}}/\mathcal{I}_g.$$

We claim that if p is injective and $g \geq 4$ then so is the map (53) at the level of vertices. Indeed, suppose D_1, D_2 are connected components of $\partial \text{Per}^{-1}(p)$ lying on the boundary component of $\Omega\bar{\mathcal{S}}^*$ corresponding to a symplectic splitting $V_1 \oplus V_2$. Since $\ker p = 0$, $\deg(p|_{V_i}) < \infty$ if and only if $\text{rank}(V_i) = 2$. The inductive hypothesis of Theorem 1.2 and Proposition 7.5 guarantee that there is only one isoperiodic irreducible component of period p in the boundary stratum of $\Omega\bar{\mathcal{S}}_g$ corresponding to $V_1 \oplus V_2$. Hence, $D_1 = D_2$. The connectedness of \mathcal{G}_p is therefore equivalent to that of the image of (53). Two vertices of the image of (53) are said to be equivalent if they lie in the same connected component of the image. Suppose that for $i = 1, 2$, $\{V_i, V_i^\perp\}$ are the modules associated to two vertices. We write $V_1 \sim V_2$ if the vertices lie in the same connected component of the image of (53). In particular $V \sim V^\perp$ for any pair $\{V, V^\perp\}$ representing a vertex in the image. In fact, more generally, if V_1, V_2 are orthogonal submodules of intermediate volume $\text{vol}_p(V_i) \in (0, \text{vol}(p))$ such that

$$0 < \text{vol}_p(V_1) + \text{vol}_p(V_2) < \text{vol}(p) \text{ then } V_1 \sim V_2.$$

Indeed, they are two of the factors of a p -admissible splitting $V_1 \oplus (V_1 \oplus V_2)^\perp \oplus V_2$. By Corollary 6.18, this splitting occurs as the period of a stable form without zero components on a curve with two separating nodes. Moreover it lies in the intersection of the closure of the components of V_1 and V_2 .

We first prove that every vertex given by a p -admissible splitting $V \oplus V^\perp$ is equivalent to another $V_1 \oplus V_1^\perp$ where $V_1 \in \mathcal{V}_p$, i.e. V_1 is of rank two. Indeed, if V is of rank strictly between 2 and $2g - 2$ then $p|_V$ is an injective Haupt homomorphism and by Proposition 6.19 there exists a $p|_V$ -admissible splitting $V_1 \oplus V_2$ of V with a factor V_1 of rank 2. Therefore $V_1 \oplus V_2 \oplus V^\perp$ is also a p -admissible splitting and $V \sim V^\perp \sim V_1$. It remains to connect the vertices corresponding to rank two p -admissible submodules.

Proposition 8.3. *If $g \geq 4$, $p \in \mathcal{H}_g$ is injective and $V, V' \in \mathcal{V}_p$, we have $V \sim V'$. Therefore the image of (53) is connected.*

We will split the proof in several Lemmas :

Lemma 8.4. *Let $g \geq 4$ and $p \in \mathcal{H}_g$ be injective. If $V, V' \in \mathcal{V}_p$ satisfy $V \cap V' \neq 0$, then*

$$V \sim V'.$$

Proof. If $V = V'$ we are done. Suppose that $V \neq V'$

First step: there is a symplectic basis $a_1, b_1, \dots, a_g, b_g$ such that $V = \mathbb{Z}a_1 + \mathbb{Z}b_1$ and $V' = \mathbb{Z}a_1 + \mathbb{Z}(b_1 + m'_2 a_2)$ for a certain integer m'_2 .

Proof of the first step. The intersection $V \cap V'$ is a primitive submodule of H_1 , since both V and V' are primitive. Being of rank 1, we have $V \cap V' = \mathbb{Z}a_1$ with a_1 primitive. Let $b_1 \in V$ (resp. $b'_1 \in V'$) such that $a_1 \cdot b_1 = 1$ (resp. $a_1 \cdot b'_1 = 1$). These elements exist since V and V' are unimodular. The element $b'_1 - b_1$ belongs to a_1^\perp . For a certain integer n , the element

$b'_1 + na_1 - b_1$ is also orthogonal to b_1 . Change b_1 to $b_1 - na_1$ if necessary. We then have that $b'_1 - b_1$ is orthogonal to $V = \mathbb{Z}a_1 + \mathbb{Z}b_1$. Write $b'_1 - b_1 = m'_2 a_2$ where a_2 is a primitive element of V^\perp . Completing a_2 into a symplectic basis $a_2, b_2, \dots, a_g, b_g$ of V^\perp gives the desired statement. \square

Second step: if the periods of $(V + V')^\perp$ do not lie in a real line of \mathbb{C} , there exists a symplectic rank two submodule $W \subset H_1(\Sigma_g)$ such that $V \perp W$, $V' \perp W$ and

$$(54) \quad 0 < \text{vol}_p(W) < \inf(\text{vol}_p(V^\perp), \text{vol}_p((V')^\perp)).$$

In particular, $V \sim W \sim V'$.

Proof of second step. In the coordinates of the first step, we have $(V + V')^\perp = \mathbb{Z}a_2 + X$ where $X := \sum_{k \geq 3} \mathbb{Z}a_k + \mathbb{Z}b_k$. We apply Proposition 6.10 to $p|_X$. If the restriction of p to X belongs to case (1) of that proposition, we are done. If it belongs to case (2), we use the

Lemma 8.5. *Let X be a unimodular symplectic module of rank 4. For every Lagrangian subspace $L \subset X$, there exists a symplectic rank two submodule $Y \subset X$ such that $L \cap Y = \{0\}$.*

Proof. We can assume that $L = \mathbb{Z}a + \mathbb{Z}a'$ is primitive. Let $a_1 = a$ and b_1 be an element of X such that $a_1 \cdot b_1 = 1$. We have $a' = m_1 a_1 + c$ where $c \in (\mathbb{Z}a_1 + \mathbb{Z}b_1)^\perp$ and $m_1 \in \mathbb{Z}$. Up to replacing a' by $a' - m_1 a_1$, we can assume that $m_1 = 0$. Since L is primitive, so is c , so that we can extend the family $a_1, b_1, a_2 = c$ to a symplectic basis of X . The symplectic submodule $Y = \mathbb{Z}(a_1 + b_2) + \mathbb{Z}b_1$ has the desired properties. \square

From now on a Greek letter will denote the period of the corresponding Latin letter. Let $l = \mathbb{R}\alpha_2$, and $L = X \cap p^{-1}(l)$. This space is either $\{0\}$ or a Lagrangian subspace of X since we assume the restriction of p to X is in case (2) of Proposition 6.10. By the preceding Lemma, there exists a symplectic rank two submodule $Y \subset X$ such that $Y \cap p^{-1}(l) = \{0\}$. Let a', b' be a symplectic basis of Y , and let

$$a = a' + Aa_2, \quad b = b' + Ba_2,$$

for some $A, B \in \mathbb{Z}$. We have $a \cdot b = a' \cdot b' = 1$, and the volume of $W = \mathbb{Z}a + \mathbb{Z}b$ is given by

$$\text{vol}_p(W) = \Im((\beta' + B\alpha_2)(\overline{\alpha' + A\alpha_2})) = \Im(\beta'\overline{\alpha'}) + \Im((A\beta' - B\alpha')\overline{\alpha_2}).$$

By construction none of the cycles of Y are mapped by p to an element of the line $l = \mathbb{R}\alpha_2$, so the linear form $(A, B) \in \mathbb{Z}^2 \mapsto \Im(A\beta' - B\alpha')\overline{\alpha_2} \in \mathbb{R}$ is injective, and thus the volume of W can approximate any real value. Since W is orthogonal to both V and V' , choosing W with $\text{vol}_p(W) \in (0, \text{vol}(p))$ gives the solution to step 2. \square

Third step: assume that the periods of $(V + V')^\perp$ lie on a real line $l \subset \mathbb{C}$. Then $V \sim V'$.

Proof of the third step. Thanks to Remark 8.2 it suffices to show that V is equivalent to V' for some w where $w \in \mathbb{C}^*$. By choosing an appropriate w of modulus one we can suppose that $wp((V + V')^\perp) \subset \mathbb{R}$. If we prove Step 3 for the case $l = \mathbb{R}$ we will be done. Recall that $X = \sum_{i \geq 3} \mathbb{Z}a_i + \mathbb{Z}b_i \subset (V + V')^\perp$. Let $c \in X$ and define $V_c := \mathbb{Z}a_1 + \mathbb{Z}(b_1 + c)$. The volume of V_c is given by

$$\text{vol}_p(V_c) = \text{vol}_p(V) + \Im(p(c)\overline{\alpha_1}).$$

If $\alpha_1 \in \mathbb{R}$, then V_c is automatically p -admissible. Otherwise $p(\alpha_1) \notin \mathbb{R}$ and V_c is p -admissible as soon as

$$-\text{vol}_p(V) < \Im(p(c)\overline{\alpha_1}) < \text{vol}_p(V).$$

Now X has rank at least 4 and so has the image of the homomorphism $X \rightarrow \mathbb{R}$

$$c \mapsto \Im(p(c)\overline{\alpha_1}).$$

Therefore, the inequality has an infinite number of solutions $c \in X \setminus \{0\}$. We can even impose a further condition that $c \cdot a_3 = 1$. Define $V'' = V_c$ for such a solution c .

We claim that $V'' \sim V$. Indeed, the space $(V + V'')^\perp$ contains the element b_2 . Observe that the period β_2 of b_2 is not real, since otherwise all the periods of V^\perp would be real, and so we would have $\text{vol}_p(V) = \text{vol}_p(V')$ which contradicts $V \in \mathcal{V}_p$. On the other hand, the submodule $c^\perp \cap X$ has rank ≥ 3 and is contained in $(V + V')^\perp$. Since the periods of X are real, this proves that some periods of $(V + V'')^\perp$ are real. We can thus apply Step 2 to the couple (V, V'') to infer $V'' \sim V$.

To prove that $V'' \sim V'$ we will show that $(V' + V'')^\perp$ does not have all its periods in a line and apply Step 2. Consider the element $b_2 + m'_2(a_1 + a_3)$. It belongs to $(V' + V'')^\perp$. If it has real period, then $b_2 + m'_2a_1$ has also real period, which implies that $(V')^\perp$ has only real periods. This is in contradiction with the fact that V' is p -admissible. Therefore $b_2 + m'_2(a_1 + a_3)$ has non-real period. On the other hand there are elements in $(V' + V'')^\perp \cap X \setminus 0$ and their periods are non-zero real numbers. \square

\square

Lemma 8.4 allows to reduce the equivalence relation \sim on submodules in \mathcal{V}_p to an equivalence relation on the elements that belong to those submodules.

Definition 8.6. Let $p \in \mathcal{H}_g$. Recall that a primitive element $w \in H_1(\Sigma_g)$ is said to be p -admissible if it is contained in some module $V \in \mathcal{V}_p$. Two p -admissible elements w, w' are equivalent and denoted $w \sim w'$ if there exist $V, V' \in \mathcal{V}_p$ containing w and w' respectively such that $V \sim V'$.

The transitivity property of this equivalence relation is proven by the use of Lemma 8.4

In particular, we already know that if $V \cap V' \neq 0$ then any pair of primitive elements in $V \cup V'$ are equivalent.

If V and W belong to \mathcal{V}_p and there exists some elements $v \in V$ and $w \in W$ such that $v \sim w$, then $V \sim W$. Indeed, we can find $V', W' \in \mathcal{V}_p$ such that $v \in V', w \in W'$ and $V' \sim W'$. By Lemma 8.4, $V \sim V'$ and $W \sim W'$, so $V \sim W$.

Let us analyze the p -admissible elements.

Lemma 8.7. Let $p : H_1(\Sigma_g) \rightarrow \mathbb{C}$ be an injective Haupt homomorphism. Given $w_1, w_2, w_3 \in H_1(\Sigma_g)$ such that

- (1) $w_i \cdot w_{i+1} = 1$ for $i = 1, 2$,
- (2) $p(w_3) \notin \mathbb{R}p(w_1)$ and
- (3) for every real line $\ell \subset \mathbb{C}$ containing 0

$$\text{rank}(p^{-1}(\ell) \cap w_1^\perp \cap w_3^\perp) < 2g - 3.$$

Then there exists $w'_2 \in H_1(\Sigma_g)$ such that $w_1 \cdot w'_2 = w'_2 \cdot w_3 = 1$ and $\mathbb{Z}w_1 \oplus \mathbb{Z}w'_2$ and $\mathbb{Z}w'_2 \oplus \mathbb{Z}w_3$ belong to \mathcal{V}_p . Therefore w_1 and w_3 are p -admissible and $w_1 \sim w_3$.

Proof. Write $w'_2 = w_2 + z$ where $z \in w_1^\perp \cap w_3^\perp$. If we show that the image of the map

$$w_1^\perp \cap w_3^\perp \rightarrow \mathbb{R}^2$$

defined by $z \mapsto (\text{vol}_p(\mathbb{Z}w_1 \oplus \mathbb{Z}(w_2 + z)), \text{vol}_p(\mathbb{Z}(w_2 + z) \oplus \mathbb{Z}w_3))$ has a point in the square $(0, \text{vol}(p)) \times (0, \text{vol}(p))$ we will be done. The previous map is affine, with linear part

$$\varphi(z) = (\Im(p(z)\overline{p(w_1)}), \Im(p(w_3)\overline{p(z)})).$$

Since $p(w_1)$ and $p(w_3)$ are not \mathbb{R} -colinear, $\text{Ker}(\varphi) = 0$ and therefore $\text{rank}(\text{Im}\varphi) = 2g - 2$. The topological closure of $\text{Im}\varphi$ in \mathbb{R}^2 is either \mathbb{R} , $\mathbb{Z} \times \mathbb{R}$ or \mathbb{R}^2 . Suppose it is not \mathbb{R}^2 . Then there exists a submodule $H \subset w_1^\perp \cap w_3^\perp$ such that $\varphi(H) \subset \ell \subset \mathbb{R}^2$ for some real line ℓ passing through the origin and $\text{rank} H \geq (2g - 2) - 1 = 2g - 3$. Write $\ell = \{(x, y) : \alpha x + \beta y = 0\}$ and then for each $z \in H$,

$$\Im(p(z)\overline{(\alpha p(w_1) - \beta p(w_3))}) = 0.$$

Hence $p(H) \subset \mathbb{R}(\alpha p(w_1) - \beta p(w_3))$ is a submodule of rank at least $2g - 3$ and we reach a contradiction with the rank hypothesis. \square

The role played by the rank of p on lines in Lemma 6.9, Lemma 8.7 and Proposition 6.10 makes it useful to introduce the following

Definition 8.8. Given a homomorphism $p : W \rightarrow \mathbb{C}$ from a \mathbb{Z} -module W we define its **line rank** as

$$r(p) = \max_{a \in \mathbb{S}^1} \text{rank}_{\mathbb{Z}}(p^{-1}(a\mathbb{R})).$$

Remark that if W is symplectic and $\text{vol}(p) > 0$ then $r(p) < 2g$. Also, if p is injective and $r(p) > g$, then the maximum is attained by a unique real line $\ell_{\max} \subset \mathbb{C}$ containing 0.

As far as Lemma 8.7 is concerned, the rank condition is automatically satisfied if $r(p) < 2g - 3$. For $g \geq 4$ and $r(p) \geq 2g - 3$ we have to check the rank condition only for ℓ_{\max} . In case $r(p) = 2g - 1$ the rank condition cannot be attained. In the other two cases it depends on whether there exists a rank two submodule of $w_1^\perp \cap w_3^\perp$ with periods outside ℓ or not.

Lemma 8.9. For $g \geq 2$, given primitive $w_1, w_4 \in H_1(\Sigma_g)$ such that $w_1 \cdot w_4 = 0$, there exists $w_2, w_3 \in H_1(\Sigma_g)$ such that

$$w_i \cdot w_{i+1} = 1 \text{ for } i = 1, 2, 3$$

Proof. Let w'_2 satisfy $w_1 \cdot w'_2 = 1$ and w'_3 satisfy $w'_3 \cdot w_4 = 1$. Since $w_1 \in w_4^\perp$ for any k we have $(w'_3 + kw_1) \cdot w_4 = 1$. Choose k as to have $w'_2 \cdot (w'_3 + kw_1) = 0$ and define $w_3 = w'_3 + kw_1$. It is primitive and we can take w''_2 satisfying $w''_2 \cdot w_3 = 1$. Since $w'_2 \in w_3^\perp$ there exists l such that $w_2 = (w'_2 + lw''_2)$ satisfies $w_1 \cdot w_2 = 1$. \square

Lemma 8.10. If $g \geq 4$, $p : H_1(\Sigma_g) \rightarrow \mathbb{C}$ is an injective homomorphism of positive volume and $r(p) < 2g - 3$, then any pair of primitive elements in $H_1(\Sigma_g)$ are equivalent.

Proof. Let $v, w \in H_1(\Sigma_g)$ be two primitive elements. By taking $z \in v^\perp \cap w^\perp$ and applying twice Lemma 8.9 we can consider a sequence $w_0, w_1, \dots, w_6 \in H_1(\Sigma_g)$ such that $w_i \cdot w_{i+1} = 1$ for $i = 0, \dots, 5$, $w_0 = v$, $w_3 = z$ and $w_6 = w$.

We claim that there exist $c_2 \in w_1^\perp \cap w_3^\perp$ and $c_4 \in w_3^\perp \cap w_5^\perp$ such that for $w'_0 = w_0$, $w'_2 = w_2 + c_2$, $w'_4 = w_4 + c_4$ and $w'_6 = w_6$ we can apply Lemma 8.7 to each of the triples w'_i, w'_{i+1}, w'_{i+2} for $i = 0, 2, 4$ and conclude. By the line rank assumption on p , we need to verify

$$(55) \quad \Im(p(w'_2)\overline{p(w_0)}) \neq 0, \quad \Im(p(w'_4)\overline{p(w'_2)}) \neq 0, \quad \Im(p(w'_6)\overline{p(w'_4)}) \neq 0,$$

for this claim to hold.

If $p(w'_2)$ is always colinear to $p(w_0)$ for any choice of $c_2 \in w_1^\perp \cap w_3^\perp$, this means that $p(w_1^\perp \cap w_3^\perp)$ is contained in the line $\mathbb{R}p(w_0)$, which contradicts the line rank assumption. Let us fix $c_2 \in w_1^\perp \cap w_3^\perp$ (and the corresponding w'_2) such that $p(w'_2)$ is not colinear with $p(w_0)$.

Now, the same argument shows that each affine subspace

$$\{c_4 \in w_3^\perp \cap w_5^\perp : \Im(p(w'_4)\overline{p(w'_2)}) = 0\} \text{ and } \{c_4 \in w_3^\perp \cap w_5^\perp : \Im(p(w'_6)\overline{p(w'_4)}) = 0\}$$

have positive corank. Hence, the union of these subspaces does not fill the whole $w_3^\perp \cap w_5^\perp$. Choosing c_4 outside the union of these subspaces gives the solution to Lemma 8.10. \square

Lemma 8.10 provides a proof of Proposition 8.3 for the case of $r(p) < 2g - 3$. In fact, a similar reasoning, but more elaborated, permits to cover also the case $r(p) = 2g - 3$. However, we will treat this case, together with the cases $r(p) \geq 2g - 3$, by different methods, in the next family of lemmata:

Lemma 8.11. *Let $g \geq 4$ and $p : H_1(\Sigma_g) \rightarrow \mathbb{C}$ an injective Haupt homomorphism with $r(p) \geq 2g - 3$. Write $I = p^{-1}(\ell_{\max})$ and suppose $v, w \in H_1(\Sigma_g) \setminus I$ are primitive elements such that $v \cdot w = 0$ and $[w] \in v^\perp / \mathbb{Z}v$ is also primitive. Then $v \sim w$.*

Proof. By applying Remark 8.2 we can suppose without loss of generality that $\ell_{\max} = \mathbb{R}$. Choose $b \in w^\perp$ such that $v \cdot b = 1$. We claim that, up to changing b by $b + e$ for some $e \in I \cap w^\perp \cap v^\perp$ we can suppose that $V = \mathbb{Z}v \oplus \mathbb{Z}b$ belongs to \mathcal{V}_p . Indeed, since $p(e) \in \mathbb{R}$,

$$\text{vol}(\mathbb{Z}v \oplus \mathbb{Z}(b + e)) = \text{vol}(\mathbb{Z}v \oplus \mathbb{Z}b) + p(e)\Im(\overline{p(v)}).$$

By hypothesis the rank of $p(I \cap w^\perp \cap v^\perp)$ is at least $2g - 5 \geq 3$ for $g \geq 4$, so the value of the volume of V can be chosen arbitrarily close to any desired value.

Next take $c \in V^\perp$ such that $w \cdot c = 1$. Given $f \in w^\perp \cap V^\perp \cap I$, we have

$$\text{vol}(\mathbb{Z}w \oplus \mathbb{Z}(c + f)) = \text{vol}(\mathbb{Z}w \oplus \mathbb{Z}c) + p(f)\Im(\overline{p(w)}).$$

Again, since the rank of $w^\perp \cap V^\perp \cap I$ is at least $2g - 6 \geq 2$ for $g \geq 4$, we can suppose that c is chosen so that $W = \mathbb{Z}w \oplus \mathbb{Z}c$ belongs to $\mathcal{V}_{p|_{V^\perp}}$. By construction $V \perp W$ and $0 < \text{vol}(V) + \text{vol}(W) < \text{vol}(p)$. Therefore $V \sim W$ and also $v \sim w$. \square

Lemma 8.12. *Let $g \geq 4$ and $p : H_1(\Sigma_g) \rightarrow \mathbb{C}$ an injective Haupt homomorphism with $r(p) \geq 2g - 3$. Define $I = p^{-1}(\ell_{\max})$. If $v, w \in H_1(\Sigma_g) \setminus I$ are primitive such that*

$$v^\perp \cap w^\perp \not\subseteq I$$

then $v \sim w$.

Proof. Take a symplectic basis a_i, b_i of $H_1(\Sigma_g)$ such that $a_1 = v$ and

$$w = m_1 a_1 + n_1 b_1 + m_2 a_2.$$

Let $X := \mathbb{Z}a_3 \oplus \mathbb{Z}b_3 \oplus \cdots \oplus \mathbb{Z}a_g \oplus \mathbb{Z}b_g$. If $X \not\subset I$ then choose $z \in X \setminus (I \cap X)$ primitive. By Lemma 8.11 $v \sim z$ and $w \sim z$ therefore $v \sim w$. If $X \subset I$ take $c \in (v^\perp \cap w^\perp) \setminus I$ and write $c = c_X + c_\perp$ where $c_\perp \in X^\perp$. Then $c_\perp \notin I$ and $c_\perp \cdot v = c_\perp \cdot w = 0$, so $v, w \in (X')^\perp$ where $X' := \mathbb{Z}(a_3 + c_\perp) \oplus \mathbb{Z}b_3 \oplus \cdots \oplus \mathbb{Z}a_g \oplus \mathbb{Z}b_g$. Restart the argument of the proof with a symplectic basis of $(X')^\perp$ whose first element is still $a_1 = v$ and complete it to a basis of $H_1(\Sigma_g)$ with the chosen basis of X' . Since $X' \not\subset I$ the argument will fall in the previous case and we will be done. \square

Lemma 8.13. *Let $g \geq 4$ and $p : H_1(\Sigma_g) \rightarrow \mathbb{C}$ an injective Haupt homomorphism with $r(p) \geq 2g - 3$. Define $I = p^{-1}(\ell_{\max})$. Let $v \in H_1(\Sigma_g) \setminus I$ and define*

$$I_v = \{z \in H_1(\Sigma_g) : z^\perp \cap v^\perp \subset I\}.$$

Then there exists a proper submodule $J \subsetneq H_1(\Sigma_g)$ such that $I_v \subset J$.

Proof. If $I_v = \emptyset$, the module $J = 0$ does the job. Otherwise take $z \in I_v$. Then $z^\perp \cap v^\perp \subset v^\perp \cap I$. We also have

$$2g - 2 \leq \text{rank}(v^\perp \cap I) < \text{rank}(v^\perp) = 2g - 1$$

where the strict inequality comes from the fact that $v \notin I$ and I is a primitive module. Therefore $\text{rank}(v^\perp \cap I) = 2g - 2$. Its primitive submodule $z^\perp \cap v^\perp$ has also rank $2g - 2$ so the only possibility is that $z^\perp \cap v^\perp = v^\perp \cap I$. Therefore $z \in (v^\perp \cap I)^\perp =: J$ \square

Lemma 8.14. *Let $g \geq 4$ and $p : H_1(\Sigma_g) \rightarrow \mathbb{C}$ an injective Haupt homomorphism with $r(p) \geq 2g - 3$. Then for any $V, W \in \mathcal{V}_p$ we have $V \sim W$.*

Proof. Again we suppose $\ell_{\max} = \mathbb{R}$ and define $I = p^{-1}(\mathbb{R})$. Since V and W are of positive volume we can find primitive elements $v \in V \cap I^c$ and $w \in W \cap I^c$. If $w^\perp \cap v^\perp \not\subset I$ we have $v \sim w$ by Lemma 8.12. Therefore $V \sim W$.

If $w^\perp \cap v^\perp \subset I$ we can consider the union $I \cup I_v \cup I_w$. Since by Lemma 8.13 it is contained in a union of proper submodules, it cannot cover the whole of $H_1(\Sigma_g)$. Take $z \in H_1(\Sigma_g) \setminus (I \cup I_v \cup I_w)$. Then by Lemma 8.12 $v \sim z \sim w$, which as before implies that $V \sim W$. \square

Proposition 8.3 is now proven for all possible ranks of an injective $p \in \mathcal{H}_g$.

8.2. Proof of Theorem 8.1 for non-injective $p \in \mathcal{H}_g$. In the case of non-injective p we also consider the dual boundary complex \mathcal{G}_p of the stratification of $\mathcal{P}er^{-1}(p)$.

Remark that each connected component of a codimension one boundary stratum in $\mathcal{P}er^{-1}(p)$ is contained in a component of the boundary of $\Omega\bar{\mathcal{S}}_g$ and again we can define a continuous map of dual complexes

$$(56) \quad \mathcal{G}_p \rightarrow \mathcal{C}_g/\mathcal{J}_g.$$

However, in the case $\ker p \neq 0$ we know that there will be some image points that correspond to Torelli classes of non-separating curves (see Lemma 6.15). In terms of the parametrization of the vertices of $\mathcal{C}_g/\mathcal{J}_g$ given in subsection 4.8, these correspond to primitive classes in $\ker p \setminus 0$ pinched by p .

In contrast with the case of injective periods, when p has large kernel, there may be boundary components of codimension one of $\Omega\bar{S}_g$ whose intersection with $\text{Per}^{-1}(p)$ is disconnected (when $\ker p$ contains symplectic submodules of large rank). For instance, the boundary component of $\Omega\bar{S}_g$ corresponding to a p -admissible splitting $V_1 \oplus V_2$ with $\text{rank } V_1 \geq 10$ and $\deg(p|_{V_1}) = 2$ provides an example, via attaching maps and Theorem 1.3. This implies that the map (56) is no longer injective. However, we can use the inductive hypothesis to identify a particular subcomplex $\mathcal{G}'_p \subset \mathcal{G}_p$ spanned by *simple vertices* where the restriction of (56) is injective:

Definition 8.15. Let $p \in \mathcal{H}_g$. A p -admissible splitting $V_1 \oplus V_2$ is simple if it satisfies $\text{rank}(V_i) \leq 6$ or $\deg(p|_{V_i}) \geq 3$ for $i = 1, 2$. A pinched class $a \in \ker p \setminus 0$ is simple if either $g = 3, 4$ or $\deg p_a \geq 3$. A vertex of $\mathcal{C}_g/\mathcal{I}_g$ is p -simple if the corresponding p -admissible splitting or pinched class is p -admissible and simple.

Lemma 8.16. Let $g \geq 3$. For any $p \in \mathcal{H}_g$ with $\ker p \neq 0$ and $\deg p \geq 3$ there exist p -simple vertices of non-compact type in $\mathcal{C}_g/\mathcal{I}_g$.

Proof. For $g = 3, 4$ the statement is equivalent to the existence of pinched classes (see Lemma 6.15). Suppose $g \geq 5$ and that there is a vertex that is not simple. If it is of compact type, it means that one of the factors of the associated p -admissible splitting is of rank at least eight and the restriction of p to it is of primitive degree two. If it is of non-compact type, there is a symplectic submodule of rank at least 8 where the restriction of p has degree two. In either case there is a symplectic submodule of $\ker p$ of rank at least two. Each primitive element in that submodule determines a simple vertex of non-compact type of \mathcal{G}_p . \square

Proof of Theorem 8.1 in the non injective case. We claim that the restriction of (56) to the subcomplex \mathcal{G}'_p —spanned by simple vertices—is injective. Indeed, suppose D_1 and D_2 are irreducible boundary components in boundary strata contained in $\text{Per}^{-1}(p)$ lying in a boundary component of $\Omega\bar{S}_g$ corresponding to a simple p -admissible splitting or primitive class in $\ker p \setminus 0$. Propositions 7.5 and 7.6 describe this set as the image of products of isoperiodic sets under attaching maps. However, by inductive hypothesis (or by Theorem 3.3 in the cases of genus two and three and degree two) the products of isoperiodic sets under consideration are connected. Hence $D_1 = D_2$. We can therefore talk about simple vertices of \mathcal{G}_p , namely, those lying over p -simple vertices of $\mathcal{C}_g/\mathcal{I}_g$. The connectedness of the complex \mathcal{G}_p in the non injective case proceeds in three basic steps:

Step 1: Each vertex of compact type is equivalent in \mathcal{G}_p to some vertex of non-compact type.

Step 2: Each vertex of non-compact type is equivalent in \mathcal{G}_p to some simple vertex of non-compact type.

Step 3: Any pair of simple vertices of non-compact type are equivalent in \mathcal{G}'_p .

8.2.1. Proof of Step 1: from compact type to non-compact type: Let $V_1 \oplus V_2 = H_1(\Sigma_g)$ be the p -admissible splitting associated to a separating simple closed curve c in Σ_g and denote $p_i = p|_{V_i}$. If one of the p_i 's is non-injective we claim that any stable form

$$(C, m, \omega) = (C_1 \vee C_2, m_1 \oplus m_2, \omega_1 \vee \omega_2)$$

of period p sitting in the boundary component associated to $V_1 \oplus V_2$ is equivalent to a stable form with a non-separating node. Indeed, suppose without loss of generality, that $\ker p_2 \neq 0$. Then V_2 has at least rank 4. Suppose first that the vertex defined by (C, m, ω) is simple.

Then, under the hypotheses of Proposition 8.1, $\text{Per}^{-1}(p_2)$ is connected, contains ω_2 and a stable form ω_3 over a curve with a non-separating node. An isoperiodic path η_t between them provides an isoperiodic path $\omega_1 \vee \eta_t$ between $\omega = \omega_1 \vee \omega_2$ and $\omega_1 \vee \omega_3$. The latter is a form in $\Omega_0^* \mathcal{S}_g$ with periods p and two nodes, one of which is non-separating. Therefore it defines an edge between the vertex defined by (C, m, ω) and a vertex of non-compact type of \mathcal{G}_p .

Next suppose that the vertex defined by (C, m, ω) is not simple. Then $\text{rank } V_2 \geq 8$ and $\deg p_2 = 2$. Integration of ω_2 along C_2 provides a branched degree two cover

$$C_2 \rightarrow \mathbb{C}/p_2(V_2) =: E_2$$

having at least two distinct critical values. The preimage of a path in E_2 joining these critical values provides a pair of twin paths for ω_2 sharing the same endpoints. The Schiffer variation along this pair of twin paths describes an isoperiodic deformation η_t joining ω_2 with a stable form ω_3 with a non-separating node. Indeed, if it would join the two critical points to a single critical point, the local degree of the branched covering around the latter would be at least three, contradicting the degree two hypothesis. If it would collapse to a separating node, the degree of the restriction of the period map to each part of the curve would be one, and since one of the parts has genus at least two it would contradict Haupt's Theorem.

It remains to treat the case where both p_1 and p_2 are injective. We will prove that the (simple) vertex is equivalent to another vertex of compact type where the restriction of p to one of the parts has non-trivial kernel.

If both p_1 and p_2 are injective, we can find p -admissible splittings of V_1 and V_2 with all factors of rank 2 by using Corollary 6.19 inductively to p_1 and p_2 . In particular the vertex associated to (C, m, ω) is equivalent to a vertex defined by a p -admissible splitting having a rank two factor, so we can suppose $\text{rank } V_1 = 2$. If p_2 is non-injective we are done. It remains to treat the case where p_2 is injective.

Take a primitive element $a \in \ker p$ and write $a = n_1 a_1 + n_2 a_2$ for primitive $a_i \in V_i$ and co-prime $n_1, n_2 \in \mathbb{N}^*$. If $a_2 \in V_2$ is contained in a factor V of a p_2 -admissible splitting of V_2 , define $V'_1 = V_1 \oplus V$. The splitting $V'_1 \oplus V_1'^\perp$ is also p -admissible and satisfies $\ker(p|_{V'_1}) \neq 0$. It defines a vertex that is equivalent to the one defined by $V_1 \oplus V_2$. In particular this argument works if a_2 is p_2 -admissible.

So suppose p_2 is injective and $a_2 \neq 0$ is not p_2 -admissible. Then by Lemma 6.9 applied to p_2 , the rank of $p_2^{-1}(\mathbb{R}p(a_2)) \cap a_2^\perp$ is at least $(2g - 2) - 3 + 1 = 2g - 4$. Completing a_2 to a symplectic basis $a_2, b_2, a_3, b_3, \dots, a_g, b_g$ of V_2 , and denoting $V_3 := \mathbb{Z}a_3 + \mathbb{Z}b_3 + \dots + \mathbb{Z}a_g + \mathbb{Z}b_g$, we conclude that $H = (p_2^{-1}(\mathbb{R}p(a_2)) \cap a_2^\perp) \cap b_2^\perp = p^{-1}(\mathbb{R}p(a_2)) \cap V_3$ is either V_3 or a co-rank one primitive submodule of V_3 .

In the latter case, by considering an element $w \in V_3$ such that $p(w)$ does not belong to $\ell := \mathbb{R}p(a_2)$, we apply Lemma 6.9 to ℓ and V_3 to construct a symplectic rank two submodule $W \subset V_3$ containing w with $0 < \text{vol}_p(W) < \text{vol}_p(V_2)$. Since p_2 is injective, this implies that $W \in \mathcal{V}_{p_2}$. The splitting $V_1 \oplus W \oplus (W^\perp \cap V_2)$ of $H_1(\Sigma_g)$ is p -admissible, and in particular so is $W \oplus W^\perp$. On the other hand $a \in W^\perp \cap \ker p$ so the restriction $p|_{W^\perp}$ is non-injective and we are reduced to one of the previous cases.

It remains to treat the case where $H = V_3$, namely $p(V_3) \subset \mathbb{R}p(a_2)$. In this case we will find a new initial p -admissible splitting $V_1 \oplus V_2$ that falls in one of the previous cases and that defines an equivalent splitting. The new V_1 that we want to construct, call it V'_1 , will be a rank two factor of a p_2 -admissible splitting of V_2 .

The candidates to V'_1 satisfy $0 < \text{vol}_p(V'_1) < \text{vol}_p(V_2)$. The initial splitting of a will change to

$$a = n''_1 a''_1 + n''_2 a''_2$$

with primitive $a''_1 \in V'_1$ and $a''_2 \in V'_2 := (V'_1)^\perp$ and co-prime n''_1 and n''_2 (the reason for this notation a''_i instead of a'_i will become clear hereafter). We claim that it is possible to find V'_1 so that either a''_2 is $p_{V'_2}$ -admissible or $p'_2 = p|_{V'_2}$ is not injective. We already explained that this would conclude the proof.

Take $b_1 \in V_1$ such that $a_1 \cdot b_1 = 1$. Up to composing p with a \mathbb{R} -linear orientation preserving equivalence from \mathbb{C} to \mathbb{C} , we can assume that

$$p(a_1) = n_2, \Im p(b_1) = 1, p(a_2) = -n_1, \Im p(b_2) < 0,$$

and that

$$p(a_k) = \alpha_k \in \mathbb{R}, p(b_k) = \beta_k \in \mathbb{R} \text{ for } k \geq 3.$$

By injectivity of p_2 , the numbers $\alpha_3, \dots, \alpha_g, \beta_3, \dots, \beta_g$ are linearly independent over \mathbb{Q} .

We are going to look for the module V'_1 as being generated by the elements a'_1 and b'_1 , where

$$a'_1 = a_2 + \sum_{k \geq 3} m_k a_k + n_k b_k, b'_1 = b_2, a'_2 = a_1, b'_2 = b_1$$

and for $k \geq 3$

$$a'_k = a_k + n_k b_2, b'_k = b_k - m_k b_2$$

form a new symplectic basis for $H_1(\Sigma_g)$. Here m_k, n_k are integers that have to be determined for $k \geq 3$. We have

$$\text{vol}_p(V'_1) = -\Im(p(b_2))(n_1 - \sum_{k \geq 3} m_k \alpha_k + n_k \beta_k).$$

Observe also that $\text{vol}_p(V_2) = -n_1 \Im(p(b_2))$. We will choose m_k, n_k multiple of n_1 , so we write $m_k = m'_k n_1, n_k = n'_k n_1$ with m'_k, n'_k integers. We then have

$$\text{vol}_p(V'_1) = \varepsilon \text{vol}_p(V_2) \text{ with } \varepsilon = 1 - \sum_{k \geq 3} m'_k \alpha_k + n'_k \beta_k.$$

Since p_2 is injective, V'_1 is p_2 -admissible if and only if $0 < \varepsilon < 1$. By rational independence of $\alpha_3, \dots, \alpha_g, \beta_3, \dots, \beta_g$ we can find $m'_k, n'_k \in \mathbb{Z}$ satisfying this condition. If for such a choice the homomorphism $p'_2 := p|_{V'_2}$ is not injective, we are done. Suppose that for all choices we have that p'_2 is injective. In this case we will refine the solution to have a''_2 to be p'_2 -admissible.

Let us understand how the class a decomposes according to the splitting $V'_1 + (V'_1)^\perp$: it is given by $a = n_2 a''_1 + n_1 a''_2$ with $a''_1 = a'_1$ and

$$a''_2 = a'_2 - n_2 \sum_{k \geq 3} m'_k a'_k + n'_k b'_k.$$

Hence, it suffices to find $n'_k, m'_k \in \mathbb{Z}$ such that a''_2 is p'_2 -admissible. The volume of the symplectic rank two submodule $\mathbb{Z}a''_2 + \mathbb{Z}b'_2 \subset V'_2$ (containing a''_2) is

$$\text{vol}_p(\mathbb{Z}a''_2 + \mathbb{Z}b'_2) = \Im(p(b'_2) \overline{p(a''_2)}) = n_2 \left(1 - \sum_{k \geq 3} m'_k \alpha_k + n'_k \beta_k\right) = n_2 \varepsilon,$$

while

$$\text{vol}_p(V'_2) = \text{vol}_p - \text{vol}_p(V'_1) = \text{vol}_p - \varepsilon \text{vol}_p(V_2).$$

Hence, as soon as $0 < \varepsilon < \frac{\text{vol}_p}{n_2 + \text{vol}_p(V_2)}$ one concludes that a''_2 is p'_2 -admissible. The rational independence of $\alpha_3, \dots, \alpha_g, \beta_3, \dots, \beta_g$ allows to find a solution, again.

8.2.2. Proof of Step 2: From non-compact type to simple of non-compact type.

Lemma 8.17. *Let (C, ω) be a stable form with a non-separating node having $(\hat{C}, \hat{\omega})$ as its normalization and $n_1, n_2 \in \hat{C}$ the points corresponding to the node in C . Let p (resp. \hat{p}) be the period homomorphism associated to ω (resp. $\hat{\omega}$). Assume that $\deg(\hat{p}) < \infty$, and let $\hat{F} : \hat{C} \rightarrow \mathbb{C}/\hat{p}(H_1(\hat{C}))$ be the map defined by integration of $\hat{\omega}$. Then*

$$\deg(\hat{p}) = \deg(p) \Leftrightarrow \hat{F}(n_1) = \hat{F}(n_2).$$

Proof. The proof is straightforward once one realizes that $\text{vol}(\hat{p}) = \text{vol}(p)$. \square

Continuing with the notations of Lemma 8.17 we consider a marked stable form (C, m, ω) with period map p of primitive degree at least three, and a non-separating node that collapses a curve c such that $\deg(p_c) = 2$. The period map \hat{p} of its normalization $(\hat{C}, \hat{\omega})$ has therefore precisely degree two. By Lemma 8.17 $\hat{F}(n_1) \neq \hat{F}(n_2)$ for the integral

$$\hat{F} : \hat{C} \rightarrow \mathbb{C}/\hat{p}(H_1(\hat{C}))$$

of $\hat{\omega}$. The map \hat{F} is a branched covering of degree two. All its branch points are therefore simple and have distinct values for \hat{F} . Choose a path in $\mathbb{C}/\hat{p}(H_1(\hat{C}))$ that avoids the two points $\hat{F}(n_1)$ and $\hat{F}(n_2)$ and joins two distinct critical values. Its preimage by \hat{F} defines a pair of twins that join two critical points in \hat{C} and also two zeros of ω in C . The closed curve formed by both twins cannot be separating. Indeed, if it was separating, it would induce a \hat{p} -admissible splitting of $H_1(\Sigma_{g-1})$. However, these do not exist for $\deg(\hat{p}) = 2$ and $g - 1 \geq 3$ (see Example 6.21). Therefore, the curve formed by both twins is non-separating. After the Schiffer variation on $(\hat{C}, \hat{\omega})$ along these twins, we obtain a new stable form $(\tilde{C}, \tilde{\omega})$ of period \hat{p} with a node \tilde{n} that is non separating. It is a degree two cover of an elliptic differential. By Lemma 8.17, identifying the points n_1 and n_2 on $(\tilde{C}, \tilde{\omega})$ produces a stable form whose period map has finite primitive degree different from $\deg(\hat{p}) = 2$ (and from one by Haupt's Theorem). Hence it has degree at least three. Again by the same Lemma, the normalization of \tilde{n} on this form has that same degree, larger than three.

8.2.3. *Proof of Step 3: connecting simple vertices of non-compact type in \mathcal{G}'_p .* Recall that each simple vertex of non-compact type of the complex \mathcal{G}_p corresponds to a unique cyclic primitive submodule $\mathbb{Z}a \subset \ker p$ such that $\deg p_a \geq 3$. We will parametrize those vertices by the primitive elements $\pm a \in \ker p \setminus 0$ generating the submodule. We say that two such primitive elements $a, a' \in \ker p \setminus 0$ are equivalent if the corresponding vertices in \mathcal{G}'_p lie in the same connected component of the subcomplex \mathcal{G}'_p of \mathcal{G}_p spanned by simple vertices.

Lemma 8.18. *Given $g \geq 4$, a homomorphism $p : H_1(\Sigma_g) \rightarrow \mathbb{C}$ with $\text{vol } p > 0$, $\deg p \geq 3$ and*

- (1) *a symplectic submodule $W \subset \ker p$ of rank two, then every pair of simple vertices of non-compact type of \mathcal{G}'_p corresponding to classes in $W \cup W^\perp$ are equivalent.*
- (2) *an isotropic primitive submodule $L \subset \ker p$ of rank two, then every pair of simple vertices of non-compact type of \mathcal{G}'_p corresponding to classes in L are equivalent.*
- (3) *a p -admissible splitting $H_1(\Sigma_g) = V_1 \oplus V_2$. The primitive elements $a \in V_i \cap \ker p$ such that $\deg((p|_{V_i})_a) \geq 2$ satisfy $\deg(p_a) \geq 3$.*

Proof. (1) First remark that the homomorphism $p_1 := p|_{W^\perp}$ is defined on a module of rank at least 6 and has the same volume and primitive degree as p . By Lemma 8.16 applied to p_1 we deduce that there exists a simple vertex for \mathcal{G}_{p_1} . Thanks to Lemma 6.13 and the fact that $W \subset \ker p$, it also determines a simple vertex for \mathcal{G}_p . Using Corollary 6.18 we construct a marked stable form (C_1, m_1, ω_1) of genus $g - 1$ and period p_1 , with a node that induces the given simple vertex and – thanks to the genus hypothesis and up to some Schiffer variations – a simple zero at some regular point. Take a pair of embedded twins with distinct endpoints starting at the simple zero and glue the endpoints, thus obtaining a stable form on a genus g curve with two nodes. After the gluing, the union of the twins forms a closed loop in the nodal curve whose period vanishes. Given any symplectic basis a, b of W we mark the obtained nodal form by collapsing a to the node, associating b to the loop obtained from the twins, and keeping the marking on W^\perp as it was for (C_1, ω_1, m_1) . The given form with two nodes defines an edge in \mathcal{G}_p joining two simple vertices. One of them corresponds to the node of the initial ω_1 and is independent of the chosen a . In particular all primitive classes in W define equivalent simple vertices. Moreover, if the chosen node for ω_1 is determined by a primitive element $\ker p \cap W^\perp$ defining a simple vertex for \mathcal{G}_p it is also a simple vertex for \mathcal{G}_{p_1} . This vertex is equivalent to any vertex defined by a primitive element of W . This shows that all simple vertices of non-compact type of \mathcal{G}_p corresponding to primitive classes in $W \cup W^\perp$ are equivalent.

(2) The module L^\perp/L has rank $2g - 4$ and the ambient symplectic form on $H_1(\Sigma_g)$ induces a symplectic form on it. Since $L \subset \ker p$, the map p induces a homomorphism

$$p_L : L^\perp/L \rightarrow \mathbb{C}.$$

If $\ker p_L$ contains a non-trivial symplectic submodule, then by item (1) of this Lemma, every pair of simple vertices in L are equivalent. Item (2) is proven for the cases $\deg p_L = 1$ (where the kernel is symplectic of corank two) or $g \geq 5$ and $\deg p_L < \infty$ (thanks to Lemma 6.14). It remains to treat the cases $g = 4$ and $\deg p_L \geq 2$ and $g \geq 5$ and $\deg p_L = \infty$.

Next suppose a_1, a_2 forms a basis of L corresponding to simple vertices and $\deg p_L \geq 2$. We claim that we can construct a marked stable form of period p with precisely two non-separating nodes that correspond to the given classes. The corresponding vertices are thus joined by an edge in \mathcal{G}_p . Indeed, take a form (C_2, m_2, ω_2) of genus $g - 2$ and periods p_L with two simple zeros. Complete the elements a_1, a_2 to a symplectic basis a_i, b_i of $H_1(\Sigma_g)$. For $i = 1, 2$, apply Lemma 6.17 to ω_2 to choose an embedded path β_i in C_2 having distinct endpoints and satisfying

$$p(b_i) = \int_{\beta_i} \omega_2.$$

We can further assume that β_1 and β_2 are disjoint. Indeed, if $p(b_1) = p(b_2) = 0$ it suffices to take pairs of short twins at distinct zeros of ω_2 . If only one of them is non-zero, we can take a very short pair of twins to realize the zero period so as to avoid the path of non-zero length. If both are non-zero and we have initially taken two paths that intersect, we change one of the paths in its homotopy class with fixed endpoints to avoid the intersections. Gluing the endpoints of the said paths and marking the form by collapsing a_1 and a_2 to the nodes, and associating b_i to the corresponding loop β_i we obtain a marked stable form of period p with two non-separating nodes as was claimed. In full generality it is not true that every primitive element of L corresponds to a simple vertex. However, if $\deg p_L \geq 3$ we have that every primitive element of L corresponds to a simple vertex. The previous argument shows

that we only have to check that for any pair of primitive elements $a, a' \in L$ we can construct a sequence $a = a_0, \dots, a_n = a'$ of primitive elements in L such that a_i, a_{i+1} form a basis of L for each $i = 0, \dots, n-1$. This is guaranteed by Gauss algorithm in L .

It remains to treat the case $g = 4$ and $\deg p_L = 2$. In this case, p_L is defined on a rank four symplectic module and has a lattice $\Lambda \subset \mathbb{C}$ as image. The normal form for periods of finite primitive degree, see Lemma 9.1 proved in Appendix 9, provides a symplectic splitting of L^\perp/L into rank two submodules $W_1 \oplus W_2$ satisfying $p_L(W_i) = \Lambda$. Let $V \subset H_1(\Sigma_g)$ be a rank four symplectic submodule containing L and consider a symplectic splitting of $V^\perp = V_1 \oplus V_2$ that induces $W_1 \oplus W_2$ on L^\perp/L . Let $V_3 = V \oplus V_1$. The symplectic splitting $H_1(\Sigma_g) = V_3 \oplus V_2$ defines a simple vertex in \mathcal{G}_p , since all factors are of rank at most 6. We claim that there is an edge of \mathcal{G}_p between this simple vertex and any simple vertex corresponding to a primitive element in L . To construct it, it suffices to construct a marked stable form with two nodes one of which induces the symplectic splitting $V_3 \oplus V_2$ and the other collapses the class $a \in L \subset V_3$ satisfying $\deg p_a \geq 3$ to a node. On the other hand we have $\text{Im}(p|_{V_3})_a = \text{Im}(p_a)$, $\text{vol}_p(V) = 0$, and $\text{vol}_p(V_2) = \text{vol}_p(V_1)$. Therefore

$$\deg((p|_{V_3})_a) = \frac{\text{vol}((p|_{V_3})_a)}{\text{vol}(\mathbb{C}/\text{Im}(p|_{V_3})_a)} = \frac{\text{vol}(p_a)/2}{\text{vol}(\mathbb{C}/\text{Im}(p_a))} = \frac{1}{2} \deg(p_a) \geq \frac{3}{2} > 1.$$

By using Corollary 6.18 we can construct a stable form with two nodes, one of which corresponds to a and the other to the splitting $V_3 \oplus V_2$.

(3) Without loss of generality suppose $a \in V_1$. Remark that by definition $\text{Im}((p|_{V_i})_a) \subset \text{Im}(p_a)$. Therefore

$$\begin{aligned} \deg p_a &= \frac{\text{vol } p_a}{\text{vol}(\mathbb{C}/\text{Im}(p_a))} \geq \frac{\text{vol } p_a}{\text{vol}(\mathbb{C}/\text{Im}(p|_{V_1})_a)} = \frac{\text{vol}((p|_{V_1})_a) + \text{vol}(p|_{V_2})}{\text{vol}(\mathbb{C}/\text{Im}(p|_{V_1})_a)} > \\ &> \deg((p|_{V_1})_a) \geq 2. \end{aligned}$$

□

Corollary 8.19. *If $c, c' \in \ker p$ are primitive elements satisfying $c \cdot c' = 0$ and $\deg(p_c), \deg(p_{c'}) \geq 3$ and then $c \sim c'$.*

Proof. Write $a_1 = c$ and choose $b_1 \in H_1(\Sigma_g)$ such that $a_1 \cdot b_1 = 1$. Define $W_1 = \mathbb{Z}a_1 \oplus \mathbb{Z}b_1$ and write

$$c' = m_1 a_1 + n_1 b_1 + m_2 a_2$$

where $a_2 \in W_1^\perp$ is a primitive element and $m_1, n_1, m_2 \in \mathbb{Z}$. Since $c \cdot c' = 0$ we have $n_1 = 0$. If $m_2 = 0$ then $c' = \pm c$ and we are done. Otherwise $a_2 \in \ker p$ and we can apply item (2) of Lemma 8.18 to $L = \mathbb{Z}a_1 \oplus \mathbb{Z}a_2$ to conclude $c \sim c'$. □

Corollary 8.20. *If there exists a non-trivial symplectic submodule $W \subset \ker p$ then every pair of primitive $c, c' \in \ker p$ such that $\deg(p_c), \deg(p_{c'}) \geq 3$ are equivalent.*

Proof. For every primitive $c \in W$ $\deg(p_c) = \deg p \geq 3$. Every pair of primitive elements in W are equivalent by (1) in Lemma 8.18. Take any primitive $c \in \ker p$ with $\deg(p_c) \geq 3$. Then $\text{rank}(c^\perp \cap W) \geq 1$ and for any primitive element $c' \in c^\perp \cap W$ we have $c' \sim c$ by Corollary 8.19 we deduce $c \sim c'$. □

Corollary 8.21. *If there exists a primitive $a \in \ker p$ such that $\deg(p_a) < \infty$ then for every pair of primitive $c, c' \in \ker p$ such that $\deg(p_c), \deg(p_{c'}) \geq 3$ we have $c \sim c'$.*

Proof. $\deg(p_a) < \infty$ implies that $\text{rank } \ker p = 2g - 2$. Since $g \geq 4$ we have that $\ker p$ contains a non-trivial symplectic submodule and conclude by Corollary 8.20. \square

The following result improves Corollary 8.19:

Corollary 8.22. *If $\text{rank}(\ker p) \geq 3$ then every pair of primitive $c, c' \in \ker p$ such that $\deg(p_c), \deg(p_{c'}) \geq 3$ satisfy $c \sim c'$.*

Proof. By Corollary 8.21 if there exists an element c such that $\deg(p_c) < \infty$ we are done. So we can suppose $\deg(p_c) = \infty$ for every primitive $c \in \ker p$. Let $c, c' \in \ker p$ be two primitive elements. Then $\text{rank}(c^\perp \cap c'^\perp \cap \ker p) \geq 1$ and a primitive element $a \in c^\perp \cap c'^\perp \cap \ker p$ will satisfy $c \sim a \sim c'$ by Corollary 8.19. \square

Lemma 8.23. *If $\text{rank}(\ker p) \leq 2$ then every pair of primitive $a, a' \in \ker p$ such that*

$$\deg(p_a), \deg(p_{a'}) \geq 3 \text{ satisfy } a \sim a'.$$

Proof. If $\ker p$ has rank one, or $a \cdot a' = 0$ or there exists a symplectic module in $\ker p$ containing a and a' we are done by Corollaries 8.19 and 8.20.

Let $a_1 = a$ and b_1 such that $a_1 \cdot b_1 = 1$. Write $W_1 = \mathbb{Z}a_1 \oplus \mathbb{Z}b_1$ and

$$(57) \quad a' = m_1 a_1 + n_1 b_1 + m_2 a_2$$

for a primitive $a_2 \in W_1^\perp$ and integers m_1, n_1, m_2 . We can assume $n_1 = a \cdot a' \neq 0$ and $m_2 p(a_2) \neq 0$ (otherwise we contradict the rank hypothesis on $\ker p$ or we fall in one of the initial cases). Define $p_1 = p|_{W_1^\perp}$.

If a_2 is p_1 -admissible, i.e., it belongs to a symplectic module of rank two $V_2 \in \mathcal{V}_{p_1}$, there exists a p -admissible splitting $W_1^\perp = V_2 \oplus V_3$. Then $a_1, a' \in V_4 := W_1 \oplus V_2$ and $\text{vol}_p(V_4) = \text{vol}_p(V_2) > 0$. The restriction of p to the orthogonal $V_4^\perp = V_3$ has infinite degree, since it has rank at least four, and the kernel of the restriction of p to V_4^\perp has rank bounded by one (otherwise the whole period would have a rank bounded from below by three). Since $a_1 \in V_4$ does not belong to any symplectic submodule contained in $\ker(p)$, $p|_{V_4}$ is a Haupt homomorphism on a rank four symplectic module. By Lemma 6.15 applied for $g = 2$, all primitive elements a'' in $\ker p|_{V_4}$ are pinched by $p|_{V_4}$. Applying Corollary 6.18 we can find a form with two nodes: one that pinches a'' and another that induces the splitting $V_4 \oplus V_4^\perp$, which is simple, hence the vertex of \mathcal{G}_p associated to a'' is equivalent to the vertex associated to $V_4 \oplus V_4^\perp$. Since a, a' are primitive elements in $V_4 \cap \ker p$ we have $a' \sim a$.

By Corollary 6.11 applied to W_1^\perp , there exists a proper submodule $I \subset W_1^\perp$ containing all elements that are not p_1 -admissible, i.e. that do not belong to any symplectic module $V_2 \in \mathcal{V}_{p_1}$. So if $a_2 \notin I$ we are done. Next we are going to show that, up to changing the initial choice of b_1 , we can guarantee that we fall in the previous case.

Indeed, suppose $a_2 \in I \setminus \ker p$ is not admissible. Denote $\ell = \mathbb{R}p(a_2)$. We know by Corollary 6.11 that $I = p^{-1}(\ell) \cap W_1^\perp$ has rank at least $2g - 4$ and for every other real line ℓ' , $\ell' \cap p(W_1^\perp)$ has rank at most 2. Since $p(a_1), p(b_1) \in \ell$, $p^{-1}(\ell)$ has rank at least $2g - 2 \geq 6$. On the other hand, for every other real line $\ell' \subset \mathbb{C}$ containing 0 we have $\text{rank}(p^{-1}(\ell') \cap W_1^\perp) \leq 2 + \text{rank}(\ker p_1) \leq 3$. Therefore $p^{-1}(\ell')$ has rank at most 5. If we manage to find a splitting as in equation (57)

where the image of the a_2 is outside ℓ we will be done. We are going to show that, up to changing the initial b_1 , we can suppose that we fall in this case or one of the previous cases.

Given $w \in a_1^\perp$ define $b'_1 = b_1 + w$ and $W'_1 = \mathbb{Z}a_1 \oplus \mathbb{Z}b'_1$. Then

$$a = m_1a_1 + n_1b'_1 + m'_2a'_2$$

where $m'_2a'_2 = m_2a_2 - n_1w$. If we manage to guarantee that

- $a'_2 \in W_1'^\perp$, or equivalently $0 = -n_1(b_1 \cdot w) + m_2(w \cdot a_2)$
- $p(a'_2) \notin \ell = \mathbb{R}p(a_2)$ or equivalently $p(w) \notin \ell$,

we will be done: $a'_2 \in W_1'^\perp$ will be $p|_{W_1'^\perp}$ -admissible.

If there exists $w \in a_2^\perp \cap W_1^\perp \setminus p^{-1}(\ell)$, it constitutes a solution. Otherwise $a_2^\perp \cap W_1^\perp \subset p^{-1}(\ell) \cap W_1^\perp$, and since W_1^\perp has positive volume, any $b_2 \in W_1^\perp$ satisfying $a_2 \cdot b_2 = 1$ satisfies $p(b_2) \notin \ell$. In this case the element $w = m_2a_1 + n_1b_2$ provides a solution. \square

\square

8.3. Proof of Theorem 1.2. The Theorem is true for $g = 2, 3$ by Theorem 3.3. We proceed by induction on the genus. Fix some $g \geq 4$ and suppose that Theorem 1.2 is true up to genus $g - 1$. Take $p \in \mathcal{H}_g$ with $\deg(p) \geq 3$. By Proposition 5.5 every connected component of $\text{Per}^{-1}(p)$ has points in the boundary. The boundary $\text{Per}^{-1}(p) \setminus \text{Per}^{-1}(p)$ is connected thanks to Theorem 8.1. Therefore $\text{Per}^{-1}(p)$ is connected. On the other hand, by Theorem 4.37, the boundary points do not locally separate $\text{Per}^{-1}(p)$, and the latter is therefore also connected.

9. APPENDIX I: PROOF OF PROPOSITION 3.10

We first begin by providing a normal form for periods of positive volume and finite primitive degree, which implies the first item of Proposition 3.10 in the case where the subspace (18) is rational.

Lemma 9.1. *Given a surjective homomorphism $p : \mathbb{Z}^{2g} \rightarrow \mathbb{Z} + i\mathbb{Z}$ of volume (and primitive degree) $d \geq 2$, there exists $M \in \text{Sp}(2g, \mathbb{Z})$ and a symplectic basis $\{a_1, b_1, \dots, a_g, b_g\}$ of \mathbb{Z}^{2g} such that*

$$p \circ M = a_1^* + i(db_1^* + a_2^*)$$

where the star denotes the symplectic dual of the given element.

Proof. Denote by $x = \Re(p)$ and $y = \Im(p)$ the elements in $(\mathbb{Z}^{2g})^*$. They satisfy $x \cdot y = d$. Choose $a_1 \in \mathbb{Z}^{2g}$ the symplectic dual of x . Choose some $b_1 \in \mathbb{Z}^{2g}$ such that $a_1 \cdot b_1 = 1$ and write

$$(58) \quad y = m_1a_1^* + db_1^* + m_2a_2^*$$

where a_2 is a primitive element satisfying $a_1 \cdot a_2 = b_1 \cdot a_2 = 0$ and m_1, d and m_2 are coprime. Complete those elements to a symplectic basis $a_1, b_1, \dots, a_g, b_g$ of \mathbb{Z}^{2g} . The image of an element $u = \sum_{k \geq 1} \alpha_k a_k + \beta_k b_k$ under p is

$$p(u) = (a_1^*(u), (m_1a_1^* + db_1^* + m_2a_2^*)(u)) = (\alpha_1, m_1\alpha_1 + d\beta_1 + m_2\alpha_2).$$

Therefore

$$\begin{aligned} p(u) = 1 &\Leftrightarrow \alpha_1 = 1 \text{ and } m_1 + d\beta_2 + m_2\alpha_2 = 0 \\ p(u) = i &\Leftrightarrow \alpha_1 = 0 \text{ and } d\beta_1 + m_2\alpha_2 = 1. \end{aligned}$$

Hence, p is surjective if and only if d and m_2 are co-prime.

To conclude, we are going to show that under the hypothesis of d and m_2 co-prime, there exists a choice of b_1 such that the coefficients m_1 and m_2 in the splitting given by (58) are 0 and 1 respectively.

Let us analyze the effect of a change of the first given b_1 . Write $\tilde{a}_1^* = a_1^*$ and

$$\tilde{b}_1^* = u_1 a_1^* + b_1^* + \sum_{k \geq 2} u_k a_k^* + v_k b_k^*$$

for some $u_k, v_k \in \mathbb{Z}$. The new decomposition $y = \tilde{m}_1 \tilde{a}_1^* + d \tilde{b}_1^* + \tilde{m}_2 \tilde{a}_2^*$ has the properties

$$(y - \tilde{m}_1 \tilde{a}_1^* - d \tilde{b}_1^*) \cdot \tilde{a}_1^* = 0 \text{ and } (y - \tilde{m}_1 \tilde{a}_1^* - d \tilde{b}_1^*) \cdot \tilde{b}_1^* = 0.$$

The first equation is automatically satisfied, and the second gives

$$(59) \quad \tilde{m}_1 = m_1 - d u_1 + m_2 v_2$$

Since d and m_2 are co-prime we can already choose u_1 and v_2 to get $\tilde{m}_1 = 0$ and restart the argument by supposing $m_1 = 0$.

For this choice, we have

$$\tilde{m}_2^* \tilde{a}_2^* = y - \tilde{m}_1 \tilde{a}_1^* - d \tilde{b}_1^* = m_2 v_2 a_1^* + (m_2 - d u_2) a_2^* - d v_2 b_2^* - d \left(\sum_{k \geq 3} u_k a_k^* + v_k b_k^* \right)$$

and from this

$$\tilde{m}_2 = \gcd(-m_2 v_2, m_2 - d u_2, -d v_2, -d u_3, -d v_3, \dots, -d u_g, -d v_g)$$

If we still want $\tilde{m}_1 = 0$ we need to impose $m_2 v_2 = d u_1$ by equation (59). The choice $v_2 = d$, $u_1 = m_2$ and all other coefficients equal to zero gives $\tilde{m}_1 = 0$ and $\tilde{m}_2 = 1$, as desired. \square

We continue the proof of Proposition 3.10, which is reminiscent of Ratner's theory.

Equip \mathbb{R}^{2g} with its canonical symplectic form $\omega(x, y) = \sum_{1 \leq k \leq g} x_{2k} y_{2k+1} - x_{2k+1} y_{2k}$. The volume of a period $p \in \mathbb{C}^{2g}$ is the symplectic product $V(p) = \omega(\Re p, \Im p)$. Since the action of Γ is linear, and that the volume is multiplicative, namely $V(\lambda p) = |\lambda|^2 V(p)$ for every $\lambda \in \mathbb{C}$ and $p \in \mathbb{C}^{2g}$, we can restrict our attention to the action of Γ on the subset $X \subset \mathbb{C}^{2g}$ whose elements have volume 1. In real and imaginary coordinates the set of periods of volume 1 is then the set of pairs $(x, y) \in \mathbb{R}^{2g} \times \mathbb{R}^{2g}$ such that $\omega(x, y) = 1$.

The simple real Lie group $G = \mathrm{Sp}(2g, \mathbb{R})$ acts transitively on the set of couples $(x, y) \in (\mathbb{R}^{2g})^2$ such that $\omega(x, y) = 1$, and that the stabilizer of the couple

$$((1, 0, \dots, 0), (0, 1, 0, \dots, 0)) \text{ is the group } \begin{pmatrix} 1 & & \\ & 1 & \\ & & \mathrm{Sp}(2g-2, \mathbb{R}) \end{pmatrix}$$

that we will denote by U in the sequel. Our set X is isomorphic to the homogeneous space G/U . The linear action of Γ on X is under the isomorphism $X \simeq G/U$ given by left multiplication on G/U .

Since the group G is simple, that U is generated by unipotent elements, and that Γ is a lattice in G , Ratner's theorem [59] tells us that the closure of the Γ -orbits on X are homogeneous in the following sense

Theorem 9.2 (Ratner). *For every $p \in X$ of the form $p = gU$, there exists a closed subgroup H of G containing $U^g = gUg^{-1}$, such that $\Gamma \cap H$ is a lattice in H , and such that $\overline{\Gamma \cdot p} = \Gamma Hp$.*

Notice that in our situation, we have $U^g = I|_W \oplus \mathrm{Sp}(W^\perp) \simeq \mathrm{Sp}(2g-2, \mathbb{R})$ where $W = \mathbb{R}\mathfrak{K}p + \mathbb{R}\mathfrak{S}p \subset \mathbb{R}^{2g}$ is the symplectic subspace associated to the volume one $p \in \mathbb{C}^2$.

Let H_0 be the connected component of H containing the identity: then $\Gamma \cap H_0$ is still a lattice in H_0 , and U^g is contained in H_0 .

If $H_0 = G$ then $\overline{\Gamma \cdot p} = G$ and we deduce that the orbit closure is dense in X . Since the closure of $\Lambda(p)$ contains all the $\Lambda(q)$ of elements $q \in \overline{\Gamma \cdot p}$, we have $\overline{\Lambda(p)} = \mathbb{C}$.

If H_0 is a proper subgroup of G , Kapovich observes that it falls into two categories

- (Semi-simple case) H_0 is of the form $S \oplus \mathrm{Sp}(W^\perp)$, where S is a Lie subgroup of $\mathrm{Sp}(W)$.
- (Non semi-simple case) H_0 is not semi-simple and preserves a line $L \subset W$.

The proof of this dichotomy can be found in [42, p. 12], and is based on Dynkin's classification of maximal connected complex Lie subgroups of $\mathrm{Sp}(2g, \mathbb{C})$, see [21]. Let L be a maximal complex Lie subgroup of $\mathrm{Sp}(2g, \mathbb{C})$ which contains H_0 . If $H_0 \neq \mathrm{Sp}(2g, \mathbb{R})$, its Zariski closure in the complex domain is a strict subgroup of $\mathrm{Sp}(2g, \mathbb{C})$, so it is contained in a maximal complex Lie (strict) subgroup of $\mathrm{Sp}(2g, \mathbb{C})$. It satisfies one of the following properties (see [27, Ch. 6, Thm 3.1, 3.2]):

- (1) $L = \mathrm{Sp}(V) \oplus \mathrm{Sp}(V^\perp)$ for some complex symplectic subspace $V \subset \mathbb{C}^{2g}$,
- (2) L is conjugated to $\mathrm{Sp}(s, \mathbb{C}) \otimes \mathrm{SO}(t, \mathbb{C})$ where $2g = st$, $s \geq 2$, $t \geq 3$, $t \neq 4$ or $t = 4$ and $s = 2$,
- (3) L preserves a line of \mathbb{C}^{2g} .

Since H_0 contains U^g , L contains the complexification of U^g , which is nothing but $\mathrm{Id}_{W_\mathbb{C}} \oplus \mathrm{Sp}(W_\mathbb{C}^\perp)$, where $W_\mathbb{C}$ denotes the complexification $W \otimes_\mathbb{R} \mathbb{C}$ of W . In case (1), the only possibility is that up to permutation of V and V^\perp , we have $W_\mathbb{C} = V$. In particular, H_0 is a subgroup of $\mathrm{Sp}(W) \oplus \mathrm{Sp}(W^\perp)$. Since it contains $\mathrm{Id}_W \oplus \mathrm{Sp}(W^\perp)$, it must be of the form $S \oplus \mathrm{Sp}(W^\perp)$, where S is a Lie subgroup of $\mathrm{Sp}(W)$. Case (2) cannot occur. In case (3), observe that the line L needs to be in $W_\mathbb{C}$, since the group $\mathrm{Id}_{W_\mathbb{C}} \oplus \mathrm{Sp}(W_\mathbb{C}^\perp)$ preserves this line. If L is defined over the reals, we are done. If not, both L and \overline{L} (the image of L by the complex conjugation) are preserved by H_0 , and thus H_0 is a subgroup of $\mathrm{Sp}(W) \oplus \mathrm{Sp}(W^\perp)$. As before, because it contains $\mathrm{Id}_W \oplus \mathrm{Sp}(W^\perp)$, it must be of the form $S \oplus \mathrm{Sp}(W^\perp)$, where S is a Lie subgroup of $\mathrm{Sp}(W)$.

Remark 9.3. If $\Gamma_W := \Gamma \cap (\mathrm{Id}_W \oplus \mathrm{Sp}(W^\perp))$ is a lattice in $\mathrm{Id}_W \oplus \mathrm{Sp}(W^\perp)$ then W is defined over \mathbb{Q} and the conclusion of the first item of Proposition 3.10 follows easily. Indeed, Γ_W acts by the identity on W^σ for every Galois automorphism σ . The Zariski closure of Γ_W being $\mathrm{Id}_W \oplus \mathrm{Sp}(W^\perp)$ (by Borel density theorem, see [73]), $\mathrm{Id}_W \oplus \mathrm{Sp}(W^\perp)$ acts by the identity on W^σ as well. This implies that $W^\sigma = W$ for every σ (otherwise $\mathrm{Id} \oplus \mathrm{Sp}(W^\perp)$ acts by the identity on the non-trivial subspace $(W + W^\sigma) \cap W^\perp$), and so W is rational.

Let us proceed to analyze the different cases.

Semi-simple case. Since the group $H_0 = S \oplus \mathrm{Sp}(W^\perp)$ contains a lattice, it must be unimodular. In particular, either S is the trivial group, or a 1-parameter subgroup, or the whole $\mathrm{Sp}(W)$. If S is trivial, then Ratner's Theorem tells us $\Gamma_W := \Gamma \cap (\mathrm{Id}_W \oplus \mathrm{Sp}(W^\perp))$ is a lattice in $\mathrm{Id}_W \oplus \mathrm{Sp}(W^\perp)$ and we fall in case (1) by Remark 9.3.

If S is 1-dimensional, $S \oplus \text{Id}_{W^\perp}$ would be the radical of H_0 , and a theorem of Wolf and Raghunathan, see [58], shows that it would intersect Γ in a lattice. This implies that the intersection of Γ with $\text{Id}_W \oplus \text{Sp}(W^\perp)$ is also a lattice. Indeed, let $\bar{\Gamma}$ denote the natural projection of Γ on $\text{Sp}(W^\perp)$. Since the sequence

$$0 \rightarrow S \cap \Gamma \rightarrow \Gamma \rightarrow \bar{\Gamma} \rightarrow 1$$

is exact, the map $\Gamma \rightarrow (S \cap \Gamma) \backslash S$ induced by the projection of $S \oplus \text{Sp}(W^\perp)$ to S induces a morphism $\bar{\Gamma} \rightarrow (\Gamma \cap S) \backslash S$. The group $(\Gamma \cap S) \backslash S$ is abelian, and since $\bar{\Gamma}$ has Kazhdan property (T), the image group in $(\Gamma \cap S) \backslash S$ is finite. We thus conclude that the image of Γ on the factor S is a lattice and therefore $\Gamma \cap (\text{Id}_W \oplus \text{Sp}(W^\perp))$ as well. We conclude as in Remark 9.3.

Finally, it remains to treat the case where $S = \text{Sp}(W)$. This case splits into two subcases, depending on the lattice $\Gamma \cap \text{Sp}(W) \oplus \text{Sp}(W^\perp)$ being reducible or irreducible. If it is reducible, this implies that $\Gamma \cap (\text{Id}_W \oplus \text{Sp}(W^\perp))$ is a lattice, and then W must be rational by the above considerations. Assume now that we are in the irreducible case. Then $g = 2$, by a theorem of Margulis [46]. Assume W is not rational, otherwise we are done. Let σ be a Galois automorphism such that $W^\sigma \neq W$. The group $\Gamma \cap \text{Sp}(W) \oplus \text{Sp}(W^\perp)$ preserves the splitting $W^\sigma \oplus (W^\sigma)^\perp$, since Γ and the symplectic form are defined over the rationals. Borel density theorem applied to the lattice $\Gamma \cap \text{Sp}(W) \oplus \text{Sp}(W^\perp)$ shows that $\text{Sp}(W) \oplus \text{Sp}(W^\perp)$ preserves the splitting $W^\sigma \oplus (W^\sigma)^\perp$. This implies that $W^\sigma = W^\perp$ and $(W^\sigma)^\sigma = W$. This being true for every Galois automorphism, this means that W is defined over a totally real quadratic field K , and we have $W^\sigma = W^\perp$ where σ is the Galois automorphism of K . This is the only situation where we fall in the last case of Proposition 3.10.

Non semi-simple case. We suppose that H_0 is not semi-simple and prove in this case that the periods p satisfy the second case of Proposition 3.10. We already know that there is a line $L \subset W$ that is invariant by the action.

For this, we will first need to understand in detail the subgroup B of $\text{Sp}(2g, \mathbb{R})$ formed by all elements that stabilize the line L , see [42, p. 10]. To unscrew the structure of B , notice that any element of B stabilizes both L and L^\perp so that we have an exact sequence

$$CH_{2g} \rightarrow B \rightarrow \text{Sp}(L^\perp/L) \simeq \text{Sp}(2g-2)$$

The group CH_{2g} is then the set of elements $M \in \text{Sp}(2g)$ which induce the identity map on L^\perp/L .

We now have another exact sequence

$$(60) \quad H_{2g-1} \rightarrow CH_{2g} \rightarrow \text{GL}(L) \simeq \mathbb{R}^*,$$

the last arrow being given by the restriction of an element $M \in CH_{2g}$ to the line L . Hence the subgroup $H_{2g-1} \subset CH_{2g}$ is the group of elements $M \in \text{Sp}(2g)$ which act as the identity on L and on L^\perp/L . Such M are easily seen to be of the form $M_{\varphi, \alpha}$, for some $\varphi \in (L^\perp/L)^*$ and $\alpha \in \mathbb{R}$, where

- the restriction of $M_{\varphi, \alpha}$ to L^\perp equals $\text{id}_{L^\perp} + \varphi a_1$
- $M_{\varphi, \alpha}(b_1) = \alpha a_1 + b_1 + \sum_{k \geq 2} \varphi(b_k) a_k - \varphi(a_k) b_k$,

where $a_1, b_1, \dots, a_g, b_g$ is a symplectic basis such that $L = \mathbb{R}a_1$. The group structure on H_{2g-1} is then given by the following relation

$$(61) \quad M_{\varphi, \alpha} M_{\varphi', \alpha'} = M_{\varphi + \varphi', \alpha + \alpha' + \omega(\varphi, \varphi')},$$

where $\omega(\varphi, \varphi')$ is the natural symplectic product induced by ω on $(L^\perp/L)^*$, namely

$$\omega(\varphi, \varphi') = \sum_{k \geq 2} \varphi(a_k) \varphi'(b_k) - \varphi'(a_k) \varphi(b_k).$$

Equation (61) is a straightforward computation. An equivalent formulation is that H_{2g-1} is the central extension

$$\mathbb{R} \rightarrow H_{2g-1} \rightarrow (L^\perp/L)^*,$$

defined by the 2-cocycle $(\varphi, \varphi') \mapsto \omega(\varphi, \varphi')$. The group H_3 is isomorphic to the classical Heisenberg group of upper triangular real matrices of size 3×3 with 1's on the diagonal.

Now CH_{2g} is a semi-direct product of \mathbb{R}^* by H_{2g-1} , see (60). To understand its structure, we introduce for every λ , one of its lift $S_\lambda \in CH_{2g}$ defined by

$$S_\lambda(a_1) = \lambda a_1, \quad S_\lambda(b_1) = \frac{1}{\lambda} b_1, \quad S_\lambda(a_k) = a_k, \quad S_\lambda(b_k) = b_k \text{ for } k \geq 2.$$

A trivial computation shows that for any $\lambda \in \mathbb{R}^*$, every $\varphi \in (L^\perp/L)^*$ and every $\alpha \in \mathbb{R}$, we have

$$(62) \quad S_\lambda M_{\varphi, \alpha} S_\lambda^{-1} = M_{\lambda\varphi, \lambda^2\alpha}.$$

This shows that CH_{2g} is not unimodular, and consequently does not contain any lattice.

By construction, our group H_0 is contained in B . We have an exact sequence $CH_{2g} \rightarrow B \rightarrow \text{Sp}(L^\perp/L, \omega)$. The image of H_0 by the right arrow is onto since H_0 contains U^g , so that H_0 itself splits as an exact sequence $CH_{2g} \cap H_0 \rightarrow H_0 \rightarrow \text{Sp}(L^\perp/L, \omega)$. The group $CH_{2g} \cap H_0$ is invariant under the action by conjugation of $\text{Sp}(L^\perp/L, \omega) \simeq \text{Sp}(W^\perp)$. The restriction of this action on H_{2g-1} can be described explicitly: for $U \in \text{Sp}(L^\perp/L) \simeq \text{Sp}(W^\perp)$ denote $s(U) = Id_W \oplus U \in H_0 \subset B$. Then

$$s(U) M_{\varphi, \alpha} s(U)^{-1} = M_{\varphi \circ s(U)^{-1}, \alpha}.$$

Lemma 9.4. *The closed non-trivial connected subgroups of CH_{2g} invariant by $\text{Sp}(W^\perp)$ are*

- (1) $Z(H_{2g-1})$
- (2) lifts of $GL^+(L)$ in CH_{2g}
- (3) H_{2g-1}
- (4) $GL^+(L) \ltimes Z(H_{2g-1})$
- (5) CH_{2g}

The proof of Lemma 9.4 is an easy consequence of the previous exact sequences and calculations.

Now $H_0 \cap CH_{2g}$ cannot fall in cases (1) and (2) of Lemma 9.4 since in either of those, H_0 would be semi-simple, contrary to hypothesis. It can neither fall in cases (4) or (5), since in those cases equation (62) does not allow H_0 to be unimodular. Hence we are left with the possibility $H_0 \cap CH_{2g} = H_{2g-1}$ (and $H_0 \simeq \text{Sp}(W^\perp) \ltimes H_{2g-1}$). In this case we will show that the invariant line L is rational, and thus we fall in the second possibility of Proposition 3.10

The theorem of Raghunathan and Wolf cited above tells us that $\Gamma \cap H_{2g-1}$ is a lattice in H_{2g-1} . By using Borel's density Theorem in [28, p.91] we deduce that its Zariski closure is H_{2g-1} . We have $L \subset K := \bigcap_{\gamma \in \Gamma \cap H_{2g-1}} \text{Ker}(\gamma - I)$ which is an intersection of rational spaces. If the inclusion is proper, then the Zariski closure of $\Gamma \cap H_{2g-1}$ would not be the whole of H_{2g-1} . This shows that $L = K$ and it is a rational one-dimensional subspace of \mathbb{R}^{2g} .

Up to a real affine change of coordinates on \mathbb{C} , we can assume that the imaginary part of p generates L , and that it is a primitive element of \mathbb{Z}^{2g} . Since the group H_{2g-1} acts transitively on the set of vectors $v \in \mathbb{R}^{2g}$ such that $v \cdot \Im p = 1$, while keeping the period $\Im p$ fixed, we see that $H \cdot p$ already contains all the periods q such that $\Im q = \Im p$ and such that $V(q) = V(p) = 1$. Since, Γ acts transitively on the set of primitive elements of \mathbb{Z}^{2g} , we infer that $\Gamma H p = \overline{\Gamma \cdot p}$ contains all the periods q with volume $V(p) = 1$ and with a primitive integer imaginary part. Since any periods of $\overline{\Gamma \cdot p}$ is of this form, we deduce that this situation is exactly the second case of the proposition. The proof of this latter is now complete.

Applying Moore's ergodic theorem in [55] to each case H above we deduce the final ergodicity part of Proposition 3.10.

10. APPENDIX II: PROOF OF LEMMA 3.9

Up to composing p by an element of $\text{GL}_2^+(\mathbb{R})$ we can assume that the image of p is the set of Gaussian integers. By Lemma 9.1, we can assume that there exists a symplectic basis $a_1, b_1, \dots, a_g, b_g$ in which the period p has the following form:

$$p(a_1) = p(a_2) = 1, \quad p(b_1) = p(b_2) = i \text{ and } p(a_k) = p(b_k) = 0 \text{ for } k \geq 3.$$

This basis permits to identify $H_1(\Sigma_g, \mathbb{Z})$ with \mathbb{Z}^{2g} equipped with the symplectic form

$$u \cdot v = u_1 v_2 - u_2 v_1 + \dots + u_{2g-1} v_{2g} - u_{2g} v_{2g-1},$$

and the group $\text{Aut}(H_1(\Sigma_g, \mathbb{Z}))$ with $\text{Sp}(2g, \mathbb{Z})$. In these coordinates we have

$$p(u_1, \dots, u_{2g}) = (u_1 + u_3) + (u_2 + u_4)i.$$

The form $u_1 + u_3 \in (\mathbb{Z}^{2g})^*$ is dual to the vector $P_1 = -(b_1 + b_2)$ (meaning that $u_1 + u_3 = -(b_1 + b_2) \cdot u$), whereas the form $u_2 + u_4 \in (\mathbb{Z}^{2g})^*$ is dual to $P_2 = a_1 + a_2$ (meaning $u_2 + u_4 = (a_1 + a_2) \cdot u$). We then have that an endomorphism of \mathbb{Z}^{2g} given by a matrix M stabilizes p if and only if $M(P_k) = P_k$ for $k = 1, 2$, and similarly an endomorphism of $(\mathbb{Z}/2\mathbb{Z})^{2g}$ given by a matrix $M[2]$ stabilizes $p[2]$ if and only if $M[2](P_k[2]) = P_k[2]$ for $k = 1, 2$. The condition can be read in the columns of the matrices. If C_k (resp. $C_k[2]$) denotes the k -th column of M (resp. $M[2]$) then

$$(63) \quad C_1 + C_3 = a_1 + a_2 \text{ and } C_2 + C_4 = b_1 + b_2.$$

$$(64) \quad (\text{resp. } C_1[2] + C_3[2] = a_1[2] + a_2[2] \text{ and } C_2[2] + C_4[2] = b_1[2] + b_2[2])$$

Let $M[2] \in \text{Stab}_{\text{Sp}(2g, \mathbb{Z}/2\mathbb{Z})}(p[2])$ a symplectic isomorphism whose columns $C_1[2], \dots, C_{2g}[2]$ satisfy (64). Let $C'_k \in \mathbb{Z}^{2g}$ be representatives of the classes $C_k[2] \in (\mathbb{Z}/2\mathbb{Z})^{2g}$, for $k = 1, \dots, 2g$, and M' be the square matrix whose columns are the C'_k . Up to changing the representative in each class, we can suppose (63) is valid for the columns of M' . We will assume in the sequel that these equations are always satisfied.

Our goal is to modify the vectors C'_k by defining

$$C_k := C'_k + 2E_k, \text{ with } E_k \in \mathbb{Z}^{2g}$$

in such a way that the matrix $M := (C_1, \dots, C_{2g})$ not only stabilizes p (satisfying (63)) but also belongs to $\text{Sp}(2g, \mathbb{Z})$. It is therefore necessary to impose

$$(65) \quad E_1 + E_3 = 0 \text{ and } E_2 + E_4 = 0.$$

$$(66) \quad C_1, \dots, C_{2g} \text{ forms a symplectic basis of } \mathbb{Z}^{2g}$$

Main step: construction of C_1, C_2, C_3, C_4 . The conditions we need to satisfy are

- (1) $C_1 \cdot C_2 = 1$
- (2) $C_1 \cdot C_3 = 0$, or equivalently $C_1 \cdot (a_1 + a_2) = 0$
- (3) $C_1 \cdot C_4 = 0$, or equivalently, knowing (1): $C_1 \cdot (b_1 + b_2) = C_1 \cdot C_2 = 1$
- (4) $C_2 \cdot C_3 = 0$, or equivalently, knowing (1): $C_2 \cdot (a_1 + a_2) = C_2 \cdot C_1 = -1$
- (5) $C_2 \cdot C_4 = 0$, or equivalently, $C_2 \cdot (b_1 + b_2) = 0$
- (6) $C_3 \cdot C_4 = 1$, or equivalently, $((a_1 + a_2) - C_1) \cdot ((b_1 + b_2) - C_2) = 1$

If (63) and (1)–(5) are true, we have that (6) is automatically satisfied

$$\begin{aligned} & ((a_1 + a_2) - C_1) \cdot ((b_1 + b_2) - C_2) = \\ & = (a_1 + a_2) \cdot (b_1 + b_2) - (a_1 + a_2) \cdot C_2 - C_1 \cdot (b_1 + b_2) + C_1 \cdot C_2 = 2 - 1 - 1 + 1 = 1. \end{aligned}$$

Let us first find C_1, C_2, C_3, C_4 so that conditions (2)–(5) are satisfied. In real and imaginary coordinates this is equivalent to

$$p(E_1) = \left(\frac{1 - C'_1 \cdot (b_1 + b_2)}{2}, \frac{C'_1 \cdot (a_1 + a_2)}{2} \right),$$

and

$$p(E_2) = \left(\frac{(b_1 + b_2) \cdot C'_2}{2}, \frac{1 + C'_2 \cdot (a_1 + a_2)}{2} \right).$$

Observe that the right hand sides of the last two equations are integers because C'_k 's are lifts of - the elements of the symplectic basis- $C_k[2]$'s and satisfy (63). By surjectivity of p there exist solutions to the equations (2)–(5), and so we can assume that the C'_k 's satisfy (2)–(5). We can now replace C'_k by $C_k = C'_k + 2E_k$ for $k = 1, 2$ with

$$E_k \in (a_1 + a_2)^\perp \cap (b_1 + b_2)^\perp \text{ for } k = 1, 2.$$

to preserve conditions (2)–(5). Condition (1) is equivalent to

$$1 = C_1 \cdot (C'_2 + 2E_2) = C_1 \cdot C'_2 + 2C_1 \cdot E_2,$$

and we know that $C_1 \cdot C'_2$ is odd. So we are done if we can choose C_1 so that the map

$$(67) \quad E_2 \in (a_1 + a_2)^\perp \cap (b_1 + b_2)^\perp \mapsto C_1 \cdot E_2 \in \mathbb{Z}$$

is onto. We have $(a_1 + a_2)^\perp \cap (b_1 + b_2)^\perp = \mathbb{Z}(a_1 - a_2) + \mathbb{Z}(b_1 - b_2) + \sum_{k \geq 3} \mathbb{Z}a_k + \mathbb{Z}b_k$. So the map (67) is onto if and only if

$$(68) \quad \gcd(C_1 \cdot (a_1 - a_2), C_1 \cdot (b_1 - b_2), C_1 \cdot a_3, C_1 \cdot b_3, \dots, C_1 \cdot a_g, C_1 \cdot b_g) = 1.$$

However, observe the following:

- $C_1 \cdot (b_1 - b_2) = C_1 \cdot (b_1 + b_2)[2] = 1[2]$ is odd, and
- If $g \geq 3$, choosing E_1 appropriately, we can assume that the value of $C_1 \cdot a_3$ is either 1 or 2.

So (68) is satisfied with these choices of E_1 . Hence the main step is achieved.

To conclude the proof of Lemma 3.9, we need to construct the columns C_5, \dots, C_{2g} . We will inductively find the pair C_5, C_6 , then the pair C_7, C_8 , etc.. Let us construct the first one. We need to find E_5, E_6 so that

$$C_5, C_6 \in (C_1, C_2, C_3, C_4)^\perp \text{ and } C_5 \cdot C_6 = 1.$$

This means that the equations

$$(69) \quad (E_k \cdot C_1, \dots, E_k \cdot C_4) = -\frac{1}{2}(C'_k \cdot C_1, \dots, C'_k \cdot C_4) \text{ for } k = 5, 6$$

and

$$(70) \quad C_5 \cdot (C'_6 + 2E_6) = 1$$

hold. We can find E_5 and E_6 so that (69) is satisfied, since C_1, C_2, C_3, C_4 is a symplectic family (i.e. $C_1 \cdot C_2 = C_3 \cdot C_4 = 1$ and other products are zero). So we can assume that C'_5 and C'_6 belong to the orthogonal of C_1, C_2, C_3, C_4 , which is a symplectic submodule of \mathbb{Z}^{2g} isomorphic to \mathbb{Z}^{2g-4} with the canonical symplectic product. In these coordinates, one of the coordinates of C'_5 is odd (since $C_5[2]$, the reduction of C'_5 modulo 2, is non zero) so by adding to C'_5 an even vector of \mathbb{Z}^{2g-4} one can assume that one of the coordinates of C'_5 is equal to 1. Hence, equation (70) being equivalent to $C_5 \cdot E_6 = \frac{-C_5 \cdot C'_6}{2}$ and the product $C_5 \cdot C_6$ being even, equation (70) can be solved for some suitable choice of E_6 . We get C_5 and C_6 in this way. For the construction of the other pairs by induction, the argument is similar.

The first part of the lemma follows, namely the surjectivity of the map $\text{Stab}_{\text{Aut}(H_1(\Sigma_g, \mathbb{Z}))}(p) \rightarrow \text{Stab}_{\text{Aut}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}))}(p[2])$.

For the second part, it suffices to remark that the stabilizer of $p[2]$ in $\text{Aut}(H_1(\Sigma_g, \mathbb{Z}/2\mathbb{Z}))$ is the stabilizer of a pair of non colinear elements of $H^1(\Sigma_g, \mathbb{Z}/2\mathbb{Z})$ that do not intersect each other. The number of elements in this group is given by the announced formula, by elementary considerations.

11. APPENDIX III: A RESULT IN PICARD-LEFSCHETZ THEORY

We prove here a result which is a well-known consequence of Picard-Lefschetz theory, but which we cannot find as such in the literature.

Let $h : S \rightarrow C$ be a holomorphic map from a compact complex surface S to a compact complex curve C . We assume both S and C are non singular, that the critical points of h are non degenerate, and that the fibers of h are connected.

Lemma 11.1. *For every $c \in C$, the sequence*

$$(71) \quad \pi_1(h^{-1}(c)) \rightarrow \pi_1(S) \rightarrow \pi_1(C)$$

where the map on the left is induced by inclusion and the one on the right by h_ , is exact.*

Proof. Let us first prove the claim for c a regular value of h . To do so, given a lift $w \in h^{-1}(c)$, we will use some useful loops $\tilde{\mu}_i : [0, 1] \rightarrow S$ starting and ending at w that do not intersect the critical fibers of h .

Denote by c_i , $i = 1, \dots, r$ the critical values of h . At any regular point p_i of the curve $h^{-1}(c_i)$, the map h induces a biholomorphism between a compact disc D_i transversal to $h^{-1}(c_i)$ at p_i and a neighborhood δ_i of c_i in C . We can assume that the discs $\delta_i := h(D_i)$ are disjoint.

Given a point $w_i \in \partial D_i$, let $\gamma_i : [0, 1] \rightarrow E \setminus \cup_i \text{Int}(\delta_i)$ be a family of paths with origin c and end point $h(w_i)$, having the property of being disjoint appart from their origin. Since h induces a C^∞ -fibration with connected fibers from $S \setminus \cup_i h^{-1}(c_i)$ to $C \setminus \{c_i\}$, given a lift $w \in h^{-1}(c)$, we can lift the path γ_i to a path $\tilde{\gamma}_i$ starting at w and ending at w_i . We denote by μ_i the concatenation $\gamma_i \star \partial \delta_i \star \gamma_i^{-1}$ and by $\tilde{\mu}_i$ its lift $\tilde{\gamma}_i \star \partial D_i \star \tilde{\gamma}_i^{-1}$. Notice the following two facts:

- (1) $\tilde{\mu}_i$ is homotopically trivial in S ,
- (2) the representatives of the loops μ_i in $\pi_1(C \setminus \{c_i\}, c)$ generate the kernel of the map

$$(72) \quad \pi_1(C \setminus \{z_i\}, c) \rightarrow \pi_1(C, c)$$

induced by inclusion.

(The first is clear, and the second comes from the fact that the paths γ_i are disjoint appart from their origin.)

We are now in a position to prove the claim in the case of a regular fiber. Consider an element in the kernel of the map $h_* : \pi_1(S, w) \rightarrow \pi_1(C, c)$ represented by a loop $\varepsilon : [0, 1] \rightarrow S$ starting and ending at w . Up to homotopy, we can assume that ε intersects none of the critical fibers of h . The image $h \circ \varepsilon$ is then a loop starting and ending at c which defines an element of the fundamental group of $\pi_1(C \setminus \{z_i\}, c)$ belonging to the kernel of the map (72). Hence, $h \circ \varepsilon$ is homotopic to a word $W(\mu_1, \dots, \mu_r)$. Consider the path $\varepsilon' = \varepsilon \star W(\tilde{\mu}_1, \dots, \tilde{\mu}_r)^{-1}$; by construction,

- (i) ε' is homotopic to ε in S
- (ii) it intersects none of the critical fibers of h ,
- (iii) its image $h \circ \varepsilon'$ with fixed extremities c is homotopic in $C \setminus \{c_i\}$ to the constant path c .

Since the restriction of h induces a locally trivial C^∞ fibration from $S \setminus \cup_i h^{-1}(c_i)$ to $C \setminus \{c_i\}$ — this is a proper submersion, so this is Ehresmann's fibration theorem — we can lift the homotopy found in (iii) to a homotopy in $S \setminus \cup_i h^{-1}(c_i)$ between ε' and a loop contained in the fiber $h^{-1}(c)$. This establishes that ε is homotopic with fixed extremities to a loop contained in the fiber $h^{-1}(c)$, and ends the proof of the claim in the case of a regular fiber.

It remains to prove the claim for the singular fibers of h . For each i , there exists a neighborhood U of $h^{-1}(c_i)$ that retracts by deformation on $h^{-1}(c_i)$ (see [3, Chapter X, Section 9]); in particular, the images of $\pi_1(h^{-1}(c_i))$ and $\pi_1(U)$ in $\pi_1(S)$ coincide. Since U contains a regular fiber, we know from the preceeding discussion that the image of $\pi_1(U)$ in $\pi_1(S)$ contains $\text{Ker}(h_*)$. On the other hand, the image of $\pi_1(h^{-1}(c_i))$ in $\pi_1(S)$ is contained in $\text{Ker}(h_*)$ since h is constant on $h^{-1}(c_i)$, so the sequence (71) is exact for the critical value $c = c_i$, and the Lemma follows. \square

Corollary 11.2. *For every $c \in C$, the sequence*

$$(73) \quad H_1(h^{-1}(c), \mathbb{Z}) \rightarrow H_1(S, \mathbb{Z}) \rightarrow H_1(C, \mathbb{Z})$$

where the map on the left is induced by inclusion and the one on the right by h_* , is exact.

Proof. This is a consequence of Lemma 11.1 and Hurewicz' theorem. \square

REFERENCES

- [1] W. ABIKOFF Degenerating families of Riemann surfaces, *Ann. of Math.* (2) 105 (1977), no. 1, pp. 29–44,
- [2] L. V. AHLFORS The complex analytic structure of the space of closed Riemann surfaces, in *Analytic functions*, Princeton Univ. Press, Princeton, N.J. (1960) pp. 45–66
- [3] E. ARBARELLO, M. CORNALBA, P. GRIFFITHS, *Geometry of Algebraic Curves*, Vol. II, Springer Verlag, (2011)
- [4] V. I. ARNOLD A remark on the branching of hyperelliptic integrals as functions of the parameters, *Funkcional. Anal. i Prilozhen.* 2 1968 no. 3, pp. 1–3.
- [5] M. BAINBRIDGE. Euler characteristics of Teichmüller curves in genus two. *Geometry and Topology* 11(4), pp. 1887–2073.
- [6] M. BAINBRIDGE, D. CHEN, Q. GENDRON, S. GRUSHEVSKY, M. MLLER Compactification of strata of Abelian differentials, *Duke Math. J.* 167 (2018), no. 12, pp. 2347–2416.
- [7] M. BAINBRIDGE, D. CHEN, Q. GENDRON, S. GRUSHEVSKY, M. MLLER The moduli of multi-scale differentials, *arXiv:1910.13492*, pp. 1–122
- [8] M. BAINBRIDGE, C. JOHNSON, C. JUDGE, AND I. PARK. Haupt’s theorem for strata of abelian differentials. *arXiv:2002.12901*
- [9] BENSON FARB AND DAN MARGALIT. A primer on Mapping Class groups, Princeton University Press, 2001.
- [10] I. BERSTEIN AND A. L. EDMONDS On the classification of generic branched coverings of surfaces, *Illinois Jr. of Math.*, Vol. 24, Nr. 1, (1984), pp. 64–82
- [11] L. BERS Spaces of degenerating Riemann surfaces, *Discontinuous groups and Riemann surfaces* (Proc. Conf., Univ. Maryland, College Park, Md., 1973), Princeton Univ. Press, Princeton, N.J., 1974, pp. 43–55. *Ann. of Math. Studies*, no. 79
- [12] L. BERS Deformations and moduli of Riemann surfaces with nodes and signatures, *Math. Scand.* 36 (1975), 12716, collection of articles dedicated to Werner Fenchel on his 70th birthday.
- [13] L. BERS Finite dimensional Teichmüller spaces and generalizations, *Bull. AMS*, Vol. 5, Nr. 2, 1981
- [14] P. BUSER *Geometry and spectra of compact Riemann surfaces*, Progress in Mathematics, vol. 106, Birkhäuser Boston Inc., Boston, MA, 1992,
- [15] G. CALSAMIGLIA, B. DEROIN, S. FRANCAVIGLIA. Branched projective structures with Fuchsian holonomy. *Geometry and Topology* 18 (2014) pp. 379–446.
- [16] A. CALDERON, N. SALTER. Relative homological representations of framed mapping class groups, to appear in *B. Lond. Math. Soc.* (2020)
- [17] A. CALDERON, N. SALTER. Connected components of strata of Abelian differentials over Teichmüller space *Comment. Math. Helv.* 95 (2): 361–420 (2020).
- [18] K. CALTA Veech surfaces and complete periodicity in genus two. *J. Amer. Math. Soc.* 17 (2004), no. 4, pp. 871–908.
- [19] C. H. CLEMENS. *A Scrapbook of Complex Curve Theory*. 1980. Plenum Press., New York and London.
- [20] P. DELIGNE AND D. MUMFORD The irreducibility of the space of curves of given genus, *Inst. Hautes Études Sci. Publ. Math.* (1969), no. 36, pp. 71–109
- [21] E. B. DYNKIN. Maximal subgroups of the classical groups. *Trudy Moskov. Mat. Obsc.* 1, 1952, pp. 39–166.
- [22] A. L. EDMONDS Deformation of maps to branched coverings in dimension two. *Ann. of Math.* (2) 110 (1979), no. 1, pp. 113–125.
- [23] A. ESKIN, H. MASUR, A. ZORICH. Moduli spaces of abelian differentials: the principal boundary, counting problems and the Siegel-Veech constants. *Publ. Math. Inst. Hautes Études Sci.* No. 97 (2003), pp. 61–179.
- [24] A. ESKIN AND M. MIRZAKHANI. Invariant and stationary measures for the action on Moduli space. *Publ. Math. Inst. Hautes Études Sci.* No. 127, (2018) pp. 95–324
- [25] B. FARB, D. MARGALIT A primer on mapping class groups, Princeton University Press, 2012
- [26] L. GERRITZEN, F. HERRLICH. The extended Schottky space. *J. Reine Angew. Math.* 389 (1988), pp. 190–208
- [27] V. GORBATSEVICH, A. ONISHCHIK, È. VINBERG. *Lie groups and Lie algebras*. III Springer-Verlag. Berlin. 1994. Structure of Lie groups and Lie algebras.
- [28] M. GROMOV Rigid transformations groups, in *Géométrie différentielle* (Paris 1986) (D. Bernard, Y. Choquet-Bruhat ed.) *Travaux en Cours*, 33, Hermann, Paris (1988), pp. 65–139.
- [29] U. HAMENSTÄDT. Dynamical properties of the absolute period foliation, *Israel Journal of Math.* 225, (2018), pp. 661–680

- [30] U. HAMENSTÄDT. Quotients of the orbifold fundamental group of strata of abelian differentials. Preprint, 2018.
- [31] U. HAMENSTÄDT. On the orbifold fundamental group of the odd component of the stratum $H(2, \dots, 2)$. Preprint, 2020.
- [32] J. HARRIS, I. MORRISON. Moduli of curves, Graduate Texts in Mathematics 187, Springer, New York (1998)
- [33] R. HARTSHORNE Algebraic Geometry, Graduate texts in mathematics, 52, 1997 Springer-Verlag, N.Y.
- [34] O. HAUPT. Ein Satz über die Abelschen integrale I. Gattung. Math. Z. 6 (1920), no. 3-4, pp. 219—237.
- [35] W.J. HARVEY Boundary Structure of the Modular Group, Riemann Surfaces and Related Topics. Proceedings of the 1978 Stony Brook Conference . 1981.
- [36] F. HERRLICH The extended Teichmüller space, Math. Z. 203 (1990), no. 2, pp. 279—291
- [37] V. HINICH, A. VAINTROB Augmented Teichmüller spaces and orbifolds , Sel. Math. New Ser. (2010) 16, pp. 533–629
- [38] HOOPER W.P., WEISS B. Rel leaves of the Arnoux-Yoccoz surfaces, Selecta Mathematica, Volume 24, pp. 875–934 (2018)
- [39] J.H. HUBBARD AND S. KOCH An analytic construction of the Deligne-Mumford compactification of the moduli space of curves, J. of Diff. Geom. 98 (2014), pp. 261–313
- [40] P. HUBERT, H. MASUR, T. A. SCHMIDT AND A. ZORICH Problems on billiards, flat surfaces and translation surfaces, in Problems on mapping class groups and related topics, Proc. Sympos. Pure Math., vol. 74, Amer. Math. Soc, 2006, pp. 233–243.
- [41] D. JOHNSON Conjugacy relations in subgroups of the mapping class group and a group-theoretic description of the Rohlin invariant, Math. Ann., 249(3), 1980, pp. 243—263 .
- [42] M. KAPOVICH. Periods of abelian differentials and dynamics. "Dynamics: Topology and Numbers" (Proceedings of Kolyada Memorial Conference), Contemporary Mathematics, AMS, vol. 744, 2020, pp. 297—315.
- [43] M. KONTSEVICH, A. ZORICH. Connected components of the moduli spaces of Abelian differentials with prescribed singularities. Invent. Math. 153 (2003), no. 3, pp. 631–678.
- [44] T. LE FILS. Periods of abelian differentials with prescribed singularities. arXiv:2003.02216
- [45] A. G. MAIER. Trajectories on the closed orientable surfaces. (In Russian) Sb. Math., 12 (54), (1943), pp. 71–83.
- [46] G. A. MARGULIS. Discrete subgroups of semi-simple Lie groups. Springer-Verlag. 1991.
- [47] H. MASUR. Closed trajectories for quadratic differentials with an application to billiards. Duke Math. J. Vol. 53 No 2 (1986) 307–314.
- [48] H. MASUR. Extension of the Weil-Petersson metric to the boundary of Teichmüller space, Duke Math. J. 43 (1976), no. 3, pp. 623–635.
- [49] H. MASUR. Interval exchange transformations and measured foliations, Ann. of Math. .2/ 115 (1982), pp. 169—200
- [50] C. McMULLEN. Moduli spaces of isoperiodic forms on Riemann surfaces. Duke Math. J. Vol. 163, Number 12 (2014), pp. 2271–2323.
- [51] C. McMULLEN. Navigating moduli space with complex twists. J. Eur. Math. Soc. 15(2013), pp. 1223–1243
- [52] C. McMULLEN. Dynamics of $SL_2(\mathbb{R})$ over moduli space in genus two. Ann. of Math. (2) 165 (2007), no. 2, pp. 397–456.
- [53] C. McMULLEN. Foliations of Hilbert Modular Surfaces. American Journal of Mathematics Vol. 129, No. 1 (Feb., 2007), pp. 183–215
- [54] G. MESS. The Torelli groups for genus 2 and 3 surfaces. Topology 31 (1992), no. 4, pp. 775–790.
- [55] C. C. MOORE. Ergodicity of flows on homogeneous spaces. Amer. J. Math. 88 (1966) pp. 154–178
- [56] H. MOVASATI. On elliptic modular foliations. Indagationes Mathematicae, Volume 19, Issue 2, June 2008, pp. 263–286
- [57] A. PUTMAN. Cutting and pasting in the Torelli group, Geometry & Topology 11 (2007), pp. 829–865
- [58] M. S. RAGHUNATHAN. Discrete subgroups of Lie groups. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 68. Springer-Verlag, New York-Heidelberg, 1972. ix+227
- [59] M. RATNER. Raghunathan’s topological conjecture and distributions of unipotent flows. Duke Math. J. 63 (1991), no. 1, pp. 235–280.
- [60] M. SCHIFFER. A method of variation within the family of simple functions. Proc. London Math. Soc. 44 (1938), pp. 450–452.

- [61] M. SCHMOLL. Spaces of elliptic differentials. Algebraic and topological dynamics. Contemp. Math. vol. 385, AMS, Providence, RI, 2006, pp. 193–206.
- [62] C. SIMPSON. Lefschetz theorems for the integral leaves of a holomorphic one-form. Compositio Math. 87 (1993) no. 1, pp. 99–113
- [63] K. STREBEL. Quadratic differentials. Ergebnisse der Mathematik. Springer-Verlag, 1984.
- [64] R. TORELLI. Sulle varietà di Jacobi. Rendiconti della Reale accademia nazionale dei Lincei. (1913) 22 (5), 98–103.
- [65] W.A. VEECH. Moduli spaces of quadratic differentials, J. Analyse Math. 55 (1990), pp. 117–171.
- [66] C. VOISIN. Hodge Theory and Complex Algebraic Geometry II, Cambridge Studies in Advanced Mathematics, 2003.
- [67] M. WOLF, S. WOLPERT. Real analytic structures on the moduli space of curves, Amer. J. Math. 114 (1992), no. 5, 1079–1102
- [68] S. WOLPERT. Geometry of the Weil-Petersson completion of Teichmüller space, Surveys in Differential Geometry, Vol. VIII (Boston, MA, 2002), Surv. Differ. Geom., VIII, Int. Press, Somerville, MA, 2003, pp. 357–393
- [69] S. WOLPERT. Families of Riemann surfaces and Weil-Petersson geometry, CBMS Regional Conference Series in Mathematics, vol. 113, published for the Conference Board of the Mathematical Sciences, Washington, DC, 2010
- [70] A. WRIGHT. Cylinder deformations in orbit closures of translation surfaces. Geometry & Topology, 19(1), 2015, pp. 414–438
- [71] S. YAMADA. On the geometry of Weil-Petersson completion of Teichmüller spaces, Math. Res. Lett. 11 (2004), no. 2-3, pp. 327–344
- [72] F. YGOUF. A criterion for density of the isoperiodic leaves in rank 1 affine invariant suborbifolds. arXiv:2002.01186
- [73] R.J. ZIMMER. Ergodic theory and semi-simple groups. Monographs in Mathematics. Vol. 81. Birkhäuser, 1984.

DEPARTAMENTO DE MATEMÁTICA APLICADA, UNIVERSIDADE FEDERAL FLUMINENSE, RUA PROFESSOR MARCOS WALDEMAR DE FREITAS REIS, S/N, CAMPUS DO GRAGUATÁ, BLOCO G, NITERÓI, 24210-201 BRAZIL
E-mail address: `gcalsamiglia@id.uff.br`

CNRS & UNIVERSITÉ CERGY-PONTOISE, UMR 8088, 2 AV. ADOLPHE CHAUVIN, 95302 CERGY-PONTOISE CEDEX, FRANCE
E-mail address: `bertrand.derooin@u-cergy.fr`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, P.ZZA PORTA S. DONATO 5, 40126 BOLOGNA, ITALY
E-mail address: `stefano.francaviglia@unibo.it`