



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE  
DELLA RICERCA

## Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Timely adjuvant chemotherapy improves outcome in dogs with non-metastatic splenic hemangiosarcoma undergoing splenectomy

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Timely adjuvant chemotherapy improves outcome in dogs with non-metastatic splenic hemangiosarcoma undergoing splenectomy / Faroni, E; Sabbatini, S; Guerra, D; Iannuzzi, C; Chalfon, C; Agnoli, C; Stefanello, D; Polton, G; Ramos, S; Aralla, M; Ciaccini, R; Foglia, A; Okonji, S; Marconato, L. - In: VETERINARY AND COMPARATIVE ONCOLOGY. - ISSN 1476-5810. - ELETTRONICO. - 21:1(2023), pp. 123-130. [10.1111/vco.12875]

*Availability:*

This version is available at: <https://hdl.handle.net/11585/921431> since: 2023-03-28

*Published:*

DOI: <http://doi.org/10.1111/vco.12875>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Faroni E, Sabattini S, Guerra D, et al. Timely adjuvant chemotherapy improves outcome in dogs with non-metastatic splenic hemangiosarcoma undergoing splenectomy. *Vet Comp Oncol.* 2023;21(1):123-130.

The final published version is available online at:

<https://doi.org/10.1111/vco.12875>

#### Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

1 **Timely adjuvant chemotherapy improves outcome in dogs with non-metastatic splenic**  
2 **hemangiosarcoma undergoing splenectomy**

3 **Word count:** 3167

4 **Number of figures and tables:** 2 figures, 4 tables

5 **Conflict of interest declaration:** The authors declare that there is no conflict of interest.

6 **Abstract**

7 Timely delivery of adjuvant chemotherapy has been shown to be advantageous in many human  
8 cancers and canine osteosarcoma. Adjuvant chemotherapy has been shown to improve outcome  
9 for canine splenic hemangiosarcoma. The aim of this retrospective study was to investigate  
10 whether timely adjuvant chemotherapy administration resulted in better outcome in dogs with  
11 non-metastatic splenic hemangiosarcoma undergoing splenectomy. Medical records were  
12 searched for dogs with non-metastatic, splenic hemangiosarcoma that received splenectomy and  
13 adjuvant chemotherapy. The number of days from surgery to the first chemotherapy dose (StoC)  
14 was evaluated to identify the cut-off value associated with the best survival advantage. StoC and  
15 other possible prognostic factors were tested for influence on time to metastasis (TTM) and  
16 overall survival (OS). Seventy dogs were included. Median StoC was 20 days (range, 4-70). The  
17 time interval associated with the greatest survival benefit was 21 days. Median TTM and OS of  
18 dogs with StoC  $\leq$ 21 days were significantly longer than those with StoC  $>$ 21 days (TTM: 163 vs. 118  
19 days,  $P=0.001$ ; OS: 238 vs. 146 days,  $P<0.001$ ). On multivariable analysis, StoC  $>$ 21 days was the  
20 only variable significantly associated with increased risk of tumor progression (HR 2.1,  $P=0.010$ )  
21 and death (HR 2.3;  $P=0.008$ ). Starting adjuvant chemotherapy within 21 days of surgery may be  
22 associated with a survival benefit in dogs with non-metastatic splenic hemangiosarcoma, possibly  
23 due to the early targeting of newly-recruited metastatic cells after surgery.

24 **Keywords**

25 Canine, doxorubicin, micrometastasis, prognosis, spleen, surgery

26 **Introduction**

27 In dogs, hemangiosarcoma (HSA) is a common and aggressive cancer of vascular origin. The spleen  
28 is the most common primary site, with nearly 50% of dogs having evidence of metastatic disease  
29 at presentation.<sup>1,2</sup> Splenectomy as the sole treatment modality is considered palliative; the  
30 addition of adjuvant doxorubicin-based chemotherapy usually only increases the average survival  
31 time by 2–4 months.<sup>3-5</sup>

32 There is accumulating evidence that surgical resection may enhance metastatic shedding of tumor  
33 cells. According to several hypotheses, surgery may promote tumor cell migration or trigger the  
34 outgrowth of resting metastatic cells through the release of inflammatory factors, catecholamines  
35 and pro-metastatic enzymes.<sup>6,7</sup> In addition, surgery-induced local and systemic  
36 immunosuppression can impair antitumor immunity (e.g., decreased number of natural killers [NK]  
37 and monocyte/macrophages, release of inflammatory cytokines such as TNF- $\alpha$  and IL-1),  
38 contributing to tumor cell survival.<sup>8-11</sup> While timely delivery of adjuvant chemotherapy may  
39 prevent these undesired effects, early administration may result in increased toxicity rates.<sup>12</sup>

40 Additionally, although never reported in dogs, there is a concern that early chemotherapy  
41 administration may increase post-surgical complications, including delayed wound healing and  
42 enhanced immunosuppression, due to cytotoxic effects on fibroblasts, thrombocytes, monocytes  
43 and leucocytes, inhibition of keratinocytes proliferation and decreased collagen synthesis.<sup>13-18</sup>

44 The timing of administration of adjuvant chemotherapy is a well-known prognostic variable for  
45 many human cancers (e.g., osteosarcoma, colon cancer), and it has gained growing interest in the  
46 veterinary community as well, with studies in mice and dogs reporting its impact.<sup>19-22</sup> Recently,

47 Marconato et al. demonstrated a survival benefit in dogs with appendicular osteosarcoma that  
48 received chemotherapy within 5 days of limb amputation.<sup>23</sup>

49 Variations in intervals between surgery and chemotherapy are common in clinical practice. This  
50 may depend on patients' medical conditions, owner's compliance and on the availability of a  
51 definitive diagnosis.

52 The aim of this study was to determine whether different time intervals between surgery and the  
53 first administration of chemotherapy would result in different outcomes, possibly identifying an  
54 optimal prognostic cut-off. Given the highly aggressive behaviour of canine splenic HSA, we  
55 hypothesized that starting chemotherapy early after splenectomy would improve outcome by  
56 targeting newly-recruited cycling metastatic cells after surgery.

## 57 **Materials and Methods**

### 58 *Study Design*

59 This was a retrospective study on client-owned dogs with spontaneous tumors seeking medical  
60 advice at a clinical facility. As the research did not influence any diagnostic or therapeutic decision,  
61 approval by an ethics committee was not required. All the examined samples and data were  
62 collected for diagnostic purposes as part of routine standard care.

63 The archives of seven veterinary oncology centers were retrospectively searched for dogs with  
64 splenic HSA undergoing splenectomy between January 2004 and December 2021. Dogs were  
65 included if they had a histologically confirmed splenic HSA, no evidence of metastases at the start  
66 of chemotherapy and received at least 1 dose of adjuvant cytotoxic chemotherapy. Dogs with  
67 incomplete staging or follow-up data were excluded. All owners provided written informed  
68 consent to the use of their dogs' medical records.

69 Collected information included signalment (i.e., breed, sex, neutering status, age, weight),  
70 presence and duration of clinical signs, clinical stage according to the World Health Organization  
71 staging system of domestic animals,<sup>24</sup> results of hematologic and coagulation analyses (i.e.,  
72 complete blood count [CBC], serum biochemical profile, coagulation profile [i.e., activated partial  
73 thromboplastin time, prothrombin time], urinalysis), and need for blood product administration.

74 Pre-chemotherapy staging was required to be performed by means of total body CT scan (TBCT) or  
75 3-view thoracic radiographs, abdominal ultrasound, and echocardiography, in addition to  
76 cytological or histologic evaluation of any suspicious visceral lesion.

77 All dogs received adjuvant chemotherapy, consisting of single agent doxorubicin (intravenous, q3  
78 weeks), or a combination of doxorubicin and dacarbazine, as previously described.<sup>25</sup> For each case,  
79 the number of days between surgery and the first administration of chemotherapy (*surgery to*  
80 *chemotherapy, StoC*) was retrieved.

81 Routine monitoring for pulmonary and abdominal metastasis with thoracic radiographs and  
82 abdominal ultrasound, respectively, occurred every 2-3 cycles of chemotherapy, unless clinical  
83 signs suspicious for metastasis were present, in which case imaging was carried out sooner. Once  
84 chemotherapy was completed, dogs were restaged every 2-3 months.

85 In case of disease progression, metronomic therapy was offered. Also, based on owners' and  
86 clinicians' preference, dogs could receive metronomic therapy at the end of the doxorubicin-based  
87 protocol.

88 Clinical, hematologic, and biochemical chemotherapy-related adverse events (AEs) were assessed  
89 based on patient history provided by the owner, physical examination and blood work. AEs were  
90 graded according to the Veterinary Cooperative Oncology Group Common Terminology Criteria for  
91 Adverse Events (VCOG-CTCAE) v2 and retrieved from medical records.<sup>26</sup>

92 *Statistical Analysis*

93 Categorical variables were summarized as frequency (percentage), while numerical variables were  
94 summarized as median (range). Non-normality of numerical data was assessed using the Shapiro–  
95 Wilk test.

96 Time to metastasis (TTM) was defined as the time interval between the first administration of  
97 chemotherapy and detection of metastasis. Metastasis was defined as the clinical and/or imaging  
98 evidence of suspected tumoral lesion(s) in any anatomic site, with cytologic confirmation (when  
99 possible) and/or progression of lesion(s) during follow-up. Overall survival (OS) was defined as the  
100 time interval between surgery and death due to any cause. Survival estimates were computed as  
101 medians and 95% CIs.

102 Cox proportional hazards regression was performed to evaluate the influence of potentially  
103 prognostic variables on TTM and OS, including sex, neutering status, age, body weight, presence of  
104 clinical signs prior to diagnosis, type of imaging used for tumor staging (TBCT vs other),  
105 hemoabdomen, anemia (hematocrit < 37%), thrombocytopenia (total platelet number <  
106 160000/ $\mu$ L), coagulation abnormalities, blood product administration, chemotherapeutic protocol  
107 (doxorubicin vs doxorubicin and dacarbazine), administration of metronomic chemotherapy, and  
108 chemotherapy-related AEs. The continuous variables of age and body weight were converted to  
109 dichotomous variables with the median value used as the cut-off point. Variables with a P value <  
110 0.2 in the univariable analyses were further tested for independence of association with outcome  
111 in a multivariable Cox proportional hazards model.

112 StoC was analyzed as a categorical variable (within 7, 14, 21, 28, or > 28 days after surgery). The  
113 risk of tumor metastasis and death of the dogs within each interval category for StoC was then  
114 compared with that of the remaining dogs (e.g.,  $\leq$  7 days vs > 7 days, and  $\leq$  14 days vs > 14 days)

115 by means of Cox proportional hazards regression, and the interval with the highest, significant ( $P <$   
116 0.05) HR was selected as the optimal interval and included in the multivariable model.  
117 The two groups identified by the optimal interval were tested for distribution of the  
118 aforementioned possible prognostic variables and time of restaging by means of chi-  
119 square/Fisher's exact test and Mann-Whitney U test for categorical and continuous variables,  
120 respectively. Survival curves were generated with the Kaplan-Meier product limit method and  
121 then compared between StoC groups with the log-rank test. One-year survival rates of different  
122 StoC-defined groups were obtained.

### 123 **Cell Line Validation Statement**

124 No cell lines were used in the current study.

### 125 **Results**

#### 126 *Patients' characteristics*

127 Seventy dogs were included in the study. There were 37 (53%) males, of which 14 were neutered,  
128 and 33 (47%) females, of which 26 were spayed. Median age was 10 years (range, 6–15), and  
129 median weight was 28 kg (range, 6–58). The most represented breeds were mixed (n=21; 30%),  
130 Labrador retriever (n=11; 16%), and German shepherd (n=5; 7%).

131 Sixty-three (90%) dogs were symptomatic at admission, with median duration of clinical signs of 2  
132 days (range, 1 – 46), while HSA was an incidental finding in 7 (10%) cases. Reported clinical signs  
133 were lethargy, reduced appetite, exercise intolerance, tachypnea, and syncope.

134 Results of CBC were available for 64 (91%) cases. Forty nine (77%) dogs were anemic, with a  
135 median hematocrit of 29% (range, 13-36). Twenty-eight (44%) dogs were thrombocytopenic, with



136 a median platelet number of 89000/ $\mu$ L (range, 16000-153000). Results of coagulation times were  
137 available for 51 (73%) cases, with abnormal results registered in 4 (8%) of those.

138 Tumors were staged by means of 3-view thoracic radiography and abdominal ultrasonography in  
139 49 (70%) dogs, whereas 21 (30%) underwent TBCT scan. At admission, hemoabdomen was present  
140 in 58 (83%) dogs. Overall, 59 (84%) and 11 (16%) dogs had stage 2 and stage 1 disease,  
141 respectively.

#### 142 *Treatment*

143 Following splenectomy, no complication in wound healing was reported.

144 Eight (11%) dogs received heterologous packed red blood cells during the surgical procedure.

145 All dogs received adjuvant chemotherapy: 66 (94%) dogs received doxorubicin as single agent

146 (median, 4 doses; range, 1 to 6 doses), while 4 (6%) received doxorubicin and dacarbazine

147 (median, 4 doses; range, 2 to 6 doses). The median doxorubicin dose was 30 mg/m<sup>2</sup> (range, 25 to

148 30 mg/m<sup>2</sup>), and the median dacarbazine dose was 200 mg/m<sup>2</sup> (range, 160 to 210 mg/m<sup>2</sup>).

149 Chemotherapy-related AEs were experienced by thirty (43%) dogs, of which nine (13%) had grade  
150 3 or 4 toxicity. Overall, there were twenty-four episodes of gastrointestinal toxic effects, including

151 four grade 3 (n=2 vomiting, n=1 diarrhea) and one grade 4 (diarrhea); there were four episodes of

152 hematologic toxicity, including three grade 4 (neutropenia in all cases). The three dogs with grade

153 4 hematologic toxicity required 1-week dose delay. The dog with grade 4 gastrointestinal toxicity

154 and one dog with grade 3 gastrointestinal toxicity (vomiting) had their chemotherapy protocol

155 stopped after a single dose of doxorubicin at the owners' request.

156 Overall, 36 (51%) dogs completed the planned course of cytotoxic chemotherapy. Twenty (29%)

157 dogs received metronomic chemotherapy as a rescue treatment at the detection of metastasis,

158 and 21 (30%) as maintenance therapy after completion of the cytotoxic chemotherapy protocol.

159 Twenty-six dogs received thalidomide, cyclophosphamide and piroxicam; 11 dogs received  
160 cyclophosphamide and firocoxib; 4 dogs received cyclophosphamide as single drug. Median  
161 thalidomide, cyclophosphamide, piroxicam and firocoxib doses were 4 mg/kg (range, 3.5-8), 10  
162 mg/m<sup>2</sup> (range, 7-13), 0.3 mg/kg (range, 0.1-0.4) and 5 mg/kg (range, 4-5), respectively.

### 163 *Outcome*

164 At data analysis closure, 63 (90%) dogs were dead, while 7 (10%) were still alive after a median  
165 follow-up time of 212 days (range, 102-1274). Among dead dogs, 60 (95%) and 3 (5%) dogs were  
166 dead for tumor-related and tumor-unrelated (2 gastric-dilatation-volvulus, 1 cardiac failure due to  
167 an end-stage myxomatous mitral valve disease) causes, respectively. One-year survival rate was  
168 9%. Median TTM was 142 days (95% CI, 103-181) and median OS was 165 days (95% C, 141-189).

169 Median StoC was 20 days (range, 4-70). Specifically, StoC was ≤ 7, ≤ 14, ≤21, and ≤ 28 days in 4, 22,  
170 39 and 57 dogs, respectively. The only cut-off significantly associated with a survival benefit was  
171 StoC ≤21 days. The median TTM of dogs with StoC ≤21 days (163 days, 95% CI 103-223) was  
172 significantly (P=0.001) longer than that of those with StoC >21 days (118 days, 95% CI 96-140;  
173 Figure 1). The median OS of dogs with StoC ≤21 days (238 days, 95% CI 184-292) was significantly  
174 (P<0.001) longer than that of those with StoC >21 days (146 days, 95% CI 129-163; Figure 2).

175 One-year survival rate of dogs with StoC ≤21 days and that of those with StoC >21 days were 16%  
176 and 0%, respectively.

177 On univariable analysis (Table 1), StoC >21 days was the only variable significantly associated with  
178 an increased risk of tumor metastasis (HR 2.5, 95% CI 1.4-4.2, P=0.001). When adjusting for other  
179 variables with P<0.2 (i.e., intact status; Table 2), StoC >21 days remained the only significant  
180 variable (HR 2.3, 95% CI 1.3-4.0, P=0.003). On univariable analysis for risk of death (Table 1), intact  
181 status (HR 1.8, 95% CI 1.1-3.1, P=0.02) and StoC >21 days (HR 2.8, 95% CI 1.6-5.1, P<0.001) were

182 both significant. When adjusting for all variables with  $P < 0.2$  (i.e., intact status, staging without  
183 TBCT, lack of maintenance metronomic chemotherapy; Table 3),  $\text{StoC} > 21$  days remained the only  
184 variable independently associated with an increased risk of death (HR 2.5, 95% CI 1.4-4.6,  
185  $P = 0.002$ ).

186 Analysis of distribution of possible prognostic variables between dogs with  $\text{StoC} \leq 21$  days and dogs  
187 with  $\text{StoC} > 21$  days is reported in Table 4. Dogs with  $\text{StoC} > 21$  days were more frequently of intact  
188 status than dogs with  $\text{StoC} \leq 21$  days ( $P = 0.029$ ). None of the remaining evaluated variable was  
189 differently distributed.

## 190 **Discussion**

191 Considering the highly aggressive behavior of splenic HSA and the limited overall benefit of  
192 medical therapy, our hypothesis was that timely administration of adjuvant chemotherapy would  
193 positively influence patients' outcome. Based on our findings,  $\text{StoC} \leq 21$  days resulted in a  
194 significantly longer median TTM (163 vs 118 days), and overall survival (238 vs 146 days) compared  
195 to dogs receiving their first dose of adjuvant chemotherapy  $> 21$  days from surgery. Notably, 1-year  
196 survival rate of dogs with  $\text{StoC} \leq 21$  days (i.e., 16%) was higher than that of dogs with  $\text{StoC} > 21$  days  
197 (i.e., 0%).

198 These findings are in line with pre-clinical and clinical studies, both in human and veterinary  
199 medicine.<sup>19-23</sup> The aim of administering adjuvant cytotoxic chemotherapy for splenic HSA is to  
200 target and kill rapidly-cycling neoplastic cells. There are many reasons to believe that early  
201 administration of adjuvant chemotherapy would result in a benefit for the patient. In 2006,  
202 Harless and Qiu proposed a mathematical model indicating that increasing tumor burden would  
203 result in decreased chemotherapy efficacy, and that the longer the time interval between tumor  
204 removal and chemotherapy delivery, the greater the tumor burden.<sup>27</sup> It is well documented that

205 quiescent micrometastases may re-enter the cell cycle after surgery and, with time,  
206 macrometastases may become clinically evident.<sup>28-29</sup> In fact, the removal of the primary tumor is  
207 associated with the release of many growth factors (e.g., catecholamines, prostaglandins), which  
208 stimulate angiogenesis and potentially trigger the outgrowth of latent distant disease; surgery-  
209 induced immunosuppression, caused by reduction in NK cell number and cytotoxic activity,  
210 impairment of monocytes/macrophages phagocytic and chemotactic function, or release of  
211 inflammatory cytokines (e.g., TNF- $\alpha$  and IL-1) into circulation, may further favor this process.<sup>6,7,9-11</sup>  
212 Additionally, a long interval between surgery and chemotherapy administration may lead to the  
213 development of a chemo-resistant phenotype due to the accumulation of spontaneous mutations  
214 in replicating tumor cells.<sup>30</sup>

215 In the present study, median StoC was 20 days; this is in line with a recent study by Finotello et  
216 al.,<sup>31</sup> whereas two studies reported shorter times.<sup>1,25</sup> This relatively long interval may be due to  
217 several reasons. In most cases, surgery for splenic HSA is an emergency procedure, as most dogs  
218 present with hemoabdomen due to mass rupture. Clinical conditions may require hospitalization  
219 in intensive care unit after surgery, thereby postponing discharge, together with the  
220 administration of the first dose of adjuvant chemotherapy. In addition, the ability to obtain a pre-  
221 operative diagnosis of HSA is limited, due to the poor sensitivity of cytology in identifying vascular  
222 neoplasms;<sup>32-34</sup> thus, the final histopathologic report is critical to guide possible adjuvant  
223 treatment decisions. Moreover, the spleen, being a blood-rich organ, generally requires longer  
224 formalin fixation times (e.g., up to 48-72 hours), increasing histologic processing and reporting  
225 time. The time interval of 21 days, which was significantly associated with the outcome in this  
226 study, is generally adequate for a patient to fully recover from surgery and for the final  
227 histopathologic report to be delivered.

228 Chemotherapy-related AEs did not increase with shorter StoC (i.e., 7, 14 days) and it is possible  
229 that shorter StoC intervals would have resulted in a survival benefit with a larger sample size  
230 available. On the other hand, one must question whether excessively early chemotherapy delivery  
231 can be counterproductive. An unacceptable level of toxicity has been reported in dogs with  
232 osteosarcoma receiving cisplatin and doxorubicin two days after surgery.<sup>12</sup> In addition, many  
233 anesthetic drugs (e.g., opioids, inhalant anesthetics) and surgery itself are known to cause  
234 transient immunosuppression.<sup>35-39</sup> A further immunosuppressive effect induced by  
235 chemotherapeutic drugs may put the dog at risk of major complications (e.g., sepsis).  
236 Furthermore, the cytotoxic effect and impaired collagen biosynthesis consequent to anthracycline  
237 and alkylating agents' administration could antagonize surgical wound repair.<sup>13,15</sup> In that scenario,  
238 waiting for the patient to recover after surgery may eventually translate in an improved outcome.  
239 Here, however, no adverse effect on wound healing was registered, even in dogs receiving  
240 adjuvant chemotherapy within 7 or 14 days from surgery.

241 StoC >21 days was the only variable independently associated with an increased risk of both tumor  
242 metastasis and death. Other factors influencing outcome in previous studies (e.g., hemoabdomen,  
243 thrombocytopenia, administration of blood products)<sup>2,40</sup> failed to demonstrate a prognostic role in  
244 the current population. Unfortunately, groups defined by some variables were not numerically  
245 homogenous. Notably, hemoabdomen was present in 83% of dogs, and we cannot exclude that  
246 this or other variables would have emerged as prognostic factors with similar number of cases in  
247 each group.

248 In disagreement with a previous study,<sup>25</sup> the chemotherapy protocol did not influence outcome in  
249 this cohort. However, only 4 dogs received doxorubicin and dacarbazine as a multidrug protocol,  
250 thus it is possible that this low number of patients may have prevented a real significance to be  
251 revealed. Again, in disagreement with the published literature,<sup>31</sup> no apparent benefit of

252 metronomic chemotherapy was observed. However, only 21 dogs received metronomic  
253 chemotherapy after completion of the cytotoxic protocol, while almost the same number received  
254 it as a rescue treatment.

255 Interestingly, HSA was an incidental finding in more than 10% of cases. This is uncommon, as  
256 splenic HSA is almost always symptomatic for active or previous hemoperitoneum due to mass  
257 rupture. It is possible that these cases, which often translate into stage I HSA, had a better  
258 outcome, but were well-distributed among the two groups, thereby minimizing a possible  
259 influence on the analysis.

260 This study has several limitations, mainly due to its retrospective nature. The population size was  
261 small and not homogeneous in terms of treatment; additionally, dogs receiving dacarbazine were  
262 underrepresented. An increased sample size would have possibly allowed for more significant  
263 findings to emerge, both in terms of StoC and other prognostic variables. There was a limited  
264 number of cases receiving chemotherapy within 7 and 14 days, but this represents daily clinical  
265 practice, as many variables (e.g., peri-operative complications, waiting for histopathology report,  
266 owners' decision-making, ordering and scheduling of treatment) may play a role in the timing of  
267 the first administration of chemotherapy. Furthermore, follow-up was not standardized,  
268 potentially affecting calculation of TTM.

269 Overall, the population was homogenous regarding common prognostic variables. Without TBCT,  
270 smaller metastases may not be revealed, increasing the risk of underestimating tumor stage, and  
271 including dogs with distant metastasis. It must be noted, however, that 70% of dogs in this cohort  
272 did not undergo TBCT, and StoC-based groups were uniform regarding this variable, too,  
273 minimizing a possible influence on inclusion and outcome.

274 Unfortunately, histologic slides were not available for re-evaluation, thus preventing inclusion of  
275 histologic variables in the analysis of prognostic factors. Mitotic count was recently associated  
276 with outcome by Moore et al.,<sup>41</sup> and may have possibly influenced prognosis in this cohort, too.

277 In conclusion, the advantages of timely administration of adjuvant chemotherapy in dogs with  
278 non-metastatic splenic HSA are supported by the current findings, being associated with longer  
279 TTM and survival. Based on this study, the authors recommend that oncologists and their teams  
280 collaborate to find strategies to shorten recovery time and interval from presentation to diagnosis  
281 to reduce the likelihood of the surgery to chemotherapy interval for these patients becoming  
282 significantly prolonged.

283 In light of our preliminary results, prospective studies on HSA and other tumor types are  
284 warranted to assess whether shorter time intervals between surgery and administration of  
285 adjuvant chemotherapy would further improve outcome.

#### 286 **Data Availability Statement**

287 The data that support the findings of this study are available from the corresponding author upon  
288 reasonable request.

#### 289 **References**

- 290 1. Ciepluch BJ, Wilson-Robles HM, Pashmakova MB, Budke CM, Ellison GW, Thieman Mankin  
291 KM. Long-term postoperative effects of administration of allogeneic blood products in 104  
292 dogs with hemangiosarcoma. *Vet Surg* 2018;47(8):1039-1045.
- 293 2. Marconato L, Chalfon C, Finotello R, et al. Adjuvant anthracycline-based vs metronomic  
294 chemotherapy vs no medical treatment for dogs with metastatic splenic

- 295 hemangiosarcoma: A multi-institutional retrospective study of the Italian Society of  
296 Veterinary Oncology. *Vet Comp Oncol* 2019;17(4):537-544.
- 297 3. Sorenmo KU, Jeglum KA, Helfand SC. Chemotherapy of canine hemangiosarcoma with  
298 doxorubicin and cyclophosphamide. *J Vet Intern Med* 1993;7(6):370-6.
- 299 4. Sorenmo KU, Baez JL, Clifford CA, et al. Efficacy and toxicity of a dose-intensified  
300 doxorubicin protocol in canine hemangiosarcoma. *J Vet Intern Med* 2004;18(2):209-13.
- 301 5. Kahn SA, Mullin CM, de Lorimier LP, et al. Doxorubicin and deracoxib adjuvant therapy for  
302 canine splenic hemangiosarcoma: a pilot study. *Can Vet J* 2013;54(3):237-42.
- 303 6. Tohme S, Simmons RL, Tsung A. Surgery for Cancer: A Trigger for Metastases. *Cancer Res*  
304 2017;77(7):1548-1552.
- 305 7. Chen Z, Zhang P, Xu Y, et al. Surgical stress and cancer progression: the twisted tango. *Mol*  
306 *Cancer* 2019;18(1):132.
- 307 8. Sawai H, Funahashi H, Yamamoto M, et al. Interleukin-1alpha enhances integrin  
308 alpha(6)beta(1) expression and metastatic capability of human pancreatic cancer. *Oncology*  
309 2003;65(2):167-73.
- 310 9. Yano S, Nokihara H, Yamamoto A, et al. Multifunctional interleukin-1beta promotes  
311 metastasis of human lung cancer cells in SCID mice via enhanced expression of adhesion-,  
312 invasion- and angiogenesis-related molecules. *Cancer Sci* 2003;94(3):244-252.
- 313 10. Jing Y, Ma N, Fan T, et al. Tumor necrosis factor-alpha promotes tumor growth by inducing  
314 vascular endothelial growth factor. *Cancer Invest* 2011;29(7):485-493.
- 315 11. Alieva M, van Rheenen J, Broekman MLD. Potential impact of invasive surgical procedures  
316 on primary tumor growth and metastasis. *Clin Exp Metastasis* 2018;35(4):319-331.
- 317 12. DeRegis CJ, Moore AS, Rand WM, Berg J. Cisplatin and doxorubicin toxicosis in dogs with  
318 osteosarcoma. *J Vet Intern Med* 2003;17(5):668-673



- 319 13. Robinson SJ, Nappi JF, Falcone RE. The effect of dacarbazine on wound healing. *J Dermatol*  
320 *Surg Oncol* 1988;14(9):975-8.
- 321 14. Kolb BA, Buller RE, Connor JP, et al. Effects of early postoperative chemotherapy on wound  
322 healing. *Obstet Gynecol* 1992;79:988–92.
- 323 15. Muszyńska A, Wolczyński S, Pałka J. The mechanism for anthracycline-induced inhibition of  
324 collagen biosynthesis. *Eur J Pharmacol* 2001;411(1-2):17-25.
- 325 16. Payne WG, Naidu DK, Wheeler CK, et al. Wound healing in patients with cancer. *Eplasty*  
326 2008;8:e9.
- 327 17. Guo S, DiPietro LA. Factors affecting wound healing. *J Dent Res* 2010;89:219–29.
- 328 18. Iodence AE, Wallace ML, Grimes JA, Schmiedt CW. Dogs undergoing surgical excision of  
329 mast cell tumors are not at increased risk of incisional complications. *J Am Vet Med Assoc*  
330 2021;260(S1):S88-S95.
- 331 19. Bell RS, Roth YF, Gebhardt MC, et al. Timing of chemotherapy and surgery in a murine  
332 osteosarcoma model. *Cancer Res* 1988;48(19):5533-8.
- 333 20. Meyers PA, Heller G, Healey J, et al. Chemotherapy for nonmetastatic osteogenic sarcoma:  
334 the Memorial Sloan-Kettering experience. *J Clin Oncol* 1992;10(1):5-15.
- 335 21. Hershman D, Hall MJ, Wang X, et al. Timing of adjuvant chemotherapy initiation after  
336 surgery for stage III colon cancer. *Cancer* 2006;107(11):2581-2588.
- 337 22. Imran H, Enders F, Krailo M, et al. Effect of time to resumption of chemotherapy after  
338 definitive surgery on prognosis for non-metastatic osteosarcoma. *J Bone Joint Surg Am*  
339 2009;91(3):604-12.
- 340 23. Marconato L, Buracco P, Polton GA, et al. Timing of adjuvant chemotherapy after limb  
341 amputation and effect on outcome in dogs with appendicular osteosarcoma without  
342 distant metastases. *J Am Vet Med Assoc* 2021;259(7):749-756.

- 343 24. Owen LN (Ed). *TNM Classification of Tumours in Domestic Animals*. Geneva, World Health  
344 Organization, 1980.
- 345 25. Finotello R, Stefanello D, Zini E, Marconato L. Comparison of doxorubicin-  
346 cyclophosphamide with doxorubicin-dacarbazine for the adjuvant treatment of canine  
347 hemangiosarcoma. *Vet Comp Oncol* 2017;15(1):25-35.
- 348 26. LeBlanc AK, Atherton M, Bentley RT, et al. Veterinary Cooperative Oncology Group-  
349 Common Terminology Criteria for Adverse Events (VCOG-CTCAE v2) following  
350 investigational therapy in dogs and cats. *Vet Comp Oncol* 2021;19(2):311-352.
- 351 27. Harless W, Qiu Y. Cancer: A medical emergency. *Med Hypotheses* 2006;67(5):1054-9.
- 352 28. Gunduz N, Fisher B, Saffer EA. Effect of surgical removal on the growth and kinetics of  
353 residual tumor. *Cancer Res* 1979;39(10):3861-5.
- 354 29. Fisher B, Gunduz N, Saffer EA. Influence of the interval between primary tumor removal  
355 and chemotherapy on kinetics and growth of metastases. *Cancer Res* 1983;43(4):1488-92.
- 356 30. Goldie JH, Coldman AJ. A mathematic model for relating the drug sensitivity of tumors to  
357 their spontaneous mutation rate. *Cancer Treat Rep* 1979;63(11-12):1727-33.
- 358 31. Finotello R, Henriques J, Sabattini S, et al. A retrospective analysis of chemotherapy switch  
359 suggests improved outcome in surgically removed, biologically aggressive canine  
360 haemangiosarcoma. *Vet Comp Oncol* 2017;15(2):493-503.
- 361 32. Watson AT, Penninck D, Knoll JS, Keating JH, Sutherland-Smith J. Safety and correlation of  
362 test results of combined ultrasound-guided fine-needle aspiration and needle core biopsy  
363 of the canine spleen. *Vet Radiol Ultrasound* 2011;52(3):317-22.
- 364 33. Patten SG, Boston SE, Monteith GJ. Outcome and prognostic factors for dogs with a  
365 histological diagnosis of splenic hematoma following splenectomy: 35 cases (2001-2013).  
366 *Can Vet J* 2016;57(8):842-6.

- 367 34. Tecilla M, Gambini M, Forlani A, Caniatti M, Ghisleni G, Roccabianca P. Evaluation of  
368 cytological diagnostic accuracy for canine splenic neoplasms: An investigation in 78 cases  
369 using STARD guidelines. *PLoS One* 2019;14(11):e0224945.
- 370 35. Yeager MP, Colacchio TA, Yu CT, et al. Morphine inhibits spontaneous and cytokine-  
371 enhanced natural killer cell cytotoxicity in volunteers. *Anesthesiology* 1995;83(3):500-508.
- 372 36. Melamed R, Bar-Yosef S, Shakhar G, Shakhar K, Ben-Eliyahu S. Suppression of natural killer  
373 cell activity and promotion of tumor metastasis by ketamine, thiopental, and halothane,  
374 but not by propofol: mediating mechanisms and prophylactic measures. *Anesth Analg*  
375 2003;97(5):1331-1339.
- 376 37. Shavit Y, Ben-Eliyahu S, Zeidel A, Beilin B. Effects of fentanyl on natural killer cell activity  
377 and on resistance to tumor metastasis in rats. Dose and timing study.  
378 *Neuroimmunomodulation* 2004;11(4):255-260.
- 379 38. Horowitz M, Neeman E, Sharon E, Ben-Eliyahu S. Exploiting the critical perioperative period  
380 to improve long-term cancer outcomes. *Nat Rev Clin Oncol* 2015;12(4):213-26.
- 381 39. Longhini F, Bruni A, Garofalo E, et al. Anesthetic Strategies in Oncological Surgery: Not Only  
382 a Simple Sleep, but Also Impact on Immunosuppression and Cancer Recurrence. *Cancer*  
383 *Manag Res* 2020;12:931-940.
- 384 40. Masyr AR, Rendahl AK, Winter AL, Borgatti A, Modiano JF. Retrospective evaluation of  
385 thrombocytopenia and tumor stage as prognostic indicators in dogs with splenic  
386 hemangiosarcoma. *J Am Vet Med Assoc* 2021;258(6):630-637.
- 387 41. Moore AS, Rassnick KM, Frimberger AE. Evaluation of clinical and histologic factors  
388 associated with survival time in dogs with stage II splenic hemangiosarcoma treated by  
389 splenectomy and adjuvant chemotherapy: 30 cases (2011-2014). *J Am Vet Med Assoc*  
390 2017;251(5):559-565.

391 **Figure legends**

392 **Figure 1.** Time to metastasis for 70 dogs with non-metastatic splenic hemangiosarcoma  
393 undergoing splenectomy and adjuvant chemotherapy. In the group with a time interval  
394 between surgery and adjuvant chemotherapy (StoC)  $\leq 21$  days, dogs had a significantly longer  
395 time to metastasis than dogs with StoC  $> 21$  days (median, 163 vs 118 days, respectively;  
396  $P=0.001$ ).

397 **Figure 2.** Overall survival for 70 dogs with non-metastatic splenic hemangiosarcoma  
398 undergoing splenectomy and adjuvant chemotherapy. In the group with a time interval  
399 between surgery and adjuvant chemotherapy (StoC)  $\leq 21$  days, dogs had a significantly longer  
400 overall survival than dogs with StoC  $> 21$  days (median, 238 vs 146 days, respectively;  $P<0.001$ ).