

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Compost improves plant and soil macronutrient content in a 14-years orchard

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Compost improves plant and soil macronutrient content in a 14-years orchard / Baldi, Elena; Mazzon, Martina; Cavani, Luciano; Quartieri, Maurizio; Toselli, Moreno; Marzadori, Claudio. - In: NUTRIENT CYCLING IN AGROECOSYSTEMS. - ISSN 1385-1314. - ELETTRONICO. - 125:(2023), pp. 425-435. [10.1007/s10705-023-10258-0]

Availability:

This version is available at: <https://hdl.handle.net/11585/923184> since: 2023-04-17

Published:

DOI: <http://doi.org/10.1007/s10705-023-10258-0>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the peer-reviewed accepted manuscript of:

Baldi, E., Mazzon, M., Cavani, L. et al. *Compost improves plant and soil macronutrient content in a 14-years orchard*.

Nutr Cycl Agroecosyst 125, 425–435 (2023).

The final published version is available online at:

<https://doi.org/10.1007/s10705-023-10258-0>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

~~The fate of generally unconsidered macronutrients in plant and soil: effect of different fertilization strategy on K, Ca, Mg and S~~

Compost improves plant and soil macronutrient content in a 14-years orchard

Elena Baldi¹, Martina Mazzon^{*1}, Luciano Cavani¹, Maurizio Quartieri¹, Moreno Toselli¹, Claudio Marzadori¹

¹Department of Agricultural and Food Sciences, Alma Mater Studiorum - University of Bologna, Bologna, Italy

^{*}Correspondence: Martina Mazzon (ORCID 0000-0001-5136-0135), martina.mazzon2@unibo.it

Abstract

Adequate plant nutritional status and soil fertility preservation ~~should~~ can be achieved through ~~the employment of~~ sustainable agricultural management techniques. The challenge of intensive orchard cultivation, besides the risk of nutrient decline, is ~~the to prevention of~~ the loss of soil fertility that could lead to soil degradation with a consequent negative impact on yield and fruit quality. The use of organic amendments could be a sustainable strategy to combine ~~elevated~~ high plant performance with soil fertility improvement.

This work aims at shedding light on the effects of compost addition with respect to an unfertilized control and a mineral fertilization treatment on macronutrient (K, Ca, Mg, and S) dynamics in plants and soil of a commercial nectarine orchard planted in 2001. Therefore, ~~the manuscript aims at shading light on macronutrient (K, Ca, Mg and S) dynamics in plants and soil of a commercial nectarine orchard planted in 2001 and in which three fertilization treatments were compared: unfertilized control, mineral fertilization, and compost.~~

In the first 0.15 m of soil, compost addition resulted in higher values (26 - 42%) of all the parameters.

Both fertilization treatments induced a 28% increase ~~of~~ in roots' S content compared to the control but did not induce macronutrients content variation in plant skeleton, pruned wood, and thinned fruits. In autumn leaves, all the macronutrients resulted in higher values (24 - 45%) with both mineral and compost fertilization, and the same was observed in fruit at harvest (increases of 15 - 31%).

~~Finally, in the first 0.15 m of soil, compost addition resulted with the higher values (+ 26 - 42%) of all the parameters.~~

In our study, the treatment with compost satisfied plants' nutrient demands as much as the mineral fertilizer. In addition, compost treatment also improved soil nutrient content while preserving yield.

Our results show that it is possible to reconcile plant nutrient needs with the preservation of soil fertility with the aim of reaching agriculture sustainability.

~~Thus, the long term use of mineral fertilizer, even if meeting plant's nutrient demand, could lead to soil macronutrient depletion. Whereas, organic amendments addition could not only satisfy plant's nutrient demand but also maintain soil quality and fertility through the higher macronutrients storage in the soil; this would lead to both higher plant performances and to economic and environmental benefits.~~

Keywords

Prunus persica, soil nutrient availability, compost, mineral fertilization, nutrient removal, soil macronutrient content.

45 1. Introduction

46 The ~~rising-increasing~~ demand for high-quality nutritional food related to ~~the constant-steady~~
47 increase ~~of in~~ population (El-Jendoubi et al., 2013) is leading to the necessity to ~~maximise-maximize~~
48 yield ~~by-while~~ minimizing ecosystem impacts deriving from agriculture. According to the report of
49 the Food and Agriculture Organization of the United Nations (FAO, 2017), the requirements for
50 adequate food supplies have to pass through sustainable agricultural management techniques. This
51 issue also deals with adequate plant nutritional status and preservation/increase of soil fertility
52 (Toselli et al., 2019a; Zhang et al., 2020). Plant nutrition ~~depends on is the result of~~ the nutrient cycle
53 in and out of the orchard ecosystem. The inputs to the soil, for example, include mineral and/or
54 organic fertilizers, atmospheric deposition, and biological nitrogen (N) fixation, while the outputs
55 are represented by harvested fruits, nutrients lost by leaching, gaseous losses, and erosion (Toselli
56 et al., 2019b). In case of a negative balance between inputs and outputs, the soil would come across
57 a nutrient depletion that, in the long-term, would lead to an unsustainable farming system (El-
58 Jendoubi et al., 2013). The challenge of intensive orchard cultivation is, besides the risk of nutrient
59 decline, the loss of soil fertility (both chemical and biological) that could lead to soil degradation (i.e.
60 loss of soil organic matter, erosion, acidification, and pollution) with a consequent negative impact
61 on plant performances (Zhang et al., 2020). Thus, the great challenge for modern farmers is to
62 maintain and/or increase soil fertility in a sustainable way. This could be reached through the
63 application of organic amendments (i.e. composts, biochar, and manures) that are widely ~~recognised~~
64 ~~recognized~~ to be ~~cheapinexpensive~~, to ~~release slowly release the~~ nutrients through mineralization
65 ~~slowly~~, and to enhance soil carbon (C) and organic matter increase (Mazzon et al., 2018; Sciubba et
66 al., 2015) thus stimulating carbon dioxide sequestration. In addition, the application of organic
67 matter ~~is able to can~~ improve soil physical properties (Chatzistathis et al., 2020) as, for example, bulk
68 density reduction and aggregate stability and water holding capacity increase (Adugna, 2016),

positively affecting soil microbial community (Fawzi et al., 2010; Safaei Khorram et al., 2019) and increasing macro- and micro-nutrient availability (Baldi et al., 2021a; Fawzi et al., 2010). Organic amendments thus represent not only a source of N, phosphorus (P), and potassium (K) but also of calcium (Ca), magnesium (Mg), and sulphur (S) ~~that-which~~ are equally considered ~~as~~-essential plant macronutrients (Barreto et al., 2021; Shiwakoti et al., 2020). In the study of Shiwakoti et al. (2020), ~~the-farmyard-manure~~ long-term (approximately 64 years) addition of farmyard manure (at the rate of 11.2 Mg ha⁻¹ year⁻¹) evidenced higher macronutrient (K, S, and Mg) content in soil than the other treatments (pea vine and wheat residues, with and without N addition) mainly due to the high amounts of these nutrients that directly contribute to soil chemical fertility. Shiwakoti et al. (2020) also highlighted that manure could have activated soil cation exchange sites releasing organic colloids and consequently adsorbing K to the exchangeable sites and increasing its availability. Macronutrient soil availability throughout the growing season is fundamental for fruit trees. In a study on pomegranate nutrient dynamics, Maity et al. (2019) demonstrated that ~~most of~~ plant needs were mostly satisfied by uptake from the soil more than from the mobilization from plant reserves; as a consequence, if soil nutrients are not replenished through appropriate fertilization management, fruit yield and quality could be severely impaired. In a different study on various fruit trees (i.e. apple, peach, and mandarin), Cruz et al. (2019) showed that adequate K supply at fruit set is of fundamental importance for the final fruit quality. Maity et al. (2019) evidenced that a great amount of Mg was remobilized from leaves to fruit at the maturity stage, while S was mainly concentrated in shoots and Ca in the woody organs of pomegranate. El-Jendoubi et al. (2013) found that K, P, and N mainly accumulate in fruits, while Mg and Ca in abscised leaves. Moreover, it is estimated that peach trees in commercial orchards have a macronutrient requirement accounting for 74 - 425 g K tree⁻¹, 25 - 518 g Ca tree⁻¹, and 9 - 74 g Mg tree⁻¹ (Baldi et al., 2021b) every year.

92 The key issue for fruit tree nutrition is the availability of nutrients in the soil during the entire
93 vegetative season, consequently, the use of organic amendments, that gradually release nutrients
94 through mineralization, could be a sustainable strategy able to combine elevated plant performance
95 with the improvement of soil fertility (Baldi et al., 2021b).
96 This manuscript follows ~~3-three~~ previous publications on C (Baldi et al., 2018), N (Toselli et al.,
97 2019b), and micronutrients ~~on-in~~ the same experiment (Baldi et al., 2021a), and aims at ~~shedding~~
98 ~~shedding~~ light on ~~other nutrients such as the macronutrients~~ K, Ca, Mg, and S. Indeed, the goal of the
99 present study was to determine the effects of the long term mineral fertilization and compost
100 addition in the long term nectarine orchard (14 years) on: i) soil macronutrient content at the end
101 of the 14-years life-time of the ~~nectarine~~ orchard, ii) macronutrient content in different plant organs,
102 and iii) the relation between plant and soil macronutrient content.

103

104 2. Materials and methods

105 2.1 Orchard description and treatments

106 The experiment was carried out on a commercial nectarine orchard [*Prunus persica*, Batsch var.
107 *nucipersica* (Bockh.) Schn.] planted in 2001 (Table 1). The orchard was located in the Po valley (Italy)
108 near Ravenna (44°27' N; 12°13' E), an area characterized by a temperate climate and a silt-loam soil
109 (Calcaric Cambisol) with a total carbonate content (% CaCO₃) of 31 ± 1 and an active carbonate
110 content (% CaCO₃) of 13 ± 1 (Baldi et al., 2021a, 2018). The planting layout and main orchard
111 management strategies are reported in Table 1. Since orchard plantation, three fertilization
112 treatments were compared with four replicates (4 trees each) according to a complete randomized
113 block design: unfertilized control (CK); mineral fertilization (MIN); and compost (COM). Specific
114 information regarding the treatments ~~are-is~~ reported in Table 1. Fertilizers were applied ~~on-to~~ the

115 tree row and tilled into the soil to a depth of 0.25 m, while pruned wood was left into the ground
 116 and chopped (Baldi et al., 2021a, 2018; Toselli et al., 2019b).

117

118 **Table 1** – Nectarine orchard main characteristics and management strategies (Baldi et al., 2021a, 2018; Toselli
 119 et al., 2019b).

Management	Description
<i>Variety and rootstock</i>	Stark RedGold, grafted on hybrid GF677 (<i>P. persica</i> × <i>P. dulcis</i>)
<i>Training system</i>	Delayed vasette, distance of 5 m between the rows and 3.8 m between trees along the row
<i>Irrigation</i>	Drip irrigation system from June to September (vegetation season).
<i>Phytosanitary treatments</i>	Done a According to Crop Management Guideline of the region Emilia-Romagna (www.regione.emilia-romagna.it).
<i>Soil management</i>	Tree row was tilled superficially for weed control. Alleys were covered with spontaneous grass managed by mowing it 3 times a year.
<i>Fertilization treatments</i>	Mineral fertilization (MIN) Nitrogen (N) applied every year at the rate of 70 kg ha ⁻¹ yr ⁻¹ (60% in May and 40% in September); in 2004, N rate was increased to 120 kg ha ⁻¹ yr ⁻¹ and from 2006 to 130 kg ha ⁻¹ yr ⁻¹ . <u>Nitrogen was applied as ammonium nitrate (N = 35 %), while P and K were applied as a binary fertilizer (P = 10 %; K = 20 %); no micro-nutrients was provided in mineral fertilization.</u> Phosphorus (P) at 100 and potassium (K) at 200 kg ha ⁻¹ (applied only at planting). Rate established according to Integrated Crop Management Guideline of the region Emilia-Romagna (www.regione.emilia-romagna.it). <u>Nitrogen was applied as ammonium nitrate (N = 35%), while P and K were applied as a binary fertilizer (P = 10%; K = 20%); no micronutrients was provided with mineral fertilization.</u>

Compost (COM)

Applied at a rate of 10 t DW ha⁻¹ yr⁻¹, equal to 240 kg N ha⁻¹ yr⁻¹.

Compost was obtained from domestic organic wastes (50%) mixed with pruning material (50%) after a 3-month stabilization. Average characteristics: DW of 73%, pH 9, EC 2.96 mS cm⁻¹, C/N ratio 10.2, and (in g kg^{DW}⁻¹) organic C 234, N 21.1, P 4.8, K 15.2.

120

121 2.2 Plant sampling and analysis

122 In 2014, after 14 years of life, 4 trees per treatment were harvested and divided into organs –as
123 described by Baldi et al. (2021a, 2018) and Toselli et al. (2019b). Briefly, thinned fruits were collected
124 in spring and weighed, and a representative sample was then oven-dried and milled (2 mm). In
125 July, a sample of 40 young fully expanded leaves was collected from the apical part of shoots, and
126 the leaf area was measured by a portable area meter (Li-3000, LiCor inc., Lincoln, Nebraska). Leaves
127 were then washed, oven-dried, and milled at 2 mm. At harvest, in August, plant yield was recorded;
128 afterwards, on a representative sample of fruits, fresh weight (FW) and dry weight (DW) of flesh
129 and kernel were measured on a representative sample of fruits; dried flesh and kernel were weighted
130 and milled. In September, one tree per plot was enclosed into a plastic net to collect autumn leaves
131 that were weighed, leaf area measured, dried, and milled. In December 2014, at the end of the
132 commercial life of the orchard, the same trees were harvested, divided into roots, trunks, branches
133 (age > 2 years), and current year shoots (twigs), and weighed. A subsample of each organ was oven-
134 dried, weighed, and milled. The sample of roots was washed with deionized water to
135 remove soil residues.

136 A sample (0.3 g) of each plant organ was mineralized according to the US EPA Method 3052
137 (Kingston and Jassie, 1988) in an Ethos TC microwave (Milestone, Bergamo, Italy), filtered
138 (Whatman 42®), and analyzed for Ca, K, Mg, and S by Inductively Inductively Coupled coupled

~~Plasma-plasma Optical-optical Emission-emission Spectrometer-spectrometer~~ (ICP-OES; Ametek Spectro, Arcos, Kleve, Germany). Blank and certified reference materials (NIST standard reference material SRM 1573a tomato leaves and SRM 1570a spinach leaves) analyses were performed. Relative uncertainty, calculated as the relative deviation of the measured element concentration to its certified value, was typically better than $\pm 5\%$.

~~Mineral~~ The mineral content in different parts of the plant (leaves, fruits, branches, trunk, and roots) was calculated by multiplying each mineral concentration by the DW of the specific organ.

The biomass of the skeleton was calculated as the sum (without pruning wood) of trunk, branch, and twigs > 2 years (identified according to their insertion into branches) measured at the end of the experiment. Skeleton and root nutrient content were divided by the age of the orchard (13) assuming a constant annual increase of weight and nutrient accumulation, and considering the increase of the first year (2001) was negligible since orchard ~~plantation~~ was done-planted at the end of the year.

2.3 Soil sampling and analysis

In December 2014, an 80 cm deep soil core (70 mm diameter) was collected in the row of each plot with a soil column cylinder auger that was inserted into the soil using a tractor. The core was carefully removed from the auger and divided into four parts according to depth: 0 - 0.15 m, 0.16 - 0.25 m, 0.26 - 0.45 m and 0.46 - 0.65 m. Soil from each depth was separately weighted and oven dried at 105°C for 24 hours to evaluate soil bulk density (BD) ~~that-which~~ was calculated as the ratio between DW and the volume of each core. In addition, soil samples were collected, always in the row, with an auger at 0-0.15 m, 0.16-0.25 m, 0.26-0.45 m, and 0.46-0.65 m ~~of~~ depth. Each sample (made of 4 sub-samples) was sieved (2 mm), cleaned from roots and visible plant residues, and air-dried or stored at 4°C. A ~~sub-sub~~ sample was then used to evaluate soil total concentration of Ca, K, Mg, and S. Briefly, samples were subjected to wet mineralization by treating 0.5 g of dry sample

163 with 6 mL of hydrochloric acid (37%), 2 mL of nitric acid (65%) and 2 mL of hydrogen peroxide
164 (30%) in an Ethos TC microwave lab station (Milestone, Bergamo, Italy) according to the methods
165 ISO 12914:2012 and 22036:2008. Solutions were filtered (Whatman 42®) and the element
166 concentration was determined by plasma spectrometer (ICP-OES; Ametek Spectro, Arcos, Kleve,
167 Germany). Blank and certified reference material (BCR reference material No 141R calcareous loam
168 soil) analyses were performed; relative uncertainty, calculated as the relative deviation of the
169 measured element concentration to its certified value, was typically better than $\pm 5\%$. Soil pseudo-
170 total mineral element content at the end of the orchard life-time was calculated by multiplying the
171 nutrient concentration by the soil bulk density at the respective depth intervals. Electrical
172 conductivity (EC) was measured on a suspension of 10 g of fresh sample and 20 mL of deionised
173 water that was stirred for 120 min at 25°C and filtered before measurement. Soil potential cation
174 exchange capacity (CEC) was estimated using the ammonium acetate method (Sumner and Miller,
175 1996).

176

177 2.4 Statistical analysis

178 After assumption verification (Shapiro-Wilk for normality and Bartlett for homogeneity of variance),
179 pPlant organ data were ~~analysed~~analyzed with a one-way ANOVA with treatment as a factor (three
180 levels: unfertilized control, mineral fertilization, and compost) ~~after assumption verification~~
181 ~~(Shapiro Wilk for normality and Bartlett for homogeneity of variance)~~. Whether necessary, data
182 were transformed using the Box-Cox procedure to fit the ANOVA assumption. Similarly, data on
183 macronutrient content for~~of~~ autumn and summer leaves ~~macronutrient content~~ were analysed with
184 a ~~two~~one-way ANOVA with ~~treatment and~~ season as a factors. When significant differences occur
185 ($P \leq 0.05$), an HSD post-hoc test (Tukey's test) was applied to separate the means. A Principal
186 Component Analysis (PCA) was carried out using plant organ data ~~that showed~~ed significant

187 differences ~~for in~~ the treatments~~ss~~. Soil data were ~~analysed-analyzed~~ using a split-plot design, with
188 treatments (three levels: unfertilized control, mineral fertilization, and compost) as the main factor
189 and sampling depths (four levels: 0 - 0.15, 0.16 - 0.25, 0.26 - 0.45, 0.46 - 0.65 m) as the sub-factor.
190 ANOVA assumption verification and means separation were ~~done-asperformed as described~~ -for
191 plant organs (R Core Team, 2021).

192

193 3. Results

194 Fertilization treatments, no matter which one, induced on average a 28% increase ~~of in~~ roots' S
195 content compared to the control (Table 2). The fertilization treatments did not induce a variation of
196 Ca, K, Mg, and S content in the plant skeleton, pruned wood_z and thinned fruits (Table 2). On the
197 contrary, all the considered macronutrients ~~considered~~ (Ca, K, Mg_z and S) ~~were~~ increased in autumn
198 leaves (in a range from 24 to 45%) by both mineral and compost fertilizations. Similar results in
199 ~~correspondence of~~ the two fertilization treatments (with increases between 15 and 31%) ~~was-were~~
200 observed in fruit at harvest with the exception of Ca content that did not show any differences
201 among treatments (Table 2).

202

203 **Table 2** – Macronutrient content (g pt⁻¹) in plant organs ± standard error (based on field replicates). Different
204 lowercase letters indicate significant (P < 0.05) differences between treatments (CK = unfertilized control, MIN
205 = mineral fertilization, COM = compost).

Plant organs	Treatments	Ca	K	Mg	S
Pruned wood	CK	26 ± 1	9.9 ± 0.7	2.2 ± 0.2	1.4 ± 0.1
	MIN	43 ± 1	17 ± 1	3.5 ± 0.3	2.5 ± 0.1
	COM	30 ± 7	13 ± 3	2.6 ± 0.6	1.9 ± 0.4
	<i>p.value</i>	0.150	0.098	0.178	0.097
Thinned fruits	CK	0.62 ± 0.14	2.6 ± 0.6	0.29 ± 0.07	0.27 ± 0.06
	MIN	0.44 ± 0.12	2.0 ± 0.6	0.22 ± 0.06	0.21 ± 0.06
	COM	0.87 ± 0.17	3.5 ± 0.7	0.38 ± 0.07	0.36 ± 0.07

	<i>p.value</i>	0.168	0.257	0.259	0.251
Fruit at harvest	CK	2.3 ± 0.2	43 ± 1 ^b	3.6 ± 0.1 ^b	2.0 ± 0.1 ^b
	MIN	2.4 ± 0.1	51 ± 2 ^{ab}	4.8 ± 0.3 ^a	2.8 ± 0.1 ^a
	COM	2.4 ± 0.2	57 ± 1 ^a	4.8 ± 0.2 ^a	2.8 ± 0.1 ^a
	<i>p.value</i>	0.867	0.010	0.012	0.006
Autumn leaves	CK	118 ± 3 ^b	43 ± 2 ^b	14 ± 0.3 ^b	2.6 ± 0.2 ^b
	MIN	181 ± 23 ^a	70 ± 7 ^a	21 ± 2 ^a	4.7 ± 0.5 ^a
	COM	156 ± 9 ^a	66 ± 4 ^{ab}	19 ± 2 ^a	4.5 ± 0.4 ^a
	<i>p.value</i>	0.010	0.026	0.012	0.013
Roots	CK	21 ± 3	5.6 ± 0.2	3.0 ± 0.5	0.32 ± 0.02 ^b
	MIN	21 ± 4	5.2 ± 0.5	3.0 ± 0.5	0.43 ± 0.03 ^a
	COM	24 ± 0.3	6.1 ± 0.7	3.6 ± 0.2	0.45 ± 0.03 ^a
	<i>p.value</i>	0.771	0.313	0.659	0.001
Skeleton	CK	45 ± 2	12 ± 1	3.8 ± 0.3	1.8 ± 0.1
	MIN	60 ± 7	16 ± 2	5.2 ± 0.7	2.4 ± 0.2
	COM	56 ± 7	17 ± 1	5.0 ± 0.6	2.5 ± 0.2
	<i>p.value</i>	0.328	0.220	0.329	0.126

206

207 No significant differences between fertilization treatments were observed in the concentration of
 208 nutrients in summer leaves (Table 3); the only exception was Mg which resulted in a 13% higher
 209 concentration in the control compared to the two fertilization strategies (Table 3).


210

211 **Table 3** – Macronutrient concentration (g 100 g⁻¹ DW) ± standard error (based on field replicates) in leaves
 212 sample in summer. Different lowercase letters indicate significant (P < 0.05) differences between treatments
 213 (CK = unfertilized control, MIN = mineral fertilization, COM = compost).

Treatments	Ca	K	Mg	S
CK	2.2 ± 0.1	1.2 ± 0.04	0.44 ± 0.01 ^a	0.16 ± 0.03
MIN	1.9 ± 0.1	1.1 ± 0.01	0.39 ± 0.01 ^b	0.15 ± 0.01
COM	1.9 ± 0.1	1.2 ± 0.04	0.39 ± 0.01 ^b	0.15 ± 0.01
<i>p.value</i>	0.088	0.088	0.013	0.947

214

215 The comparison of macronutrients content in autumn and summer leaves highlighted the significant
 216 impact of the season-phenological state for all ~~the~~ four macronutrients considered in this study
 217 (Figure 1). Specifically, higher concentrations of Ca and K were observed in autumn than in summer
 218 leaves; the opposite was observed for Mg and S.



219
 220 **Figure 1**– Means of summer and autumn leaves content of calcium (Ca), potassium (K), magnesium (Mg) and
 221 sulphur (S). Error bar represent the standard error (based on four leaf samples collected in field replicates) and
 222 the significant differences ($P \leq 0.05$) are reported.

223
 224
 225 A PCA (Figure 2) was ~~done-performed using with-macronutrients-content in data from~~ those organs
 226 whose macronutrient content was that were significantly affected by the fertilization treatments
 227 (Table 2). The treatments clusterized were split ~~resulted into~~ two main cluster-groups (according to
 228 PC1 ANOVA results): the unfertilized control in-on the left side of the plot, and the two fertilization
 229 strategies (mineral and compost) in-on the right side of the plot. These two groups showed to be

clearly defined ~~with by the~~ Mg content in summer leaves that characterized the unfertilized control group (Figure 2). A clear separation in plot space was also evident between macronutrients content in the autumn leaves (bottom-right side of the plot) and in the fruit at harvest and roots (upper-right side of the plot).

230
231
232
233
234
235
236
237
238
239
240
241
242
243




Figure 2– Principal component (PC) analysis with macronutrients content (calcium (Ca), potassium (K), magnesium (Mg) and sulfur/sulphur (S)) in the most relevant plant organs: harvested fruits (HF), roots (R), autumn leaves (AL), and summer leaves (SL). ~~In the table are reported reports~~ the statistical output of the ANOVA done on the PC (Treatments: CK = unfertilized control, MIN = mineral fertilization, COM = compost) with t-he significant differences (P < 0.05).

244
245
246
247
248
249
250
251
252
253
254

In the first 0.15 m of soil, the addition of compost increased CEC, EC, Ca, K, Mg, and S compared with control and mineral fertilization (Fig. 3). With the exception of CEC and Ca, the positive effect of compost was observed also in other soil layers (Fig. 3) and in particular EC was higher in all soil

255 profiles, while the concentration of K, Mg and S were higher than in the control and mineral
256 fertilization between 0 and 45 cm of depth. The effect of compost decreased with depth for all the
257 investigated parameters ~~investigated~~; however, while the concentration of K and Mg decreased
258 constantly with depths, the EC decreased until 25 cm and then ~~was~~ remained steady from 26 to 65
259 cm, and the concentration of Ca, S and CEC decreased until 15 cm and then ~~was~~ remained steady
260 from 16 to 65 cm (Fig. 3). Unfertilized control and mineral fertilization did not show significant
261 differences in nutrient concentrations and in both treatments only a slightly decreasing trend with
262 depth was observed for K and Mg concentration. In control and mineral fertilization, S concentration
263 was higher in the deepest layer than in the ~~superficial~~ shallowest one (Fig. 3).




Figure 3 – Means of soil cation exchange capacity (CEC), electrical conductivity (EC), and soil calcium (Ca), potassium (K), magnesium (Mg) and sulphur (S) content at four sampling depth (0 - 0.15, 0.16 - 0.25, 0.26 - 0.45, 0.46 - 0.65 m) for the three fertilization treatments (CK = unfertilized control, MIN = mineral fertilization, COM = compost). Error bars represent the standard error (based on data on field replicates) and the significant differences ($P \leq 0.05$) between “Treatment”, “Depth”, and “Treatment*Depth” interaction are reported.

269 4. Discussion

270 ~~In this study and as already observed previously (El-Jendoubi et al., 2013) plants macronutrients~~
271 ~~were mainly allocated in autumn leaves and in fruits at harvest. However, while in roots, skeleton,~~
272 ~~pruned wood, autumn leaves, and summer leaves Ca contents were highest among all the other~~
273 ~~macronutrients, K was the most important macronutrient in thinned fruit and fruit at harvest.~~
274 ~~Similarly, El-Jendoubi et al. (2013) showed that each nutrient was characterized by a precise~~
275 ~~allocation pattern: fruits were the largest sink for K, while Mg and Ca were mainly accumulated in~~
276 ~~abscised leaves.~~

277 4.1 Calcium and potassium returned to soil with leaf abscission ~~abundance and allocation~~

278 In this study and as already observed previously (El-Jendoubi et al., 2013) plants macronutrients
279 were mainly allocated in autumn leaves and in fruits at harvest. However, while in roots, skeleton,
280 pruned wood, autumn leaves, and summer leaves Ca contents were highest among all the other
281 macronutrients, K was the most important macronutrient in thinned fruit and fruit at harvest.
282 Similarly, El-Jendoubi et al. (2013) showed that each nutrient was characterized by a precise
283 allocation pattern: fruits were the largest sink for K, while Mg and Ca were mainly accumulated in
284 abscised leaves.

285 The difference between the amount of nutrients in summer leaves (sampled in summer July) and at
286 natural abscission gives an estimation of the fraction of nutrients remobilized at the end of the
287 vegetative season and stored inside the woody part of the plant. In the present experiment, Ca and
288 K showed higher concentrations in autumn leaves than in those sampled in summer indicating no
289 net remobilization through the season. ~~SA~~ similar behavior was reported in almond trees
290 (Muhammad et al., 2015). As a consequence, Ca and K allocated ~~in autumn to~~ leaves returned ~~into to~~
291 the soil after abscission and ~~degradation~~ decomposition, thus ~~becoming again~~ returning partially
292 available for root uptake (Baldi et al., 2021b) after mineralization. ~~On the other hand, little quantities~~

293 ~~of Ca and K reservation are available for spring new growth.~~ However, despite what was expected
294 and what was observed in other ~~study studies~~ (Dang et al., 2022; do Carmo et al., 2016), in ~~this our~~
295 case, soil Ca and K concentration increased in the surface horizon and in the whole soil profile
296 respectively, only when compost was applied (CaO and K mean content of compost ~~was were~~ of
297 ~~8.25.8~~ ± 1.0 % and 1.5 ± 0.2 %, respectively), and not in control and mineral fertilized plots, meaning
298 that the source of additional Ca was the organic fertilizer rather than the litter ~~made of~~ formed by
299 abscised leaves. ~~This The lack of effect~~ fact that abscised leaves and/or mineral fertilizer have no such
300 effect on soil Ca content was probably in relation to the natural soil's high abundance in total and
301 active carbonate content (CaCO_3) (Baldi et al., 2018), indicating a large soil endowment in carbonates
302 thus making it impossible to observe variation in soil Ca content even in the long term.

303

304 4.2 ~~K and Mg in plant and soil~~ Compost contributed to soil and plants K and Mg content

305 Differently from Ca, ~~the fertilization treatments~~ (both mineral fertilizer and compost) ~~induced an~~
306 increase ~~of~~ K and Mg content ~~also~~ in fruits at harvest (Table 2). This effect was also observed by
307 ~~Increase of K content in fruits was observed also in another study on nectarine~~ (Delian et al., (2012),
308 who reported a higher concentration of potassium than magnesium in nectarine fruits at
309 harvest ~~where at harvest macronutrients concentration in fruits followed the order $K > Ca > Mg$.~~
310 Potassium ~~is not an essential element for organic molecules and it is not considered a structural~~
311 ~~element~~ (Delian et al., 2012); however, ~~it~~ is involved in many physiological and biochemical
312 processes related to plant growth, crop quality, and plant response to stress factors (Delian et al.,
313 2012; Wang et al., 2018). An excess in available P potassium is ~~also~~ known to induce a Mg deficiency
314 in the plant due to the unidirectional competition for uptake ~~between K and Mg for which an~~
315 ~~increase in K concentration reduces Mg uptake~~ (Xie et al., 2021). In ~~this our~~ study, independently
316 from the fertilization treatment (mineral or organic), the ~~lower lowest~~ K/Mg ratio (approx. 1.8) was

determined-measured in plant roots with respect to the other plant organs, (approx. 1.8) thus indicating that the increase in K concentration in plant organs did not inhibit plants' efficiency to uptake Mg (Xie et al., 2021) ~~was not limited by the high plant K concentrations~~. The apparently similar plant uptake of K and Mg is confirmed also by the content of these elements in soil samples at the end of the commercial orchard life-time. Indeed, soil samples were not depleted in K nor Mg with values that ranged between 150 – 400 mg kg dw⁻¹, considering that optimum soil concentrations lie between 240 and 300 mg kg dw⁻¹ for K (Xie et al., 2021) and between 25 and 180 mg kg dw⁻¹ for Mg (Fox and Piekielek, 1984). Moreover, at the end of the experiment, significantly higher values of both K and Mg content were measured with compost fertilization treatment (compost ~~K₂O and MgO~~ mean concentration was ~~of 1.4 ± 0.2 % and 0.72 ± 0.07 % respectively~~); this may indicating indicate a its greater contribution of compost to soil nutrient content with respect to the ~~other-mineral fertilization~~ treatments. Similar results ~~for these macronutrients~~ were also observed in previous studies (Acharya et al., 2019; do Carmo et al., 2016) ~~and could be also related to compost chemical characteristics~~.

331

4.3 ~~Sulfur~~ Sulphur content had a different trend with compost and mineral fertilization ~~destiny in plant roots and soil: fertilization treatment effects~~

Mineral and compost fertilization enhanced S content in fruits s at harvest and in summer leaves more than in autumn leaves. At the end of the nectarine orchard commercial life-time, S content in roots was still high with the two both treatments indicating a potential availability for bud sprout break in the next ~~vegetation-vegetative~~ season. ~~Indeed, S is a component of some secondary metabolites used for plant's physiological functions and development (Narayan et al., 2022). This p~~Plant activity production of secondary metabolites is ~~supposed to be~~ supported by soil S content which ~~indeed~~ resulted particularly high in correspondence of compost addition treatment. Compost ~~as an organic~~

341 ~~matter supply increased~~increases soil organic matter, the largest reservoir of S (in organic form) ~~into~~
342 soil ~~-, and~~Thus compost (in our study characterized by a ~~0.44-18~~0.44-18 ± 0.04 % of SO_3 mean content) or soil
343 organic matter decomposition could result in organic sulfur mineralization into the SO_4^{2-} , which is
344 available to plants (do Carmo et al., 2016; Narayan et al., 2022). ~~However, while s~~Soil S-sulfur content
345 decreased~~d~~ with depth in ~~correspondence of the plots treated with~~ compost ~~addition~~, while it slightly
346 increased or did not change with depth in the control and mineral fertilized plots. This different
347 trend could be ascribed to the different S forms present in the soil: the organic one related to
348 compost, ~~and~~ and thus mainly present in the upper soil layers ~~-, and~~ and the inorganic one, which ~~accumulate~~
349 moves deeper in the soil profile, and is probably less available to plants and more subjected to
350 leaching and/or co-precipitation as calcium, magnesium or sodium ~~sulphate-sulfate~~ (Scherer, 2001).

351

352 4.4 Compost ~~effects on soil chemistry~~increased soil CEC and EC

353 Compost contributes to the increase of exchangeable cations (i.e. Ca^{2+} , K^+ , and Mg^{2+}) creating
354 favorable conditions for cation exchange (Acharya et al., 2019; Dang et al., 2022). Fourteen years of
355 compost addition significantly increased soil CEC and EC not only in the first (0 - 0.15 m) but also
356 in the deeper soil layers. Changes in the CEC of soils are directly linked to the negative charges in
357 the SOM and in the humified compounds (do Carmo et al., 2016) and to the colloidal nature of
358 organic matter (Kumar Bhatt et al., 2019). Similarly, soil EC increase can be explained by the inputs
359 of nutrients and salts contained in the compost and by the soil organic matter mineralization rate.
360 Notwithstanding, the increase ~~of in~~ soil EC values needs to be considered carefully, since ~~-above a~~
361 critical range of 750 - 3490 $\mu\text{S cm}^{-1}$ plant growth could be damaged (do Carmo et al., 2016).




Figure 4 - Macronutrient dynamics in the soil-plant system. Ca, K, Mg and S content removed from the plant (solid grey arrow) and recycled from plant to soil (dashed brown arrow) are expressed in kg ha⁻¹, while Ca, K, Mg and S soil content in soil are expressed in mg over kg dry weight.

5. Conclusion

Considering the goals of this study we have found that: (i) Long-term compost addition facilitates macronutrients storage in the soil and thus indicating that it would favour soil macronutrients availability for the next crops (Figure 4). Moreover, organic amendment addition, not only maintains this is functional both for the current crops and for future ones (Fig. 4); (ii) The supply of compost addition, besides maintaining soil quality and fertility, but also meets plant's nutrient demand thus leading to higher plant performances and to economic and environmental benefits.; (iii) The synchronization between plant needs and nutrient soil availability is fundamental for a correct fertilization management since it avoid plants' nutritional imbalance and,

377 at the same time, reduce the risk of nutrient leaching. The use of compost makes it difficult to
378 guarantee the mentioned synchronization, but significantly contributes to soil nutrient content and
379 soil quality increase. On the other hand, tThe exclusive use of mineral fertilizer, if not carefully
380 managed, even if meeting the plant's nutrient demand, could lead to macronutrient depletion due
381 to scarce reserve creation in the soil; on the other hand, mineral fertilizer represents a source of
382 nutrients readily available to plant uptake. Therefore, the choice of the fertilizer to be used needs to
383 be calibrated on soil and plant requirements taking into account their potential effects (either
384 positive or negative, i.e. the increase of soil organic matter content or the increase of nutrient
385 leaching) on the environment.
386 Future studies should take into consideration the effects of other organic amendments not only in
387 relation to soil and plant nutrient availability, but also on the effects that compost decomposition
388 could have on the amount of CO₂ emitted or sequestered by the orchard, and the impact of
389 macronutrient dynamics on soil microbial communities structure and activity.

390
391

392 **Acknowledgments**

393 We would like to thank Dr. Andrea Simoni and Dr. Giampaolo Di Biase for performing the ICP-AES
394 measurements.

395

396 **Authors Contribution**

397 **Conceptualization:** Elena Baldi, Moreno Toselli; **Methodology:** Elena Baldi; **Formal analysis and**
398 **investigation:** Elena Baldi, Martina Mazzon, Maurizio Quartieri, Luciano Cavani; **Writing – original**
399 **draft preparation:** Martina Mazzon; **Writing – review and editing:** Martina Mazzon, Elena Baldi,

400 Luciano Cavani; **Resources:** Moreno Toselli, Claudio Marzadori; **Supervision:** Moreno Toselli,
401 Claudio Marzadori.

402

403 **References**

404 Acharya, P., Ghimire, R., Cho, Y., 2019. Linking soil health to sustainable crop production: Dairy
405 compost effects on soil properties and sorghum biomass. *Sustain.* 11, 1–13.
406 <https://doi.org/10.3390/su11133552>

407 Adugna, G., 2016. A review on impact of compost on soil properties, water use and crop
408 productivity. *Agric. Sci. Res. J.* 4, 93–104. <https://doi.org/10.14662/ARJASR2016.010>

409 Baldi, E., Cavani, L., Margon, A., Quartieri, M., Sorrenti, G., Marzadori, C., Toselli, M., 2018. Effect
410 of compost application on the dynamics of carbon in a nectarine orchard ecosystem. *Sci. Total*
411 *Environ.* 637–638, 918–925. <https://doi.org/10.1016/j.scitotenv.2018.05.093>

412 Baldi, E., Cavani, L., Mazzon, M., Marzadori, C., Quartieri, M., Toselli, M., 2021a. Fourteen years of
413 compost application in a commercial nectarine orchard: effect on microelements and potential
414 harmful elements in soil and plants. *Sci. Total Environ.* 752, 141894.
415 <https://doi.org/10.1016/j.scitotenv.2020.141894>

416 Baldi, E., Quartieri, M., Sorrenti, G., Toselli, M., 2021b. Evaluation of nutrients removed and recycled
417 in a commercial peach orchard over a 14-years-production cycle. *Italus Hortus* 28, 1–12.
418 <https://doi.org/10.26353/j.itahort/2021.3.0112>

419 Barreto, C.F., Antunes, L.E.C., Toselli, M., Baldi, E., Sorrenti, G., Quartieri, M., 2021. Organic
420 fertilization and crop load in yield and quality of organic nectarines in Italy. *Rev. Bras. Frutic.*
421 43, 0–3. <https://doi.org/10.1590/0100-29452021016>

422 Chatzistathis, T., Papadakis, I.E., Papaioannou, A., Chatzissavvidis, C., Giannakoula, A., 2020.
423 Comparative study effects between manure application and a controlled-release fertilizer on

the growth, nutrient uptake, photosystem II activity and photosynthetic rate of *Olea europaea* L. (cv. 'Koroneiki'). *Sci. Hortic.* (Amsterdam). 264, 109176.
<https://doi.org/10.1016/j.scienta.2020.109176>

Cruz, A.F., de Almeida, G.M., Salvador Wadt, P.G., de Carvalho Pires, M., Gerosa Ramos, M.L., 2019. Seasonal Variation of Plant Mineral Nutrition in Fruit Trees. *Brazilian Arch. Biol. Technol.* 62.
<https://doi.org/10.1590/1678-4324-2019180340>

Dang, L. Van, Ngoc, N.P., Hung, N.N., 2022. Effects of Biochar, Lime, and Compost Applications on Soil Physicochemical Properties and Yield of Pomelo (*Citrus grandis* Osbeck) in Alluvial Soil of the Mekong Delta. *Appl. Environ. Soil Sci.* 2022. <https://doi.org/10.1155/2022/5747699>

Delian, E., Chira, L., Dumitru, L., Badulescu, L., Chira, A., Petcuci, A., 2012. Mineral content of nectarines fruits in relation to some fertilization practices. *Horticulture LVI*, 73–80.

do Carmo, D.L., de Lima, L.B., Silva, C.A., 2016. Soil fertility and electrical conductivity affected by organic waste rates and nutrient inputs davi lopes. *Rev. Bras. Cienc. do Solo* 40, 1–17.
<https://doi.org/10.1590/18069657rbcs20150152>

El-Jendoubi, H., Abadía, J., Abadía, A., 2013. Assessment of nutrient removal in bearing peach trees (*Prunus persica* L. Batsch) based on whole tree analysis. *Plant Soil* 369, 421–437.
<https://doi.org/10.1007/s11104-012-1556-1>

FAO, 2017. The future of food and agriculture - Trends and challenges.

Fawzi, M.I.F., Shahin, .F.M., Elham, A.D., Kandil, E.A., 2010. Effect of organic and biofertilizers and magnesium sulphate on growth yield, chemical composition and fruit quality of "Le-Conte" pear trees. *Nat. Sci.* 8, 1–7.

Fox, R.H., Piekielek, W.P., 1984. Soil magnesium level, corn (*Zea mays* L.) yield, and magnesium uptake. *Commun. Soil Sci. Plant Anal.* 15, 109–123.

Kingston, H.M., Jassie, L.B., 1988. Journal of Research of the National Bureau of Standards Accuracy

in Trace Analysis Microwave Acid Sample Decomposition for Elemental Analysis 93, 1–6.

Kumar Bhatt, M., Labanya, R., Joshi, H.C., 2019. Influence of Long-term Chemical fertilizers and Organic Manures on Soil Fertility - A Review. *Univers. J. Agric. Res.* 7, 177–188. <https://doi.org/10.13189/ujar.2019.070502>

Maity, A., Babu, K.D., Sarkar, A., 2019. Guidelines for fertilizer use in pomegranate orchards based on seasonal uptake and partitioning of nutrients. *Sci. Hortic. (Amsterdam)*. 252, 138–148. <https://doi.org/10.1016/j.scienta.2019.03.047>

Mazzon, M., Cavani, L., Margon, A., Sorrenti, G., Ciavatta, C., Marzadori, C., 2018. Changes in soil phenol oxidase activities due to long-term application of compost and mineral N in a walnut orchard. *Geoderma* 316. <https://doi.org/10.1016/j.geoderma.2017.12.009>

Narayan, O.P., Kumar, P., Yadav, B., Dua, M., Johri, A.K., 2022. Sulfur nutrition and its role in plant growth and development. *Plant Signal. Behav.* 00. <https://doi.org/10.1080/15592324.2022.2030082>

R Core Team, 2021. R: A language and environment for statistical computing.

Safaei Khorram, M., Zhang, G., Fatemi, A., Kiefer, R., Maddah, K., Baqar, M., Zakaria, M.P., Li, G., 2019. Impact of biochar and compost amendment on soil quality, growth and yield of a replanted apple orchard in a 4-year field study. *J. Sci. Food Agric.* 99, 1862–1869. <https://doi.org/10.1002/jsfa.9380>

Scherer, H.W., 2001. Sulphur in crop production - Invited paper. *Eur. J. Agron.* 14, 81–111. [https://doi.org/10.1016/S1161-0301\(00\)00082-4](https://doi.org/10.1016/S1161-0301(00)00082-4)

Sciubba, L., Cavani, L., Grigatti, M., Ciavatta, C., Marzadori, C., 2015. Relationships between stability, maturity, water-extractable organic matter of municipal sewage sludge composts and soil functionality. *Environ. Sci. Pollut. Res.* 22, 13393–13403. <https://doi.org/10.1007/s11356-015-4611-7>

472 Shiwakoti, S., Zheljazkov, V.D., Gollany, H.T., Kleber, M., Xing, B., Astatkie, T., 2020. Macronutrient
 473 in soils and wheat from long-term agroexperiments reflects variations in residue and fertilizer
 474 inputs. *Sci. Rep.* 10, 1–9. <https://doi.org/10.1038/s41598-020-60164-6>

475 Sumner, M.E., Miller, W.P., 1996. Cation Exchange Capacity and Exchange Coefficients, in: Sparks,
 476 D., Page, A., Helmke, P., Loeppert, R., Soltanpour, P.N., Tabatabai, M.A., Johnston, C.T.,
 477 Sumner, M.E. (Eds.), *Methods of Soil Analysis*.
 478 <https://doi.org/https://doi.org/10.2136/sssabookser5.3.c40>

479 Toselli, M., Baldi, E., Cavani, L., Sorrenti, G., 2019a. Nutrient management in fruit crops: An organic
 480 way, in: *Fruit Crops: Diagnosis and Management of Nutrient Constraints*. Elsevier Inc., pp.
 481 379–392. <https://doi.org/10.1016/B978-0-12-818732-6.00027-7>

482 Toselli, M., Baldi, E., Cavani, L., Mazzon, M., Quartieri, M., Sorrenti, G., Marzadori, C., 2019b. Soil-
 483 plant nitrogen pools in nectarine orchard in response to long-term compost application. *Sci.*
 484 *Total Environ.* 671, 10–18. <https://doi.org/10.1016/j.scitotenv.2019.03.241>

485 Wang, X., Hao, L., Zhu, B., Jiang, Z., 2018. Plant calcium signaling in response to potassium
 486 deficiency. *Int. J. Mol. Sci.* 19, 1–16. <https://doi.org/10.3390/ijms19113456>

487 Xie, K., Cakmak, I., Wang, S., Zhang, F., Guo, S., 2021. Synergistic and antagonistic interactions
 488 between potassium and magnesium in higher plants. *Crop J.* 9, 249–256.
 489 <https://doi.org/10.1016/j.cj.2020.10.005>

490 Zhang, Y., Wang, X., Liu, B., Liu, Q., Zheng, H., You, X., Sun, K., Luo, X., Li, F., 2020. Comparative
 491 study of individual and Co-Application of biochar and wood vinegar on blueberry fruit yield
 492 and nutritional quality. *Chemosphere* 246, 125699.
 493 <https://doi.org/10.1016/j.chemosphere.2019.125699>

494