

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Forming and preserving aragonite in shear zones: First report of blueschist facies metamorphism in the Jabal Akhdar Dome, Oman Mountains

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Forming and preserving aragonite in shear zones: First report of blueschist facies metamorphism in the Jabal Akhdar Dome, Oman Mountains / Zuccari, C.; Vignaroli, G.; Callegari, I.; Nestola, F.; Novella, D.; Giuntoli, F.; Guillong, M.; Viola, G.. - In: GEOLOGY. - ISSN 0091-7613. - ELETTRONICO. - 51:5(2023), pp. 454-459. [10.1130/G51079.1]

Availability:

This version is available at: <https://hdl.handle.net/11585/927683> since: 2023-11-14

Published:

DOI: <http://doi.org/10.1130/G51079.1>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Zuccari, C.; Vignaroli, G.; Callegari, I.; Nestola, F.; Novella, D.; Giuntoli, F.; Guillong, M.; Viola, G.: Forming and preserving aragonite in shear zones: First report of blueschist facies metamorphism in the Jabal Akhdar Dome, Oman Mountains

GEOLOGY Vol. 51 ISSN 0091-7613

DOI: 10.1130/G51079.1

The final published version is available online at:

<https://dx.doi.org/10.1130/G51079.1>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Forming and preserving aragonite in shear zones: first report of blueschist facies metamorphism in the Jabal Akhdar Dome, Oman Mountains

Tracking no: G51079

Authors:

Costantino Zuccari (Università degli Studi di Bologna), Gianluca Vignaroli (Università degli Studi di Bologna), Ivan Callegari (German University of Technology in Oman - GUTech), Fabrizio Nestola (Università degli Studi di Padova), Davide Novella (Università degli Studi di Padova), Francesco Giuntoli (Università degli Studi di Bologna), Marcel Guillong (ETH Zurich), and Giulio Viola (University of Bologna)

Abstract:

We report the first occurrence of high-pressure metamorphic aragonite in Precambrian carbonates of the Jabal Akhdar Dome in the Oman Mountains. We offer a model for its formation at blueschist facies metamorphic conditions and its subsequent preservation to the surface within the tectonic framework of the late Cretaceous obduction of the Semail Ophiolite. Aragonite formed at $T \sim 350^{\circ}\text{C}$ and $P \geq 0.9$ GPa and is preserved within mylonitic shear zones and in stretched fiber dilational veins where the necessary conditions for its formation and preservation, such as plastic strain accommodation, fluid-enhanced mineralogical reactions, and an anisotropic permeability structure, were preferentially met with respect to the surrounding rock. High-strain structural domains are thus documented to potentially be ideal sites where to look for- and study pro- and retrograde high-pressure metamorphic histories in deeply subducted and exhumed terrains.

1 Forming and preserving aragonite in shear zones: first report of
2 blueschist facies metamorphism in the Jabal Akhdar Dome,
3 Oman Mountains

4
5 Zuccari C.^{1*}, Vignaroli G.¹, Callegari I.², Nestola F.³, Novella D.³, Giuntoli F.¹, Guillong M.⁴
6 and Viola G.^{1*}

7
8 ¹*Alma Mater Studiorum, University of Bologna, Department of Biological, Geological and*
9 *Environmental Sciences - BiGeA, Bologna, Italy.*

10 ²*German University of Technology in Oman – GUTech, Department of Applied Geosciences,*
11 *Muscat, Oman.*

12 ³*University of Padua, Department of Geosciences, Padova, Italy.*

13 ⁴*Department of Earth Sciences, Institute of Geochemistry and Petrology, ETH Zürich, Zurich,*
14 *Switzerland.*

15
16 ***Corresponding authors**
17
18
19
20
21
22
23
24
25
26

ABSTRACT

We report the first occurrence of high-pressure metamorphic aragonite in Precambrian carbonates of the Jabal Akhdar Dome in the Oman Mountains. We propose a model for both its formation at blueschist facies conditions and its subsequent preservation to the surface within the tectonic framework of the late Cretaceous obduction of the Semail Ophiolite. Aragonite formed at $T \sim 350^\circ\text{C}$ and $P \geq 0.9\text{GPa}$ and is preserved within mylonitic shear zones and in stretched fiber dilational veins where the necessary conditions for its formation and preservation, such as plastic strain accommodation, fluid-enhanced mineralogical reactions, and an anisotropic permeability structure were preferentially met with respect to the surrounding rock. High-strain structural domains are ideal sites where to look for and study pro- and retrograde high-pressure metamorphic histories in deeply subducted and exhumed terrains.

1. INTRODUCTION

The identification and characterization of relic high-pressure (HP) assemblages in exhumed rocks are key to constraining the thermobaric conditions and the deformation processes typical of otherwise inaccessible subduction zones (e.g., [Agard et al., 2009](#); [Rubatto et al., 2011](#)). Even though exhumed terrains may be affected by retrograde metamorphism, relic assemblages may locally survive within favourable structural sites that act as sheltering capsules to specific HP phases. Formation and later preservation of such phases therein is favoured or inhibited by the interplay of numerous factors, such as the exhumation rate and geothermal gradient, which may inhibit retrograde mineralogical re-equilibrations, presence and composition of fluids, which can favour element transport and enhance mineralogical reactions during pro- and retrograde histories, amount of strain and type of involved mineralogical phases (e.g., [Goffé and Velde, 1984](#); [Bucher and Frey, 2002](#)). The documentation of index minerals, such as lawsonite and carpholite, has facilitated the study of blueschist facies conditions within metabasite and

metapelite (e.g., [Evans and Brown, 1986](#)). Aragonite, on the other hand, is a metastable polymorph of calcite that represents a reliable blueschist facies indicator for “pure” carbonates that otherwise lack other P-sensitive mineral phases (e.g., [Hacker et al., 1992](#); [Stöckhert et al., 1999](#); [Giuntoli et al., 2020](#)). Aragonite formation and preservation are mostly limited to <10°C/km thermal gradients and rapid exhumation in dry conditions (e.g., [Carlson, 1980](#); [Hacker et al., 1992](#)).

We report the first record of HP aragonite in the Precambrian carbonates of the Jabal Akhdar Dome (Oman Mountains), universally acknowledged as a non-subducted portion of the Arabian Plate (e.g., [Breton et al., 2004](#)). We demonstrate that aragonite is therein selectively preserved within narrow mylonitic shear zones that formed due to cyclic brittle-ductile deformation under blueschist facies conditions during the Cretaceous obduction of the Semail Ophiolite. We propose a model wherein the identified mylonitic shear zones acted as sheltering structural capsules, within which aragonite initially formed at HP metamorphic conditions and through which it survived exhumation all the way to the surface.

2. GEOLOGICAL SETTING

The Jabal Akhdar Dome is the largest tectonic window in the NW-SE Oman Mountains ([Fig. 1A-B](#)). It is composed of a pre-mid Carboniferous series overlain by a 2.5km-thick Permian to Cretaceous carbonate succession (Autochthonous A and B, respectively). This stack was overthrust by the Hawasina Nappe and the Semail Ophiolite during the Cretaceous (e.g., [Searle, 2007](#); [Fig. 1B](#)). The autochthonous series is part of the Arabian passive margin, which is reported as having experienced subduction, exhumation and obduction only in its northeasternmost external portion (e.g., [Lippard, 1983](#)). Therefore, deformation in the Precambrian JAD has so far been predominantly related to a complex sequence of contractional and extensional phases from the late Ediacaran to the early Cambrian (e.g., [Callegari et al., 2020](#); [Scharf et al., 2021](#); [Fig. 1C](#)), with evidence of Cretaceous continental subduction

exclusively confined to the Saih Hatat Dome (Fig. 1B), where rocks experienced up to eclogite facies conditions (e.g., Warren et al., 2003; Agard et al., 2010). The JAD in the inner part of the Arabian margin is thus considered to belong to the un-subducted portion of the Arabian passive margin, and to have experienced only anchizonal deformation conditions during the Semail Ophiolite obduction (e.g., Searle, 2007). Exhumation of the subducted Arabian margin through a still ill-defined mechanism started at the end of the Cretaceous (e.g., Hansman et al., 2021), leading to rapid cooling of the whole region (e.g., Grobe et al., 2019).

3. METHODS AND RESULTS

3.1 ARAGONITE-BEARING MYLONITIC SHEAR ZONES

In the study area (Fig. 1C), contractional calcmylonitic shear zones cut across the Precambrian organic-matter-rich carbonate of the Hajir Formation (Fig. 2; Fig. S1). Strain localised in 1) discrete, up to 20cm thick mylonitic to ultramylonitic shear zones (Figs. 2A-B; S1) that discordantly cut across the bedding and 2) thinner shears along bed-bed interfaces within NE-verging folds in response to a flexural slip folding mechanism (Fig. 2). In the former, the mylonitic foliation is penetrative and defined by lenses and laterally continuous films of graphite (Figs. 2C; S1), dips to the SW (Fig. 2B) and is invariably associated with a SW-plunging stretching lineation and thrusting-related top-to-NE S/C-structures and oblique foliation (Fig. 2B). The mylonitic foliation commonly transposes quartz and calcite veins (Fig. 2B). Carbonates exhibit a strong shape preferred orientation (SPO, Fig. 2C) with grains elongated parallel to the transport direction and contained within the foliation. They preserve a peculiar “rod-like” shape (Fig. 2C), which is commonly reported as typical for calcite pseudomorphs after aragonite (e.g., Brady et al., 2004; Seaton et al., 2009; Giuntoli et al., 2020), and as ensuing during shearing at the calcite to aragonite phase transformation. Such rod-shaped fabric is not found within the less deformed Hajir Fm outside of the shear zones (Fig. S1).

Shear zones related to flexural-slip folding along bed-bed interfaces are extremely localised and thin (<5cm; Fig. 2D-E) and contain sheared quartz and calcite-aragonite veins and fibres (Fig. 2E). They exhibit a strong SPO with calcite and aragonite grains aligned parallel to the mylonitic foliation, embedding and transposing lenses and films of graphite (Fig. 2F). All shear zones document broadly coeval brittle and ductile deformation, with the mylonitic fabric cut across by mode-I veins with opening directions subparallel to the regional SW-NE shear direction. These veins contain aragonite and quartz fibres (Fig. 2F; Fig. S2A-B) and are, themselves transposed along the mylonitic foliation, suggesting cyclic brittle-ductile deformation (Figs. 2B-F; S2C-D; S3A-B).

3.2 *P-T CONSTRAINTS*

Raman Spectroscopy on Carbonaceous Material (RSCM) spectra and high-resolution micro-Raman maps (see Supplementary Material) were acquired to discriminate calcite from aragonite in both mylonites and mode-I veins. Representative RSCM spectra from graphitic films and patches, which outline the mylonitic foliation, constrain synkinematic deformation temperatures in the 336-347°C range (Fig. 3A). Micro-Raman maps and spectra document aragonite (only partially retrogressed to calcite) in both mylonitic shear zones and mode-I veins (Fig. 3B-C), thus confirming inferences from the mylonitic rod-shaped fabric in the shear zones (Fig. 2C and F).

Based on the calcite-aragonite stability field (e.g., Johannes and Puhon, 1971), the obtained deformation temperature range requires 0.8-0.9GPa (Fig. 3D) as minimum pressure during shearing and coeval brittle deformation, thus within the range of blueschist facies conditions. Trace element analysis on aragonite fibres and grains in mylonites constrains Sr to <1wt% (Table S1; Fig. S4 and S5), thus excluding lower P values for the calcite-aragonite transformation (e.g., Carlson, 1980).

3.3 AGE OF DEFORMATION

In order to temporally place the formation of the studied shear zones in the evolution of northeastern Oman, we dated by U-Pb the described tectonic carbonates. Ages from mode-I carbonate fibres cluster at 96.5 ± 31.6 Ma (2σ ; Fig. S6A-C, E). Dating of the calcmylonitic shear zones defines two different clusters at 97.5 ± 23.3 Ma and 573.3 ± 27.6 Ma, respectively (2σ ; Fig. S6D-F). The large analytical uncertainties notwithstanding, we interpret the new dates as related to the Cretaceous subduction-obduction of the Semail Ophiolite, whereas the Precambrian date as the age of the Hajir Fm protolith, which is ascribed to the early-middle Ediacaran (e.g., Scharf et al., 2021).

4. DISCUSSION AND CONCLUSIONS

This first documentation of blueschist facies conditions from genetically connected shear zones and mode-I veins within the JAD and the interpretation of this data require considering the possible role of non-lithostatic pressure during deformation. Tectonic overpressure may promote HP metamorphism, such that analytically obtained P estimates cannot be converted to depth. Several conditions are discussed as leading to tectonic overpressure, among which the following might affect our results: i) significant viscosity contrast between juxtaposed rock types, ii) heterogenous pressure conditions in folds and iii) fluid release (dehydration, decarbonation) in a closed system (e.g., Mancktelow, 2008; Schmalholz and Podladchikov, 2013; Luisier et al., 2019).

We tend to exclude a significant contribution of tectonic overpressure during the evolution of the studied shear zones in the JAD based on the following:

- Mode-I fracturing causes instantaneous pressure drops in dilatant fractures to conditions corresponding to the local lithostatic load. Coeval fracture infilling is, therefore, at equilibrium with the lithostatic pressure and aragonite in veins thus reflects the overburden.

- Mylonitic shearing occurred in the absence of a significant viscosity contrast between shear zone and the undeformed host rock, as evidenced by their similar composition and by aragonite occurring in both shear zones and mode-I veins.
- Shear zones locally formed by flexural slip along bed-bed interfaces on fold limbs, which are not prone to develop significant overpressure (e.g., [Mancktelow, 2008](#)).
- Shearing occurred at too low a temperature to release CO₂-rich fluids from decarbonation (e.g., [Samtani et al., 2002](#)).

Theoretically, aragonite might form even at low-pressure conditions in highly-strained domains, where shear strain would increase the internal energy of calcite crystals by densifying defects and dislocations in the lattice ([Newton et al., 1969](#)). This process, however, has been described for late calcite veins cutting across mylonitic fabrics and is thought to be rather unlikely in natural shear zones ([Gillet and Goffé, 1988](#)).

We therefore consider our minimum P estimate of 0.8-0.9GPa as indicative of at least 25-30km depth, which attests to reaching blueschist facies conditions in the dynamic context of the Oman Cretaceous subduction. This is consistent with the new U-Pb results, which, despite the large analytical errors (24-30%; [Fig. S6E-F](#)), fit available age constraints for the known HP and subduction-related metamorphism in northeastern Oman ([Fig. S6G](#)). We also exclude an Ediacaran sedimentary origin for aragonite because we deem highly unlikely its preservation over such a long time span due to the rapid kinetics of the aragonite/calcite transformation (e.g., [Theye and Seidel, 1993](#); [Hacker et al., 2005](#)).

Cyclic brittle-ductile deformation under HP conditions in the JAD finds analogies with descriptions of similar deformation styles under blueschist facies conditions elsewhere in orogenic belts (e.g., [Molli et al., 2017](#); [Giuntoli and Viola, 2022](#)) and allows us to present a conceptual model for the formation and preservation of aragonite within mylonitic shear zones ([Fig. 4](#)). Formation of aragonite is known to be both favoured and limited by the i) amount of strain, ii) dynamic evolution of the permeability tensor through time and fluid-rock interaction,

183 iii) temperature changes, iv) overall composition of the deforming system and v) exhumation
184 modes (Fig. 4A). The interplay of these factors within the studied shear zones would have
185 made it possible for aragonite to first form and then be preserved. We suggest that the
186 mylonitic fabric triggered transient chemical/physical processes that were, instead, unable to
187 operate in the less deformed Hajir Fm outside of the shear zones (as shown by the lack of
188 aragonite therein; Fig. S1).

189 Shear zones cyclically accommodated broadly coeval ductile and brittle deformation,
190 both taking place under blueschist facies conditions, as demonstrated by aragonite occurring in
191 both coeval mylonites and mode-I veins. Progressive strain localisation within the initially
192 undeformed Hajir Fm under increasing pressure and temperature (Fig. 4B) caused calcite
193 recrystallization and grain size reduction (Fig. 4C). Fluids enhanced the calcite-aragonite
194 transformation at favourable P-T conditions that were met as the Hajir Fm reached
195 progressively greater depth during subduction (Carlson, 1980). We see this process as being
196 favoured by enhanced permeability along the mylonitic foliation due to the progressive
197 alignment of ordered graphite films along the mylonitic foliation planes (Fig. 4C; Upton and
198 Craw, 2008). Fluid ingress and flow within shear zones was facilitated by broadly coeval
199 fracturing and viscous deformation cyclically alternating under HP conditions (e.g., Molli et al.,
200 2017) and by the establishment of a large foliation-parallel permeability, with shear zones thus
201 acting as conduits (Fig. 4D).

202 Close to or at P-T peak conditions, rod-shaped aragonite grains formed (Fig. 4C-D)
203 growing an interwoven network with graphite crystals to develop laterally continuous and
204 extensive layers (Fig. 4E). This would have sealed the shear zones, eventually leading to lower
205 permeability conditions compared to the less deformed hosting Hajir Fm. Low-permeability
206 and sealed shear zones characterised the exhumation retrograde path of this part of the
207 subducted slab (Fig. 4E). Finally, the Hajir Fm re-entered the stability field of calcite at
208 $T \leq 200^\circ\text{C}$ (e.g., Hacker et al., 1992; Fig. 4A and F), with shear zones remaining relatively

209 sealed and dry, thus fulfilling a key condition for aragonite preservation (e.g., [Gillet and Goffé,](#)
210 [1988](#)). When coupled with a low geothermal gradient ($<10^{\circ}\text{C}/\text{km}$) typical of cold subduction
211 zones and fast exhumation ($\sim 1\text{-}3\text{mm/a}$), as indeed reported for NE Oman (e.g., [Grobe et al.,](#)
212 [2019](#)), our proposed structural evolution explains the seldomly observed crystallographic and
213 morphologic preservation of aragonite in the stability field of calcite.

214 In conclusion, we document for the first time the occurrence of metamorphic aragonite
215 in the JAD, attesting to blueschist facies conditions during subduction to at least 25-30 km
216 depth. Its preservation calls for a model based on structural capsules where HP phases
217 selectively formed and were later preserved by the concomitant effects of regional (rapid
218 exhumation) and local factors (low permeability horizons). Our results provide new insights
219 into both the regional evolution of the JAD and the processes allowing for the preservation of
220 HP phases in deeply subducted and exhumed carbonate(meta)sedimentary successions.

221 222 **ACKNOWLEDGEMENTS**

223 We thank Alberto Vitale Brovarone, Urs Schaltegger and Maria Ovtcharova for the discussions
224 about the aragonite-calcite transformation and the dating methodology. The manuscript
225 benefited from the fruitful comments of the editor Rob Strachan, Thomas Lamont and five
226 anonymous reviewers.

REFERENCES

- Agard, P., Searle, M.P., Alsop, G.I., and Dubacq, B., 2010, Crustal stacking and expulsion tectonics during continental subduction: P-T deformation constraints from Oman: *Tectonics*, v. 29, p. 1–19, [doi:10.1029/2010TC002669](https://doi.org/10.1029/2010TC002669).
- Agard, P., Yamato, P., Jolivet, L., and Burov, E., 2009, Exhumation of oceanic blueschists and eclogites in subduction zones: Timing and mechanisms: *Earth-Science Reviews*, v. 92, p. 53–79, [doi:10.1016/j.earscirev.2008.11.002](https://doi.org/10.1016/j.earscirev.2008.11.002).
- Brady, J.B., Markley, M.J., Schumacher, J.C., Cheney, J.T., and Bianciardi, G.A., 2004, Aragonite pseudomorphs in high-pressure marbles of Syros, Greece: *Journal of Structural Geology*, v. 26, p. 3–9, [doi:10.1016/S0191-8141\(03\)00099-3](https://doi.org/10.1016/S0191-8141(03)00099-3).
- Breton, J.P., Béchenec, F., le Métour, J., Moen-Maurel, L., and Razin, P., 2004, Eoalpine (Cretaceous) evolution of the Oman Tethyan continental margin: insights from a structural field study in Jabal Akhdar (Oman Mountains): *GeoArabia*, v. 9, p. 41–58, [doi:10.2113/GEOARABIA090241](https://doi.org/10.2113/GEOARABIA090241).
- Bucher, K., and Frey, M., 2002, Petrogenesis of Metamorphic Rocks: Petrogenesis of Metamorphic Rocks, [doi:10.1007/978-3-662-04914-3](https://doi.org/10.1007/978-3-662-04914-3).
- Callegari, I., Scharf, A., Mattern, F., Bauer, W., Pinto, A.J., Rarivoarison, H., Scharf, K., and al Kindi, M., 2020, Gondwana accretion tectonics and implications for the geodynamic evolution of eastern Arabia: First structural evidence of the existence of the Cadomian Orogen in Oman (Jabal Akhdar Dome, Central Oman Mountains): *Journal of Asian Earth Sciences*, v. 187, [doi:10.1016/j.jseaes.2019.104070](https://doi.org/10.1016/j.jseaes.2019.104070).
- Carlson, W.D., 1980, The calcite–aragonite equilibrium: effects of Sr substitution and anion orientational disorder: *American Mineralogist*, v. 65, p. 1252–1262.
- Clark Jr., S.P., 1957, A Note on Calcite-Aragonite Equilibrium*: *American Mineralogist*, v. 42, p. 564–566.
- Evans, B.W., and Brown, E.H., 1986, Blueschists and Eclogites: v. 164, [doi:10.1130/MEM164](https://doi.org/10.1130/MEM164).
- Gillet, P., and Goffé, B., 1988, On the significance of aragonite occurrences in the Western Alps: Contributions to Mineralogy and Petrology, v. 99, p. 70–81, [doi:10.1007/BF00399367](https://doi.org/10.1007/BF00399367).
- Giuntoli, F., and Viola, G., 2022, A likely geological record of deep tremor and slow slip events from a subducted continental broken formation: *Scientific Reports*, v. 12, p. 4506, [doi:10.1038/s41598-022-08489-2](https://doi.org/10.1038/s41598-022-08489-2).

- Giuntoli, F., Vitale Brovarone, A., and Menegon, L., 2020, Feedback between high-pressure genesis of abiogenic methane and strain localization in subducted carbonate rocks: *Scientific Reports*, v. 10, p. 9848, [doi:10.1038/s41598-020-66640-3](https://doi.org/10.1038/s41598-020-66640-3).
- Goffé, B., and Velde, B., 1984, Contrasted metamorphic evolutions in thrust cover units of the Briançonnais zone (French Alps): A model for the conservation of HP-LT metamorphic mineral assemblages: *Earth and Planetary Science Letters*, v. 68, p. 351–360, [doi:10.1016/0012-821X\(84\)90166-3](https://doi.org/10.1016/0012-821X(84)90166-3).
- Grobe, A., von Hagke, C., Littke, R., Dunkl, I., Wübbeler, F., Muchez, P., and Urai, J.L., 2019, Tectono-Thermal evolution of Oman's Mesozoic passive continental margin under the obducting Semail Ophiolite: A case study of Jebel Akhdar, Oman: *Solid Earth*, v. 10, p. 149–175, [doi:10.5194/se-10-149-2019](https://doi.org/10.5194/se-10-149-2019).
- Hacker, B.R., Kirby, S.H., and Bohlen, S.R., 1992, Time and metamorphic petrology: Calcite to aragonite experiments: *Science*, v. 258, p. 110–112, [doi:10.1126/science.258.5079.110](https://doi.org/10.1126/science.258.5079.110).
- Hacker, B.R., Rubie, D.C., Kirby, S.H., and Bohlen, S.R., 2005, The calcite → aragonite transformation in low-Mg marble: Equilibrium relations, transformations mechanisms, and rates: *Journal of Geophysical Research: Solid Earth*, v. 110, p. 1–16, [doi:10.1029/2004JB003302](https://doi.org/10.1029/2004JB003302).
- Hansman, R.J., Ring, U., Scharf, A., Glodny, J., and Wan, B., 2021, Structural architecture and Late Cretaceous exhumation history of the Saih Hatat Dome (Oman), a review based on existing data and semi-restorable cross-sections: *Earth-Science Reviews*, v. 217, p. 103595, [doi:10.1016/j.earscirev.2021.103595](https://doi.org/10.1016/j.earscirev.2021.103595).
- Johannes, W., and Puhon, D., 1971, The Calcite-Aragonite Transition , Reinvestigated: *Contributions to Mineralogy and Petrology*, v. 38, p. 28–38.
- Lin, S.J., and Huang, W.L., 2004, Polycrystalline calcite to aragonite transformation kinetics: Experiments in synthetic systems: *Contributions to Mineralogy and Petrology*, v. 147, p. 604–614, [doi:10.1007/s00410-004-0574-2](https://doi.org/10.1007/s00410-004-0574-2).
- Lippard, S.J., 1983, Cretaceous high pressure metamorphism in NE Oman and its relationship to subducted and ophiolite nappe emplacement.: *Journal of the Geological Society*, v. 140, p. 97–104, [doi:10.1144/GSJGS.140.1.0097](https://doi.org/10.1144/GSJGS.140.1.0097).
- Luisier, C., Baumgartner, L., Schmalholz, S.M., Siron, G., and Vennemann, T., 2019, Metamorphic pressure variation in a coherent Alpine nappe challenges lithostatic pressure paradigm: *Nature Communications*, v. 10, [doi:10.1038/s41467-019-12727-z](https://doi.org/10.1038/s41467-019-12727-z).
- Mancktelow, N.S., 2008, Tectonic pressure: Theoretical concepts and modelled examples: *Lithos*, v. 103, p. 149–177, [doi:10.1016/j.lithos.2007.09.013](https://doi.org/10.1016/j.lithos.2007.09.013).

- 294 Molli, G., Menegon, L., and Malasoma, A., 2017, Switching deformation mode and
295 mechanisms during subduction of continental crust: a case study from Alpine Corsica:
296 Solid Earth, v. 8, p. 767–788, [doi:10.5194/se-8-767-2017](https://doi.org/10.5194/se-8-767-2017).
- 297 Newton, R.C., Goldsmith, J.R., and Smith, J. v, 1969, Aragonite crystallization from strained
298 calcite at reduced pressures and its bearing on aragonite in low-grade metamorphism:
299 Contributions to Mineralogy and Petrology, v. 22, p. 335–348, [doi:10.1007/BF00400128](https://doi.org/10.1007/BF00400128).
- 300 Rubatto, D., Regis, D., Hermann, J., Boston, K., Engi, M., Beltrando, M., and McAlpine,
301 S.R.B., 2011, Yo-yo subduction recorded by accessory minerals in the Italian Western
302 Alps: Nature Geoscience, v. 4, p. 338–342, [doi:10.1038/ngeo1124](https://doi.org/10.1038/ngeo1124).
- 303 Samtani, M., Dollimore, D., and Alexander, K.S., 2002, Comparison of dolomite
304 decomposition kinetics with related carbonates and the effect of procedural variables on its
305 kinetic parameters: Thermochimica Acta, v. 392–393, p. 135–145, [doi:10.1016/s0040-
306 6031\(02\)00094-1](https://doi.org/10.1016/s0040-6031(02)00094-1).
- 307 Scharf, A., Mattern, F., Al-Wardi, M., Frijia, G., Moraetis, D., Pracejus, B., Bauer, W., and
308 Callegari, I., 2021, The Geology and Tectonics of the Jabal Akhdar and Saih Hatat
309 Domes, Oman Mountains: Geological Society, London, Memoirs, v. 54, p. 113–115,
310 [doi:10.1144/M54.7](https://doi.org/10.1144/M54.7).
- 311 Schmalholz, S.M., and Podladchikov, Y.Y., 2013, Tectonic overpressure in weak crustal-scale
312 shear zones and implications for the exhumation of high-pressure rocks: Geophysical
313 Research Letters, v. 40, p. 1984–1988, [doi:10.1002/grl.50417](https://doi.org/10.1002/grl.50417).
- 314 Searle, M.P., 2007, Structural geometry, style and timing of deformation in the Hawasina
315 Window, Al Jabal al Akhdar and Saih Hatat culminations, Oman Mountains: GeoArabia,
316 v. 12, p. 99–130, [doi:10.2113/geoarabia120299](https://doi.org/10.2113/geoarabia120299).
- 317 Seaton, N.C.A., Whitney, D.L., Teyssier, C., Toraman, E., and Heizler, M.T., 2009,
318 Recrystallization of high-pressure marble (Sivrihisar, Turkey): Tectonophysics, v. 479, p.
319 241–253, [doi:10.1016/j.tecto.2009.08.015](https://doi.org/10.1016/j.tecto.2009.08.015).
- 320 Stöckhert, B., Wachmann, M., Küster, M., and Bimmermann, S., 1999, Low effective viscosity
321 during high pressure metamorphism due to dissolution precipitation creep: The record of
322 HP-LT metamorphic carbonates and siliciclastic rocks from Crete: Tectonophysics, v.
323 303, p. 299–319, [doi:10.1016/S0040-1951\(98\)00262-5](https://doi.org/10.1016/S0040-1951(98)00262-5).
- 324 Theye, T., and Seidel, E., 1993, Uplift-related retrogression history of aragonite marbles in
325 Western Crete (Greece): Contributions to Mineralogy and Petrology, v. 114, p. 349–356,
326 [doi:10.1007/BF01046537](https://doi.org/10.1007/BF01046537).

Upton, P., and Craw, D., 2008, Modelling the role of graphite in development of a mineralised mid-crustal shear zone, Macraes mine, New Zealand: *Earth and Planetary Science Letters*, v. 266, p. 245–255, [doi:10.1016/j.epsl.2007.10.048](https://doi.org/10.1016/j.epsl.2007.10.048).

Warren, C.J., Parrish, R.R., Searle, M.P., and Waters, D.J., 2003, Dating the subduction of the Arabian continental margin beneath the Semail ophiolite, Oman: *Geology*, v. 31, p. 889–892, [doi:10.1130/G19666.1](https://doi.org/10.1130/G19666.1).

CAPTIONS TO FIGURES

Fig. 1. A) Geographical setting and B) Geological setting of the southern Oman Mountains with the Jabal Akhdar (JAD) and Saih Hatat (SHD) Domes; from [Searle, 2007](#). C) Geological map of the study area and local stratigraphy (from [Callegari et al., 2020](#)).

Fig. 2. A) Calcmylonitic shear zone cutting a NE-verging fold. B) Detail of mylonitic shear zone in (A). C) Ultramylonitic shear zone in (B), with rod-shaped aragonite crystals interleaved with semi-continuous graphitic films defining the foliation. D) NE-verging fold with flexural-slip related shear zones along bed-bed interfaces. Men for scale. E) Detail of shear zone in (D) deforming fibrous quartz and aragonite. F) Aragonite-calcite mylonitic foliation cut across by mode-I stretched fiber quartz and aragonite veins.

Fig. 3. A) Raman spectra from carbonaceous material (RSCM) on rock matrix with (black) and without (blue) polishing. B-C) High-resolution micro-Raman spectroscopy spectra, related microphotograph and phase map. Bright yellow aragonite in blue ellipses. D) Calcite-aragonite stability field. Different experimental stability curves are shown (from Clark Jr., 1957; Johannes and Puhon, 1971; Lin and Huang, 2004; Hacker et al., 2005). The estimated P-T conditions for studied JAD carbonates are indicated.

354 Fig. 4. Model for the formation/preservation of aragonite. A) P-T retrograde path. B)
355 Undeformed Hajir Fm carbonate. C) Calcite dynamic recrystallization forming new grains. D)
356 Shear zone structuring at P-T peak conditions and formation of rod-shaped aragonite grains.
357 with Quartz and aragonite fibre mode-I veins locally transposed by mylonitic fabrics constrain
358 brittle-ductile cyclicity under HP conditions. E) Shear zone sealing by high-ordered graphite
359 coalescence, forming thin, permeable and laterally continuous layers. F) Present configuration
360 of the JAD mylonitic shear zones, acting as sheltering structural capsules preserving aragonite.







