



ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Influence of high-pressure homogenization treatments combined with lysozyme activated packaging on microbiological and technological quality of vegetable smoothie during shelf life

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Influence of high-pressure homogenization treatments combined with lysozyme activated packaging on microbiological and technological quality of vegetable smoothie during shelf life / Siroli, Lorenzo; Glicerina, Virginia; Capelli, Filippo; Patrignani, Francesca; Fiorini, Maurizio; Andrisano, Vincenza; Minelli, Matteo; Gherardi, Matteo; Lanciotti, Rosalba; Colombo, Vittorio; Romani, Santina. - In: FOOD PACKAGING AND SHELF LIFE. - ISSN 2214-2894. - ELETTRONICO. - 37:(2023), pp. 101093.1-101093.9. [10.1016/j.fpsl.2023.101093]

This version is available at: <https://hdl.handle.net/11585/929953> since: 2023-06-08

Published:

DOI: <http://doi.org/10.1016/j.fpsl.2023.101093>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

(Article begins on next page)

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

This is the final peer-reviewed accepted manuscript of:

Lorenzo Siroli, Virginia Glicerina, Filippo Capelli, Francesca Patrignani, Maurizio Fiorini, Vincenza Andrisano, Matteo Minelli, Matteo Gherardi, Rosalba Lanciotti, Vittorio Colombo, Santina Romani, Influence of high-pressure homogenization treatments combined with lysozyme activated packaging on microbiological and technological quality of vegetable smoothie during shelf life, Food Packaging and Shelf Life, Volume 37, 2023, 101093, ISSN 2214-2894.

The final published version is available online at:
<https://doi.org/10.1016/j.fpsl.2023.101093>.

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

1 **Influence of high-pressure homogenization treatments combined with lysozyme activated**
2 **packaging on microbiological and technological quality of vegetable smoothie during shelf life**

3
4 **Lorenzo Siroli^{ab}, Virginia Glicerina^{*c}, Filippo Capelli^d, Francesca Patrignani^{ab}, Maurizio**
5 **Fiorini^e, Vincenza Andrisano^f, Matteo Minelli^e, Matteo Gherardi ^{dg}, Rosalba Lanciotti^{ab},**
6 **Vittorio Colombo^{dg} and Santina Romani^{ab}**

7 ^a Department of Agricultural and Food Sciences, University of Bologna, 40127, Bologna, Italy

8 ^b Interdepartmental Centre for Agri-Food Industrial Research, Alma Mater Studiorum, University of
9 Bologna, Campus of Food Science, 47521, Cesena (FC), Italy

10 ^c Department of Agriculture, Forest and Food Sciences, University of Turin, 10095 Grugliasco (TO),
11 Italy

12 ^d Department of Industrial Engineering, Bologna, 40131, Italy

13 ^e Department of Civil, Chemical, Environmental and Materials Engineering, Alma Mater
14 Studiorum, University of Bologna, 40136, Bologna, Italy

15 ^f Department for Life Quality Studies, Alma Mater Studiorum – University of Bologna, 47921, Rimini,
16 Italy

17 ^g Advanced Mechanics and Materials, Interdepartmental Center for Industrial Research (AMM-13
18 ICIR), 40131, Bologna, Italy

19
20
21 ^{*} Corresponding author: virginiateresa.glicerina@unito.it

22

23

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

Abstract

In this study the effect of a high pressure homogenization (HPH) treatment, alone and in combination with a biodegradable polylactic acid (PLA) packaging activated with lysozyme on a smoothie made with apple, carrot and rice beverage was evaluated. In particular, the effects of the treatments on the microbiological and quality characteristics of the smoothies were assessed during storage at 4 and 10 °C in climatic chambers at RH of 50%. Obtained results showed the efficacy of HPH treatment at 100 MPa to reduce the initial cell load of mesophilic and psychotrophic aerobic bacteria, lactic acid bacteria, yeasts and aerobic spore-forming bacteria. Moreover, the combination of HPH and lysozyme activated packaging resulted in an increased antimicrobial effect against mesophilic, psychotropic and lactic acid bacteria during product storage at 10 °C. Yeast and spore forming bacteria showed, instead, lower cell loads independently from the samples. Samples packed in active packaging, both treated and not, showed also a better color retention in terms of lightness and red index, compared to the control (C) ones both when stored at 4 and 10 °C. The HPH treated smoothies, both activated and not, stored at 4 °C remained microbiologically stable for more than 20 days. However, the use of active packaging has also allowed to reduce the microbiological proliferation during storage at 4 °C. Moreover, few differences in color indices were highlighted between samples stored at 4 °C. Obtained results indicate that the combination of HPH treatment and active packaging may represent a useful strategy to increase safety and shelf-life of vegetable smoothies.

49

50 **Keywords: High Pressure, lysozyme, active packaging, cold plasma, shelf- life, vegetable juices,**
51 **combined treatments, quality, safety**

52

53

54 **1. Introduction**

55 Vegetable smoothies are currently very popular beverages, which are consumed by people of all ages
56 for their nutritional and sensory properties. They are perceived by consumers as "healthy" foods due
57 to their low sodium, cholesterol, fat content and because they are rich in vitamin C, polyphenols and
58 flavonoids. The latter compounds contribute to the good antioxidant properties of these products
59 (Kumar et al. 2009; Patrignani, Vannini, Kamdem, Lanciotti, & Guerzoni, 2009; Patrignani et al.
60 2019). Furthermore, the current lifestyle has led consumers to look for more practical and ready-to-
61 eat products (Andrés, Villanueva, & Tenorio, 2016). For these reasons, one of the fastest growing
62 sectors in the fruit juice market in the last 10 years is represented by smoothies (Nieva, Jagus, Agüero
63 & Fernandez, 2022). Smoothies are blended beverages that can contain fruit pulp, fruit juice,
64 vegetables, yoghurt, milk or vegetable beverage (Nunes et al. 2016). **The smoothies are generally**
65 **minimally processed and not subjected to heat treatments compared to juices that are generally**
66 **pasteurized this can favor the nutrients bioavailability and the presence of bioactive compounds**
67 **(Bestwick et al., 2020)**. Furthermore, mixing multiple ingredients represents a good strategy to
68 incorporate non-traditional and under-utilized vegetables, such as beet leaves and stems, increasing
69 the added value of such products (Jayachandran, Chakraborty, & Rao, 2015). The growing variety of
70 products launched on the market has amplified the issues concerning their safety and stability. Many
71 microorganisms have been associated to these typologies of soft drinks as consequence of
72 environmental or raw material contaminations (Siroli, Camprini, Pisano, Patrignani, & Lanciotti,
73 2019). In fact, smoothies are a suitable substrate for the growth of yeasts and lactic acid bacteria and
74 if the final pH is not low enough, smoothies can favour the persistence of pathogenic microorganisms,

75 for this reason are increasingly associated to foodborne illnesses (Bevilacqua et al. 2018). Moreover,
76 the overall quality of these products must remain unchanged over the whole shelf life (Palgan et al.
77 2012). Therefore, currently are being sought technological solutions aimed to guarantee the safety
78 and stability of the product while preserving its nutritional and functional characteristics and
79 consequently reducing the thermal damage associated with traditional heat treatments (Patrignani &
80 Lanciotti, 2016; Gul, Saricaoglu, Mortas, Atalar, & Yazici, 2017; Yi et al., 2017). Among non-
81 thermal technologies, high pressure homogenization (HPH) is widely reported as a very interesting
82 alternative to traditional heat treatments for the stabilization, **in terms of microbial inactivation and**
83 **quality improvement, of fruit and vegetable beverages including juices based on tomato, apple, carrot,**
84 **mango, orange, kiwi, sugarcane, apricot, and pomegranate (Donsì Esposito, Lenza, Senatore, &**
85 **Ferrari, 2009; Patrignani et al., 2009; Patrignani, Tabanelli, Siroli, Gardini, & Lanciotti, 2013; Zhao**
86 **et al., 2014; Betoret, Calabuig-Jiménez, Patrignani, Lanciotti, & Dalla Rosa, 2017; Patrignani et al.,**
87 **2019; Singh et al., 2022).** Due to the phenomena of cavitation, shear stress, turbulence, and
88 impingement that take place during the food treatment with HPH, a strong antimicrobial activity
89 (Zamora & Guamis, 2015; Patrignani & Lanciotti, 2016), a modulation of some enzyme activities
90 and the maintenance of colour, flavour, and nutritional/functional properties can be observed in the
91 treated matrices (Patrignani et al. 2013; Patrignani & Lanciotti, 2016; Błaszczak, Amarowicz, &
92 Górecki, 2017). **In addition, HPH is considered a green and mild technology with a lower impact on**
93 **the environment, more sustainable, saving energy, time and additional costs since it can be used as a**
94 **continuous processing treatment over a batch process such as high hydrostatic pressure (HHP)**
95 **(Patrignani & Lanciotti, 2016; Singh et al., 2022).** Several studies have reported that the efficacy of
96 **HPH treatments is increased when high-pressure is combined with other sublethal stresses. For**
97 **example, the combination of HPH treatments with H₂O₂ or low pH is reported to increase the**
98 **effectiveness against spore forming bacteria such as *Clostridium* spp. and *Bacillus* spp. (Patrignani &**
99 **Lanciotti, 2016; Chaves-López et al. 2009).** Other authors reported that the combined use of natural
100 **antimicrobial based nanoemulsions and HPH is able to increase safety and shelf-life of apple juice**

101 due to the improved antimicrobial activity against spoiling *Saccharomyces cerevisiae* and
102 *Lactiplantibacillus plantarum* and pathogens *Staphylococcus aureus*, *Listeria monocytogenes* and
103 *Escherichia coli* (Patrignani, Siroli, Braschi & Lanciotti, 2020).

104 Another proposed solution to increase shelf-life and prevent microbial spoilage on food products is
105 represented by active packaging (Mastromatteo, Mastromatteo, Conte, & Del Nobile, 2010).
106 Different types of active packaging, based on the controlled release of the antimicrobials added either
107 in the packaging itself or in additional elements positioned inside the package, are currently studied.
108 The polymeric materials that can be used are different but, among the biodegradable polymers one of
109 the most interesting is the polylactic acid (PLA), produced mainly from renewable agricultural
110 resources, thanks to starch fermentation and the condensation of lactic acid (Krishnamurthy, Demirci,
111 Puri, & Cutter, 2004). Recently, PLA activated with antimicrobial compounds has been proposed in
112 food packaging (Jin & Zhang, 2008; Glicerina et al. 2021). Among antimicrobial molecules employed
113 in active packaging, lysozyme showed interesting properties (Glicerina et al., 2021). In fact, lysozyme
114 is a polypeptide with antimicrobial activity especially against Gram-positive bacteria (Iucci,
115 Patrignani, Vallicelli, Guerzoni, & Lanciotti, 2007) and is one of the most studied natural
116 antimicrobial agents for application in packaging to be used for food preservation (Barbiroli et al.
117 2012; Corradini et al. 2013; Ozer, Uz, Oymaci, & Altinkaya, 2016). It is classified as "generally
118 recognized as safe" (GRAS) by the Food and Drug Administration (FDA) and as a food additive by
119 the European Union (E1105) with bacteriostatic, bacteriolytic and bactericidal activity (Muriel-Galet
120 Talbert, Hernandez-Munoz, Gavara, & Goddard, 2013) currently it is widely used in the wine sector
121 and in the production of Grana Padano (Wu et al. 2019). Recently, Glicerina et al. (2021) showed a
122 strong *in-vitro* antimicrobial activity of PLA biodegradable packaging activated with lysozyme by
123 cold plasma against several spoilage and pathogenic microorganisms associated to the food industry.
124 The authors showed that this innovative packaging was able to prevent the growth of *Listeria*
125 *monocytogenes* in rice based smoothie stored at 4 and 10 °C, allowing a better colour retention during
126 storage compared to the controls.

127 In this context, the aim of this study was to evaluate the effect of an HPH treatment (100 MPa for 6
128 times), alone and in combination with the packaging in polylactic acid (PLA), activated with
129 lysozyme by cold plasma, on the shelf-life of a smoothie consisting of a mixture of apple, carrot and
130 rice based beverage, during storage at 4 °C and 10 °C, at 50% RH, in order to simulate real and
131 accelerated storage conditions.

132

133 **2. Material and methods**

134 **2.1. Smoothie preparation**

135 The smoothie used in this study was made with apple, carrot and rice based beverage. Carrots and
136 apples were purchased from a local market on the same day of the trial, sanitized in a solution of 120
137 ppm of sodium hypochlorite for two minutes and dried with a tissue paper; after cutting, their juice
138 was extracted by a juice extractor (R.G.V. - Juice Art, Como, Italy). To avoid negative phenomena
139 due to enzymatic browning, citric acid (Sigma Aldrich) was added at a concentration of 1 g/L. The
140 rice-based drink used in the smoothie preparation was a commercial Ultra High Temperature (UHT)
141 rice beverage (Alce nero, Bologna, Italy). Carrot extract, apple extract and rice beverage were finally
142 mixed in a 3:1:1 ratio to obtain the final smoothie mixture. The final pH of the smoothie was 5.25.

143

144 **2.2. Selection of the appropriate high-pressure homogenization treatment (HPH)**

145 The prepared carrot/apple/rice milk smoothie was immediately subjected to HPH treatment by using
146 a continuous high-pressure homogenizer PANDA (Gea, Parma, Italy) provided of a thermal
147 exchanger and a PS-type valve. Prior to the use, the device was sterilized according to the
148 manufacture's suggestions: 1% NaOH water solution, hot sterilized water, and finally refrigerated
149 sterilized water. **A total of 3 L of smoothie was prepared and treated by HPH at 100 MPa; after each
150 passage at 100 MPa, an aliquot was collected. In fact, 150 mL were collected in sterile flask
151 respectively after 1, 3, 5 and 6 passes at 100MPa. An aliquot of smoothie was treated at 0.1 MPa
152 (control samples).** Immediately after treatment, samples were microbiologically analysed for

153 mesophilic and psychotropic aerobic bacteria, yeasts, lactic acid bacteria, aerobic spore forming
154 bacteria and total coliforms according to the methodology reported below (2.5).

155

156 **2.3. PLA film activation and pouches realization**

157 The polymeric material and the supporting gel used for this study were respectively polylactic acid
158 (PLA) and polyvinyl alcohol (PVOH). PLA films were purchased by Taghleef Industries, S.p.A (San
159 Giorgio di Nogaro, Italy) while polyvinyl alcohol (PVOH) (MW = 88,000–97,000) was purchased
160 by Sigma–Aldrich (Gallarate, Italy). The active compound was lysozyme from hen egg white
161 (Sigma–Aldrich, Gallarate, Italy).

162 The PLA activation and packaging production were carried out according with the previous study of
163 Glicerina et al. (2021), by using a cold plasma treatment in order to activate one surface of the PLA
164 and promoting the adhesion between PLA and PVOH that forming a coating, entraps enzyme causing
165 its immobilization.

166

167 **2.4. High pressure homogenization (HPH) treatment and packaging**

168 Twelve litres of carrot/apple/rice smoothie were divided into two batches, one was subjected to 0.1
169 MPa treatment (used as control) and the other to 100 MPa for 6 cycles. This HPH treatment was
170 selected after preliminary trials aimed to evaluate the microbiological quality after different HPH
171 treatment cycles, as previous described in section 2.2. Half of each treated smoothie was used to fill
172 the PLA activated pouches and half was filled in PLA not activated pouches (control). A total of 204
173 pouches were obtained. Each pouch was filled with 50 mL of smoothie. The filling of the pouches
174 was performed in sterile conditions by using an automatic pipette, under a laminar flow cabinet, and
175 then sealed off on the upper part by using an automatic sealer according with the previous study of
176 Glicerina et al., (2021). Four different smoothie samples were obtained:

- 177 - Smoothies treated at 0.1 MPa and packed in not activated PLA pouches (C 0.1)
- 178 - Smoothies treated at 0.1 MPa and packed in lysozyme activated PLA pouches (L 0.1)

- 179 - Smoothies treated at 100 MPa for 6 cycles and packed in not activated PLA pouches (C 100)
- 180 - Smoothies treated at 100 MPa for 6 cycles and packed in lysozyme activated PLA pouches (L
- 181 100).

182 Obtained samples were stored at two different temperatures, 4 and 10 °C, in climatic chambers at

183 50% RH, for respectively 24 and 13 days, in order to simulate real and accelerated storage conditions.

184 Smoothies stored at 4 °C were analysed in triplicate at 0, 2, 4, 6, 8, 10, 13, 15 and 24 days for pH and

185 colour determinations and at 0, 4, 8, 13, 15 and 20 days for microbiological ones. At the same time,

186 smoothies stored at 10 °C were analysed in triplicate at 0, 1, 2, 3, 4, 6, 8, 10 and 13 days for pH and

187 colour parameters and at 0, 2, 4, 6, 8, 10 and 13 days for microbiological ones.

188 A total of 240 pouches were obtained.

189 In figure 1 a picture of an obtained pouch has been reported, by way of explanation.

190

191

192 **Analytical determinations**

193 **2.5. Microbiological analysis**

194 The cell loads of natural occurring mesophilic aerobic bacteria, psychotropic aerobic bacteria, lactic

195 acid bacteria, yeasts, aerobic spore forming bacteria and total coliforms were detected by plate

196 counting on Plate Count Agar (PCA) (Oxoid Ltd., Basingstoke, United Kingdom), de Man Rogosa

197 and Sharpe (MRS) (Oxoid Ltd., Basingstoke, United Kingdom), Yeast Extract Peptone Dextrose

198 (YPD) (Oxoid Ltd., Basingstoke, United Kingdom), PCA and Violet Red Bile Agar (VRBA) (Oxoid

199 Ltd., Basingstoke, United Kingdom) according to Siroli et al. (2015). PCA and YPD plates were

200 incubated at 30 °C for 48h. MRS and VRBA plates were incubated at 37 °C for 24h while PCA agar

201 plates for psychotropic bacteria were stored at 10 °C for 7 days. The presence of *L. monocytogenes*,

202 *Salmonella enteritidis*, and *E. coli* in all samples was investigated according to the ISO methods 6579-

203 1:2017 (2017); ISO 11290-1:2017 (2017); and ISO 16649-3:2015 (2015), respectively. **According to**

204 **literature data, *L. monocytogenes*, *Salmonella enteritidis* and *E. coli* must be absent in 25g of product,**

205 while the limit of acceptability of yeasts and mesophilic aerobic bacteria is reported to be 6.0 log
206 CFU/g.

207

208 **2.6. pH and colour**

209 pH was measured immediately after treatments and at each storage time by using a pH-meter Basic
210 20 (Crison Instruments, Barcelona, Spain).

211 Colour of samples treated at 0.1 and 100 MPa and packed in both activated (L) and not activated
212 pouches (C) at 10 and 4 °C, was determined by using a tristimulus spectrophotometer (mod.
213 A60-1010-615 ColorFlex, HunterLab, USA) equipped with a sample holder (12 mm diameter).
214 Colour was measured using the CIE $L^*a^*b^*$ colour space and illuminant D65, and expressed as
215 lightness (L^*), and red index (a^*) values calculated as reported by Mc Guire, (1993).

216

217 **2.7. Statistical analysis**

218 At each time of storage, the data obtained represent the means of three independents replicates.
219 Analysis of variance (ANOVA) and the test of mean comparisons according to Fisher's least
220 significant difference (LSD) with a 0.05 level of significance were applied to find out significant
221 differences among the different samples. The statistical package STSG Statistica for Windows,
222 version 6.0 (Statsoft Inc., Tulsa, USA) was used.

223

224 **3. Results and discussion**

225 **3.1. Effect of HPH treatments on microbiological quality of apple-carrot-rice smoothie**

226 In table 1 the cell loads of mesophilic aerobic bacteria, psychotrophic aerobic bacteria, lactic acid
227 bacteria, yeasts, aerobic spore forming bacteria and total coliforms, immediately after the HPH

228 treatments, are reported. In particular, the effect of the 100 MPa treatment, for a different number of
229 cycles (1, 3, 5 and 6), on the initial microbial population of the considered smoothies was evaluated.
230 As expected, the HPH treatment performed lead to a strong inactivation effect as the number of HPH
231 cycles increased. One HPH cycle at 100 MPa was enough to reduce the cell loads of lactic acid
232 bacteria (LAB) and yeasts below the detection limit. The increase of the number of HPH passes
233 significantly reduced the cell load of total mesophilic and psychotropic aerobic bacteria. One cycle
234 at 100 MPa reduced of more than 1.5 logarithmic cycles the total aerobic population, while 3 and 5
235 100 MPa cycles increased the inactivation to 3.06 and 3.23 logarithmic cycles, respectively. Finally,
236 6 passes at 100 MPa (C 100) reduced the mesophilic and psychotropic population of 3.74 and 3.93
237 logarithmic cycles, respectively. The observed microbial inactivation suggests an additive effect of
238 each HPH cycle but without a linearity in terms of reduction of the microbial load. Generally, **the**
239 **effectiveness of HPH treatments is major against gram-negative bacteria and is also dependent on the**
240 **initial microbial load (Patrignani & Lanciotti, 2016; Patrignani et al., 2019). Regarding** the additive
241 effect of HPH cycles on microbial deactivation, **literature** data are contradictory. In fact, some authors
242 indicate a non-additive behaviour of multiple HPH cycles on microbial population, attributing this
243 trend to the physiological diversity of microbial populations and to the presence of resistant cells from
244 the starting microbiota capable to survive the cycles at the applied pressure (Donsi et al., 2009;
245 Patrignani et al. 2013, Patrignani & Lanciotti, 2016). On the basis of the results obtained, the
246 treatment at 100 MPa for 6 cycles was selected as the most suitable for the further shelf-life trials
247 **because it showed a significant reduction of microbial population (mesophilic and psychotropic**
248 **aerobic bacteria), compared to treatment for 5 cycles.**

249

250 **3.2. Effect of HPH treatments combined or not with active packaging on smoothies** 251 **microbiological growth during storage**

252 The effect of the selected hyperbaric treatment (100 MPa for 6 cycles), combined or not with
253 lysozyme activated PLA packaging, on the microbiological shelf-life of apple, carrot and rice

254 smoothie samples was evaluated. Several microbiological groups associated to the spoilage of this
255 kind of products were considered over storage at 4 and at 10 °C.

256 ***Storage at 10 °C***

257 Microbiological results obtained during storage at 10 °C indicate the mesophilic aerobic bacteria and
258 psychotropic aerobic bacteria as mainly responsible for the microbiological spoilage of the smoothies
259 (Table 2). On the contrary, yeasts and spore forming bacteria were not present at spoilage levels for
260 all the 13 days of storage considered. The pathogenic microorganisms *L. monocytogenes*, *Salmonella*
261 spp., and *E. coli* were never detected in the samples stored at 10 °C, independently on the treatment
262 applied (data not shown).

263 Moreover, the stabilizing treatment performed by HPH (6 steps at 100 MPa) allowed a significant
264 reduction of the initial microbial load of all the microbiological groups considered. In fact, the starting
265 load of yeasts, lactic bacteria and spore forming bacteria was below the detection limit (0.5 log
266 CFU/mL). Only total mesophilic aerobic bacteria and psychotropic ones were present in the HPH
267 treated samples respectively at 1.67 and 1.75 log CFU/mL, but with a cell load significantly lower
268 compared to the C.01 samples (total mesophilic and psychotropic ranged between 5.4 and 5.7 log
269 CFU/mL). LAB, yeasts and aerobic spore-forming bacteria which were present at a level ranging
270 between 1.1 and 1.9 log CFU/mL in untreated samples, were found to be below the detection limit
271 following hyperbaric treatment. On the other hand, HPH technology is widely reported to
272 significantly reduce the spoilage microbiota in fruit juices (Patrignani et al., 2013, 2019; Mesa et al.,
273 2020). For example, Patrignani et al. (2019) observed a reduction in yeast load of kiwi juice by more
274 than 2 logarithmic cycles following an HPH treatment at 200 MPa for two cycles. In addition,
275 Patrignani, Siroli, Braschi, & Lanciotti (2020) showed a decrease of the cell load of *S. cerevisiae*,
276 deliberately inoculated in apple juice, of almost 3 log cycles following an HPH treatment at 200 MPa
277 x 2 cycles.

278 During the storage, mesophilic and psychotropic bacteria quickly overcome the level of 7.0 log
279 CFU/mL within 2 days in smoothie samples not subjected to hyperbaric treatment and packed in not

280 activated pouches (C 0.1) and after 6 days in samples packed in activated pouches (L 0.1). Due to the
281 rapid spoilage, the not HPH treated samples have been analysed until day 4 and day 6 when packed
282 in traditional or active packaging respectively.

283 Regarding the HPH treated samples, mesophilic and psychotropic bacteria showed a slower growth
284 kinetic in samples L 100 packed in activated pouches compared to the ones packed in not activated
285 (C 100). In fact, the cell load of mesophilic bacteria, starting from the second day of storage, resulted
286 significantly lower in L 100 samples respect to the C 100 samples. These differences were further
287 increased in the following storage days and after 3 days, samples L 100 showed a mesophilic load
288 lower of more than 2.0 log cycles compared to the controls (C 100). Also, LAB cell load was affected
289 by the lysozyme activated packaging. On the other hand, lysozyme is an enzyme active mainly against
290 Gram-positive bacteria because of its specific activity is to lyse the cell walls by hydrolyzing the
291 linkage β (1-4) between the monomers of peptidoglycan (Iucci et al. 2007).

292 In fact, the samples packed in activated pouches (L 100), starting from the sixth day of storage,
293 presented a significantly lower cell load compared to the C 100 samples. Yeasts and spore forming
294 bacteria showed limited cell loads independently on the sample. However, in the case of yeasts, in
295 samples L 100, they resulted below the detection limit (0.5 log CFU/mL) for the whole period of
296 storage. On the contrary, starting from the eighth day of storage, the yeasts were found in the C 100
297 samples highlighting how the HPH treatment applied induced sub-lethal damages on the yeast
298 population. In fact, as a result of HPH treatment part of the most resistant microbial population can
299 be viable but not culturable (VBNC) and recovers their metabolic activity under favourable
300 environmental conditions (Patrignani et al. 2019). Finally, no significant differences were found
301 between the samples regarding spore forming bacteria that showed cell loads below 2.0 log CFU/mL
302 for the whole period of storage independently to the sample considered.

303 As reported in literature, in vegetable matrices the threshold of total aerobic mesophilic bacteria
304 which leads to spoilage of the product is generally considered 6.0 log CFU/g (Stannard, 1997; FSA
305 of Ireland, 2016). As previous stated, in these trials, the not HPH treated samples (L 0.1 and C 0.1)

306 exceeded this limit after 2 days of storage at 10 °C, independently on the packaging type. However,
307 L 0.1 samples resulted borderline showing a total mesophilic cell load of 6.11 log CFU/mL while C
308 0.1 samples showed a cell load of 7.46 log CFU/mL. Contrarily, in HPH treated samples, the
309 packaging showed a higher effect, since the control samples (C100) exceeded the threshold limit of
310 total aerobic mesophilic bacteria after 8 days of storage while the samples in active pouches, at the
311 eighth day of storage still had a relatively low mesophilic load (3.49 log CFU/mL). In activated and
312 treated pouches the limit of 6.0 log CFU/mL was exceeded only at the thirteenth day of storage. This
313 represents a very interesting result, indicating the possibility of increasing the shelf-life of vegetable
314 smoothies by combining active packaging and HPH technologies and preventing the negative effects
315 of thermal abuse during the refrigerated storage. Other authors reported a microbial stabilization of
316 mango, orange and apple juices after HPH treatments at 200-300 MPa for 1-3 cycles with inlet
317 temperatures ranging between 20 and 60 °C in products stored at 12-20 °C up to 28 days (Suárez-
318 Jacobo et al. 2010; McKay, Linton, Stirling, Mackle, & Patterson, 2011; Guan et al. 2016). However,
319 all the reported authors have considered fruit juices with an extremely acidic pH ranging between 3.2
320 and 4.1, that, as known, represents a very strong hurdle for microbial growth. In this context, the
321 microbial stabilization of a vegetable smoothie characterized by higher pH values as in the present
322 study (5.35 for C100 and L 100 respectively at 8 and 13 days) represents a promising result. The use
323 of the lysozyme activated packaging allowed an increase of 5 shelf-life days of the smoothie. On the
324 other hand, lysozyme is effective against different Gram-positive bacteria including spoiling lactic
325 acid bacteria in wine and beer fermentation (Silvetti, 2010; Azzolini, Tosi, Lorenzini, Finato, &
326 Zapparoli, 2015). In addition, Conte, Buonocore, Bevilacqua, Sinigaglia, & Del Nobile (2006)
327 showed that lysozyme immobilized on polyvinyl alcohol-based film was able to inhibit the growth of
328 viable vegetative cells of *Alicyclobacillus acidoterrestris* including spores. Glicerina et al. (2021),
329 showed that PLA biodegradable packaging activated with lysozyme was able to inhibit several
330 spoiling and pathogenic microorganisms both *in-vitro* and in real systems and was characterized by
331 a gradual release of lysozyme into a milk-based smoothie during refrigerated storage at 10 °C.

332 In addition, it is widely reported that microbial cells may be only partially damaged by HPH
333 treatment, resulting viable but not culturable, and can easily recover their metabolic activity during
334 the storage in particular in a nutrient-rich medium (Patrignani et al. 2009; Patrignani & Lanciotti,
335 2016; Patrignani et al. 2019). The combination of the HPH treatment with an active packaging that
336 slowly releases lysozyme during the storage, may increase the antimicrobial activity of the peptide
337 since the partially damaged microbial cells are more sensitive; the result is an effective control of
338 microbial spoilage on the product during storage. On the other hand, extensive literature reports that
339 the effects of combined stresses (chemical and physical) on microbial growth and survival may be
340 synergistic (Patrignani & Lanciotti, 2016). In fact, even if the antimicrobial efficacy of lysozyme is
341 almost exclusively against gram-positive bacteria, the damage of the cell wall of gram-negative
342 bacteria caused by HPH treatment results in an increase in the antimicrobial activity of lysozyme also
343 against these last microorganisms.

344

345 *Storage at 4 °C*

346 In tables 3 the cell loads of total mesophilic aerobic bacteria and psychotropic aerobic bacteria of
347 samples stored at 4 °C are reported in relation to the treatment applied.

348 Microbiological results of smoothie samples stored at 4 °C highlighted the good efficacy of the HPH
349 pre-treatment to stabilize the initial vegetable matrix. In fact, with the exception of mesophilic and
350 psychotropic aerobic bacteria, present in C 100 and L 100 samples at a very low level (1.67 log
351 CFU/mL and 1.75 CFU mL), all the other microbiological groups (lactic acid bacteria, yeasts, total
352 coliforms and aerobic spore forming bacteria) were below the detection limit (data not shown). The
353 not HPH treated samples reached the spoilage threshold within 8 days when packed in not activated
354 pouches (C 0.1) and 13 days in activated pouches (L 0.1). On the contrary, HPH treated samples
355 stored at 4 °C never exceeded the level of 3.0 log CFU/mL for total mesophilic and psychotropic
356 bacteria over the 20th day of storage at 4 °C.

357 At the storage temperature of 4 °C, no particular differences were found between the HPH treated
358 control samples and those packed in active pouches. In both cases, the microbiological groups were
359 present at very low levels throughout the whole period of storage considered.

360 In HPH treated samples, the mesophilic aerobic bacteria did not reach the spoilage threshold of the
361 product and never exceed the load of 2.7 log CFU/mL for all the storage periods. The pathogenic
362 microorganisms *L. monocytogenes*, *Salmonella* spp., and *E. coli* were never detected in the samples
363 stored at 4 °C independently to the treatment applied (data not shown).

364 The obtained results suggest that the HPH treatment may provide a microbial stabilization of
365 vegetable smoothies at 4 °C. A comparable effect was observed by McKay et al. (2011) that reported
366 a total aerobic count ranging between 2 and 3 log CFU/mL in HPH treated apple juice at 300 MPa
367 for 1 cycle during a storage of 35 days at 4 °C. Other authors reported a microbial stabilization of
368 apple juice treated at 200-300 MPa for 1-3 cycles and stored at 4 °C up to 28-35 days (Suárez-Jacobo
369 et al. 2010; Maresca, Donsì, & Ferrari, 2011). Guan et al. (2016) showed that HPH treatments at 190
370 MPa for 1 and 3 cycles at an inlet temperature of 60 °C provided a long shelf-life up to 60 days of
371 mango juice stored at 4 °C and no microorganisms increased during storage. Also, Patrignani et al.
372 (2019), showed that the application of a treatment at 200 MPa for 3 cycles allowed to obtain a stable
373 kiwifruit juice for more than 40 days under refrigerated storage and to extend the shelf-life of 1 week
374 at room temperature with respect to the control, increasing at the same time the polyphenols
375 availability and its antioxidant activity, and allowing to better retain the colour.

376

377 **3.3. Effect of HPH treatments combined or not with active packaging on smoothies' pH and** 378 **colour during storage**

379 The pH values of smoothies treated with HPH and stored at 4 °C were 5.08±0.03 for both C and L
380 samples and did not change during storage. On the contrary, in not HPH treated samples a rapid
381 acidification was observed during the storage at 10 °C. In fact, the pH resulted below 5.0 after 2 and
382 4 days in samples packed in not activated and activated pouches respectively. In the not HPH treated

383 smoothies stored at 4 °C there was a pH decrease below 5.0 after 4 and 8 days of storage in C 0.1 and
384 L 0.1 samples, respectively.

385 As for colour of smoothies stored at 10 °C, it is possible to highlight that both C samples presented a
386 greater lightness decrease (Figure 2a) and red index increase (Figure 2b), than samples packed in
387 activated pouches, during all storage times, despite of the HPH treatment. This means that the colour
388 of smoothie samples packed in not activated pouches showed a greater and faster darkening during
389 storage, probably as a result of browning reactions due to oxidation phenomena of polyphenols,
390 carotenoids and other phytochemicals present in carrot and apple juice (Riganakos, Karabagias,
391 Gertzou, & Stahl, 2017). In both L samples, independently by HPH treatment, a general lower
392 browning was observed during storage compared to C ones. According with literature, these results
393 can be explained considering that the lysozyme tend to bind with phenolic compounds present into
394 the smoothie that probably, protecting phenols from oxidation, reduce browning phenomena (Green,
395 1995; Bartowsky, Costello, Villa, & Hensche, 2004; Rawduken, Suthiluk, Kamhangwong, &
396 Benjakul, 2012; Liburdi, Benucci & Esti, 2014). Moreover, in all four samples an increase in the lightness
397 was observed after 3 days of storage subsequently to its reduction, probably attributable to the presence of
398 unstable, suspended particles in juice that tends to settle during storage (Cortes et al., 2008; Cao et al., 2012).
399 Figure 3 (a, b) present the changes in lightness (L^*) and red index (a^*) colour values of all smoothie
400 samples during storage at 4 °C. Also, in this case, it was observed a similar trend as for samples stored
401 at 10 °C. Both L samples, packed in activated pouches and pre-treated at 0.1 and 100 MPa, presented
402 respectively higher lightness and lower red index values than C ones, corresponding to a general
403 greater colour maintenance. However, the colour values between L and C samples, mainly the L^*
404 ones, did not always differ significantly during storage.

405

406 **4. Conclusions**

407 The results obtained in the present research highlighted the potential of HPH treatment combined to
408 a lysozyme active biodegradable packaging for the stabilization of a vegetable smoothie characterized

409 by a low-acidic pH (5.25). In fact, the applied HPH treatments allowed to reduce the initial microbial
410 load of the smoothie thus slowing down the **deterioration** of the product stored at both 4 and 10 °C.
411 **The increased antimicrobial activity observed by the active packaging is related to the initial reduction**
412 **of the load of microbial population that was more sensitive to the released lysozyme. The combination**
413 **of these two technologies allowed** the smoothie to reach a shelf-life of 13 days at 10 °C, compared to
414 the only HPH treated product that showed a shelf-life of 8 days. Moreover, the HPH treated smoothies
415 stored at 4 °C remained microbiologically stable for more than 20 days, independently from the
416 packaging. However, the use of active packaging has also allowed to reduce the microbiological
417 proliferation. In addition, L samples packed in active packaging (both HPH treated and not) showed
418 a general better color retention, compared to C ones. In conclusion, the combination of the
419 technologies used in this experiment could be an innovative strategy for increasing the safety and
420 shelf-life of low-acid innovative drinks; the use of lysozyme active packaging could be also a solution
421 to prevent the negative effects of thermal abuse during the refrigerated storage. However, further
422 analyses are necessary to verify the perception of the consumer towards this type of minimally
423 processed product and to quantify its nutritional and functional benefit.

424

425 **Acknowledgments**

426 The present work is part of the research activity developed in the project “ECOPACKLAB - 412
427 ADVANCED TECHNOLOGIES FOR ACTIVE AND ECO-FRIENDLY PACKAGING” 413
428 financed by POR FESR 2014-2020 Region Emilia Romagna, Axis 1, Activity 1.2.2.

429

430 **Declaration of Competing Interest**

431 The authors report no declarations of interest.

432

433

434

435

436

437

438

439 **References**

440

441 Andrés, V., Villanueva, M. J., & Tenorio, M. D. (2016). Influence of high pressure processing on
442 microbial shelf life, sensory profile, soluble sugars, organic acids, and mineral content of milk-and
443 soy-smoothies. *LWT-Food Science and Technology*, 65, 98-105. [https://doi:
444 10.1016/j.lwt.2015.07.066.](https://doi.org/10.1016/j.lwt.2015.07.066)

445

446 Azzolini, M., Tosi, E., Lorenzini, M., Finato, F., & Zapparoli, G. (2015). Contribution to the aroma
447 of white wines by controlled *Torulasporea delbrueckii* cultures in association with *Saccharomyces*
448 *cerevisiae*. *World Journal of Microbiology and Biotechnology*, 31(2), 277-293. [https://doi:
449 10.1007/s11274-014-1774-1.](https://doi.org/10.1007/s11274-014-1774-1)

450

451 Barbiroli, A., Bonomi, F., Capretti, G., Iametti, S., Manzoni, M., Piergiovanni, L., & Rollini, M.
452 (2012). Antimicrobial activity of lysozyme and lactoferrin incorporated in cellulose-based food
453 packaging. *Food Control*, 26(2), 387-392. <https://doi.org/10.1016/j.foodcont.2012.01.046>

454

455 Bartowsky, B. E. J., Costello, P. J., Villa, A., & Henschel, P. A. (2004). The chemical and sensorial
456 effects of lysozyme addition to red and white wines over six months' cellar storage. *Australian*

457 *Journal of Grape and Wine Research*, 10(2), 143-150. <https://doi.org/10.1111/j.1755->
458 [0238.2004.tb00017.x](https://doi.org/10.1111/j.1755-0238.2004.tb00017.x)

459

460 Bestwick, C., Scobbie, L., Milne, L., Duncan, G., Cantlay, L., & Russell, W. (2020). Fruit-based
461 beverages contain a wide range of phytochemicals and intervention targets should account for the
462 individual compounds present and their availability. *Foods*, 9(7), 891. doi: 10.3390/foods9070891.

463

464 Betoret, E., Calabuig-Jiménez, L., A., Patrignani, F., Lanciotti, R., & Dalla Rosa, M. (2017). Effect
465 of high pressure processing and trehalose addition on functional properties of mandarin juice enriched
466 with probiotic microorganisms. *LWT-Food Science and Technology*, 85, 418-422. [https://doi:](https://doi:10.1016/j.lwt.2016.10.036)
467 [10.1016/j.lwt.2016.10.036](https://doi:10.1016/j.lwt.2016.10.036)

468

469 Bevilacqua, A., Petruzzi, L., Perricone, M., Speranza, B., Campaniello, D., Sinigaglia, M., & Corbo,
470 M. R. (2018). Nonthermal technologies for fruit and vegetable juices and beverages: overview and
471 advances. *Comprehensive Reviews in Food Science and Food Safety*, 17(1), 2-62. [https:// doi:](https://doi:10.1111/1541-4337.12299)
472 [10.1111/1541-4337.12299](https://doi:10.1111/1541-4337.12299)

473

474 Błaszczak, W., Amarowicz, R., & Górecki, A. R. (2017). Antioxidant capacity, phenolic composition
475 and microbial stability of aronia juice subjected to high hydrostatic pressure processing. *Innovative*
476 *Food Science & Emerging Technologies*, 39, 141-147. <https://doi.org/10.1016/j.ifset.2016.12.005>
477 rights and content

478

479 Cao, X., Bi, X., Huang, W., Wu, J., Hu, X., & Liao, X. (2012). Changes of quality of high hydrostatic
480 pressure processed cloudy and clear strawberry juices during storage. *Innovative Food Science &
481 Emerging Technologies, 16*, 181-190.

482

483 Conte, A., Buonocore, G. G., Bevilacqua, A., Sinigaglia, M., & Del Nobile, M. A. (2006).
484 Immobilization of lysozyme on polyvinylalcohol films for active packaging applications. *Journal of
485 Food Protection, 69*(4), 866-870. [https:// doi: 10.4315/0362-028x-69.4.866](https://doi.org/10.4315/0362-028x-69.4.866)

486

487 Corradini, C., Alfieri, I., Cavazza, A., Lantano, C., Lorenzi, A., Zucchetto, N., & Montenero, A.
488 (2013). Antimicrobial films containing lysozyme for active packaging obtained by sol-gel technique.
489 *Journal of Food Engineering, 119*(3), 580-587. [https://doi: 10.1016/j.jfoodeng.2013.05.046](https://doi.org/10.1016/j.jfoodeng.2013.05.046)

490

491 Cortés, C., Esteve, M. J., & Frígola, A. (2008). Color of orange juice treated by high intensity pulsed
492 electric fields during refrigerated storage and comparison with pasteurized juice. *Food control, 19*(2),
493 151-158.

494

495 Donsì, F., Esposito, L., Lenza, E., Senatore, B., & Ferrari, G. (2009). Production of shelf-stable
496 annurca apple juice with pulp by high pressure homogenization. *International Journal of Food
497 Engineering, 5*(4). [https://doi: 10.2202/1556-3758.1602](https://doi.org/10.2202/1556-3758.1602)

498

499 Food Safety Authority of Ireland. (FSA) (2016). Monitoring & Surveillance Series, Chemical. Report
500 on a total diet study carried out by the food safety authority of Ireland in the period 2012-2014.
501 Dublin.

502

503 Glicerina, V., Siroli, L., Canali, G., Chinnici, F., Capelli, F., Lanciotti, R., Colombo, V., & Romani,
504 S. (2021). Efficacy of biodegradable, antimicrobial packaging on safety and quality parameters
505 maintenance of a pear juice and rice milk-based smoothie product. *Food Control*, *128*, 108170.
506 <https://doi.org/10.1016/j.foodcont.2021.108170>

507

508 Green, J. L. (1995). The use of lysozyme in winemaking: the interaction of lysozyme with wine and
509 efficacy in preventing malolactic fermentation in Oregon Pinot noir and Chardonnay. Oregon State
510 University.

511

512 Guan, Y., Zhou, L., Bi, J., Yi, J., Liu, X., Chen, Q., Wu, X., & Zhou, M. (2016). Change of microbial
513 and quality attributes of mango juice treated by high pressure homogenization combined with
514 moderate inlet temperatures during storage. *Innovative Food Science and Emerging Technologies*,
515 *36*, 320-29. <https://dx.doi.org/10.1016/j.ifset.2016.07.009>

516

517 Gul, O., Saricaoglu, F. T., Mortas, M., Atalar, I., & Yazici, F. (2017). Effect of high pressure
518 homogenization (HPH) on microstructure and rheological properties of hazelnut milk. *Innovative
519 Food Science and Emerging Technologies*, *41*, 411–420. doi: 10.1016/j.ifset.2017.05.002

520

521 Iucci, L., Patrignani, F., Vallicelli, M., Guerzoni, M. E., & Lanciotti, R. (2007). Effects of high
522 pressure homogenization on the activity of lysozyme and lactoferrin against *Listeria monocytogenes*.
523 *Food Control*, *18*, 558–565. <https://doi: 10.1016/j.foodcont.2006.01.005>

524

525 Jayachandran, L. E., Chakraborty, S., & Rao, P. S. (2015). Effect of high pressure processing on
526 physicochemical properties and bioactive compounds in litchi based mixed fruit beverage. *Innovative*
527 *Food Science & Emerging Technologies*, 28, 1-9. <https://doi.org/10.1016/j.ifset.2015.01.002>

528

529 Jin, T., & Zhang, H. (2008). Biodegradable polylactic acid polymer with nisin for use in antimicrobial
530 food packaging. *Journal of Food Science*, 73(3), M127-M134. [https://doi: 10.1111/j.1750-](https://doi:10.1111/j.1750-3841.2008.00681.x)
531 [3841.2008.00681.x](https://doi:10.1111/j.1750-3841.2008.00681.x)

532

533 Krishnamurthy, K., Demirci, A., Puri, V. M., & Cutter, C. N. (2004). Effect of packaging materials
534 on inactivation of pathogenic microorganisms on meat during irradiation. *Transactions of the ASAE*,
535 *47(4)*, 1141. [https://doi: 10.13031/2013.16546](https://doi:10.13031/2013.16546)

536

537 Kumar, S., Thippareddi, H., Subbiah, J., Zivanovic, S., Davidson, P. M., & Harte, F. (2009).
538 Inactivation of Escherichia coli K-12 in apple juice using combination of high-pressure
539 homogenization and chitosan. *Journal of Food Science*, 74(1), M8-M14.
540 <https://doi.org/10.1111/j.1750-3841.2008.00974.x>

541

542 Liburdi, K., Benucci, I., & Esti, M. (2014). Lysozyme in wine: an overview of current and future
543 applications. *Comprehensive Reviews in Food Science and Food Safety*, 13(5), 1062-1073.
544 <https://doi.org/10.1111/1541-4337.12102>

545

546 Maresca, P., Donsì, F., & Ferrari, G. (2011). Application of a multi-pass high-pressure
547 homogenization treatment for the pasteurization of fruit juices. *Journal of Food Engineering*, *104*(3),
548 364–372. [https://doi: 10.1016/j.jfoodeng.2010.12.030](https://doi.org/10.1016/j.jfoodeng.2010.12.030)

549

550 Mastromatteo, M., Mastromatteo, M., Conte, A. & Del Nobile, M.A., 2010. Advances in controlled
551 release devices for food packaging applications. *Trends in Food Science & Technology*, *21*(12),
552 pp.591-598. <https://doi.org/10.1016/j.tifs.2010.07.010>

553

554 McGuire, R. G. (1992). Reporting of objective color measurements. *HortScience*, *27*(12), 1254-1255.
555 <https://doi.org/10.21273/HORTSCI.27.12.1254>

556

557 Mesa, J., Hinestroza-Córdoba, L. I., Barrera, C., Seguí, L., Betoret, E., & Betoret, N. (2020). High
558 homogenization pressures to improve food quality, functionality and sustainability. *Molecules*, *25*,
559 3305. [https://doi: 10.3390/molecules25143305](https://doi.org/10.3390/molecules25143305)

560

561 McKay, A. M., Linton, M., Stirling, J., Mackle, A., & Patterson, M. F. (2011). A comparative study
562 of changes in the microbiota of apple juice treated by high hydrostatic pressure (HHP) or high
563 pressure homogenisation (HPH). *Food Microbiology*, *28*(8), 1426–1431.
564 <https://doi.org/10.1016/j.fm.2011.06.013>

565

566 Muriel-Galet, V., Talbert, J. N., Hernandez-Munoz, P., Gavara, R., & Goddard, J. M. (2013).
567 Covalent immobilization of lysozyme on ethylene vinyl alcohol films for nonmigrating antimicrobial

568 packaging applications. *Journal of Agricultural and Food Chemistry*, 61(27), 6720-6727. [https://doi:](https://doi:10.1021/jf401818u)
569 10.1021/jf401818u

570

571 Nieva, S. G., Jagus, R. J., Agüero, M. V., & Fernandez, M. V. (2022). Fruit and vegetable smoothies
572 preservation with natural antimicrobials for the assurance of safety and quality. *LWT-Food Science*
573 *and Technology*, 154, 112663. <https://doi.org/10.1016/j.lwt.2021.112663>

574

575 Nunes, M. A., Barreira, J. C. M., Costa, A. S. G., Vinha, A. F., Alves, R. C., Rocha, A., & Oliveira,
576 M. B. P. P. (2016). Influence of packing material on the quality parameters and bioactivity of
577 smoothies. Livro de Atas do XIII Encontro de Química dos Alimentos, 170-173, Porto (Lisbon).

578

579 Ozer, B. B. P., Uz, M., Oymaci, P., & Altinkaya, S. A. (2016). Development of a novel strategy for
580 controlled release of lysozyme from whey protein isolate based active food packaging films. *Food*
581 *Hydrocolloids*, 61, 877-886. <https://doi.org/10.1016/j.foodhyd.2016.07.001>

582

583 Palgan, I., Muñoz, A., Noci, F., Whyte, P., Morgan, D. J., Cronin, D. A., & Lyng, J. G. (2012).
584 Effectiveness of combined Pulsed Electric Field (PEF) and Manothermosonication (MTS) for the
585 control of *Listeria innocua* in a smoothie type beverage. *Food Control*, 25(2), 621-625.
586 <https://doi.org/10.1016/j.foodcont.2011.11.009>

587

588 Patrignani, F., Vannini, L., Kamdem, S. L. S., Lanciotti, R., & Guerzoni, M. E. (2009). Effect of high
589 pressure homogenization on *Saccharomyces cerevisiae* inactivation and physico-chemical features in

590 apricot and carrot juices. *International Journal of Food Microbiology*, 136(1), 26-31. [https://doi:](https://doi.org/10.1016/j.ijfoodmicro.2009.09.021)
591 10.1016/j.ijfoodmicro.2009.09.021

592

593 Patrignani, F., Tabanelli, G., Siroli, L., Gardini, F., & Lanciotti, R. (2013). Combined effects of high
594 pressure homogenization treatment and citral on microbiological quality of apricot juice.
595 *International Journal of Food Microbiology*, 160(3), 273-281.
596 <https://doi.org/10.1016/j.ijfoodmicro.2012.10.021>

597

598 Patrignani, F., & Lanciotti, R. (2016). Applications of high and ultra high pressure homogenization
599 for food safety. *Frontiers in Microbiology*, 7, 1132. [https://doi: 10.3389/fmicb.2016.01132](https://doi.org/10.3389/fmicb.2016.01132)

600

601 **Patrignani, F., Siroli, L., Braschi, G., & Lanciotti, R. (2020). Combined use of natural antimicrobial**
602 **based nanoemulsions and ultra high pressure homogenization to increase safety and shelf-life of apple**
603 **juice. *Food Control*, 111, 107051. <https://doi.org/10.1016/j.foodcont.2019.107051>**

604

605 Patrignani, F., Mannozi, C., Tappi, S., Tylewicz, U., Pasini, F., Castellone, V., Riciputi, Y., Rocculi,
606 P., Romani, S., Caboni, M.F., Gardini, F., Lanciotti, R., & Dalla Rosa, M. (2019). (Ultra) High
607 pressure homogenization potential on the shelf-life and functionality of kiwifruit juice. *Frontiers in*
608 *Microbiology*, 10, 246. <https://doi.org/10.3389/fmicb.2019.00246>

609

610 Patrignani, F., Siroli, L., Braschi, G., & Lanciotti, R. (2020). Combined use of natural antimicrobial
611 based nanoemulsions and ultra high pressure homogenization to increase safety and shelf-life of apple
612 juice. *Food Control*, 111, Article 107051. <https://doi.org/10.1016/j.foodcont.2019.107051>

613

614 Rawdkuen, S., Suthiluk, P., Kamhangwong, D., & Benjakul, S. (2012). Mechanical, physico-
615 chemical, and antimicrobial properties of gelatin-based film incorporated with catechin-
616 lysozyme. *Chemistry Central Journal*, 6(1), 1-10. [https://doi: 10.1186/1752-153X-6-131](https://doi.org/10.1186/1752-153X-6-131)

617

618 Riganakos, K. A., Karabagias, I. K., Gertzou, I., & Stahl, M. (2017). Comparison of UV-C and
619 thermal treatments for the preservation of carrot juice. *Innovative Food Science & Emerging
620 Technologies*, 42, 165-172. <https://doi.org/10.1016/j.ifset.2017.06.015>

621

622 Silvetti, T. (2010). Development of innovative techniques for studying microbial population in milk
623 and dairy products. Doctoral Thesis.

624

625 Singh, S. V., Singh, R., Singh, A., Chinchkar, A. V., Kamble, M. G., Dutta, S. J., & Singh, S. B.
626 (2022). A review on green pressure processing of fruit juices using microfluidization: Quality, safety
627 and preservation. *Applied Food Research*, 2, 100235. <https://doi.org/10.1016/j.afres.2022.100235>

628

629 Siroli, L., Camprini, L., Pisano, M. B., Patrignani, F., & Lanciotti, R. (2019). Volatile molecule
630 profiles and anti-*Listeria monocytogenes* activity of nisin producers *Lactococcus lactis* strains in
631 vegetable drinks. *Frontiers in Microbiology*, 10, 563. <https://doi.org/10.3389/fmicb.2019.00563>

632

633 Stannard, C. (1997). Development and use of microbiological criteria for foods. *Food Science and
634 Technology Today*, 11(3), 137-177. [https:// doi: 10.2903/j.efsa.2017.5052](https://doi.org/10.2903/j.efsa.2017.5052)

635

636 Suárez-Jacobo, A., Gervilla, R., Guamis, B., Roig-Sagués, A. X., & Saldo, J. (2010). Effect of UHPH
637 on indigenous microbiota of apple juice: A preliminary study of microbial shelf-life. *International*
638 *Journal of Food Microbiology*, 136(3), 261–267. [https://doi: 10.1016/j.ijfoodmicro.2009.11.011](https://doi.org/10.1016/j.ijfoodmicro.2009.11.011)

639

640 Wu, T., Jiang, Q., Wu, D., Hu, Y., Chen, S., Ding, T., Ye, X., Liu, D., & Chen, J. (2019). What is
641 new in lysozyme research and its application in food industry? A review. *Food Chemistry*, 274, 698-
642 709. <https://doi.org/10.1016/j.foodchem.2018.09.017>

643 Yi, J., Kebede, B., Kristiani, K., Grauwet, T., Van Loey, A., & Hendrickx, M. (2017). Minimizing
644 quality changes of cloudy apple juice: the use of kiwifruit puree and high pressure homogenization.
645 *Food Chemistry*, 249, 202–212. <https://doi:10.1016/j.foodchem.2017.12.088>

646

647 Zamora, A. and Guamis, B., 2015. Opportunities for ultra-high-pressure homogenisation (UHPH) for
648 the food industry. *Food Engineering Reviews*, 7(2), pp.130-142. [https:// doi 10.1007/s12393-014-](https://doi.org/10.1007/s12393-014-9097-4)
649 [9097-4](https://doi.org/10.1007/s12393-014-9097-4)

650

651 Zhao, L., Wang, Y., Wang, S., Li, H., Huang, W., and Liao, X. (2014). Inactivation of naturally
652 occurring microbiota in cucumber juice by pressure treatment. *International Journal of Food*
653 *Microbiology*, 174, 12–18. [https://doi: 10.1016/j.ijfoodmicro.2013.12.023](https://doi.org/10.1016/j.ijfoodmicro.2013.12.023)

654

655 Zhou, C. L., Liu, W., Zhao, J., Yuan, C., Song, Y., Chen, D., Ni & Li, Q. H. (2014). The effect of
656 high hydrostatic pressure on the microbiological quality and physical–chemical characteristics of

657 Pumpkin (*Cucurbita maxima* Duch.) during refrigerated storage. *Innovative food science & Emerging*
658 *Technologies*, 21, 24-34. <https://doi.org/10.1016/j.ifset.2013.11.002>