

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Leiomyosarcoma of the Larynx: A Complex Diagnosis

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Molteni G., Soloperto D., Fabbris C., Gazzini L., De Rossi S., Valotto G., et al. (2020). Leiomyosarcoma of the Larynx: A Complex Diagnosis. *EAR, NOSE & THROAT JOURNAL*, 99(5), 300-302 [10.1177/0145561319840204].

Availability:

This version is available at: <https://hdl.handle.net/11585/944719> since: 2023-10-12

Published:

DOI: <http://doi.org/10.1177/0145561319840204>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

LEIOMYOSARCOMA OF THE LARYNX: A COMPLEX DIAGNOSIS.

Gabriele Molteni MD¹, Davide Soloperto MD PhD¹, Cristoforo Fabbris MD¹, Luca Gazzini MD¹, Stefano De Rossi MD¹, Giovanni Valotto MD², Daniele Marchioni MD Prof¹

¹: Otolaryngology Department, University Hospital of Verona, Piazzale Aristide Stefani 1, Verona, 37126, Italy

²: Pathology Department, University Hospital of Verona, Piazzale Aristide Stefani 1, Verona, 37126, Italy

FUNDING AND CONFLICT OF INTERESTS

The authors have no conflict of interest. They have no funding to disclose. The human participant involved in this research signed an informed consent before any further treatment. All procedures performed were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

CORRESPONDING AUTHOR

Cristoforo Fabbris, MD. Otolaryngology Department, University Hospital of Verona, Piazzale Aristide Stefani 1, Verona, Italy. Email: cristoforo.fabbris@gmail.com. Work phone: +39 045 8122330. Fax: +39 045 8122070. ORCID 0000-0002-4476-9802.

KEYWORDS

Leiomyosarcoma, purulent chondritis, laryngeal tumor, laryngectomy.

WORD COUNT: 638 words.

CASE REPORT

An 84-year-old man was referred to our institution for dysphonia present for 2 months. Fiberoptic laryngoscopy demonstrated a polypoid lesion of the right true vocal fold with normal motility.

Excisional biopsy of a submucosal thickening of the right vocal fold with vocal ligament calcification was performed. Histology revealed squamous cell epithelium without malignancy. Post-operative endoscopic follow up was regular.

Six months after surgery, dysphonia and dyspnea occurred. Flexible laryngoscopy demonstrated right hemilaryngeal paralysis, edema and partial airway obstruction. Necrotic tissue and purulent secretions were visible at the anterior commissure, right vocal fold and ipsilateral vestibule. A second microlaryngoscopic surgery included multiple biopsies and debridement of the necrotic tissue. Histological report was negative for malignancy, and purulent chondritis was diagnosed. Intravenous antibiotic therapy was administered with improvement of infection, inflammation, dysphonia, and dyspnea. The patient was discharged.

Two months later, a new sudden and severe dyspnea occurred. He was admitted to the E.R., and oropharyngeal intubation was necessary because of respiratory failure.

CT-scan showed thickening of the right laryngeal wall with wide fluid collection (48-43 mm diameter) that almost completely obstructed the airway, with erosion of the right thyroid and cricoid cartilage (Figure 1).

Debridement of the necrotic tissue and drainage of the purulent collection was performed with direct laryngoscopy. Culture was positive for *Staphylococcus epidermidis* and *Candida albicans*. Intravenous antibiotic, antimicotic and corticosteroid therapy was administered, and other 3 surgical debridements and drainage were performed, but without clinical or

radiological improvements. Moreover, patient developed signs of sepsis. Since no improvement had been achieved, total laryngectomy was planned even without a positive histologic biopsy. Intraoperative findings showed trans-glottic extension of the ulcerated material with full-thickness infiltration of the laryngeal wall and external spread through the thyroid cartilage, with invasion of pre-laryngeal tissues (Figure 2). The patient's clinical condition improved, and he started oral feeding without pharyngo-cutaneous fistula. Morphological and immunohistochemical features were consistent with leiomyosarcoma (FNCLCC Grade 3). (Figure 3).

Multidisciplinary consultation suggested adjuvant radiation therapy (RT), but pre-irradiation staging CT-scan showed rapid local recurrence and lung metastasis. The poor clinical condition of the patient contraindicated 1st line chemotherapy. At 3-month-follow-up, the patient was alive and receiving best supportive care.

DISCUSSION

The first case of laryngeal leiomyosarcoma (LLMS) was reported by Jackson et al. in 1939.[1] After that, fewer than 50 cases of LLMS have been reported in international literature,[2] including an 87-year-old man with laryngopyoceles and LLMS who was treated with total laryngectomy.[3]

Simultaneous presence of laryngeal carcinoma, laryngopyocoele, and laryngomucocoele was described by Cassano et al[4], and Mitro et al[5], but no leiomyosarcoma was found in these patients or in cases reviewed in similar articles.

Purulent chondritis of laryngeal cartilages (PCLC) is a very uncommon abscess that grows between inner and outer perichondria of the laryngeal framework. To date, only 5 cases have been described in the literature[6-8], and none of them was associated with neoplasm.

In our case, radiological suspicion of laryngeal abscess and no clear radiological signs of malignant tumor were detectable. Laryngeal biopsies were negative for malignancy, and cultures were consistent with PCLC, but combined antibiotic therapy, surgical drainage and debridement did not improve the clinical condition, and the patient developed sepsis. Fortunately, after total laryngectomy, his clinical condition improved quickly with complete sepsis resolution.

Etiology of LLMS is still unclear. A possible factor is an abnormal post-operative healing process.[9] In our case, the patient developed worsening dysphonia and dyspnea, six months after his first vocal fold biopsy with negative findings.

Treatment guidelines for LLMS are lacking due to its rarity. First line treatment should be tumor resection with wide margins, reserving radiotherapy as adjuvant treatment or in case of recurrence or persistent tumor.[10,11] Chemotherapy usually is suggested for distant metastases[12].

Even though if LLMS is a very rare laryngeal malignancy, it should be considered when dealing with laryngeal abscess, especially with cartilage involvement.

REFERENCES

- [1]. Jackson C, Jackson CL. Sarcoma of the larynx. In: Jackson C. (editors). Cancer of the larynx. Philadelphia: W. B. Saunders; 1939:164-8.
- [2]. Jagveer Yadav, Jaimanti Bakshi, Mahendra Chouhan, Rahul Modi. Head and Neck Leiomyosarcoma. Indian J Otolaryngol Head Neck Surg. July 2013;65(Suppl 1):S1–5.
- [3]. Helmberger RC, Croker BP, Mancuso AA. Leiomyosarcoma of the Larynx Presenting as a Laryngopyoce. AJNR. Jun 1996;17:1112–4.
- [4]. Cassano L, Lombardo P, Marchese-Ragona R, Pastore A. Laryngopyoce: three new clinical cases and review of the literature. Eur Arch Otorhinolaryngol. 2000;257:507–11.
- [5]. Mitroi M, Căpitănescu A, Popescu FC, Popescu C, Mogoantă CA, Mitroi G, Surlin C. Laryngocele associated with laryngeal carcinoma. Rom J Morphol Embryol. 2011;52(1):183–5.
- [6]. Eliashar R, Gross M, Goldfarb A, Sichel JY. Purulent chondritis of the laryngeal framework cartilages. Ann Otol Rhinal Laryngol. 2005 Mar;114(3):219-22.
- [7]. Lee JK, Cho HH, Kim HD, Lim SC. Purulent chondritis of the thyroid cartilage. Am J Otolaryngol. 2007 Jan-Feb;28(1):64-6.
- [8]. Leiva-Salinas C, Mukherjee S. Purulent chondritis of the cricoid. Eur Ann Otorhinolaryngol Head Neck Dis. 2016 Apr;133(2):137-9.
- [9]. Khadivi E, Taziky MH, Jafarian AH, Nasser Sadr M. Laryngeal leiomyosarcoma, a case report and review of articles. Iran J Otorhinolaryngol. 2013;25:253–8.
- [10]. Abbas A, Ikram M, Yagoob N. Leiomyosarcoma of the larynx: A case report. Ear Nose Throat J. 2005;84(7):435-6.
- [11]. Kleinsasser O, Glanz H. Myogenic tumors of the larynx. Arch Otorhinolaryngol. 1979.225:107-8.

- [12]. Ismi O, Arpaci RB, Puturgeli T, Ozcan C, Gorur K. Leiomyosarcoma of Larynx as a Metachronous Tumor of Squamous Cell Carcinoma: An Unusual Case. J Voice. 2016 Mar;30(2):242-5.

FIGURE LEGENDS

Figure 1. CT image showing an almost complete erosion of the laryngeal framework cartilages due to LLMS. Residual airway is minimal, therefore an emergency oropharyngeal intubation is needed.

Figure 2. Intraoperative picture showing the almost complete erosion of laryngeal cartilages with necrotic material substituting normal laryngeal tissue.

Figure 3. Histopathology of the lesion: SMA (Smooth Muscle specific Actin) positivity demonstrates the smooth muscle nature of the neoplastic population. Written histological report shows a malignant neoplasm with spindle cells, high mitotic index (10-14 mitosis x 10HPF) and large areas of tissue necrosis (>50%).