



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Rapid weight gain predicts fight success in mixed martial arts - evidence from 1,400 weigh-ins

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Faro, H., de Lima-Junior, D., Machado, D.G.d.S. (2023). Rapid weight gain predicts fight success in mixed martial arts - evidence from 1,400 weigh-ins. EUROPEAN JOURNAL OF SPORT SCIENCE, 23(1), 8-17 [10.1080/17461391.2021.2013951].

Availability:

This version is available at: <https://hdl.handle.net/11585/945188> since: 2024-06-20

Published:

DOI: <http://doi.org/10.1080/17461391.2021.2013951>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)



Each pound matters: the influence of weight regain on fight success in mixed martial arts - an analysis of 1,474 weigh-in

Journal:	<i>European Journal of Sports Science</i>
Manuscript ID	TEJS-2020-1010
Manuscript Type:	Original Paper
Keywords:	Body composition, Competition, Performance, Recovery

SCHOLARONE™
Manuscripts

1
2
3 1 **Each pound matters: the influence of weight regain on fight success in mixed**
4
5 2 **martial arts - an analysis of 1,474 weigh-in**
6
7
8
9 3

10
11 4 **Abstract:**
12
13

14 5 We aimed to verify whether the weight regained (%WR) between the official weigh-in
15 6 and the time of the fight is associated with fight success in MMA. A total of 1,474 MMA
16 7 weigh-ins/fights from 21 MMA companies regulated by the California State Athletic
17 8 Commission were analyzed. Logistic regression with the odds ratio (OR) performed using
18 9 the %WR as the independent variable and the fight outcome (win and loss) was used as
19 10 the dependent variable (controlled for sex, division, and weight difference from the
20 11 opponent) with $p < 0.05$. Bantamweight division presents the highest median of %WR
21 12 (9.72%) and one athlete from the lightweight division presented the highest individual
22 13 %WR (20.44%). The %WR was a significant predictor of fight outcome ($\beta = 0.044$;
23 14 $OR = 1.045$; $95\%CI = 1.014 - 1.077$; $p = 0.004$) so that for each 1%WR increased the chance
24 15 of winning by 4.6%. The results suggest that the magnitude of weight regain is linked to
25 16 the chance to win. Considering the regulatory commissions, confederations, and events'
26 17 organizers should consider defining limits for weigh-regain, considering the potential
27 18 advantage with high weight regain found. Otherwise, the athletes might invest in RWL
28 19 and RWR after official weight-in, to increase their chance of winning.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51
52 21 **Keywords:** Martial arts; Weight loss; Weight gain.
53
54
55
56
57
58
59
60

24 INTRODUCTION

25 Mixed Martial Arts (MMA) is a combat sport that uses mixed techniques from
26 different martial arts (12). The combats are divided into three rounds (five rounds for title
27 fights) of five minutes with a one-minute interval. The fight might be interrupted if there
28 is a submission or knockout (5). The victory in MMA is determined by actual or technical
29 knockout (i.e. KO/TKO), submission, or by judges' decision (i.e., by points) (6). MMA
30 gained popularity in the early 1990s when The Ultimate Fighting Championship (UFC)
31 was created (7). The first events lacked weight-divisions, but after rules reformulation,
32 divisions based on weight were created, similarly to other combat sports (6,12). This rule
33 was created to equalize strength between opponents (1,6,11) so that the only difference
34 between them would be their technical skills and/or strategy.

35 Despite the divisions by weight, most athletes choose to fight in a weight-division
36 in which the weight limit is several pounds below their habitual weight (11). To ensure
37 qualification, rapid weight loss (RWL) followed by rapid weight regain (RWR) strategies
38 are adopted in order to get an advantage related to weight, size, and strength (5,11,17).
39 The use of RWL strategies generally begins approximately seven days before the official
40 weigh-in and has the characteristics to be acute and aggressive (7,14). The use of
41 laxatives, diuretics, plastic or rubber clothing, and sauna are the most frequently reported
42 method for RWL in combat sports (11). Specifically, in MMA, the most common
43 strategies self-reported by the fighters are caloric and hydric restriction, sauna, diuretics,
44 clothes that increase sweat production, high volume of low-intensity exercise, and water
45 load (17,19). Moreover, some of these methods are used in combination (17,19). These
46 results indicate that dehydration is the main strategy of RWL. Dehydration decreases
47 blood volume, plasma volume, free testosterone, and blood creatine concentration
48 (1,11,17). In this sense, Jetton et al. (12) compared the hydration status by urinary analysis

1
2
3 49 of 40 MMA fighters at 24h and two hours before the combat. The results showed that
4
5 50 even after ~4.4% of weight regain (WR), 39% of fighters were still significantly
6
7 51 dehydrated and 11% were seriously dehydrated two hours before the combat (12).
8
9 52 Therefore, considering that the RWL strategy may impair performance and negatively
10
11 53 impact the athletes' health, ways of avoiding it are warranted.
12
13
14

15 54 Despite the risk involved in the RWL, several athletes still opt to compete in
16
17 55 divisions below their habitual weight, given that with RWR they may gain advantages
18
19 56 over shorter and lighter fighters on the day of the fight. In this regard, Brechney et al. (3)
20
21 57 investigated if the fight outcome was related to the weight loss magnitude in 75 MMA
22
23 58 fighters and found that athletes who lost the fight lose more weight than the winners
24
25 59 (10.6% vs 8.6%). However, most athletes included in their sample were amateurs (n =
26
27 60 59) (3). Considering that aggressive RWL is repeated throughout one's athletic career,
28
29 61 there is a fear of the consequences of the athletes' health (1). Most importantly, there are
30
31 62 reports of deaths that were caused as the consequence of RWL (7,14). On the other hand,
32
33 63 the magnitude of WR has been suggested to be a decisive factor for fight success in MMA,
34
35 64 more influential than the magnitude of weight loss (6,14). Thus, considering that the
36
37 65 majority of events have the official weigh-in from 24h to 36h before the combat (4), the
38
39 66 athletes consume high quantities of high energy macronutrients (i.e. carbohydrates) and
40
41 67 use intravenous and oral rehydration methods, to recover as much as possible the weight
42
43 68 lost (1,6,7,14).
44
45
46
47
48
49

50 69 MMA athletes lose ~10% of body weight in approximately seven days, with ~5%
51
52 70 in the last 24h before the official weigh-in, to fit the division's weight limit
53
54 71 (2,4,5,7,14,15). Hence, it has suggested that regulatory agencies create ways to inhibit the
55
56 72 RWL process in combat sport for both athlete safety and fairness of sport. However,
57
58 73 whether WR impacts on the outcome of combat is less discussed. Currently, there is no
59
60

1
2
3 74 formal consensus regarding the limit of weight loss or regain in MMA. However, some
4
5 75 commissions have been monitoring the magnitudes of WR in order to create new rules in
6
7 76 this regard. For instance, since 2016 the California State Athletic Commission (CSAC)
8
9 77 recommends that athletes who exceed 10% of WR compete in the division above their
10
11 78 current division (8). In 2019, CSAC approved the new rule stating that a fight must be
12
13 79 canceled if one of the athletes regained >15% of WR (9). Although previous studies
14
15 80 showed that the WR magnitude is a fundamental element for increasing the chances of
16
17 81 winning, few studies verified this fact on a large scale. Thus, we aimed to verify whether
18
19 82 the WR between the official weigh-in and the fight influences the chances of winning.
20
21
22
23
24 83 We hypothesize that greater WR would positively associate with greater chances of
25
26 84 winning.
27
28
29
30
31

85

32 **METHODS**

33 34 35 **Study Design**

36
37
38 88 To test whether the WR between the official weigh-in and the fight influence the
39
40 89 chances of winning the present study analyzed the data of 1,474 weight-ins in professional
41
42 90 MMA events from 20 MMA companies under the California State Athletic Commission
43
44 91 regulation (CSAC). The CSAC regulates, licentiate, and supervise combat sports events
45
46 92 in the California State, USA. The CSAC provided datasheets containing all information
47
48 93 used in the present study. These data were acquired in 21 different locations, from national
49
50 94 and international professional MMA events from January 2015 and August 2019 at the
51
52 95 California state.
53
54
55
56
57
58
59
60

96

97 **Subjects**

1
2
3 98 The report from 1,550 fights was provided, involving both sexes. The events
4
5 99 involved professionals MMA fighters divided into 11 weight divisions (Table 1). To be
6
7 100 included, sufficient information regarding the study variables such as the fight outcome
8
9 101 (win or lose, not draw), weight at weight-in day within the category limit should be
10
11 102 reported. The data of 76 fighters were excluded: 31 did not present the weight at the fight
12
13 103 day, 44 fight outcomes as a draw, one athlete exceeded his/her weight division limit.
14
15 104 Therefore, the final analysis included the data on 1,474 fights (Female = 110; Males =
16
17 105 1,364).

18
19
20
21
22 106 The data were collected from the following MMA companies: BAMMA, Bellator
23
24 107 MMA, Cage Fury Fighting Championship (CFFC), California Extreme Fight (CXF),
25
26 108 California Fight Championship (CFC), California Fight League (CFL), Combat
27
28 109 Americas, Dragon House MMA, Extreme Fighters, In Sync Promotions, Invicta MMA,
29
30 110 King of the Cage (KOTC), Legacy Fighting Alliance (LFA), No Limit/Goodman, PR
31
32 111 Promotions, Roy Englebrecht Events, Smash Global, Tachi Palace Fight, Titans Cage
33
34 112 MMA, TS-1 Combate, and Ultimate Fight Championship (UFC).

35
36
37
38
39 113 The CSAC provided datasheet including the official weight-in date, event dates,
40
41 114 official body weight at the weigh-in, at the fight day, and the difference between them
42
43 115 (i.e., WR), and the difference in weight from the opponent (WD), sex, and divisions of
44
45 116 the athletes. The official weigh-in (OW) was measured between 24-36h before the
46
47 117 combat, while the re-weighing (RW) was performed ~3h before de fight. From these data
48
49 118 was possible calculated the WR ($WR = RW - OW$), and the percentage of WR: [%WR =
50
51 119 $(WR/OW) \times 100$] (12).

52
53
54
55
56 120

57 58 59 121 **Statistical Analyses**

60

1
2
3 122 Descriptive data are presented as median and interquartile range as well as
4
5 123 minimum and maximum values. Logistic regression was performed using as the
6
7 124 independent variables all measures with a theoretical relationship with the dependent
8
9 125 variable, namely %WR, sex, division, and WD from the opponent. The outcome of the
10
11 126 fight (win and loss) was used as the dependent variable. The Odds Ratio (OR) with a 95%
12
13 127 confidence interval (95%CI) was calculated for each independent variable. Accordingly,
14
15 128 the probability of winning was calculated as the value exceeding OR of 1.00 multiplied
16
17 129 by 100, only for variables that achieved a significant level of 5%.

20
21
22 130

23 24 131 **RESULTS**

25
26
27 132

28
29
30 133 Table 1 describes the information about weight categories (separated by sex) and
31
32 134 %WR in each weight category. The highest %WR was found for the Bantamweight
33
34 135 division and the highest individual %WR was from one athlete of the Lightweight
35
36 136 division (20.4%). On the other hand, the heaviest categories (i.e. Heavy and Super
37
38 137 Heavyweight divisions) presented the smallest median %WR.

39
40
41 138

42
43
44 139

45 46 **Insert Table 1**

47
48
49 140

50
51
52 141 Table 2 presents the results of the logistic regression analysis. Only the %WR was
53
54 142 significantly associated with the fight outcome ($p = 0.004$). For each 1% of WR the
55
56 143 probability of winning increased by 4.6%.

57
58
59 144
60

Insert Table 2**DISCUSSION**

This study aimed to verify the influence of weight regain on fight success in MMA, analyzing 1,474 fights in events that occurred between 2015 and 2019 regulated by the CSAC. The main finding was that %WR was a predictor of fight success, independently of sex, division, and WD from the opponent. In fact, for each 1% of WR increased the probability of winning by 4.6%, which confirmed our hypothesis. Previous studies have suggested that the magnitude of WR, but not weight loss, was the main factor for fighters' performance (6,14). However, to the best of the authors' knowledge, this is the first study to investigate the influence of WR on fight success in an ecological and large-scale study with professional MMA athletes.

Our findings corroborate with Artioli et al. (1) statement that the RWL might be an indirect method to gain an advantage in the fight. These authors suggested that this method should be banned in combat sports and listed by World Anti-Doping Agency (WADA) as doping because it might enhance sports performance indirectly, and put athletes' health at risk and violating the spirit of the sport (1). One possible explanation for RWL to enhance sports performance is because it has small or no impact in physical capacities (1). This maintenance of performance has been associated with WR after the official weight-in, with reports that the weight of the athlete in the day of the fight corresponds to three weight divisions above the one he/she is fighting (14,15), which reinforce the idea of violation of the sports spirit. Indeed, the Coswig et al. (6) study concluded that WR was significantly higher in winners compared to losers. Interestingly, WR was positively correlated with time spent at high intensity during the fight, lower

1
2
3 169 limb blows, and ground and pound actions of the athletes. It is noteworthy that WR did
4
5 170 not compensate for the weight loss (6), and only WR but not weight loss was correlated
6
7 171 with measures of performance, which reinforce the idea that WR is more important than
8
9 172 weight loss. Our study corroborates these findings, by showing that the magnitude of WR
10
11 173 has high relevance for fight success in MMA.
12
13
14

15 174 In recent years, RWL has been highly criticized for the dangers it poses to the
16
17 175 athletes. One fighter from Brazil and one from China died during the RWL process (7,14).
18
19 176 Moreover, kidney injury (13) and rhabdomyolysis (16) cases were reported as a
20
21 177 consequence of the RWL process. Despite that, few actions have taken to inhibit RWL
22
23 178 methods. The Asian event ONE Championship banned since 2015 the weight loss by
24
25 179 dehydration and has increased weight limits for each category compared to other events
26
27 180 (14). On the other hand, UFC increased the time between official weigh-in and fight time
28
29 181 to 24h to 36h in 2016, justifying as a protection for the athletes. However, it may have
30
31 182 influenced the athletes to adopt even more aggressive methods for RWL, because the time
32
33 183 window for regain is larger (14). In 2019, UFC together with U.S. Anti-doping Agency
34
35 184 (USADA), prohibited the use of intravenous solutions for WR (20). This is important
36
37 185 because athletes have been using this method to speed up the WR process (7). Whilst all
38
39 186 these measures are important, the lack of control and limit about the magnitude of WR
40
41 187 remains. In this sense, some researchers reported that superior values of WR comparing
42
43 188 weight loss are influenced by the time available for WR (14,15). Thus, a decrease in the
44
45 189 use of aggressive RWL methods could occur only when the time available for WR
46
47 190 decreases considerably or if the official weigh-in is performed very close to the fight (4).
48
49 191 In the “real-world”, athletes still use RWL methods because a high percentage of weight
50
51 192 loss can be regained after the weight-in. In fact, seven out 11 divisions presented a median
52
53
54
55
56
57
58
59
60

1
2
3 193 %WR of >8%. Furthermore, one extreme example was the athlete from the lightweight
4
5 194 division that showed 20.44% WR.
6
7

8 195 In order to restrict both RWL and high WR, some changes to the rules of MMA
9
10 196 competitions should be made. First, the creation of intermediate weight division, between
11
12 197 the existent categories, which would avoid severe RWL, as seen at the present (7). This
13
14 198 is important because the difference of the limit of weight between some divisions reaches
15
16 199 20 pounds (i.e. between middle and light heavyweight divisions) and some athletes could
17
18 200 achieve the weight to be characterized in 2-3 divisions above of the one they are
19
20 201 competing in the fight day (15). Second, the official weigh-in should occur a few hours
21
22 202 before the combat, as it occurs currently in Brazilian Jiu-Jitsu competitions (10), which
23
24 203 prevents high weight loss and high WR. Some amateur events have adopted a similar
25
26 204 approach, weighting the athletes in the morning of the fight day (4). Third, make the re-
27
28 205 weighting official and put limits for the percentage of WR, under penalty of cancellation
29
30 206 of the combat and/or punishment to the athlete. This rule was adopted in judo, where the
31
32 207 extra weight-in has a limit of 5% of body weight (18). Fourth, events
33
34 208 promoters/companies obligate a change of division if the athlete exceeds 5% of WR on
35
36 209 the extra weight-in. This suggestion is limited because it depends on the voluntary
37
38 210 adherence of the companies. Finally, organizations such as USADA and WADA could
39
40 211 consider high levels of WR as doping with its corresponding punishment, as it has been
41
42 212 suggested by other authors (1,7).
43
44
45
46
47
48
49

50 213 Our findings should be interpreted with caution, considering the limitations of the
51
52 214 present study. Considering the nature of data, it was not possible to include other variables
53
54 215 in the regression analysis such as age, off-season weight, time of experience, and cartel,
55
56 216 which could also influence the results. The data analyzed in our study came from only
57
58 217 one athletic commission of the USA. This is important because the CSAC has a specific
59
60

1
2
3 218 orientation when the athletes exceed 10% of WR (8). Lastly, the time variability between
4
5 219 official weight-in and re-weighing (i.e., between 24h and 36h) also is relevant and is was
6
7 220 no specified in the datasheet. For instance, in the study by Coswig et al. (6) the weight
8
9 221 loss (~6%) was not compensated by the WR, while in Matthews et al., (14) weight loss
10
11 222 (~7%) were surpassed by WR. The main difference between the two studies was the
12
13 223 measurement time of the re-weighing (24h vs 36, respectively).
14
15
16

17 224 Considering that the athletes regain ~8-10% of body weight and our findings
18
19 225 conclude that a single 1% of regaining increase de probability of winning by 4,5%, the
20
21 226 magnitude of WR is a strong variable linked to the chance to win. In this way,
22
23 227 commissions, confederations, and events' organizers should consider changing the rules
24
25 228 related to weigh-in, not only equalize the chances of athletes in the combat, mainly linked
26
27 229 to body size and strength level, but also for safety reasons considering the risks involved
28
29 230 in RWL. Until the rules are not changed, the athletes should invest in good ways to regain
30
31 231 weight fast after official weight-in, to increase their chance of a win.
32
33
34
35

36 232
37
38

39 233 **ACKNOWLEDGMENTS**

40
41
42 234 The authors would like to appreciate to California State Athletic Commission for
43
44 235 kindly providing us with the data used in the present study. We also would like to thank
45
46 236 André Werneck for helping with data analysis
47
48
49

50 237
51
52

53 238 **DECLARATIONS**

54
55
56 239 The authors do not present any conflict of interest.
57
58
59
60

241 **REFERENCES**

- 242 1. Artioli, GG, Saunders, B, Iglesias, RT, and Franchini, E. It is Time to Ban Rapid
243 Weight Loss from Combat Sports. *Sport Med* 46: 1579–1584, 2016.
- 244 2. Barley, OR, Chapman, DW, and Abbiss, CR. The Current State of Weight-
245 Cutting in Combat Sports. *Sports* 7: 123, 2019.
- 246 3. Brechney, GC, Chia, E, and Moreland, AT. Weight-Cutting Implications for
247 Competition Outcomes in Mixed Martial Arts Cage Fighting. *J Strength Cond*
248 *Res* 1: 1, 2019.
- 249 4. Connor, J and Egan, B. Prevalence, Magnitude and Methods of Rapid Weight
250 Loss Reported by Male Mixed Martial Arts Athletes in Ireland. *Sports* 7: 206,
251 2019.
- 252 5. Coswig, VS, Fukuda, DH, and Del Vecchio, FB. Rapid weight loss elicits
253 harmful biochemical and hormonal responses in mixed martial arts athletes. *Int J*
254 *Sport Nutr Exerc Metab* 25: 480–486, 2015.
- 255 6. Coswig, VS, Miarka, B, Pires, DA, Silva, LM da, Bartel, C, and Del Vecchio,
256 FB. Weight Regain, But Not Weight Loss, Is Related to Competitive Success in
257 Real-life Mixed Martial Arts Competition. *Int J Sport Nutr Exerc Metab* , 2018.
- 258 7. Crighton, B, Close, GL, and Morton, JP. Alarming weight cutting behaviours in
259 mixed martial arts: A cause for concern and a call for action. *Br J Sports Med* 50:
260 446–447, 2016.
- 261 8. CSAC, CSAC. Commission Meeting materials - May 16, 2017. 2017. pp. 1–350
- 262 9. CSAC, CSAC. Commission Meeting Minutes - October 15, 2019. 2019. pp. 1–18

- 1
2
3 263 10. Faro, HKC, Tavares, MPM, Gantois, PMD, Pereira, DC, Lima, MNM De, Lima-
4
5 264 junior, D De, et al. Comparison of competitive anxiety and self-confidence in
6
7 265 Brazilian Jiu-Jitsu skills levels : a cross-sectional in-event study. *Ido Mov Cult*
8
9 266 20: 30–39, 2020.
- 11
12
13 267 11. Franchini, E, Brito, CJ, and Artioli, GG. Weight loss in combat sports:
14
15 268 Physiological, psychological and performance effects. *J Int Soc Sports Nutr* 9: 2–
16
17 269 7, 2012.
- 19
20 270 12. Jetton, A, Lawrence, M, Meucci, M, Haines, T, Collier, S, Morris, D, et al.
21
22 271 Dehydration and acute weight gain in mixed martial arts fighters before
23
24 272 competition. *J Strength Cond Res* 27: 1322–1326, 2013.
- 26
27
28 273 13. Kasper, Andreas, M, Crighton, B, Langan-Evans, C, Riley, P, Sharma, A, Close,
29
30 274 GL, et al. Case Study: Extreme weight making causes relative energy deficiency,
31
32 275 dehydration and acute kidney injury in a male mixed martial arts athlete. *Int J*
33
34 276 *Sport Nutr Exerc Metab* 29: 331–338, 2019.
- 36
37
38 277 14. Matthews, JJ and Nicholas, C. Extreme rapid weight loss and rapid weight gain
39
40 278 observed in UK mixed martial arts athletes preparing for competition. *Int J Sport*
41
42 279 *Nutr Exerc Metab* 27: 122–129, 2017.
- 44
45 280 15. Matthews, JJ, Stanhope, EN, Godwin, MS, Holmes, MEJ, and Artioli, GG. The
46
47 281 magnitude of rapid weight loss and rapid weight gain in combat sport athletes
48
49 282 preparing for competition: A systematic review. *Int J Sport Nutr Exerc Metab* 29:
50
51 283 441–452, 2019.
- 53
54
55 284 16. Murugappan, KR, Cocchi, MN, Bose, S, Neves, SE, Cook, CH, Sarge, T, et al.
56
57 285 Case Study: Fatal Exertional Rhabdomyolysis Possibly Related to Drastic Weight
58
59 286 Cutting. *Int J Sport Nutr Exerc Metab* 11: 1–4, 2018.

- 1
2
3 287 17. Park, S, Alencar, M, Sassone, J, Madrigal, L, and Ede, A. Self-reported methods
4
5 288 of weight cutting in professional mixed-martial artists: How much are they losing
6
7 289 and who is advising them? *J Int Soc Sports Nutr* 16: 1–8, 2019.
- 9
10 290 18. Reale, R, Cox, RG, Slater, G, and Burke, LM. Regain in Body Mass After
11
12 291 Weigh-In is Linked to Success in Real Life Judo Competition Authors: *Int J*
13
14 292 *Sport Nutr Exerc Metab* 26: 525–530, 2016.
- 15
16
17
18 293 19. Santos-Junior, RB, Utter, AC, McAnulty, SR, Bernardi, BRB, Buzzachera, CF,
19
20 294 Franchini, E, et al. Weight loss behaviors in Brazilian mixed martial arts athletes.
21
22 295 *Sport Sci Health* , 2019. Available from: <https://doi.org/10.1007/s11332-019->
23
24 296 00581-x
- 25
26
27
28 297 20. UFC and USADA. Prohibited list and guide to the prohibited list. 2019.
- 29
30
31 298
32
33
34 299
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Distribution of athletes by division, sex, and percentage of weight regained by category.

Weight Class	% of the sample	Sex (n)			Percentage of WR (%)	
		M	F	All	Median (IQR)	Range (min-max)
Atomweight	1.00	-	15	15	9.35 (6.48 – 11.12)	0.85 – 12.39
Strawweight	3.30	-	48	48	8.06 (5.23 – 11.04)	-2.51 – 15.96
Flyweight	6.80	83	17	100	9.42 (7.30 – 11.77)	1.13 – 16.86
Bantamweight	16.1	224	13	237	9.72 (7.83 – 12.45)	-0.23 – 17.78
Featherweight	20.9	291	17	308	8.95 (6.46 – 10.80)	-0.14 – 16.58
Lightweight	16.3	240	-	240	8.45 (6.02 – 10.60)	-2.61 – 20.44
Welterweight	16.4	242	-	242	8.24 (5.79 – 10.41)	-1.89 – 16.78
Middleweight	10.6	156	-	156	5.82 (4.24 – 8.45)	-6.15 – 14.27
Light Heavyweight	4.40	65	-	65	5.69 (4.19 – 7.67)	-1.09 – 13.19
Heavyweight	3.90	57	-	57	0.75 (0.08 – 1.88)	-6.93 – 8.56
Super Heavyweight	0.40	6	-	6	1.05 (-0.38 – 1.89)	-0.44 – 3.27
Overall	100	1,374	100	1,474	8.26 (5.48 – 10.62)	-6.93 – 20.44

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Note: IQR; interquartile range; WR = weight regain

Table 2. Logistic regression of independent variables to predict fight outcome (win) (n=1,474).

Variable	Fight outcome		
	β (95%CI)	OR (95%CI)	p
%WR	0.044 (0.014 – 0.074)	1.046 (1.015 – 1.078)	0.004
Sex	0.007 (-0.450 – 0.465)	1.007 (0.637 – 1.592)	0.974
Official weight	-0.000 (-0.013 – 0.012)	0.999 (0.986 – 1.012)	0.954
WD from de opponent	0.001 (-0.010 – 0.012)	1.000 (0.990 – 1.012)	0.874
Division	0.024 (-0.140 – 0.190)	1.025 (0.869 – 1.209)	0.767
Constant	-0.601 (-1.833 – 0.630)	0.547 (0.159 – 1.889)	0.339

Note: CI = confidence interval; OR = odds ratio; WD = weight difference; WR = weight regain.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Table 1. Distribution of athletes by division, sex, and percentage of weight regained by category.

Weight Class	% of the sample	Sex (n)			Percentage of WR (%)	
		M	F	All	Median (IQR)	Range (min-max)
Atomweight	1.00	-	15	15	9.35 (6.48 – 11.12)	0.85 – 12.39
Strawweight	3.30	-	48	48	8.06 (5.23 – 11.04)	-2.51 – 15.96
Flyweight	6.80	83	17	100	9.42 (7.30 – 11.77)	1.13 – 16.86
Bantamweight	16.1	224	13	237	9.72 (7.83 – 12.45)	-0.23 – 17.78
Featherweight	20.9	291	17	308	8.95 (6.46 – 10.80)	-0.14 – 16.58
Lightweight	16.3	240	-	240	8.45 (6.02 – 10.60)	-2.61 – 20.44
Welterweight	16.4	242	-	242	8.24 (5.79 – 10.41)	-1.89 – 16.78
Middleweight	10.6	156	-	156	5.82 (4.24 – 8.45)	-6.15 – 14.27
Light Heavyweight	4.40	65	-	65	5.69 (4.19 – 7.67)	-1.09 – 13.19
Heavyweight	3.90	57	-	57	0.75 (0.08 – 1.88)	-6.93 – 8.56
Super Heavyweight	0.40	6	-	6	1.05 (-0.38 – 1.89)	-0.44 – 3.27
Overall	100	1,374	100	1,474	8.26 (5.48 – 10.62)	-6.93 – 20.44

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Note: IQR; interquartile range; WR = weight regain

Table 2. Logistic regression of independent variables to predict fight outcome (win) (n=1,474).

Variable	Fight outcome		
	β (95%CI)	OR (95%CI)	p
%WR	0.044 (0.014 – 0.074)	1.046 (1.015 – 1.078)	0.004
Sex	0.007 (-0.450 – 0.465)	1.007 (0.637 – 1.592)	0.974
Official weight	-0.000 (-0.013 – 0.012)	0.999 (0.986 – 1.012)	0.954
WD from de opponent	0.001 (-0.010 – 0.012)	1.000 (0.990 – 1.012)	0.874
Division	0.024 (-0.140 – 0.190)	1.025 (0.869 – 1.209)	0.767
Constant	-0.601 (-1.833 – 0.630)	0.547 (0.159 – 1.889)	0.339

Note: CI = confidence interval; OR = odds ratio; WD = weight difference; WR = weight regain.