

SUPPLEMENTARY MATERIALS

Changes of labile, stable and water-soluble fractions of biochar after two years in a vineyard soil

Alessandro G. Rombolà^{a*}, Nicolas Greggio^{b*}, Daniele Fabbri^c, Andrea Facchin^a, Cristian Torri^a, Roberta Pulcher^b, Carlotta Carlini^d, Enrico Balugani^d, Diego Marazza^d, Denis Zannoni^b, Alessandro Buscaroli^b

^a *Department of Chemistry “Giacomo Ciamician”, Alma Mater Studiorum-University of Bologna, Campus of Ravenna, Via Sant’Alberto 163, 48123, Ravenna, Italy*

^b *Department of Biological, Geological, and Environmental Sciences, Alma Mater Studiorum-University of Bologna, Campus of Ravenna, Interdepartmental Centre for Research in Environmental Sciences (CIRSA), Via Sant’Alberto 163, 48123, Ravenna, Italy*

^c *Department of Chemistry “Giacomo Ciamician” and C.I.R.I. FRAME Tecnopolo di Rimini, Alma Mater Studiorum-University of Bologna, Via Dario Campana 71, 47192, Rimini, Italy*

^d *Department of Physics and Astronomy, University of Bologna, 40126 Bologna, Italy*

*Corresponding authors.

E-mail address: alessandro.rombola@unibo.it (A.G. Rombolà)

E-mail address: nicolas.greggio@unibo.it (N. Greggio)

Number of pages: 3 (including this page)

Number of figures: 3



Figure SM1. Example of furrows filled with biochar, before of shoot transplanting.




Figure SM2. Thermogravimetric (TG) curves obtained from fresh and aged biochar applied in the field experimental.




Figure SM3. Derivative thermogravimetric (DTG) curves obtained from fresh and aged biochar applied in the field experimental.