

- Mandic A, Sakac M, Benedettelli S, Dinelli G, 2012. Health-promoting phytochemicals of Italian common wheat varieties grown under low-input agricultural management. *J. Sci. Food Agr.* 92:2800-10.
- Dinelli G, Segura-Carretero A, Di Silvestro R, Marotti I, Arráz-Román D, Benedettelli S, Ghiselli L, Fernandez-Gutierrez A, 2011. Profiles of phenolic compounds in modern and old common wheat varieties determined by liquid chromatography coupled with time-of-flight mass spectrometry. *J. Chromatogr. A* 1218:7670-81.
- Döring TF, Knapp S, Kovacs G, Murphy K, Wolfe MS, 2011. Evolutionary plant breeding in cereals: into a new era. *Sustainability-Basel* 3:1944-71.
- Döring TF, Annicchiarico P, Clarck S, Haigha Z, Jonese HE, Pearcea H, Snapef J, Zhang J, Wolfe MS, 2015. Comparative analysis of performance and stability among compositecross populations, variety mixtures and pure lines of winter wheat inorganic and conventional cropping systems. *Field Crops Res.* 183:235-45.
- EU, 2013. Regulation (EU) 1305/2013 of the European Parliament and of the Council of 17 December 2013.
- EU, 2018. Regulation (EU) 2018/848 of the European Parliament and of the Council of 30 May 2018.
- Folch J, Lees M, Sloane Stanley GH, 1957. A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.* 226:497-509.
- Food FF, 2003. Nutrition paper 77: food energy - methods of analysis and conversion factors. FAO: Rome, Italy.
- Frutos E, Galindo MP, Leiva V, 2014. An interactive biplot implementation in R for modeling genotype-by-environment interaction. *Stoch. Environ. Res. Risk Assess.* 28:1629-41
- Johnson VA, Biever KJ, Haunold A, Schmidt JW, 1966. Inheritance of plant height, yield of grain, and other plant and seed characteristics in a cross of hard red winter wheat (*Triticum aestivum* L.). *Crop Sci.* 6: 336-8.
- Lazzaro M, Costanzo A, Farag DH, Bärberi P, 2017. Grain yield and competitive ability against weeds in modern and heritage common wheat cultivars are differently influenced by sowing density. *Ital. J. Agron.* 12:901.
- Lee SC, Prosky L, Vries JWD, 1992. Determination of total, soluble, and insoluble dietary fiber in foods. Enzymatic-gravimetric method, MES-TRIS buffer: collaborative study. *J. Aoac Int.* 75:395-416.
- Li S, Li X, 2017. Global understanding of farmland abandonment: a review and prospects. *J. Geogr. Sci.* 27:1123-50.
- Liu H, Shi Z, Ma F, Xu Y, Han G, Zhang J, Liu D, An D, 2022. Identification and validation of plant height, spike length and spike compactness loci in common wheat (*Triticum aestivum* L.). *BMC Plant. Biol.* 22:568.
- MacDonald D, Crabtree JR, Wiesinger G, Dax T, Stamou N, Fleury P, Gutierrez Lazpita J, Gibon A, 2000. Agricultural abandonment in mountain areas of Europe: environmental consequences and policy response. *J. Environ. Manag.* 59:47-69.
- Marcos-Barbero EL, Pérez P, Martínez-Carrasco R, Arellano JB, Morcuende R, 2021. Genotypic variability on grain yield and grain nutritional quality characteristics of wheat grown under elevated CO₂ and high temperature. *Plants* 10:1043.
- Migliorini P, Spagnolo S, Torri L, Arnoulet M, Lazzarini G, Ceccarelli S, 2016. Agronomic and quality characteristics of old, modern and mixture wheat varieties and landraces for organic bread chain in diverse environments of northern Italy. *Eur. J. Agron.* 79:131-41.
- Mitrovic B, Treski S, Stojakoviã M, Ivanoviã M, Bekavac G, 2012. Evaluation of experimental maize hybrids tested in multi-location trials using AMMI and GGE biplot analyses. *Turk. J. Field Crops* 17:35-40.
- Morris ML, Bellon MR, 2004. Participatory plant breeding research: opportunities and challenges for the international crop improvement system. *Euphytica*, 136:21-35.
- Nuttall JG, O'Leary GJ, Panozzo JF, Walker CK, Barlow KM, Fitzgerald GJ, 2017. Models of grain quality in wheat - a review. *Field Crops Res.* 202:136-45.
- Porfiri O, 2014. I frumenti: dalle varietà al campo. Pentàgora.
- Prosky L, Asp NG, Schweizer TF, Devries JW, Furda I, 1988. Determination of insoluble, soluble, and total dietary fiber in foods and food products: interlaboratory study. *J. Assoc. Off. Ana. Chem.* 71:1017-23.
- R Core Team, 2021. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Url: <https://www.R-project.org/>.
- Regione Emilia-Romagna, 2022. Ricerca, qualità e sostenibilità delle politiche regionali in agricoltura e nell'agroalimentare. Available from: <https://agricoltura.regione.emilia-romagna.it>
- Shewry PR, Piironen V, Lampi AM, Edelmann M, Kariluoto S, Nurmi T, Aman P, 2010. The HEALTHGRAIN wheat diversity survey: effects of genotype and environment on phytochemical and dietary fiber components, *J. Agric. Food Chem.* 48:9291-8.
- Singleton VL, Orthofer R, Lamuela-Raventós RM, 1999. Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. In: *Method. Enzymol.* (Vol. 299, pp. 152-178). Academic press.
- Tellarini S, 2017. Grani e gente. Tutto quello che avreste voluto sapere sui vecchi grani e sui loro uomini. Stilgraf.
- Triboi E, Martre P, Girousse C, Ravel C, Triboi-Blondel A, 2006. Unravelling environmental and genetic relationships between grain yield and nitrogen concentration for wheat. *Eur. J. Agron.* 25:108-18.
- Vallega A, 1992. Sea management: a theoretical approach. CRC Press.
- Van der Meulen A, Chauhan BS, 2017. A review of weed management in wheat using crop competition. *Crop Protection*, 95:38-44.
- Varotto M, 2006. Montagne deserte: l'abbandono delle 'terre alte' visto attraverso la cartografia. *Bollettino dell'Associazione Italiana di Cartografia* 117-118-119, Trieste, EUT Edizioni Università di Trieste, pp. 165-177.
- Villa TCC, Maxted N, Scholten M, Ford-Lloyd B, 2005. Defining and identifying crop landraces. *Plant Gen. Res.* 3:373-84.
- Zi Y, Ding J, Song J, Humphreys G, Peng Y, Li C, Zhu X, Wenshan G, 2018. Grain yield, starch content and activities of key enzymes of waxy and non-waxy wheat (*Triticum aestivum* L.). *Sci. Rep.* 8:4548.