



Performance of a seed amplification assay for misfolded alpha-synuclein in cerebrospinal fluid and brain tissue in relation to Lewy body disease stage and pathology burden

Giuseppe Mario Bentivenga¹ · Angela Mammana² · Simone Baiardi¹ · Marcello Rossi² · Alice Ticca¹ · Franco Magliocchetti² · Andrea Mastrangelo¹ · Anna Poleggi³ · Anna Ladogana³ · Sabina Capellari^{1,2} · Piero Parchi^{1,2}

Received: 18 September 2023 / Revised: 19 November 2023 / Accepted: 20 November 2023
© The Author(s) 2024

Abstract

The development of in vitro seed amplification assays (SAA) detecting misfolded alpha-synuclein (α Syn) in cerebrospinal fluid (CSF) and other tissues has provided a pathology-specific biomarker for Lewy body disease (LBD). However, α Syn SAA diagnostic performance in early pathological stages or low Lewy body (LB) pathology load has only been assessed in small cohorts. Moreover, the relationship between SAA kinetic parameters, the number of α Syn brain seeds and the LB pathology burden assessed by immunohistochemistry has never been systematically investigated. We tested 269 antemortem CSF samples and 138 serially diluted brain homogenates from patients with and without neuropathological evidence of LBD in different stages by the α Syn Real-Time Quaking-Induced Conversion (RT-QuIC) SAA. Moreover, we looked for LB pathology by α Syn immunohistochemistry in a consecutive series of 604 Creutzfeldt–Jakob disease (CJD)-affected brains. α Syn CSF RT-QuIC showed 100% sensitivity in detecting LBD in limbic and neocortical stages. The assay sensitivity was significantly lower in patients in early stages (37.5% in Braak 1 and 2, 73.3% in Braak 3) or with focal pathology (50% in amygdala-predominant). The average number of CSF RT-QuIC positive replicates significantly correlated with LBD stage. Brain homogenate RT-QuIC showed higher sensitivity than immunohistochemistry for the detection of misfolded α Syn. In the latter, the kinetic parameter lag phase (time to reach the positive threshold) strongly correlated with the α Syn seed concentration in serial dilution experiments. Finally, incidental LBD prevalence was 8% in the CJD cohort. The present results indicate that (a) CSF RT-QuIC has high specificity and sufficient sensitivity to detect all patients with LB pathology at Braak stages > 3 and most of those at stage 3; (b) brain deposition of misfolded α Syn precedes the formation of LB and Lewy neurites; (c) α Syn SAA provides “quantitative” information regarding the LB pathology burden, with the lag phase and the number of positive replicates being the most promising variables to be used in the clinical setting.

Keywords RT-QuIC · α -Synuclein · Lewy bodies · Parkinson’s disease · Diagnosis · Amygdala

Introduction

Lewy body disease (LBD), the second most prevalent neurodegenerative disorder after Alzheimer’s disease (AD), is characterized by the misfolding and aggregation of alpha-synuclein (α Syn), resulting in the formation of cytoplasmic neuronal inclusion in cell bodies (i.e., Lewy bodies [LBs]) and nerve terminals (i.e., Lewy neurites [LNs]) [36, 61]. Besides being the pathological hallmark of dementia with Lewy bodies (DLB) and the majority of Parkinson’s disease (PD) cases, LB pathology often manifests as co-pathology in patients with AD, and it also affects a variable proportion of aged individuals lacking clinical signs of

Giuseppe Mario Bentivenga and Angela Mammana have contributed equally to the study.

✉ Piero Parchi
piero.parchi@unibo.it

- 1 Department of Biomedical and Neuromotor Sciences, University of Bologna, Bologna, Italy
- 2 IRCCS, Istituto Delle Scienze Neurologiche di Bologna, Bologna, Italy
- 3 Department of Neurosciences, Istituto Superiore di Sanità, Rome, Italy

motor disturbances or cognitive decline. The latter condition is known as “incidental LBD” (iLBD) and likely represents the pathological preclinical stage of PD or DLB [22, 33, 35]. As it is increasingly recognized for misfolded proteins linked to neurodegeneration, misfolded α Syn can act as a seed to induce the conversion of the normal cellular α Syn, allowing for the transcellular propagation of LB pathology in a prion-like manner [24, 67]. In the CNS, LBD most often starts in the medulla oblongata and then spreads intracerebrally, following an ascending course from the brainstem to the cerebral cortex, passing through the limbic areas [12, 13, 28, 63]. Accordingly, three major disease stages have been identified, i.e., the brainstem (I), limbic or transitional (II), and neocortical (III) [37, 43]. A second, more detailed staging scheme, proposed by Heiko Braak [12, 13], recognizes six stages, among which the first three, corresponding to the brainstem stage, define in more detail the progressive involvement of the brainstem from the medulla oblongata to the pons and the midbrain reaching the basal forebrain and the hypothalamus before involving the limbic areas. There are well-established exceptions to the LB pathology progression described above [2, 50, 69], the most common being the amygdala-predominant variant, in which a widespread α Syn deposition is found in the amygdala, associated with only mild and focal LB pathology in the brainstem, not precisely following the Braak staging, often in association with moderate or severe AD pathology [19, 29, 40, 41, 59, 63, 65].

The development of *in vitro* seed amplification assays (SAA) that indirectly detect misfolded α Syn in the cerebrospinal fluid (CSF) and other tissues has recently provided a pathology-specific biomarker for LBD, thus representing a breakthrough in the field [14]. However, only a few studies in limited cohorts have correlated the *in vivo* outcome based on α Syn SAA result against the gold standard of the post-mortem LB pathology assessment [3, 7, 10, 21, 27, 55]. At the same time, none has, to date, specifically correlated the kinetic parameter values of α Syn SAA with the LB pathology burden.

In this study, we took advantage of a unique collection of CSF and brain pairs from the same patients collected within a limited time frame due to a rapidly progressive disease to address critical questions regarding the sensitivity of our α Syn SAA in detecting LB pathology at different stages and its overall specificity.

Moreover, to provide new insights into the relationship between the kinetic parameters of the α Syn SAA and the number of α Syn seeds, and into the correlation between LB pathology burden assessed by immunohistochemistry and the α Syn SAA outcome, we performed the α Syn SAA on tissue homogenates from different brain areas affected by LB pathology at different stages.

Finally, we used a cohort of consecutive well-characterized patients with Creutzfeldt–Jakob disease (CJD) lacking

any clinical evidence of cognitive decline and/or motor disturbances before the rapid onset of symptoms related to CJD to estimate the frequency of iLBD pathology in the elderly population.

Materials and methods

Case selection

A series of 862 consecutive brains have been collected at the Neuropathology Laboratory (NP-Lab) at the Istituto delle Scienze Neurologiche di Bologna (ISNB) between 2003 and 2022. The cohort (NP cohort) mainly included cases of progressive dementia, often with a rapidly progressive course, or of atypical parkinsonism, primarily belonging to the following diagnostic categories: CJD, LBD, AD, frontotemporal lobar degeneration, multiple system atrophy (MSA), progressive supranuclear palsy, encephalitis, lymphoma, and vascular dementia. Neuropathological diagnosis of neurodegenerative diseases was made according to established criteria [2, 17, 18, 23, 38, 45, 48]. All brains have been classified into α Syn LB+ and α Syn LB– subgroups based on protein aggregate assessment by immunohistochemistry (see below for further details).

For the study's primary aims, we selected from the NP cohort all the participants with antemortem CSF available and those with a final CJD diagnosis. The first group comprised 269 patients (CSF cohort), including 210 subjects lacking LB pathology (α Syn LB– group) and 59 with LB pathology (α Syn LB+ group). In the α Syn LB– group, we excluded CSF samples for which the time between lumbar puncture (LP) and death was more than one year, while we included all available samples in the α Syn LB+ group (of note, only 7/59 had a time LP-death > 1 year). The second group included 604 brains with a CJD diagnosis (CJD cohort).

Neuropathological studies

Neuropathological examination was performed at NP-Lab using standardized procedures as described [39]. Briefly, each brain was divided sagittally, and the left hemibrain fixed in 10% buffered formalin while the right one was sectioned coronally and then immediately frozen at -80°C in sealed plastic bags. The formalin-fixed left hemibrain was serially sectioned in 1 cm slices, and tissue blocks from 24 regions were processed routinely to obtain paraffin-embedded brain tissue blocks [49].

Seven μm thick sections from each block were stained with hematoxylin–eosin for screening. Also, immunohistochemistry with antibodies specific for α Syn (LB509, dilution 1:100, Thermo Fisher Scientific, and KM51, dilution 1:500,

Novocastra), phospho-tau (*p*-tau) (AT8, dilution 1:100, Innogenetics), and A β (4G8, dilution 1:5000, Signet Labs), was applied to all cases using several brain regions, mainly following established consensus criteria [1, 2, 16, 45, 48]. An experienced neuropathologist (PP) formulated the final diagnosis, assigned the stage of LB pathology according to consensus criteria [2, 4], and classified each α Syn LB+ case according to the level of AD neuropathologic change (ABC score) [1, 45].

For the semiquantitative assessment of LB pathology, brains with α Syn positive LBs or LNs in at least one section and available CSF samples were examined independently by two evaluators (PP and GMB). Specifically, α Syn immunoreactivity was evaluated semi-quantitatively (0-no immunoreactivity; 1-mild; 2-moderate; 3-prominent immunoreactivity) in the following regions: medulla oblongata (vagus dorsomedial nucleus, intermediate reticular zone), pons (locus coeruleus, raphe), substantia nigra, hypothalamus, basal forebrain (Meynert's basal nucleus and surrounding gray matter), amygdala, hippocampus (CA2–CA4 sectors), parahippocampal gyrus, cingulate gyrus, middle temporal gyrus, middle frontal gyrus, occipital cortex, and cerebellum. We scored the neuronal LBs and LNs separately. Each case was given a combined “LB score” (range 0–78).

Evaluation of α Syn seeding activity in CSF and brain homogenates by SAA

All CSF samples were obtained by LP at the L3/L4 or L4/L5 level following a standard procedure. They were subsequently divided into 400–500 μ l aliquots and stored in polypropylene tubes at -80°C until analysis. We performed the CSF and brain homogenates α Syn Real-Time Quaking-Induced Conversion (RT-QuIC) SAA, including the purification of recombinant wild-type human α Syn, as previously described with minor analysis modifications [26, 53–55]. Samples and controls were deemed positive after the first run when at least three out of four replicates reached a threshold arbitrarily set at 30% of the median of the maximum fluorescence intensity (*I*_{max}) reached by the positive control (positive) replicates. To minimize the risk of false-positive results, we repeated three times the analysis of samples showing seeding activity in only one or two of four replicates in the first run. Then, we considered the result “positive” only when at least 4 of the 12 total replicates reached the threshold.

Tenfold serial dilutions were prepared for the α Syn seeds quantification in brain homogenates, and each sample was run in quadruplicates. The dilution series was analyzed until the α Syn RT-QuIC SAA yielded ≤ 2 out of 4 positive replicates. Then, we used the Spearman–Kärber algorithm, which has been previously applied for the quantification of prion seeding activity in analogy to a bioassay's lethal dose

(LD₅₀) calculation [20, 30, 68], to estimate a seeding dose or dilution at which only 50% (e.g., 2 of 4) of replicates are positive (50% seeding doses or SD₅₀). Next, to estimate the α Syn seed concentration in each brain area, a logSD₅₀ per mg of tissue was calculated. Finally, we obtained a combined logSD₅₀ score for each brain by summing all brain areas' logSD₅₀/mg values to compare the overall α Syn seeding activity between brains.

We used eight batches of α Syn recombinant protein throughout the study for the brain homogenate analyses. Each batch underwent a quality control test before use. We ran at least one positive and negative control on each plate. The positive controls were chosen from patients with definite LBD whose brain homogenates or CSF samples gave four positive replicates during screening. In each validated experiment (plate) included in the final analysis, the positive controls showed at least three positive replicates. The average inter-batch coefficient of variation (CV) of the lag phase (i.e., the time required for the α Syn RT-QuIC SAA reaction to reach the above-mentioned “positivity” threshold) and the *I*_{max} in positive controls were 10.7% and 22.5%, respectively. All α Syn RT-QuIC SAA experiments were performed at the NP-Lab by personnel blinded to neuropathological diagnosis.

Statistical analysis

α Syn RT-QuIC SAA relative fluorescence response data were analyzed and plotted using the software GraphPad Prism 8.3. The *I*_{max}, the lag phase and the area under curve (AUC) were extracted. The normality of the distribution of these variables from each group was assessed using the Kolmogorov–Smirnov test. The protein aggregation rate (PAR) was calculated as the inverse of the lag phase (1/lag phase). Mann–Whitney U test, one-way analysis of variance (followed by Tukey's multiple comparisons post hoc test), and Kruskal–Wallis test (followed by Dunn's post hoc analysis) were used to compare the continuous variables between the groups. The Spearman's rank correlation coefficient was computed to determine significant associations between parameters. *P* values < 0.05 were considered statistically significant.

Results

Neuropathological analysis of LB pathology

The neuropathological analysis of the NP cohort revealed LB pathology in the brains of 114 (13.2%) patients. Forty-eight of 604 (7.9%) participants in the CJD group showed LB pathology, whereas 66 of 258 (25.6%) individuals had LB pathology in the non-CJD group.

Of the 114 patients with LB pathology in the NP cohort, 29 showed a neocortical stage, 37 showed a limbic stage, 42 a brainstem stage, and only 6 belonged to the amygdala-predominant variant. The relative proportion of subjects in each LBD stage of LB pathology in the whole cohort and the major subgroups is shown in Table 1.

Rates of AD pathology were high across the cohort with LB pathology: 30 of 66 (45.5%) participants with limbic or neocortical LBD stages had intermediate or high degrees of AD neuropathological changes, as did almost all (5/6, 83.3%) cases with the amygdala-predominant variant.

Sensitivity and specificity of the α Syn RT-QuIC SAA using CSF Samples

We performed the α Syn RT-QuIC SAA in 269 patients with antemortem CSF available, including 210 subjects lacking LB pathology and 59 with LB pathology assessed by α Syn immunostaining (see Supplementary Table 1 for further information about the pathological entities in the α Syn LB+ group). Demographic characteristics and α Syn RT-QuIC SAA results in the CSF cohort are shown in Table 2. The CSF α Syn RT-QuIC SAA yielded an overall specificity against the gold standard (i.e., neuropathologic examination) of 98.6%. Unexpected positive results were limited to three cases with a primary neuropathological diagnosis of Wernicke's encephalopathy, AD, and sporadic CJD (sCJD), which showed no detectable α Syn brain deposits, despite the

Table 1 Demographic characteristics and neuropathological assessment results in the whole NP cohort and in the main diagnostic groups

Main NP-diagnosis	N	Age at death	F (%)	α Syn LB+				α Syn LB-
				Neocortical	Limbic	Brainstem	Amygdala-predominant	
Whole NP cohort	862	69.1 \pm 9.8	49.4	29 (3.4%)	37 (4.3%)	42 (4.9%)	6 (0.7%)	748 (86.8%)
CJD	604	68.4 \pm 9.4	50.0	2 (0.3%)	19 (3.1%)	27 (4.5%)	-	556 (92.1%)
Non-CJD ^a	258	71.0 \pm 10.5	48.0	27 (10.5%)	18 (6.9%)	15 (5.8%)	6 (2.3%)	192 (74.4%)

Age at death is expressed as mean \pm SD

CJD, Creutzfeldt–Jakob disease; LB, Lewy body pathology; NP, neuropathological

^amain diagnostic groups: Alzheimer's disease, Lewy body disease, subcortical arteriosclerotic encephalopathy

Table 2 Demographic characteristics and CSF analysis results in the CSF cohort

	N	F (%)	Age at death (years)	Time LP-death (months)	N. α Syn RT-QuIC SAA-positive	N. α Syn RT-QuIC SAA-negative	Sensitivity (%)	Specificity (%)
α Syn LB-	210	47.1	70.1 \pm 9.4	1.8 \pm 2.7	3	207		98.6
α Syn LB+	59	42.3	73.0 \pm 7.9	5.0 \pm 12.4	48	11	81.4	
McKeith stage								
Neocortical	15	53.3	77.5 \pm 4.7	8.2 \pm 12.2	15	0	100	
Limbic	17	47.0	72.8 \pm 7.1	3.5 \pm 11.9	17	0	100	
Brainstem	23	39.1	70.3 \pm 8.5	1.9 \pm 5.4	14	9	60.9	
Braak stage								
1	4	50.0	66.5 \pm 5.9	0.5 \pm 0.5	2	2	50.0	
2	4	50.0	69.7 \pm 9.2	1.0 \pm 0.8	1	3	25.0	
3	15	33.3	71.4 \pm 9.0	2.5 \pm 6.7	11	4	73.3	
4	5	60.0	73.6 \pm 9.1	0.4 \pm 0.5	5	0	100	
5	12	41.6	72.5 \pm 6.6	4.9 \pm 14.2	12	0	100	
6	15	53.3	77.5 \pm 4.7	8.2 \pm 12.2	15	0	100	
Amygdala-predominant	4	0	72.7 \pm 11.9	16.7 \pm 31.5	2	2	50	
LB score (0–78)					33.0 \pm 15.4	7.0 \pm 4.6		

Age at death, time LP-death, and LB score are expressed as mean \pm SD

LB, Lewy body; LP, lumbar puncture

positivity by α Syn RT-QuIC SAA. Of note, the assay yielded negative results also in the two MSA cases.

Of the 59 individuals with LB pathology, 48 tested positive by the α Syn RT-QuIC SAA, resulting in an assay sensitivity to detect α Syn pathology of 81.4% (Table 2). However, we observed significant differences in the test sensitivity after stratification by the extent and load of LB pathology. The α Syn RT-QuIC SAA had a sensitivity of 100% in detecting α Syn seeds in limbic and neocortical LBD stages (i.e., Braak stages > 3), 60.8% in the brainstem stage (73.3% and 37.5% in Braak stages 3 and 1/2, respectively) and 50% in detecting the amygdala-predominant variant. Interestingly, there was a clear association between the quantitative score of LB pathology and the result of the α Syn RT-QuIC SAA (Fig. 1). LB score in CSF α Syn+ individuals was significantly higher than in CSF α Syn— ones ($p < 0.0001$) (Table 2). Indeed, virtually all (44/45) individuals with a score > 10 were positive, whereas most of those (10/14) with a score below this threshold were negative (Fig. 1).

Specifically, the “negative” group included two individuals showing only focal LBs and LNs in the medulla oblongata (Braak stage 1), three with a Braak stage 2 with sparse LBs and LNs in the medulla oblongata and pons, four in Braak stage 3 (among whom, one showed an atypical distribution with sparse LBs and LNs in the medulla oblongata and substantia nigra skipping the pons) and two with the amygdala-predominant variant showing moderate LB pathology in the amygdala associated with a mild pathology limited to the medulla oblongata.

Among the positive cases, we found a strong association between the number of positive replicates and the LB score ($r = 0.7715$, 95% confidence interval [CI] 0.6378 to




Fig. 1 CSF α Syn RT-QuIC SAA results in α Syn LB+ patients according to Braak LBD stage and LB score. Red dots represent α Syn RT-QuIC SAA negative patients, while blue dots indicate the α Syn RT-QuIC SAA-positive ones. Almost all negative patients exhibited a low LBD (or amygdala-predominant) stage or low LB scores. AP, amygdala-predominant; CSF, cerebrospinal fluid; LBD, Lewy body disease

0.8601; $p < 0.0001$), and the Braak stage ($r = 0.7554$, 95% CI 0.6143–0.8497; $p < 0.0001$). Specifically, 80% and 20% of patients in neocortical stages showed 4/4 and 3/4 positive replicates, respectively. In the limbic stage, 47%, 35%, and 18% of cases had 4/4, 3/4, and 2/4 positive replicates, respectively. Among those in the brainstem stage, 7%, 42%, and 51% showed 4/4, 3/4, and 2/4 positive replicates, respectively. Finally, the two α Syn RT-QuIC SAA-positive amygdala-predominant cases showed 2/4 positive replicates (Supplementary Fig. 1).

Detection of α Syn seeds in brain samples from α Syn LB+ and α Syn LB-/ CSF α Syn RT-QuIC SAA negative patients

Next, we analyzed a subset of 20 patients who had frozen brain tissue available, including 5 brainstem, 5 limbic, 3 neocortical, 3 amygdala-predominant, and 4 α Syn LB-/CSF α Syn RT-QuIC negative cases. Tenfold serial dilutions of brain homogenates were obtained from the medulla oblongata, substantia nigra, amygdala, basal forebrain, cingulate gyrus, temporal cortex, frontal cortex, cerebellum, and occipital cortex. The 4 α Syn LB— participants were negative by the α Syn RT-QuIC SAA in all examined brain regions, consistent with the results obtained with the antemortem CSF. The opposite was true for the limbic and neocortical cases, in which we detected α Syn seeding activity in all tested brain areas over a varying dilution range (from 10^{-4} to 10^{-21}), depending on the case and the brain area. In contrast, participants showing the brainstem stage or the amygdala-predominant patients showed significant heterogeneity, mainly depending on the LB pathology load. Specifically, patients #51 (Braak 1) exhibited α Syn seeding activity only in the medulla oblongata (dilution range 10^{-4} – 10^{-6}), patient #48 (Braak 3) was α Syn RT-QuIC SAA-positive in the medulla oblongata and the substantia nigra (dilution range 10^{-4} – 10^{-13}), patient #50 (Braak 3) was α Syn RT-QuIC SAA-positive in brainstem areas, basal forebrain, amygdala, and cingulate gyrus (dilution range 10^{-4} – 10^{-6}), and patients #43 and #54 (both Braak 3) showed α Syn seeds in all brain areas (dilution range of 10^{-4} – 10^{-14} , with the highest dilutions needed in brainstem areas and basal forebrain). All three subjects with amygdala-predominant variant (#37, #41, and #57) were positive in the amygdala (dilution range 10^{-4} – 10^{-14}), basal forebrain and medulla (dilution range 10^{-4} – 10^{-7}). Moreover, α Syn seeding activity was variably detected in a few additional brain areas, such as the substantia nigra and cingulate gyrus (cases #37 and #41) or the temporal and occipital cortices (case #41) (dilution range 10^{-4} – 10^{-6}).

Notably, 47 of the 138 brain samples showed α Syn seeding activity despite the negative immunohistochemical assessment, highlighting the increased sensitivity of the

α Syn RT-QuIC SAA for the detection of misfolded α Syn compared to the standard immunohistochemical approach.

To better characterize the relationship between the kinetic parameters of the α Syn RT-QuIC SAA and the number of α Syn seeds, we computed the protein aggregation rate (PAR) (1/lag phase), a measure of amyloid formation rate. In all cases, a progressive decrease in PAR along the dilution series (i.e., as the number of α Syn seeds in the sample decreased) was observed in each brain area (Fig. 2). In contrast, I_{max} and AUC did not show a regular decline over the dilution series, exhibiting an often irregular and unpredictable pattern in numerous brain areas (data not shown).

Quantification of α Syn seeds in brain samples from α Syn LB+ patients

We then estimated each area's average α Syn seed concentration using the Spearman-Kärber algorithm. We expressed it as $\log SD_{50}$ (seed concentrations giving 50% positive replicate reactions) per mg of tissue (Table 3). $\log SD_{50}/mg$ values were highly variable depending on the case and the brain area, ranging from 7.2 to 23.9, with the highest values detected in the substantia nigra, the basal forebrain, and the amygdala. When stratifying patients according to the McKeith stage, we reported no significant differences between the average seed concentrations of the four groups in the different brain areas (considering only the cases in which α Syn seeding activity was detected). However, the results showed

a trend for the amygdala-predominant LBD cases to have lower average $\log SD_{50}/mg$ values than the other groups in all brain areas where α Syn positivity was detected (medulla oblongata, substantia nigra, basal forebrain, cingulate gyrus, and temporal cortex), except in the amygdala, where the values were in the brainstem cases' range (Fig. 3a).

To evaluate the association between the amount of LB pathology estimated through α Syn RT-QuIC SAA and that estimated through α Syn immunohistochemistry, we gave a combined $\log SD_{50}$ score to each case by summing the $\log SD_{50}/mg$ values of all tested brain areas. $\log SD_{50}$ score significantly correlated with LB score ($r=0.7903$, 95% CI 0.4719–0.9263; $p<0.001$) (Fig. 3b).

We also exploited α Syn seeds quantification analyses to compare the performance of α Syn RT-QuIC SAA and immunohistochemistry in detecting LB pathology. As previously mentioned, in 47/138 examined brain regions α Syn RT-QuIC SAA detected α Syn seeding activity notwithstanding a negative immunohistochemical assessment (Table 3). We found that all but 6 areas with $\log SD_{50}/mg$ values ≤ 8.7 were negative at immunohistochemical evaluation, while only 12 above this threshold were negative.

Differences in α Syn seeding activity in brain samples at SD_{50} dilution

Next, we tested the hypothesis that a different α Syn conformational strain would sustain the amygdala-predominant




Fig. 2 α Syn RT-QuIC SAA results in brain homogenates from α Syn LB+ patients. **a** Progressive reduction of seeding activity in brain homogenates of cingulate gyrus from case #24 along the dilution series. Each curve represents the average RFU values of positive quadruplicates. Standard deviation (SD) was hidden to improve the image's readability. **b–f** Progressive PAR reduction of α Syn RT-QuIC SAA reactions in different brain areas along the dilution series. The

dilution coefficient is expressed in a logarithmic scale. Each point represents the average PAR value of positive quadruplicates at each dilution. Standard deviation (SD) was hidden to improve the image's readability. Linear regression lines were applied to values from patients in the same McKeith stage only when at least three values were available. AP, amygdala-predominant; PAR, protein aggregation rate; RFU, relative fluorescent units; SAA, seed amplification assay

Table 3 LogSD₅₀/mg of brain homogenates and LB score results in different brain areas

Case		43	48	50	51	54	13	18	24	47	49	3	26	44	37	41	57	
Braak stage		3	3	3	1	3	5	5	5	4	5	6	6	6	6			
McKeith stage		Brainstem				Limbic				Neocortical				AMG-predominant				
Brain areas																		
ME	SD ₅₀ /mg	15.7	8.9	8.9	8.7	16.4	17.4	16.7	9.7	10.4	15.4	15.4	22.9	9.9	7.7	8.7	7.2	
	IHC	6	3	4	1	6	6	6	6	6	6	6	6	6	3	Neg	Neg	
SN	SD ₅₀ /mg	11.4	14.7	7.9	Neg	13.7	23.9	20.2	8.9	9.2	10.4	11.7	20.9	9.7	7.2	8.9	Neg	
	IHC	5	5	1	Neg	2	6	5	4	6	5	6	6	6	3	3	Neg	
BF	SD ₅₀ /mg	13.2	Neg	7.7	Neg	12.2	12.5	18.5	9.2	9.7	11.7	18.2	12.0	9.7	8.5	9.2	9.2	
	IHC	6	Neg	1	Neg	6	6	6	6	6	6	6	6	4	Neg	3	Neg	
AMG	SD ₅₀ /mg	9.2	Neg	7.7	Neg	8.9	10.2	15.7	10.4	9.4	11.9	14.9	19.9	9.7	8.7	12.2	9.4	
	IHC	Neg	Neg	Neg	Neg	Neg	3	2	4	4	6	5	2	6	6	5	3	
CI	SD ₅₀ /mg	9.7	Neg	7.2	Neg	8.7	9.2	12.7	10.2	11.2	8.9	10.7	18.2	9.9	7.2	7.4	Neg	
	IHC	Neg	Neg	Neg	Neg	Neg	2	1	1	Neg	2	5	2	5	Neg	Neg	Neg	
TC	SD ₅₀ /mg	8.2	Neg	NA	NA	8.7	8.7	10.7	8.2	NA	8.7	9.4	10.9	NA	NA	7.9	Neg	
	IHC	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	2	1	4	Neg	Neg	Neg	
FC	SD ₅₀ /mg	9.9	Neg	Neg	Neg	8.4	11.7	8.7	7.2	7.5	7.4	8.7	10.2	11.4	Neg	Neg	Neg	
	IHC	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	1	1	4	Neg	Neg	Neg	
CE	SD ₅₀ /mg	7.2	Neg	NA	Neg	8.4	9.8	7.4	7.7	Neg	7.9	7.9	7.7	7.5	Neg	Neg	Neg	
	IHC	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	
OCC	SD ₅₀ /mg	9.7	Neg	Neg	Neg	7.4	8.2	9.2	7.2	7.4	7.7	8.2	8.2	9.2	Neg	7.2	Neg	
	IHC	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	

AMG, amygdala; BF, basal forebrain; CI, cingulate gyrus; CE, cerebellum; FC, frontal cortex; IHC, immunohistochemical LB score; ME, medulla oblongata; NA, not available; Neg, negative; OCC, occipital cortex; SN, substantia nigra; TC, temporal cortex. SD₅₀/mg values are expressed in logarithmic scale




Fig. 3 αSyn seeds quantification in brain samples **a** Average SD₅₀/mg values in the main McKeith stages in different brain areas. Standard deviation (SD) was hidden to improve the readability of the image. **b** Correlation analysis between logSD₅₀ score and LB score ($r=0.7903$, 95% CI 0.4719 to 0.9263; $p<0.001$). AMG, amygdala; BF, basal

forebrain; CI, cingulate gyrus; CE, cerebellum; FC, frontal cortex; LB, Lewy body; ME, medulla oblongata; OCC, occipital cortex; SN, substantia nigra; TC, temporal cortex. SD₅₀/mg values are expressed in logarithmic scale

variant than that associated with the typical LBD stages. For this purpose, we looked for differences between αSyn seeding activity in the medulla oblongata and amygdala, assuming that an equal seed dose would be present in the sample at the dilution corresponding to the SD₅₀. More specifically, we extrapolated the kinetic data (lag phase, I_{max}, and AUC) of the αSyn RT-QuIC SAA curves of the

dilution related to the SD₅₀. We compared the mean values across the brainstem, limbic, neocortical, and amygdala-predominant stages and found no significant differences in αSyn RT-QuIC SAA kinetic parameters among the four groups in either the medulla oblongata or the amygdala (data not shown).

Analysis of LBD prevalence in the CJD cohort

In the last part of the study, we investigated the prevalence of *i*LBD in CJD patients, the largest and best-characterized subgroup of our NP cohort. Of the 604 examined CJD cases, 48 (7.9%) showed LB pathology. More in detail, 2 (0.3%) showed a neocortical, 19 (3.1%) a limbic, and 27 (4.5%) a brainstem stage (Table 1). Of note, no case exhibited an amygdala-predominant involvement. When stratifying patients by age, we found a prevalence of LBD in 3% of individuals aged 50–59, rising to 8–9% and staying stable in the following three decades (Fig. 4).

Discussion

The performance against the gold standard, i.e., the neuropathologic diagnosis, represents a key issue for validating novel pathology-specific biomarkers. Studies of this kind are often limited by the long latency between the *in vivo* and postmortem assessments, especially in the case of usually long-standing pathological processes (e.g., AD or LBD). In this context, our collection of CSF-brain pairs from patients with rapidly progressive disease who died within a limited time frame after LP provides the best scenario for evaluating the specificity of an α Syn SAA assay to detect LB pathology. In the present study, we analyzed a comprehensive cohort of CSF samples and postmortem brain tissues, including a good representation of all LBD pathological stages.

Consistent with previous results [3, 7, 10, 21, 27, 55], we confirmed the assay's high overall specificity (98.6%) and sensitivity (81.4%) for LB pathology. Only three cases with a primary neuropathological diagnosis of Wernicke's encephalopathy, AD, and sCJD yielded a positive test result without evidence of LB brain pathology. The finding of no

positive α Syn seeding activity in brain homogenates from two of these cases (data not shown) excluded the hypothesis of prodromal situations in which the accumulation of misfolded α Syn preceded the appearance of LB pathology. The presence of LB pathology limited to the spinal cord and/or peripheral ganglia or a CSF contamination favoring the recombinant α Syn spontaneous aggregation may explain these incongruent positive results. Additional studies on larger sample cohorts are needed to address this critical issue further.

Regarding the assay sensitivity, there was a clear correlation with the LBD stage. The sensitivity was 100% in patients with neocortical or limbic LBD stages but decreased significantly in patients with the amygdala-predominant variant (50%) or a brainstem stage (60.9%), mainly when α Syn accumulation was sparse and limited to the medulla oblongata or the pons. Specifically, the sensitivity was 73.3% in those with Braak stage 3 and 37.5% in Braak stages 1 and 2. Of note, 13 of 14 patients in the brainstem stage with a negative α Syn SAA showed a low LB pathology load at the semiquantitative assessment. Furthermore, one exhibited an atypical LB pathology distribution showing sparse LBs and LNs in the medulla oblongata and substantia nigra skipping the pons.

To date, only a few studies in small cohorts have investigated the CSF α Syn SAA diagnostic accuracy in individuals with the amygdala-predominant form, while no detailed data are available for those with the brainstem stage of LBD [3, 21, 27]. Moreover, one of these studies [27] only analyzed postmortem ventricular CSF, which cannot be easily compared with the CSF obtained by LP at an early disease stage [57]. Fairfoul et al. reported that 31% (4/13) of AD brains with incidental LBD cases (with no details on the LBD stage) tested positive at CSF α Syn SAA [21]. More recently, Hall et al. detected α Syn seeding activity in the postmortem ventricular CSF of 0% (0/2) individuals with "brainstem only", 71% (5/7) with "amygdala only", and 40% (2/5) of those with "LB pathology restricted to amygdala and brainstem" [27]. Finally, Arnold et al. reported a low (3/21, 14.3%) α Syn SAA sensitivity using antemortem CSF in the amygdala-predominant LBD variant [3].

There is no definite explanation for the reduced sensitivity of the α Syn SAA in the brainstem and amygdala-predominant stages. Hypothetically, this could be due to either the lower brain LB pathology loads compared to full-blown LBD in limbic and neocortical stages or, especially for the amygdala-predominant variant, the presence of a distinct pathological α Syn conformer, exhibiting no aggregation at all or a different kinetic profile compared to the α Syn associated with typical LBD. Our demonstration of a clear threshold effect, with the assay yielding negative results in all but four cases with a LB score < 10 out of 78, and the evidence of a significant correlation between the number of positive




Fig. 4 Age-specific *i*LBD prevalence rates in the CJD cohort. The green area shows the number of LB-positive cases at α Syn immunohistochemistry in each decade. The red line expresses these data in percentage. *i*LBD, incidental Lewy body disease, CJD, Creutzfeldt Jakob disease

replicates and the LB score strongly support the hypothesis of an association between the assay sensitivity and the overall brain LB pathology load.

The data we obtained by performing the α Syn SAA in brain homogenates from different regions further supports the positive association between LB pathology burden and assay sensitivity. First, we demonstrated for the first time a robust positive correlation between the brain α Syn seeding activity and the LB pathology load evaluated by α Syn immunostaining, thus excluding that a significant proportion of the α Syn aggregates forming LBs and LNs may lack seeding properties. Second, in each examined brain area showing LB pathology, we did not find significant differences in the mean $\log SD_{50}/mg$ values among brains stratified according to the McKeith stage, thus excluding the hypothesis of misfolded α Syn isoforms contributing differently to the overall seeding activity depending on the LBD stage. As the only possible exception, we found a trend toward lower average $\log SD_{50}/mg$ values for the amygdala-predominant cases compared to those with typical LBD. However, although the small sample size does not allow us to draw firm conclusions, we did not find differences in amygdala seeding activity between the amygdala-predominant cases and those in other stages. Further studies should investigate the structural, physicochemical, and seeding properties of α Syn in this peculiar histopathological variant compared to typical LBD.

Our results, combined with those of previous studies [3, 27], indicate that the α Syn SAA have enough sensitivity to detect all symptomatic patients with DLB or PD related to LB pathology. Indeed, DLB patients and most of those with PD are expected to have reached the limbic LBD stage already at clinical onset. Moreover, in PD patients, a high LB pathology load in the brainstem is expected to explain the significant neurodegeneration in the substantia nigra. However, CSF α Syn SAAs have shown high diagnostic performance in confirming LB pathology not only in DLB and PD patients but also in most of those affected by prodromal LBD syndromes, such as Isolated Rapid Eye Movement Sleep Behavior Disorder (iRBD) or mild cognitive impairment due to LBD (MCI-LB) and even in asymptomatic individuals [15, 32, 54, 55]. In this context, our finding of an incomplete sensitivity of the α Syn SAA for the brainstem stage of LBD might be interpreted as partially diverging from the results of the above clinical studies and deserves specific comment. The issue mainly concerns the asymptomatic individuals and those with iRBD since the 90–95% sensitivity demonstrated for MCI-LB by our SAA in the clinical setting [54] is entirely in line with the present results given that the LBD Braak stage expected in these patients is > 3 [33]. Regarding iRBD, we recently demonstrated a 75% sensitivity of our CSF α Syn SAA in the largest clinical cohort tested to date [32]. The three other studies with significant cohorts of iRBD patients conducted to date

showed 90% [31] and 84–94% [15, 58] assay sensitivities. Possible explanations for these differences are the different SAA protocols, including the recombinant protein and differences in patients' characteristics. Regarding the latter, it must be emphasized that iRBD patients often manifested other signs of "prodromal" LBD, including MCI and motor deficits, suggesting a relatively high LB pathology burden and spread, thus justifying, in line with our current results, a higher than expected sensitivity of the SAA assay. For example, the percentage of patients with iRBD with coexistent MCI or motor signs was significantly higher in the study reporting an α Syn SAA sensitivity of 90% compared to ours in which the sensitivity was 75%. In summary, the results of the present study, in line with recent imaging findings [35], strongly suggest that iRBD patients harbor, on average, a Braak stage 3 of LB pathology, with significant variations from case to case mainly depending on the prodromal LBD score determined by the associated neurological signs. Finally, our recent finding of 8% α Syn SAA positivity in a large cohort of cognitively unimpaired individuals [47] suggests that even the asymptomatic "incidental" LBD status may be associated with a variable LB pathology load, likely including Braak stages 3 and 4. The fact that all α Syn SAA-positive patients with CJD or a non-degenerative etiology included in the present study did not have any anamnestic evidence of motor signs or cognitive decline before the onset of a rapidly progressive disease and yet showed LB pathology up to Braak stage 4 supports this conclusion.

In brain sample experiments, we sought to analyze how the kinetic parameters of the α Syn SAA fluorescence signal vary depending on the number of α Syn seeds in the sample. For this purpose, we extrapolated each run's I_{max} , AUC, and lag times and studied their values' distribution along the dilution series. We computed the PAR ($1/\text{lag time}$) as a measure of the amyloid formation rate to deal with negative reactions (which show a lag phase virtually approaching infinity so that the rate of amyloid formation can be assumed equal to 0), as reported [30]. I_{max} and AUC did not show a regular decline over the dilution series, exhibiting an unpredictable pattern, likely depending not only on the amount of seed but also heavily on the experimental conditions. Conversely, following the amyloid seeding model, which predicts a linear relationship between the initial seed concentration and the rate of amyloid formation [30, 68], we found a linear dependence of the PAR (and thus the lag phase) on seed concentration. This paves the way for using the lag phase as a surrogate for the number of seeds in the sample. These results have significant implications for evaluating and validating the reliability of the "quantitative" information provided by the α Syn SAA. Developing a quantitative assay will allow the evaluation of disease severity and progression and may take the α Syn SAA to the next level. Preliminary evidence indicates that the assay might provide quantitative

data indirectly revealing the number of protein seeds in the biosample and the associated LB pathology burden in addition to the well-established reliable qualitative dichotomous (positive/negative) response. The data we obtained in the present study indicate that both the number of positive replicates and the lag phase are closely related to the α Syn burden in the CNS and likely represent the two most reliable variables to pursue the clinical validation of a quantitative α Syn SAA.

As one of the main findings, we demonstrated that in brain tissue α Syn SAA detect misfolded α Syn with higher sensitivity than immunohistochemistry. Additionally, we showed a strong correlation between the overall seed quantity, estimated through the $\log SD_{50}/mg$ score and the brain LB pathology load, assessed through immunohistochemistry. The formation of LBs and LNs is likely a multistep process starting from monomeric misfolded proteins and involving intermediate soluble species (i.e., oligomers and protofibrils), which further aggregate in insoluble fibrils, eventually forming immunohistochemically detectable deposits [60]. Our results suggest that seeds triggering α Syn SAA reactions likely include pre-fibrillary aggregates. The finding supports the numerous pieces of evidence suggesting that pre-fibrillary species are those mediating α Syn pathology spread and toxic effects [8]. In this perspective, it is possible that α Syn SAA may not only have diagnostic applications but also a potential role in monitoring the response to therapy in case future treatments targeting the early stages of the aggregation process will be available.

In the last part of the study, we reported the data obtained in our cohort of CJD patients screened for the presence of LB pathology. Consistent with previous evidence [25], we found that almost 8% of CJD patients showed LB pathology (with 4.5% cases in the brainstem, 3.1% in limbic, 0.3% in neocortical stages, and none in the amygdala-predominant variant). Based on the assumption that there is no association between prion disease and LBD, which is compatible with the current literature, we speculate that the prevalence of LB pathology in our wide case series of consecutive well-characterized CJD patients lacking clinical evidence of cognitive decline and/or motor disturbances before the rapid onset of symptoms related to CJD, can be bona fide considered an estimate of the frequency of iLBD in the elderly population. Previous studies estimated iLBD prevalence ranging between 8 and 15% in neurologically unimpaired individuals [9, 11, 22, 34, 42, 44, 51, 52, 64, 66]. In line with our results, in most cases, iLBD follows the Braak progression model [33], although it can sometimes be limited to limbic or neocortical areas [22, 33]. In our cohort, iLBD prevalence increased significantly in the transition between the sixth and seventh decades, consistent with previous results [52, 66]. However, at variance with previous studies [52, 66], which reported a steep and progressive increase from the sixth

decade onwards (reaching up to 15–18% of cognitively normal subjects > 90 years old), iLBD prevalence in our cohort remained substantially stable across decades (around 8–9%). Interestingly, similar results have been obtained in recent studies employing CSF α Syn SAA to estimate the prevalence of iLBD in cognitively unimpaired individuals [15, 47, 56]. Lastly, although amygdala LB pathology has been reported to occur frequently in concomitance with other protein misfolding disorders (e.g., AD and other tauopathies [46, 62]), we found no cases of the amygdala-predominant LBD variant in our CJD cohort, not even in sCJD subtypes with significant pathological involvement of the amygdala (i.e., VV2, VV1, and MV2K) [5, 6], supporting the prominent role of AD-related tau pathology in determining this subtype of focal LBD pathology.

The main study limitation is the low number of amygdala-predominant and Braak stage 1/2 cases. Furthermore, we are aware that the results on brain homogenates are strongly conditioned by the sampling accuracy, with some areas more accessible than others and harboring more widespread pathology. However, we are confident that including many areas from a relatively large number of subjects at each stage has minimized the inherent variability associated with the sampling procedure. We are also aware that our cohort mainly includes patients who, in most cases, do not have LBD as their primary diagnosis, so we could not correlate the assay results with clinical variables (e.g., motor or cognitive scores). As a further limitation, our cohort comprised mostly patients with a rapidly progressive disease course, which therefore may not be representative of the overall population affected by LB pathology. Finally, the inclusion, albeit in a small number, in the α Syn LB+ cohort of cases with a timespan LP-death > 1 year is a further limitation of the study.

In conclusion, taking advantage of a large neuropathologic cohort including LBD cases in various stages, our results confirm the virtually full accuracy of our ante mortem CSF α Syn SAA as a marker of LB pathology in limbic and neocortical stages, making it a revolutionary tool for LBD diagnosis. However, we observed lower α Syn SAA sensitivity in patients in the early stages (especially in cases of pathology limited to the medulla and/or pons) or with focal pathology (e.g., amygdala-predominant variant). Together with the CSF data, the results obtained by performing α Syn SAA on brain homogenates from different regions suggest that the LB pathology burden is the factor that likely mostly affects the assay sensitivity. Furthermore, kinetic analysis of the fluorescence signal along the dilution series of brain samples suggests that α Syn SAA can provide "quantitative" information, with the lag phase and the number of positive replicates being the most promising and reliable variables to develop a "quantitative" α Syn SAA prospectively. Finally, taking advantage of our well-characterized CJD cohort, we

estimated the prevalence of iLBD in the elderly population to be about 8%, thus emphasizing the importance of developing a reliable method for early and accurate LBD diagnosis.

Future efforts should focus on improving the assay sensitivity in cases with low LB pathology load, standardizing the protocol across laboratories, evaluating the impact of its inclusion in PD and DLB diagnostic criteria, and fully developing its “quantitative” potential.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00401-023-02663-0>.

Author contributions Conceptualization and design of the study: GMB, AM, and PP. Drafting/revision of the manuscript for content, including medical writing for content GMB, AM, and PP. Neuropathological analyses: GMB, SB, and PP. α Syn RT-QuIC SAA experiments: AM, MR, AT, FM. Major role in the acquisition, analysis, or interpretation of data: SB, AMas, AL, SC, and PP. Statistical analysis: GMB, AM, and AT. Critical review of the manuscript and approval of the final version: all authors.

Funding Open access funding provided by Alma Mater Studiorum - Università di Bologna within the CRUI-CARE Agreement. The study was supported by the grant Ricerca Finalizzata-2021–12374386 and “Ricerca Corrente” funded by the Ministero della Salute, and the #NextGenerationEU (NGEU), funded by the Ministry of University and Research (MUR), National Recovery and Resilience Plan (NRRP), project MNESYS (PE0000006).

Availability of data and materials The datasets used and analyzed during the current study are available from the corresponding author upon reasonable request.

Declarations

Conflict of interest The authors declare there are no competing interests.

Ethical approval The study was conducted according to the revised Declaration of Helsinki and Good Clinical Practice guidelines and approved by the local ethics committee (approval number AVEC:18025, 113/2018/OSS/AUSLBO).

Informed consent Written informed consent was given by study participants or the next of kin.

Consent for publication Not applicable.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Alafuzoff I, Arzberger T, Al-Sarraj S, Bodi I, Bogdanovic N, Braak H et al (2008) Staging of neurofibrillary pathology in Alzheimer's disease: a study of the BrainNet Europe consortium. *Brain Pathol* 18(4):484–496. <https://doi.org/10.1111/j.1750-3639.2008.00147.x>
- Alafuzoff I, Ince PG, Arzberger T, Al-Sarraj S, Bell J, Bodi I et al (2009) Staging/typing of lewy body related α -synuclein pathology: a study of the BrainNet Europe consortium. *Acta Neuropathol* 117:635–652. <https://doi.org/10.1007/s00401-009-0523-2>
- Arnold MR, Coughlin DG, Brumbach BH, Smirnov DS, Concha-Marambio L, Farris CM et al (2022) α -Synuclein seed amplification in CSF and brain from patients with different brain distributions of pathological α -synuclein in the context of co-pathology and non-LBD diagnoses. *Ann Neurol* 92:650–662. <https://doi.org/10.1002/ana.26453>
- Attems J, Toledo JB, Walker L, Gelpi E, Gentleman S, Halliday G et al (2021) Neuropathological consensus criteria for the evaluation of Lewy pathology in postmortem brains: a multi-centre study. *Acta Neuropathol* 141:159–172. <https://doi.org/10.1007/s00401-020-02255-2>
- Baiardi S, Magherini A, Capellari S, Redaelli V, Ladogana A, Rossi M et al (2017) Towards an early clinical diagnosis of sporadic CJD VV2 (ataxic type). *J Neurol Neurosurg Psychiatry* 88:764–772. <https://doi.org/10.1136/jnnp-2017-315942>
- Baiardi S, Mammana A, Dellavalle S, Rossi M, Redaelli V, Colaizzo E et al (2023) Defining the phenotypic spectrum of sporadic Creutzfeldt-Jakob disease MV2K: the kuru plaque type. *Brain* 146(8):3289–3300. <https://doi.org/10.1093/brain/awad074>
- Bargar C, Wang W, Gunzler SA, LeFevre A, Wang Z, Lerner AJ et al (2021) Streamlined alpha-synuclein RT-QuIC assay for various biospecimens in Parkinson's disease and dementia with Lewy bodies. *Acta Neuropathol Commun* 9:62. <https://doi.org/10.1186/s40478-021-01175-w>
- Bengoa-Vergniory N, Roberts RF, Wade-Martins R, Alegre-Abarrategui J (2017) Alpha-synuclein oligomers: a new hope. *Acta Neuropathol* 134:819–838. <https://doi.org/10.1007/s00401-017-1755-1>
- Bennett DA, Wilson RS, Boyle PA, Buchman AS, Schneider JA (2012) Relation of neuropathology to cognition in persons without cognitive impairment. *Ann Neurol* 72:599–609. <https://doi.org/10.1002/ana.23654>
- Bongianni M, Ladogana A, Capaldi S, Klotz S, Baiardi S, Cagnin A et al (2019) α -Synuclein RT-QuIC assay in cerebrospinal fluid of patients with dementia with Lewy bodies. *Ann Clin Transl Neurol* 6:2120–2126. <https://doi.org/10.1002/acn3.50897>
- Boyle PA, Yu L, Wilson RS, Schneider JA, Bennett DA (2013) Relation of neuropathology with cognitive decline among older persons without dementia. *Front Aging Neurosci* 5:50. <https://doi.org/10.3389/fnagi.2013.00050>
- Braak H, Del Tredici K, Rüb U, de Vos RAI, Jansen Steur ENH, Braak E (2003) Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol Aging* 24:197–211. [https://doi.org/10.1016/s0197-4580\(02\)00065-9](https://doi.org/10.1016/s0197-4580(02)00065-9)
- Braak H, Ghebremedhin E, Rüb U, Bratzke H, Del Tredici K (2004) Stages in the development of Parkinson's disease-related pathology. *Cell Tissue Res* 318:121–134. <https://doi.org/10.1007/s00441-004-0956-9>
- Candelise N, Baiardi S, Franceschini A, Rossi M, Parchi P (2020) Towards an improved early diagnosis of neurodegenerative diseases: the emerging role of in vitro conversion assays for protein amyloids. *Acta Neuropathol Commun* 8:117. <https://doi.org/10.1186/s40478-020-00990-x>

15. Concha-Marambio L, Weber S, Farris CM, Dakna M, Lang E, Wicke T et al (2023) Accurate detection of α -synuclein seeds in cerebrospinal fluid from isolated rapid eye movement sleep behavior disorder and patients with parkinson's disease in the DeNovo parkinson (DeNoPa) cohort. *Mov Disord* 38:567–578. <https://doi.org/10.1002/mds.29329>
16. Crary JF, Trojanowski JQ, Schneider JA, Abisambra JF, Abner EL, Alafuzoff I et al (2014) Primary age-related tauopathy (PART): a common pathology associated with human aging. *Acta Neuropathol* 128:755–766. <https://doi.org/10.1007/s00401-014-1349-0>
17. Dickson DW (2012) Parkinson's disease and parkinsonism: neuropathology. *Cold Spring Harb Perspect Med* 2(8):a009258. <https://doi.org/10.1101/cshperspect.a009258>
18. Dickson DW, Bergeron C, Chin SS, Duyckaerts C, Horoupian D, Ikeda K et al (2002) Office of rare diseases neuropathologic criteria for corticobasal degeneration. *J Neuropathol Exp Neurol* 61:935–946. <https://doi.org/10.1093/jnen/61.11.935>
19. Dickson DW, Uchikado H, Fujishiro H, Tsuboi Y (2010) Evidence in favor of Braak staging of Parkinson's disease. *Mov Disord* 1:S78–82. <https://doi.org/10.1002/mds.22637>
20. Dougherty T (1964) Animal virus titration techniques. In: Harris RJC (ed) *Techniques in experimental virology*. Academic Press, New York, pp 183–186
21. Fairfoul G, McGuire LL, Pal S, Ironside JW, Neumann J, Christie S et al (2016) Alpha-synuclein RT-QuIC in the CSF of patients with alpha-synucleinopathies. *Ann Clin Transl Neurol* 3:812–818. <https://doi.org/10.1002/acn3.338>
22. Frigerio R, Fujishiro H, Ahn T-B, Josephs KA, Maraganore DM, DelleDonne A et al (2011) Incidental Lewy body disease: do some cases represent a preclinical stage of dementia with Lewy bodies? *Neurobiol Aging* 32:857–863. <https://doi.org/10.1016/j.neurobiolaging.2009.05.019>
23. Gilman S, Wenning GK, Low PA, Brooks DJ, Mathias CJ, Trojanowski JQ et al (2008) Second consensus statement on the diagnosis of multiple system atrophy. *Neurology* 71:670–676. <https://doi.org/10.1212/01.wnl.0000324625.00404.15>
24. Goedert M, Masuda-Suzukake M, Falcon B (2017) Like prions: the propagation of aggregated tau and α -synuclein in neurodegeneration. *Brain* 140:266–278. <https://doi.org/10.1093/brain/aww230>
25. Grau-Rivera O, Gelpi E, Nos C, Gaig C, Ferrer I, Saiz A et al (2015) Clinicopathological correlations and concomitant pathologies in rapidly progressive dementia: a brain bank series. *Neurodegener Dis* 15:350–360. <https://doi.org/10.1159/000439251>
26. Groveman BR, Orrù CD, Hughson AG, Raymond LD, Zanusso G, Ghetti B et al (2018) Rapid and ultra-sensitive quantitation of disease-associated α -synuclein seeds in brain and cerebrospinal fluid by α Syn RT-QuIC. *Acta Neuropathol Commun* 6:7. <https://doi.org/10.1186/s40478-018-0508-2>
27. Hall S, Orrù CD, Serrano GE, Galasko D, Hughson AG, Groveman BR et al (2022) Performance of α Synuclein RT-QuIC in relation to neuropathological staging of Lewy body disease. *Acta Neuropathol Commun* 10:90. <https://doi.org/10.1186/s40478-022-01388-7>
28. Halliday GM, Holton JL, Revesz T, Dickson DW (2011) Neuropathology underlying clinical variability in patients with synucleinopathies. *Acta Neuropathol* 122:187–204. <https://doi.org/10.1007/s00401-011-0852-9>
29. Hamilton RL (2006) Lewy bodies in Alzheimer's disease: a neuropathological review of 145 cases using α -synuclein immunohistochemistry. *Brain Pathol* 10:378–384. <https://doi.org/10.1111/j.1750-3639.2000.tb00269.x>
30. Henderson DM, Davenport KA, Haley NJ, Denkers ND, Mathiason CK, Hoover EA (2015) Quantitative assessment of prion infectivity in tissues and body fluids by real-time quaking-induced conversion. *J Gen Virol* 96:210–219. <https://doi.org/10.1099/vir.0.069906-0>
31. Iranzo A, Fairfoul G, Ayudhaya ACN, Serradell M, Gelpi E, Vilaseca I et al (2021) Detection of α -synuclein in CSF by RT-QuIC in patients with isolated rapid-eye-movement sleep behaviour disorder: a longitudinal observational study. *Lancet Neurol* 20:203–212. [https://doi.org/10.1016/S1474-4422\(20\)30449-X](https://doi.org/10.1016/S1474-4422(20)30449-X)
32. Iranzo A, Mammana A, Muñoz-Lopetegi A, Dellavalle S, Mayà G, Rossi M et al (2023) Misfolded α -synuclein assessment in the skin and CSF by RT-QuIC in isolated REM sleep behavior disorder. *Neurology* 100:e1944–e1954. <https://doi.org/10.1212/WNL.000000000207147>
33. Jellinger KA (2019) Neuropathology and pathogenesis of extrapyramidal movement disorders: a critical update—I. Hypokinetic-rigid movement disorders. *J Neural Transm* 126:933–995. <https://doi.org/10.1007/s00702-019-02028-6>
34. Knopman DS, Parisi JE, Salviati A, Floriach-Robert M, Boeve BF, Ivnik RJ et al (2003) Neuropathology of cognitively normal elderly. *J Neuropathol Exp Neurol* 62:1087–1095. <https://doi.org/10.1093/jnen/62.11.1087>
35. Koeglsperger T, Rumpf S-L, Schließer P, Struebing FL, Brendel M, Levin J et al (2023) Neuropathology of incidental Lewy body and prodromal Parkinson's disease. *Mol Neurodegener* 18:32. <https://doi.org/10.1186/s13024-023-00622-7>
36. Koga S, Sekiya H, Kondru N, Ross OA, Dickson DW (2021) Neuropathology and molecular diagnosis of Synucleinopathies. *Mol Neurodegener* 16:83. <https://doi.org/10.1186/s13024-021-00501-z>
37. Kosaka K, Yoshimura M, Ikeda K, Budka H (1984) Diffuse type of Lewy body disease: progressive dementia with abundant cortical Lewy bodies and senile changes of varying degree—a new disease? *Clin Neuropathol* 3:185–192
38. Kovacs GG, Lukic MJ, Irwin DJ, Arzberger T, Respondek G, Lee EB et al (2020) Distribution patterns of tau pathology in progressive supranuclear palsy. *Acta Neuropathol* 140:99–119. <https://doi.org/10.1007/s00401-020-02158-2>
39. Lattanzio F, Abu-Rumeileh S, Franceschini A, Kai H, Amore G, Poggiolini I et al (2017) Prion-specific and surrogate CSF biomarkers in Creutzfeldt-Jakob disease: diagnostic accuracy in relation to molecular subtypes and analysis of neuropathological correlates of p-tau and A β 42 levels. *Acta Neuropathol* 133:559–578. <https://doi.org/10.1007/s00401-017-1683-0>
40. Leverenz JB, Fishel MA, Peskind ER, Montine TJ, Nochlin D, Steinbart E et al (2006) Lewy body pathology in familial Alzheimer disease. *Arch Neurol* 63:370–376. <https://doi.org/10.1001/archneur.63.3.370>
41. Lippa CF, Fujiwara H, Mann DMA, Giasson B, Baba M, Schmidt ML et al (1998) Lewy bodies contain altered α -synuclein in brains of many familial Alzheimer's disease patients with mutations in presenilin and amyloid precursor protein genes. *Am J Pathol* 153:1365–1370
42. Markesbery WR, Jicha GA, Liu H, Schmitt FA (2009) Lewy body pathology in normal elderly subjects. *J Neuropathol Exp Neurol* 68:816–822. <https://doi.org/10.1097/NEN.0b013e3181ac10a7>
43. McKeith IG, Dickson DW, Lowe J, Emre M, O'Brien JT, Feldman H et al (2005) Diagnosis and management of dementia with Lewy bodies: third report of the DLB Consortium. *Neurology* 65:1863–1872. <https://doi.org/10.1212/01.wnl.0000187889.17253.b1>
44. Mikolaenko I, Pletnikova O, Kawas CH, O'Brien R, Resnick SM, Crain B et al (2005) Alpha-synuclein lesions in normal aging, Parkinson disease, and Alzheimer disease: evidence from the Baltimore Longitudinal Study of Aging (BLSA). *J Neuropathol Exp Neurol* 64:156–162. <https://doi.org/10.1093/jnen/64.2.156>
45. Montine TJ, Phelps CH, Beach TG, Bigio EH, Cairns NJ, Dickson DW et al (2012) National Institute on Aging–Alzheimer's Association guidelines for the neuropathologic assessment of

- Alzheimer's disease: a practical approach. *Acta Neuropathol* 123:1–11. <https://doi.org/10.1007/s00401-011-0910-3>
46. Nelson PT, Abner EL, Patel E, Anderson S, Wilcock DM, Kryscio RJ et al (2018) The amygdala as a locus of pathologic misfolding in neurodegenerative diseases. *J Neuropathol Exp Neurol* 77:2–20. <https://doi.org/10.1093/jnen/nlx099>
 47. Palmqvist S, Rossi M, Hall S, Quadalti C, Mattsson-Carlgrén N, Dellavalle S et al (2023) Cognitive effects of Lewy body pathology in clinically unimpaired individuals. *Nat Med* 29:1971–1978. <https://doi.org/10.1038/s41591-023-02450-0>
 48. Parchi P, de Boni L, Saverioni D, Cohen ML, Ferrer I, Gambetti P et al (2012) Consensus classification of human prion disease histotypes allows reliable identification of molecular subtypes: an inter-rater study among surveillance centres in Europe and USA. *Acta Neuropathol* 124:517–529. <https://doi.org/10.1007/s00401-012-1002-8>
 49. Parchi P, Strammiello R, Notari S, Giese A, Langeveld JPM, Ladogana A et al (2009) Incidence and spectrum of sporadic Creutzfeldt-Jakob disease variants with mixed phenotype and co-occurrence of PrPSc types: an updated classification. *Acta Neuropathol* 118:659–671. <https://doi.org/10.1007/s00401-009-0585-1>
 50. Parkkinen L, Pirttilä T, Alafuzoff I (2008) Applicability of current staging/categorization of alpha-synuclein pathology and their clinical relevance. *Acta Neuropathol* 115:399–407. <https://doi.org/10.1007/s00401-008-0346-6>
 51. Parkkinen L, Soininen H, Alafuzoff I (2003) Regional distribution of alpha-synuclein pathology in unimpaired aging and Alzheimer disease. *J Neuropathol Exp Neurol* 62:363–367. <https://doi.org/10.1093/jnen/62.4.363>
 52. Parkkinen L, Soininen H, Laakso M, Alafuzoff I (2001) Alpha-synuclein pathology is highly dependent on the case selection. *Neuropathol Appl Neurobiol* 27:314–325. <https://doi.org/10.1046/j.0305-1846.2001.00342.x>
 53. Quadalti C, Palmqvist S, Hall S, Rossi M, Mammana A, Janelidze S et al (2023) Clinical effects of Lewy body pathology in cognitively impaired individuals. *Nat Med*. <https://doi.org/10.1038/s41591-023-02449-7>
 54. Rossi M, Baiardi S, Teunissen CE, Quadalti C, van de Beek M, Mammana A et al (2021) Diagnostic value of the CSF α -synuclein real-time quaking-induced conversion assay at the prodromal mci stage of dementia with Lewy bodies. *Neurology* 97:e930–e940. <https://doi.org/10.1212/WNL.0000000000012438>
 55. Rossi M, Candelise N, Baiardi S, Capellari S, Giannini G, Orrù CD et al (2020) Ultrasensitive RT-QuIC assay with high sensitivity and specificity for Lewy body-associated synucleinopathies. *Acta Neuropathol* 140:49–62. <https://doi.org/10.1007/s00401-020-02160-8>
 56. Russo MJ, Orru CD, Concha-Marambio L, Giaisi S, Groveman BR, Farris CM et al (2021) High diagnostic performance of independent alpha-synuclein seed amplification assays for detection of early Parkinson's disease. *Acta Neuropathol Commun* 9:179. <https://doi.org/10.1186/s40478-021-01282-8>
 57. Saijo E, Metrick MA, Koga S, Parchi P, Litvan I, Spina S et al (2020) 4-Repeat tau seeds and templating subtypes as brain and CSF biomarkers of frontotemporal lobar degeneration. *Acta Neuropathol* 139:63–77. <https://doi.org/10.1007/s00401-019-02080-2>
 58. Siderowf A, Concha-Marambio L, Lafontant D-E, Farris CM, Ma Y, Urenia PA et al (2023) Assessment of heterogeneity among participants in the Parkinson's Progression Markers Initiative cohort using α -synuclein seed amplification: a cross-sectional study. *Lancet Neurol* 22:407–417. [https://doi.org/10.1016/S1474-4422\(23\)00109-6](https://doi.org/10.1016/S1474-4422(23)00109-6)
 59. Sorrentino ZA, Goodwin MS, Riffe CJ, Dhillon J-KS, Xia Y, Gorion K-M et al (2019) Unique α -synuclein pathology within the amygdala in Lewy body dementia: implications for disease initiation and progression. *Acta Neuropathol Commun* 7:142. <https://doi.org/10.1186/s40478-019-0787-2>
 60. Soto C, Pritzkow S (2018) Protein misfolding, aggregation, and conformational strains in neurodegenerative diseases. *Nat Neurosci* 21:1332–1340. <https://doi.org/10.1038/s41593-018-0235-9>
 61. Spillantini MG, Schmidt ML, Lee VM, Trojanowski JQ, Jakes R, Goedert M (1997) Alpha-synuclein in Lewy bodies. *Nature* 388:839–840. <https://doi.org/10.1038/42166>
 62. Spires-Jones TL, Attems J, Thal DR (2017) Interactions of pathological proteins in neurodegenerative diseases. *Acta Neuropathol* 134:187–205. <https://doi.org/10.1007/s00401-017-1709-7>
 63. Toledo JB, Gopal P, Raible K, Irwin DJ, Brettschneider J, Sedor S et al (2016) Pathological α -synuclein distribution in subjects with coincident Alzheimer's and Lewy body pathology. *Acta Neuropathol* 131:393–409. <https://doi.org/10.1007/s00401-015-1526-9>
 64. Tsuboi Y, Ahlskog JE, Apaydin H, Parisi JE, Dickson DW (2001) Lewy bodies are not increased in progressive supranuclear palsy compared with normal controls. *Neurology* 57:1675–1678. <https://doi.org/10.1212/wnl.57.9.1675>
 65. Uchikado H, Lin W-L, DeLucia MW, Dickson DW (2006) Alzheimer disease with amygdala lewy bodies: a distinct form of α -synucleinopathy. *J Neuropathol Exp Neurol* 65:685–697. <https://doi.org/10.1097/01.jnen.0000225908.90052.07>
 66. Wakisaka Y, Furuta A, Tanizaki Y, Kiyohara Y, Iida M, Iwaki T (2003) Age-associated prevalence and risk factors of Lewy body pathology in a general population: the Hisayama study. *Acta Neuropathol* 106:374–382. <https://doi.org/10.1007/s00401-003-0750-x>
 67. Walker LC, Jucker M (2015) Neurodegenerative diseases: expanding the prion concept. *Annu Rev Neurosci* 38:87–103. <https://doi.org/10.1146/annurev-neuro-071714-033828>
 68. Wilham JM, Orrù CD, Bessen RA, Atarashi R, Sano K, Race B et al (2010) Rapid end-point quantitation of prion seeding activity with sensitivity comparable to bioassays. *PLoS Pathog* 6:e1001217. <https://doi.org/10.1371/journal.ppat.1001217>
 69. Zaccai J, Brayne C, McKeith I, Matthews F, Ince PG, MRC Cognitive Function, Ageing Neuropathology Study (2008) Patterns and stages of alpha-synucleinopathy: Relevance in a population-based cohort. *Neurology* 70:1042–1048. <https://doi.org/10.1212/01.wnl.0000306697.48738.b6>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.