

Supplementary information 1. Diagnostic schedule for study dogs.

<b>Time point</b>	<b>CBC</b>	<b>PCV</b>	<b>Blood lactate</b>	<b>SAT exam</b>	<b>Serum biochemistry</b>	<b>CRP</b>	<b>AT activity</b>	<b>Urine dipstick</b>	<b>UPC §</b>
T0 (day 0)	X	X	X	X	X	X	X	X	
T2 (day 2)	X	X	X	X				X	
T3 (day 3)	X	X	X	X	X		X	X	
T4 (day 4)	X	X	X	X				X	
T5 (day 5)	X	X	X	X				X	
T6 (day 6)	X	X	X	X				X	

---

T7 (day 7)	X	X	X	X	X	X	X*	X*	X		
T14 (day 14)	X		X	X*	X	X*	X*	X*	X	X*	
T30 (day 30)	X		X	X*	X	X*	X*	X*	X	X*	
T60 (day 60)	X		X	X*	X	X*	X*	X*	X*	X*	
T120 (day 120)	X		X	X*	X	X*	X*	X*	X*	X*	
T180 (day 180)	X					X					
T365 (day 365)	X					X					

---

\*Clinicopathological variables were measured out until normalization within the reference range.

§Urine protein-to-urine creatinine ratio quantification was performed from the disappearance of pigmented urine onwards at different time points.

AT activity, antithrombin activity; CRP, C-reactive protein; SAT, saline autoagglutination test; UPC, urine protein-to-urine creatinine ratio.